
Zinc

The aesthetic objective for zinc is ≤ 5.0 mg/L. Zinc is an essential element and is generally considered to be non-toxic. Intake of zinc from food is more than sufficient to satisfy the recommended daily requirements. Drinking water is not regarded as an important nutritional source of this element. Water containing zinc at concentrations above 5.0 mg/L tends to be opalescent, develops a greasy film when boiled, and has an undesirable astringent taste.

General

Zinc is an abundant element and constitutes approximately 0.004 percent of the Earth's substance.⁽¹⁾ The most common zinc mineral is sphalerite (ZnS), which is often associated with the sulphides of other metallic elements, e.g., lead, copper, cadmium, and iron.⁽²⁾ Zinc is also found as calamine (ZnCO₃) in carbonate sediments; other forms of zinc are usually products of the oxidation of sphalerite.^(2,3)

During 1984, the zinc content of ores and concentrates produced in Canada was 1 207 098 tonnes; domestic zinc consumption was 150 528 tonnes.⁽⁴⁾ Canadian zinc reserves in known deposits have been estimated at 20 492 700 tonnes.⁽⁴⁾ The main use for zinc in Canada is for galvanizing iron and steel, which accounted for 40 percent of the zinc consumption in 1984. A further 20 percent went towards the production of brass plumbing and heating fittings, and 15 percent was used in the die-casting industry to produce builders' hardware and automobile fittings.⁽⁵⁾ The remaining consumption of zinc is divided among many industries. Zinc oxide is used in the manufacture of paints, rubber, rayons, yarns, inks, matches, and chemicals; zinc "dust" is used in printing, textile dyeing, and fat purification, and for cementing metals in metallurgical processing; rolled zinc is used in ornamental work and batteries; and zinc salts are used as wood preservatives and pesticides.

Occurrence

Industrial and domestic emissions contribute a considerable quantity of zinc to the air and water environments. In Canada during 1972, primary iron and steel production accounted for the emission of 2690 tonnes of zinc (in the form of zinc oxide) to the

atmosphere. Primary copper and nickel production contributed a similar quantity (2290 tonnes). Total industrial emissions of zinc amounted to 6340 tonnes. Fuel combustion from stationary sources (coal and heavy oils) added a further 134 tonnes. Transportation, solid waste incineration, and pesticide application contributed 794, 124, and 1.8 tonnes, respectively. Emissions from all sources for 1972 totalled 7400 tonnes.⁽⁵⁾

In a 1980 study, it was estimated that the annual contribution of zinc to the Canadian environment from natural and anthropogenic sources was 768 000 and 414 000 tonnes, respectively.⁽⁶⁾

Analyses for zinc in airborne particulate matter over urban areas showed concentrations ranging from 0.32 mg/g in Toronto to 5.40 mg/g in Vancouver.⁽⁷⁾ Concentrations as high as 6 mg/g were found in dust along Toronto's waterfront.⁽⁸⁾

Analyses of the total suspended particulates in the air over Edmonton during 1978 and 1979 showed a mean concentration of 0.085 $\mu\text{g}/\text{m}^3$ with little seasonal variation. During one event in November 1978, the concentration reached 0.6 $\mu\text{g}/\text{m}^3$, possibly as a result of fugitive emissions from incinerators.⁽⁹⁾

In a 1982 survey across Ontario, the spatial pattern of precipitation and air quality regarding trace metal content, including zinc, was monitored.⁽¹⁰⁾ The mean air concentration of zinc ranged from 0.019 $\mu\text{g}/\text{m}^3$ in the south to 0.007 $\mu\text{g}/\text{m}^3$ in the north. Similarly, the annual dry deposition of zinc ranged from 1.51 $\mu\text{g}/\text{m}^2$ in the south to 0.51 $\mu\text{g}/\text{m}^2$ in the north. The mean concentration of zinc in the annual precipitation ranged from 0.0083 mg/L in the south to 0.0055 mg/L in the north.⁽⁸⁾ The trace metal concentration across Ontario showed a general decreasing trend from south to north. The quantity of zinc found in precipitation reflects the presence of zinc-emitting operations. The mean concentration of zinc in precipitation near Sudbury, Ontario, during 1972 to 1973 was 0.028 mg/L, but reached 120 mg/L during one rainfall.⁽¹¹⁾ The average concentration of zinc in rain falling in the United States was 0.107 mg/L.⁽¹²⁾

In unpolluted snow sampled in a remote area of northern Manitoba, the concentrations of zinc were less than 0.001 mg/L.⁽¹¹⁾

The average soil concentration of zinc in Canada is 90 mg/kg.⁽⁷⁾ A similar value was found in soils not associated with sulphide ores in New Brunswick.⁽¹³⁾ The zinc content of representative soils from industrial locations in Canada ranged from 106 to 2095 mg/kg (the maximum value was found in Toronto).⁽⁸⁾ Urban soil samples from Vancouver and Toronto contained 395 and 220 mg/kg, respectively.⁽⁷⁾

The carbonates, oxides, and sulphides of zinc are sparingly soluble, and therefore zinc is present in natural waters at low concentrations. The highly soluble chloride and sulphate salts are hydrolyzed to form the hydroxide and carbonate. Adsorption onto hydrolysate sediments further depletes the levels of dissolved zinc.⁽³⁾

The environmental concentration of zinc in Canadian surface waters ranged from 0.001 mg/L to a high of 1.17 mg/L over a 1980 to 1985 sampling period.⁽¹⁴⁾ Analysis of several lakes in Ontario (used as sources of drinking water) showed that the mean concentration of zinc was less than 0.010 mg/L.⁽¹⁵⁾

In surveys of river water in central and western Canada, it was found that the level of zinc varied widely with both the location and the season. The range was 0.001 to 0.096 mg/L, with maximum levels observed in the Slave River in the Northwest Territories; normally, the concentrations in river water do not exceed 0.04 mg/L.⁽¹⁶⁾

The concentration of zinc can be considerably higher in tap water than in surface water because of the leaching of zinc from galvanized pipes, hot water tanks, and brass fittings.⁽¹⁷⁾

In a national survey of Canadian drinking water supplies, it was found that the median zinc content in the raw, treated, and distributed water samples rarely exceeded 0.01 mg/L. However, zinc was frequently picked up while passing through the distribution network. In New Brunswick and Saskatchewan, pickup was negligible, whereas in British Columbia and Quebec, pickup was found in over half the distribution systems.⁽¹⁸⁾

Recent data from Ontario have corroborated these findings, with zinc in water withdrawn after an overnight stand sometimes reaching levels 100 times greater than that of the treated water. Thus, in Gananoque on the St. Lawrence River, the concentration of zinc after treatment was less than 0.01 mg/L in 1986, whereas the mean concentration at seven stations after an overnight stand was 0.309 mg/L (range: 0.03 to 1.17 mg/L). Water sampled after a five-minute flush had a mean zinc concentration of 0.014 mg/L.⁽¹⁵⁾

Because of the nutritional value of zinc, several studies have been undertaken to determine the zinc

content of foods and total diets. Two surveys have been carried out to establish the zinc content of representative Canadian diets. These showed that meat, fish, poultry, cereals, oils, and fats have the highest zinc content, with meat, fish, and poultry contributing most to the daily dietary intake.^(19,20)

Canadian Exposure

The average daily intake from a “representative Canadian diet” has been estimated to be between 15.2 and 19.9 mg.^(19,20) The average daily intake for a “normal” man was reported to be 12 mg,⁽⁷⁾ whereas other estimates of the normal daily intake range from 13.0 to 16.1 mg.⁽²¹⁾

Based on a 1981 national survey of trace metals in drinking water supplies, it was estimated that the average daily intake of zinc from drinking water for Canadian adults is $\leq 13.0 \mu\text{g/day}$.⁽¹⁸⁾ Based on a 1984 study, this value ranged from 33.8 to 97.5 $\mu\text{g/day}$ and was found to be highly dependent on the sampling strategy.⁽²²⁾

If the average concentration of zinc in air in Canada is assumed to be $0.035 \mu\text{g/m}^3$ and the daily respiratory volume 20 m^3 , then the daily intake from air would be 0.7 μg .

Based on the above considerations, the total daily intake of zinc from food, air, and water is estimated to be about 16 mg, with food contributing over 99 percent of the intake.

Health Considerations

Essentiality

Zinc is an essential element for all living things, including man. Zinc-containing proteins and enzymes are involved in every aspect of metabolism, including the replication and translation of genetic material.⁽²³⁾ Nearly 200 zinc-containing enzymes have been identified from all species.⁽²⁴⁾ A few of the well-characterized zinc-containing enzymes are carbonic anhydrase, aspartase, transcarbamylase, and alcohol dehydrogenase.

The Recommended Dietary Intake (RDI) for Canadians is 2 mg/day for young infants, 3 to 7 mg/day for children to age 12, and, from the age of 13, 9 mg/day for males and 8 mg/day for females.⁽²⁵⁾ The United States Recommended Daily Allowance (RDA) is 15 mg/day for adults.⁽²⁶⁾

Zinc deficiency is most often found in countries where the primary source of protein is cereal grains. Zinc absorption from these grains may be limited by high phosphate and phytate content.⁽²⁷⁾ Zinc deficiency, however, has also been identified in some children in the United States.⁽²⁸⁾ Zinc deficiency is most likely to

occur in segments of the population with high zinc requirements, such as infants, adolescents, and pregnant women.⁽²⁹⁾

General symptoms of zinc deficiency in humans include retarded growth, hypogonadism, anorexia, mental lethargy, skin changes, and night blindness.⁽²⁷⁾ Zinc deficiency may also impair the immune system,^(30–32) slow wound healing,^(27,33) and lead to eczema⁽³⁴⁾ and acne.^(35,36) The importance of zinc in brain development and function has also been emphasized.⁽³⁷⁾ Recently it was reported that hyperactive children showed some degree of zinc deficiency.⁽³⁸⁾

Absorption

In humans, zinc is absorbed in the small intestine by a carrier-mediated mechanism.⁽³⁹⁾ The fraction of zinc absorbed is difficult to determine because zinc is also secreted into the gut. Generally, 33 percent is accepted as the average zinc absorption in humans,^(25,40,41) although zinc may be more bioavailable from drinking water than from food.⁽⁴²⁾ Zinc absorption increases with increasing dietary zinc up to a maximum rate, indicating a saturated carrier-mediated mechanism.⁽⁴³⁾ In addition, zinc status may influence zinc absorption. Zinc-deprived humans absorb this element with increased efficiency, whereas humans on a high-zinc diet show a reduced efficiency of absorption.

Almost 20 regulatory and dietary factors have been reported to influence zinc absorption,⁽³⁹⁾ one of the most important of which is phytate (myoinositol hexaphosphate).⁽⁴⁴⁾ The bioavailability of zinc from cereals and legumes is greatly reduced by the phytate content of these foods,^(40,45) although some components of dietary fibre may also reduce zinc absorption.^(46,47) Phytate, particularly when ingested with calcium,⁽⁴⁸⁾ reduces zinc absorption by forming insoluble precipitates.

At high doses, zinc and iron appear to antagonize each others' absorption.^(49,50) In addition, excessive levels of both tin and copper in the diet may reduce zinc absorption.^(51–54)

Distribution

Zinc absorbed from the intestine is carried to the liver bound to albumin.⁽⁵⁵⁾ Within the circulatory system, between 80 and 90 percent of the zinc found in whole blood is in the erythrocytes,⁽⁵⁶⁾ where it is largely bound to carbonic anhydrase.⁽⁴⁹⁾ The remaining plasma zinc is primarily bound to albumin, although other proteins and free amino acids also bind zinc.⁽⁵⁷⁾ Only a small fraction of plasma zinc is in the ionic form.

The body tissues high in zinc include bone, liver, kidney, pancreas, retina, and prostate.^(58,59) Redistribution of zinc can occur under stress.⁽⁶⁰⁾

Excretion

Zinc is primarily excreted via the gastrointestinal tract; zinc excreted in this way consists of unabsorbed dietary zinc, endogenous zinc secreted into the gastrointestinal tract, and zinc from mucosal cell desquamation.⁽⁶¹⁾ Excretory mechanisms contribute to zinc homeostasis; however, regulation of zinc absorption is the major homeostatic mechanism. Under normal circumstances, about 0.5 mg of zinc may be lost in perspiration,⁽⁶²⁾ and about an equal amount is lost in daily urine. Although the kidney nephrons appear capable of secreting and reabsorbing zinc, urinary zinc does not appear to fluctuate with diet.⁽⁶³⁾ High-protein diets, however, are reported to increase urinary zinc.⁽⁶⁴⁾

Toxicity

Toxicity from dietary zinc has not been reported, although occupational exposure⁽⁶⁵⁾ or pharmacological interventions⁽⁶⁶⁾ can produce symptoms. There have been reports of teratogenic effects in sheep⁽⁶⁷⁾ and disrupted cholesterol metabolism in man,^(68,69) both thought to be due to the adverse effects of high zinc concentrations on copper metabolism. The copper status of individuals consuming high-zinc diets, therefore, should be monitored. The immune response in humans is reported to be impaired by excessive zinc intake.⁽⁷⁰⁾

Other Considerations

Taste threshold tests have shown that 5 percent of the population can taste zinc at about 5 mg/L.⁽⁷¹⁾ At a concentration of 40 mg/L, zinc gives water a milky appearance.⁽⁷²⁾

Rationale

1. Zinc is an essential element for human nutrition. The daily requirement is between 4 and 10 mg depending on age and sex, but pregnant women and new mothers may require up to 16 mg/day. Food constitutes the most important source of zinc. Long-term ingestion of quantities considerably in excess of these amounts has not resulted in adverse effects. Furthermore, because of efficient homeostatic control mechanisms, the occurrence of chronic zinc toxicity is extremely unlikely. A maximum acceptable concentration (MAC) for zinc in drinking water has therefore not been set.

2. Although surface waters seldom have zinc concentrations greater than 0.1 mg/L, levels in tap water can be considerably higher because of the use of zinc in plumbing materials. Water containing zinc at concentrations in excess of 5.0 mg/L has an undesirable astringent taste and may be opalescent and develop a greasy film on boiling.

3. The aesthetic objective for zinc in drinking water is therefore ≤ 5.0 mg/L.

References

1. Browing, E. Toxicity of industrial metals. 2nd edition. Butterworths, London. p. 348 (1969).
2. U.S. Environmental Protection Agency. Quality criteria for water. Washington, DC. p. 481 (1976).
3. Hem, J.K. Zinc. In: Study and interpretation of the chemical characteristics of natural water. Geological Survey Water Supply Paper No. 1473, Washington, DC. p. 125 (1970).
4. Gauvin, M.J. Zinc. In: Canadian minerals yearbook. Mineral Resources Branch, Department of Energy, Mines and Resources (1985).
5. Environment Canada. National inventory of sources and emissions of zinc (1972). Internal Report APCD 76-1, Air Pollution Control Directorate, June (1976).
6. Taylor, M.C. and Demayo, A. Zinc. In: Guidelines for surface water quality. Vol. 1. Inorganic chemical substances. Inland Waters Directorate, Water Quality Branch, Environment Canada, Ottawa (1980).
7. Warren, H.V. Some trace element concentrations in various environments. In: Environmental medicine. G.M. Howe and J.A. Loraine (eds.). William Heinemann Medical Books Ltd., London. p. 9 (1973).
8. Van Loon, J.C. Toronto's precipitation analysed for heavy metal content. Water Pollut. Control, 111: 38 (1973).
9. Klemm, R.F. and Gray, J.M.L. A study of the chemical composition of particulate matter and aerosols over Edmonton. Report RMD 82/9, prepared for the Research Management Division by the Alberta Research Council. p. 125 (1982).
10. Chan, W.H., Tang, A.J.S., Chung, D.H.S. and Lusic, M.A. Concentrations and deposition of trace metals in Ontario — 1982. Water Air Soil Pollut., 29: 373 (1986).
11. Beamish, R.J. and Van Loon, J.C. Precipitation loading of acid and heavy metals to a small acid lake near Sudbury, Ontario. J. Fish. Res. Board Can., 34: 649 (1977).
12. Lazrus, A.L., Lorange, E. and Lodge, J.P. Lead and other metal ions in United States precipitation. Environ. Sci. Technol., 4: 55 (1970).
13. Present, E.W. and Tupper, W.M. Trace elements in some New Brunswick soils. Can. J. Soil Sci. 45: 305 (1965).
14. National Water Quality Data Bank (NAQUADAT). Water Quality Branch, Inland Waters Directorate, Environment Canada, Ottawa (1985).
15. Ontario Ministry of the Environment. Survey of water quality in the distribution systems, Ontario 1981–1986. Unpublished data (1987).
16. Environment Canada. Detailed surface water quality data, Northwest Territories 1980–1981, Alberta 1980–1981, Saskatchewan 1980–1981, Manitoba 1980–1981. Inland Waters Directorate (1984).
17. National Academy of Sciences. Water quality criteria. Vol. 93. Washington, DC (1972).
18. Méranger, J.C., Subramanian, K.S. and Chalifoux, C. Metals and other elements. Int. J. Assoc. Anal. Chem., 64: 44 (1981).
19. Kirkpatrick, D.C. and Coffin, D.E. The trace metal content of representative Canadian diets in 1970 and 1971. J. Inst. Can. Sci. Technol. Aliment., 7: 56 (1974).
20. Méranger, J.C. and Smith, D.C. The heavy metal content of a typical Canadian diet. Can. J. Public Health, 62: 53 (1972).
21. Hamilton, E.I. Relations between metal elements in man's diet and environmental factors. Trace Subst. Environ. Health, 13: 3 (1979).
22. Méranger, J.C., Subramanian, K.S., Langford, C.H. and Umbrasas, R. Use of an on-site integrated pump sampler for estimation of total daily intake of some metals from tap water. Int. J. Environ. Anal. Chem., 17: 307 (1984).
23. Galdes, A. and Vallee, B.L. Categories of zinc metalloenzymes. Metal Ions Biol. Syst., 15: 2 (1983).
24. O'Dell, B.L. History and status of zinc in nutrition. Fed. Proc., 43: 2821 (1984).
25. Department of National Health and Welfare. Recommended nutrient intakes for Canadians. Ottawa (1983).
26. National Academy of Sciences. Recommended daily allowances. 8th edition. Washington, DC (1974).
27. Prasad, A.S. Clinical biochemical and nutritional spectrum of zinc deficiency in human subjects: an update. Nutr. Rev., 41: 197 (1983).
28. Hambridge, K.M. and Walraevens, P.A. In: Trace elements in human health and disease. Vol. 1. A.S. Prasad (ed.). Academic Press, New York, NY. p. 21 (1976).
29. Pleban, P.A., Numerof, B.S. and Wirth, F.H. Trace element metabolism in the fetus and neonate. Clin. Endocrinol. Metab., 14: 545 (1985).
30. Fraker, P.S., Haas, S.M. and Luecke, R.W. Effect of zinc deficiency on the immune response of young adult A/J mouse. J. Nutr., 107: 1889 (1977).
31. De Pasquale-Jardieu, P. and Fraker, P.J. Interference in the development of a secondary immune response in mice by zinc deprivation: persistence of effects. J. Nutr., 114: 1762 (1984).
32. Baer, M.T., King, J.C., Tamura, T., Morgen, S., Bradfield, R.B., Weston, W.L. and Daugherty, N.A. Nitrogen utilization, enzyme activity, glucose intolerance and leukocyte chemotaxis in human zinc depletion. Am. J. Clin. Nutr., 41: 1220 (1985).
33. Senapati, A. and Thompson, R.P.H. Zinc deficiency and the prolonged accumulation of zinc in wounds. Br. J. Surg., 72: 583 (1985).
34. David, T.J., Wells, F.E., Sharpe, T.C. and Gibbs, A.C.C. Low serum zinc in children with atopic eczema. Br. J. Dermatol., 111: 597 (1984).
35. Fritzerherbert, J.C. Acne vulgaris — zinc deficiency? Med. J. Aust., 1: 848 (1976).
36. Pohit, J., Saha, K.C. and Pal, B. Zinc status of acne vulgaris patients. J. Appl. Nutr., 37: 18 (1985).
37. Sandstead, H.H. Zinc: essential for brain development and function. Nutr. Rev., 43: 129 (1985).
38. Barlow, P.J. and Sidani, S.A. Metal imbalance and hyperactivity. Acta Pharmacol. Toxicol., 59: 458 (1986).
39. Cousins, R.J. Absorption, transport and hepatic metabolism of copper and zinc: special reference to metallothionein and ceruloplasmin. Physiol. Rev., 65: 238 (1985).
40. Turmlund, J.R., King, J.C., Keyes, W.R., Gong, B. and Michel, M.C. A stable isotope study of zinc absorption in young men: effects of phytate and α -cellulose. Am. J. Clin. Nutr., 40: 1071 (1984).
41. Honstead, J.F. and Brady, D.N. The uptake and retention of ^{32}P and ^{65}Zn from the consumption of Columbia River fish. Health Phys., 13: 455 (1967).

42. Van Barneveld, A.A. and Van Den Hamer, C.J.A. Influence of isotope administration mode and of food consumption on absorption and retention of ⁶⁵Zn in mice. *Trace Subst. Environ. Health*, 16: 196 (1983).
43. Steel, L. and Cousins, R.J. Kinetics of zinc absorption by lumenally and vascularly perfused rat intestine. *Am. J. Physiol.*, 248: G46 (1985).
44. House, W.A., Welch, R.M. and Van Campen, D.R. Effect of phytic acid on the absorption, distribution and endogenous excretions of zinc in rats. *J. Nutr.*, 112: 941 (1982).
45. O'Dell, B.L. and Savage, J.E. Effect of phytate on zinc availability. *Proc. Soc. Exp. Biol. Med.*, 103: 304 (1960).
46. Reinhold, J.G., Faradji, B., Abade, P. and Ismail-Beigi, F. Decreased absorption of calcium, magnesium, zinc and phosphorus by humans due to increased fiber and phosphorus consumption as wheat bread. *J. Nutr.*, 106: 493 (1976).
47. Kelsay, J.L., Jacob, R.A. and Prather, E.S. Effect of fiber from fruits and vegetables on metabolic responses of human subjects: III — Zinc, copper and phosphorus balances. *Am. J. Clin. Nutr.*, 32: 2307 (1979).
48. Sandstead, J.J., Dintzis, F. and Johnson, L. Influence of diet fiber, phytate and calcium on human zinc absorption. *Fed. Proc.*, 43: 851 (1984).
49. Crofton, R.W., Gvozdanovic, D. and Agget, P.J. A study of the effect of zinc on iron absorption in Man. *Proc. Nutr. Soc.*, 41: 17 (abstr.) (1962).
50. Solomons, N.W. and Jacob, R.A. Studies on the bioavailability of zinc in humans: effect of heme and non-heme iron on absorption of zinc. *Am. J. Clin. Nutr.*, 34: 475 (1981).
51. Valberg, L.S., Flanagan, P.R. and Chamberlain, M.J. Effects of iron, tin and copper on zinc absorption in humans. *Am. J. Clin. Nutr.*, 40: 536 (1984).
52. Johnson, M.A., Baier, J.M. and Greger, J.L. Effect of dietary tin on zinc, copper, iron, manganese and magnesium metabolism of adult males. *Am. J. Clin. Nutr.*, 35: 1332 (1982).
53. Van Campen, D.R. Copper interference with the intestinal absorption of zinc-65 by rats. *J. Nutr.*, 97: 104 (1969).
54. Van Campen, D.R. and Scaife, P.U. Zinc interference with coffee absorption in rats. *J. Nutr.*, 91: 473 (1967).
55. Smith, K.T., Failla, M.L. and Cousins, R.J. Identification of albumin as the plasma carriers for zinc absorption by perfused rat intestine. *Biochem. J.*, 184: 627 (1979).
56. Scott, K.T. and Bradwell, A.R. Identification of the serum binding proteins for iron, zinc, cadmium, nickel and calcium. *Clin. Chem.*, 29: 629 (1983).
57. Vallee, B.L., Lewis, H.D., Attschule, M.P. and Gibson, J.G. The relationship between carbonic anhydrase activity and zinc content of erythrocytes in normal, in anemic and other pathological conditions. *Blood*, 4: 467 (1949).
58. Underwood, E.J. In: *Trace elements in human and animal nutrition*. 4th edition. E. Underwood (ed.). Academic Press, New York, NY. p. 196 (1977).
59. Halsted, J.A., Smith, J.C. and Irwin, M.I. Conspectus of research on zinc requirements of man. *J. Nutr.*, 104: 345 (1974).
60. Beisel, W.R., Pekarek, R.S. and Wannemacher, R.W., Jr. In: *Trace elements in human health and disease*. Vol. 1. A.S. Prasad (ed.). Academic Press, New York, NY. p. 87 (1976).
61. Kirchgessner, M. and Weigand, E. Zinc absorption and excretion in nutrition. *Metal Ions Biol. Syst.*, 15: 319 (1983).
62. Prasad, A.S. Clinical, biochemical and nutritional spectrum of zinc deficiency in human subjects: an update. *Nutr. Rev.*, 41: 197 (1983).
63. Abu-Hamdan, D.K., Migdal, S.O., Whitehorse, R., Rabbani, P., Prasad, A.S. and McDonald, F.D. Renal handling of zinc: effect of cysteine infusion. *Am. J. Physiol.*, 241(F): 487 (1981).
64. Greger, J.L. and Snedeker, S.M. Effect of dietary protein and phosphorus levels on the utilization of zinc, copper and manganese by adult males. *J. Nutr.*, 110: 243 (1980).
65. Papp, J.P. Metal fume fever. *Postgrad. Med.*, 43: 160 (1968).
66. Prasad, A.S., Brewer, G.J., Schoemaker, E.B. and Rabbani, P. Hypocupremia induced by zinc therapy in adults. *J. Am. Med. Assoc.*, 240: 2166 (1978).
67. Campbell, J.K. and Mills, C.F. Toxicity of zinc to pregnant sheep. *Environ. Res.*, 20: 1 (1979).
68. Klevay, L.M. Interactions of copper and zinc in cardiovascular disease. *Ann. N.Y. Acad. Sci.*, 355: 140 (1980).
69. Katya-Katya, M., Ensminger, A., Mejean, L. and Derley, G. The effect of zinc supplementation on plasma cholesterol levels. *Nutr. Res.*, 4: 663 (1984).
70. Chandra, R.K. Excessive intake of zinc impairs immune responses. *J. Am. Med. Assoc.*, 252: 1443 (1984).
71. Cohen, J.M., Kamphake, L.J., Harris, E.K. and Woodward, R.L. Taste threshold concentrations of metals in drinking water. *J. Am. Water Works Assoc.*, 52: 660 (1960).
72. Anderson, E., Reinhard, C. and Hammel, W. The corrosion of zinc in various waters. *J. Am. Water Works Assoc.*, 26: 49 (1934).