

Micro-PIXE Analysis of Arctic Char, *Salvelinus alpinus*, Otoliths from Quttinirpaaq National Park, Nunavut

J.A. Babaluk, C.L. Evans, and J.D. Reist

Central and Arctic Region
Fisheries and Oceans Canada
Winnipeg, Manitoba R3T 2N6

2002

Canadian Data Report of Fisheries and Aquatic Sciences 1092



Fisheries and Oceans
Canada Pêches et Océans
Canada

Canada

**Canadian Data Report of
Fisheries and Aquatic Sciences**

Data reports provide a medium for filing and archiving data compilation where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of data reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was introduced with the publication of report number 161.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

**Rapport statistique canadien des
sciences halieutiques et aquatiques**

Les rapports statistiques servent a archiver les compilations de donnees pour les quelles il y a peu ou point d'analyse. Ces compilations auront d'ordinaire ete preparees a l'appui d'autres publications ou rapports. Les sujets des rapports statistiques refletent la vaste gamme des interets et des politiques du ministere des Peches et des Oceans, c'est-a-dire les sciences halieutiques et aquatiques.

Les rapports statistiques ne sont pas destine' a une vaste distribution et leur contenu ne doit pas etre mentionne dans une publication sans autorisation ecrite prealable de l'établissement auteur. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numeros 1 a 25 de cette serie ont ete publies a titre de releves statistiques, Services des peches et de la mer. Les numeros 26 a 160 ont ete publies a titre de rapports statistiques du Service des peches et de la mer, ministere des Peches et de l'Environnement. Le nom actuel de la serie a ete etabli lors de la parution du numero 161.

Les rapports statistiques sont produits a l'echelon regional, mais numerotes a l'echelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports epuisés seront fournis contre retribution par des agents commerciaux.

Canadian Data Report of
Fisheries and Aquatic Sciences 1092

2002

**MICRO-PIXE ANALYSIS OF ARCTIC CHAR, *SALVELINUS ALPINUS*,
OTOLITHS FROM QUTTINIRPAAQ NATIONAL PARK, NUNAVUT**

by

J.A. Babaluk, C.L. Evans, and J.D. Reist

Central and Arctic Region
Fisheries and Oceans Canada
Winnipeg, MB R3T 2N6

© Her Majesty the Queen in Right of Canada, 2002

Cat. No. Fs 97-13/1092E ISSN 0706-6465

Correct citation for this publication is:

Babaluk, J.A., C.L. Evans, and J.D. Reist. 2002. Micro-PIXE analysis of Arctic char, *Salvelinus alpinus*, otoliths from Quttinirpaaq National Park, Nunavut. Can. Data Rep. Fish. Aquat. Sci. 1092: xiv + 143 p.

ERRATA

Babaluk, J.A., C.L. Evans, and J.D. Reist. 2002. Micro-PIXE analysis of Arctic char, *Salvelinus alpinus*, otoliths from Quttinirpaaq National Park, Nunavut. Can. Data Rep. Fish. Aquat. Sci. 1092: xiv + 143 p.

On pages xi and 111, Fig. 106 caption should read “Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (**285 mm, 511 g, female, 14 yr**) caught in Heintzelman Lake, May/June 1995...”.

On pages xi and 112, Fig. 107 caption should read “Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (**252 mm, 314 g, male, 11 yr**) caught in Heintzelman Lake, May/June 1995...”.

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT/RÉSUMÉ	xiv
INTRODUCTION.....	1
MATERIALS AND METHODS	2
Description of the study area	2
Otolith collection	2
Otolith preparation.....	2
Micro-PIXE analysis	2
DATA PRESENTATION	3
ACKNOWLEDGMENTS.....	3
REFERENCES.....	3

LIST OF TABLES

Table

1	Summary of micro-PIXE point analysis data from Arctic char otoliths from Quttinirpaaq National Park, Nunavut	5
---	--	---

LIST OF FIGURES

Figure

1	Map of the Lake Hazen area, Quttinirpaaq National Park, Nunavut showing collection locations of Arctic char used for micro-PIXE analysis and dates (year) fish were collected	6
2	(a) Lateral side of the external surface of an Arctic char otolith: broken line (---) indicates transverse cut to expose the internal surface, (b) diagram of the exposed surface of an otolith showing typical areas (dots) where point analyses and transect (---) along which line-scan were conducted and (c) diagram showing the general features and arrangement of the University of Guelph scanning proton microprobe.	7
3	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (291 mm, 160 g, female, 13 yr) caught in Lake A, May 1996. Point analysis results are also indicated.....	8
4	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (588 mm, 1730 g, male, 28 yr) caught in Lake A, May 1996. Point analysis results are also indicated.....	9

<u>Figure</u>		<u>Page</u>
5	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (539 mm, 1246 g, male, 20 yr) caught in Lake A, May 1996. Point analysis results are also indicated	10
6	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (541 mm, 1678 g, male, 17 yr) caught in Lake A, May 1996. Point analysis results are also indicated	11
7	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (582 mm, 1965 g, male, 17 yr) caught in Lake A, May 1996. Point analysis results are also indicated	12
8	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (524 mm, 1352 g, female, 20 yr) caught in Lake A, May 1996. Point analysis results are also indicated	13
9	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (289 mm, 210 g, female, 15 yr) caught in Lake A, May 1996. Point analysis results are also indicated	14
10	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (290 mm, 189 g, female, 18 yr) caught in Lake A, May 1996. Point analysis results are also indicated	15
11	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (259 mm, 163 g, female, 10 yr) caught in Lake A, May 1996. Point analysis results are also indicated	16
12	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (286 mm, 171 g, male, 16 yr) caught in Lake A, May 1996. Point analysis results are also indicated	17
13	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (366 mm, 869 g, female, 19 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	18
14	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (310 mm, 618 g, male, 21 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	19
15	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (519 mm, 3242 g, female, 14 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	20
16	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (201 mm, 156 g, male, 18 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	21
17	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (222 mm, 184 g, male, 19 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	22
18	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (177 mm, 101 g, female, 18 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	23

<u>Figure</u>		<u>Page</u>
19	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (220 mm, 100 g, male, 21 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	24
20	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (219 mm, 87 g, female, 14 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	25
21	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (262 mm, 150 g, female, 13 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.....	26
22	Strontium (top) and zinc (bottom) profiles from two scanning proton microprobe line-scans of an otolith from an Arctic char (519 mm, 3242 g, 14 yr) caught in Alexandra Lake, May 1995. Point analysis was conducted in conjunction with one of the line-scans and the results are indicated	27
23	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (257 mm, 119 g, female, 14 yr) caught in Lake B, May 1996. Point analysis results are also indicated	28
24	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (245 mm, 131 g, female, 10 yr) caught in Lake B, May 1996. Point analysis results are also indicated	29
25	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (480 mm, 916 g, male, 26 yr) caught in Lake B, May 1996. Point analysis results are also indicated	30
26	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (376 mm, 348 g, male, 27 yr) caught in Lake B, May 1996. Point analysis results are also indicated	31
27	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (262 mm, 123 g, female, 19 yr) caught in Lake B, May 1996. Point analysis results are also indicated	32
28	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (257 mm, 137 g, female, 15 yr) caught in Lake B, May 1996. Point analysis results are also indicated	33
29	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (269 mm, 169 g, male, 14 yr) caught in Lake B, May 1996. Point analysis results are also indicated	34
30	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (244 mm, 121 g, female, 11 yr) caught in Lake B, May 1996. Point analysis results are also indicated	35
31	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (247 mm, 143 g, male, 11 yr) caught in Lake B, May 1996. Point analysis results are also indicated	36
32	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (272 mm, 150 g, male, 16 yr) caught in Lake B, May	

<u>Figure</u>		<u>Page</u>
	1996. Point analysis results are also indicated	37
33	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (243 mm, 141 g, female, 12 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.....	38
34	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (244 mm, 124 g, male, 11 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.....	39
35	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (255 mm, 148 g, male, 11 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.....	40
36	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (238 mm, 123 g, male, 11 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated	41
37	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (232 mm, 123 g, female, 10 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated	42
38	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (223 mm, 95 g, male, 10 yr) caught in Craig Lake, June 1992. Point analysis results were also indicated	43
39	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (281 mm, 184 g, male, 16 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated	44
40	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (274 mm, 184 g, male, 13 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated	45
41	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (259 mm, 175 g, female, 13 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated	46
42	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (173 mm, 44 g, female, 20 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	47
43	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (178 mm, 54 g, male, 14 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	48
44	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (180 mm, 58 g, female, 24 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	49
45	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (123 mm, 22 g, female, 15 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	50
46	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan	

<u>Figure</u>	<u>Page</u>
of an otolith from an Arctic char (131 mm, 21 g, male, 13y r) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	51
47 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (126 mm, 22 g, male, 13 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	52
48 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (218 mm, 90 g, female, 20 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	53
49 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (216 mm, 103 g, female, 21 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	54
50 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (489 mm, 1540 g, male, 20 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	55
51 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (170 mm, 46 g, female, 21 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated	56
52 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (633 mm, 2450 g, male, 20 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	57
53 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (347 mm, 450 g, male, not aged) caught in Lake Hazen, August 1981. Point analysis results are also indicated	58
54 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (574 mm, 1475 g, male, 25 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	59
55 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (420 mm, 625 g, male, 18 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	60
56 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (717 mm, 4800 g, male, 22 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	61
57 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (610 mm, 2250 g, male, 19 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	62
58 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (390 mm, 600 g, female, 23 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	63
59 Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (664 mm, 3250 g, male, 21 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	64

<u>Figure</u>		<u>Page</u>
60	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (557 mm, 1700 g, male, 21 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	65
61	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (472 mm, 1000 g, female, 18 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	66
62	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (417 mm, 700 g, male, 10 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	67
63	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (470 mm, 1000 g, female, 14 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	68
64	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (365 mm, 450 g, male, 17 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	69
65	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (353 mm, 400 g, male, 21 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	70
66	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (435 mm, 725 g, male, 18 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated	71
67	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (357 mm, 333 g, male, 22 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated	72
68	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (348 mm, 416 g, female, 26 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated	73
69	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (326 mm, 365 g, male, 19 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated	74
70	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (328 mm, 347 g, female, 21 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated	75
71	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (362 mm, 448 g, male, 22 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated	76
72	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (370 mm, 482 g, male, 22 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated	77
73	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (391 mm, 533 g, male, 23 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	78

<u>Figure</u>		<u>Page</u>
74	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (362 mm, 483 g, male, 25 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	79
75	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (366 mm, 472 g, male, 24 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	80
76	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (611 mm, 2170 g, male, 22 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	81
77	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (635 mm, 2216 g, male, 24 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	82
78	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (478 mm, 979 g, female, 22 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	83
79	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (578 mm, 1956 g, male, 21 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	84
80	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (618 mm, 2050 g, male, 24 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	85
81	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (570 mm, 1768 g, male, 27 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	86
82	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (560 mm, 1733 g, male, 27 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	87
83	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (452 mm, 836 g, male, 19 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	88
84	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (569 mm, 2037 g, male, 21 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	89
85	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (301 mm, 236 g, female, 16 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	90
86	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (565 mm, 1645 g, male, 21 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	91
87	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (672 mm, 1836 g, male, 25 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	92

<u>Figure</u>		<u>Page</u>
88	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (518 mm, 1275 g, female, 19 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated	93
89	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (435 mm, 810 g, female, 16 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated	94
90	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (533 mm, 1270 g, female, 25 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated	95
91	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (630 mm, 1820 g, male, 21 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated	96
92	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (177 mm, 48 g, male, 11 yr) of unknown form caught in Lake Hazen, July 1998. Point analysis results are also indicated	97
93	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (644 mm, 2410 g, male, 21 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated	98
94	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (172 mm, 42 g, female, 13 yr) of unknown form caught in Lake Hazen, July 1998. Point analysis results are also indicated	99
95	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (524 mm, 1190 g, female, 22 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated	100
96	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (549 mm, 1805 g, female, 20 yr) caught in Lake Hazen, July 1998. Point analysis results are also indicated	101
97	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (531 mm, 1803 g, female, 18 yr) caught in Lake Hazen, July 1998. Point analysis results are also indicated	102
98	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (397 mm, 520 g, male, 21 yr) caught in Lake Hazen, July 1998. Point analysis results are also indicated.....	103
99	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (174 mm, 126 g, female, 9 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	104
100	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (243 mm, 261 g, male, 8 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	105
101	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (231 mm, 227 g, female, 17 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	106

<u>Figure</u>		<u>Page</u>
102	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (187 mm, 146 g, male, 12 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	107
103	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (198 mm, 164 g, female, 13 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	108
104	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (224 mm, 209 g, male, 24 yr) caught in Heintzelman Lake, June/May 1995. Point analysis results are also indicated.....	109
105	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (246 mm, 292 g, female, 20 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	110
106	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (252 mm, 314 g, male, 11 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	111
107	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (285 mm, 511 g, female, 14 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	112
108	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (336 mm, 565 g, female, 11 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.....	113
109	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (268 mm, 167 g, female, 17 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	114
110	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (292 mm, 165 g, female, 22 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	115
111	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (275 mm, 197 g, female, 15 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	116
112	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (308 mm, 283 g, male, 22 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	117
113	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (310 mm, 252 g, male, 19 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	118
114	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (277 mm, 200 g, female, 20 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	119
115	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (293 mm, 218 g, male, 19 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	120

<u>Figure</u>		<u>Page</u>
116	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (283 mm, 223 g, male, 18 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	121
117	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (264 mm, 177 g, female, 16 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	122
118	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (260 mm, 171 g, male, 13 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.....	123
119	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (158 mm, 54 g, male, 10 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	124
120	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (124 mm, 26 g, male, 6 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	125
121	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (134 mm, 33 g, female, 5 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	126
122	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (161 mm, 47 g, male, 7 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	127
123	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (145 mm, 42 g, female, 5 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	128
124	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (153 mm, 46 g, male, 11 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	129
125	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (539 mm, 1949 g, male, 18 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	130
126	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (168 mm, 47 g, female, 11 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	131
127	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (176 mm, 53 g, female, 7 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	132
128	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (153 mm, 46 g, female, 9 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated	133
129	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (590 mm, 1946 g, female, 24 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	134

<u>Figure</u>		<u>Page</u>
130	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (630 mm, 2617 g, female, 20 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	135
131	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (196 mm, 63 g, female, 15 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	136
132	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (669 mm, 2639 g, male, 30 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	137
133	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (529 mm, 1522 g, female, 27 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	138
134	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (635 mm, 1928 g, male, 27 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	139
135	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (227 mm, 83 g, female, 17 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	140
136	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (182 mm, 59 g, male, 14 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	141
137	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (689 mm, 3316 g, male, 27 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	142
138	Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (168 mm, 46 g, male, 13 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated	143

ABSTRACT

Babaluk, J.A., C.L. Evans, and J.D. Reist. 2002. Micro-PIXE analysis of Arctic char, *Salvelinus alpinus*, otoliths from Quttinirpaaq National Park, Nunavut. Can. Data Rep. Fish. Aquat. Sci. 1092: xiv + 143 p.

Micro-PIXE analysis (line-scans) was used to determine the distribution of strontium (Sr) and zinc (Zn) in otoliths of Arctic char, *Salvelinus alpinus*, from 10 lakes in Quttinirpaaq National Park, Ellesmere Island, Nunavut. Profiles of these distributions are presented. Trace element (Sr and Zn) concentrations in otolith cores from these fish were also determined by micro-PIXE analysis (point analysis) and are tabulated.

Key words: anadromy; life history; otolith microchemistry; scanning proton microprobe; trace elements.

RÉSUMÉ

Babaluk, J.A., C.L. Evans, and J.D. Reist. 2002. Micro-PIXE analysis of Arctic char, *Salvelinus alpinus*, otoliths from Quttinirpaaq National Park, Nunavut. Can. Data Rep. Fish. Aquat. Sci. 1092: xiv + 143 p.

Nous avons utilisé la micro-analyse PIXE par balayage pour déterminer la répartition du (Sr) et du zinc (Zn) dans des otolithes d'omble chevalier (*Salvelinus alpinus*) capturés dans dix lacs du parc national Quttinirpaaq, situé sur l'île d'Ellesmere (Nunavut). Nous présentons des profils de ces répartitions. Nous avons aussi mesuré les concentrations des éléments traces Sr et Zn dans le centre des otolithes de ces poissons par micro-analyse PIXE ponctuelle : les résultats sont présentés dans un tableau.

Mots clés: anadromie; cycle vital; microchimie des otolithes; microsonde à proton à balayage; éléments traces.

INTRODUCTION

A basic understanding of the biology and diversity of the Arctic char, *Salvelinus alpinus*, populations in Quttinirpaaq National Park, formerly Ellesmere Island National Park Reserve, is fundamental to Parks Canada's long-term conservation and management plans for these fish (Parks Canada 1994). Arctic char is the only freshwater fish species in the Canadian High Arctic including Quttinirpaaq National Park (Scott and Crossman 1973; Parks Canada 1994). However, the species exhibits great diversity at a number of levels below that of species including: a) life history type (e.g., anadromous or non-anadromous), b) ecological type (e.g., pelagic or benthic forms), c) trophic type (e.g., planktivore or piscivore), d) evolutionary lineages (e.g., subspecies, biological stocks), as well as, e) variants within many of the above types (e.g., life history variability). This diversity can be observed through a number of techniques, but is best understood using several complementary approaches such as: a) morphology (e.g., body form and function), b) genetics (e.g., delineation of groups using DNA chemistry), c) stable isotopes and diet (e.g., trophic pattern), d) population dynamics (e.g., age and size structure), and e) otolith microchemistry (e.g., life history types, stock discrimination).

Fisheries and Oceans Canada (DFO), in collaboration with Parks Canada, has assessed the diversity in Quttinirpaaq National Park Arctic char populations since 1990. In 1990, a study on contaminants in Lake Hazen sediments was conducted (Muir et al. 1995). At the same time, Arctic char were also collected for a study on contaminants in muscle tissue (Muir and Lockhart 1992; Muir and Lockhart 1993). Between 1992 and 1998, DFO, assisted by Parks Canada, intermittently assessed Arctic char populations within Quttinirpaaq National Park, in particular Lake Hazen Arctic char. Reist et al. (1995) showed that there were at least two morphotypes or forms of Arctic char in Lake Hazen (a small and a large form). Some results of other studies have been reported by Brown Gladden et al. (1995), Reist et al. (1995), Babaluk et al. (1997), Babaluk et al. (2001), Babaluk et al. (2002) and Guiquier et al. (2002). Before this area of northern Ellesmere Island became initially, a national park reserve and then, a national park, it had only been visited by two other groups of fisheries researchers: 1) in 1958, a group under the auspices of the Defence Research Board of Canada collected biological

information on the Arctic char in Lake Hazen (Hunter 1960; Johnson 1983), and 2) in 1981, DFO biologists collected additional information on Lake Hazen Arctic char (Johnson 1983).

Otoliths ("ear stones") are small, paired structures found in a fish's inner ear used to sense orientation and acceleration. They are composed mainly of a calcium carbonate (aragonite) and protein matrix; but trace elements similar to calcium (Ca), such as strontium (Sr) and zinc (Zn), can be incorporated into or replace Ca in the otolith (Degens et al. 1969). The Ca and trace elements are derived mainly from the waters that the fish inhabits (Ichii and Mugiya 1983). Otoliths do not undergo resorption during the life of the fish (Simkiss 1974) and thus provide an elemental record of the fish's life.

Sea water contains, on average, 8.0 mg L^{-1} Sr whereas freshwater contains, on average, 0.1 mg L^{-1} Sr (Rosenthal et al. 1970). Local freshwater Sr concentrations also can vary. For example, in Quttinirpaaq National Park lakes, Sr concentrations range from $0.033 - 0.329 \text{ mg L}^{-1}$ (Babaluk et al. 1999). Sr concentrations in otoliths closely reflect environmental levels of Sr (Babaluk et al. 1998) and thus provide information on the types of environments occupied at particular times of life. This can be used to reconstruct life history and also to differentiate specific environments occupied.

Previous studies using micro-Proton-Induced X-ray Emission (micro-PIXE) analysis of Sr in Arctic char otoliths showed that differences in Sr concentrations can retrospectively indicate anadromous behaviour or, in the case of Lake Hazen Arctic char, indicated non-andromous behaviour (Babaluk et al. 1997) and, in the case of Dolly Varden, *S. malma*, provided details regarding the habitats occupied early in a fish's life (Babaluk et al. 1998).

Another micro-PIXE study, incorporating data published in this report, showed that Zn has a typically oscillatory variation that corresponds to the annular structure of Arctic char otoliths (Halden et al. 2000). Incorporation of Zn is linked to nutrient uptake in the fish so the concentration of Zn in the otolith must, in part, be a proxy for nutrient availability in the environment and perhaps the productivity of a particular environment. Zinc is an important trace nutrient that is necessary for the development of fish gonads (Lall 1989). The onset of sexual maturity

and the development of gonads will likely alter the metabolic distribution of Zn and this may be recorded in the otolith.

As part of the ongoing assessment of Arctic char populations in Quttinirpaaq National Park, we extended the otolith microchemistry study to: a) determine the extent of anadromy (if any) existing within Quttinirpaaq National Park Arctic char populations and b) provide additional trace element information, in particular Sr and Zn, to assist in discriminating stocks or populations. A summary of this work was presented at the Ninth International Conference on Particle-Induced X-ray Emission and its Analytical Applications, June 8 - 12, 2001 at the University of Guelph, Guelph, Ontario (Babaluk et al. 2002). This report presents detailed otolith microchemistry data from the Quttinirpaaq National Park Arctic char samples examined to date.

MATERIALS AND METHODS

DESCRIPTION OF THE STUDY AREA

Quttinirpaaq National Park is located at the northern end of Ellesmere Island, Nunavut (Fig. 1, left inset). The most northern area of the Park, Cape Aldrich at 83°06'N, is also the most northern point in the Canadian Arctic Archipelago. The park, with an area of 37 775 km², is the second largest national park in Canada and is described in detail in Parks Canada (1994).

There are relatively few lakes in Quttinirpaaq National Park and the most productive ones are located on the Hazen Plateau or along the coastal areas of the Grant Land Mountains (Fig. 1, right inset). Most of these lakes provide suitable habitat for some aquatic flora and invertebrates but are too shallow to support Arctic char (Parks Canada 1994). Lakes for this study were chosen because they were known to contain populations of Arctic char (R. Wissink, Parks Canada, Pangnirtung, pers. comm. 1992).

OTOLITH COLLECTION

Otoliths were collected from Arctic char from 10 lakes in Quttinirpaaq National Park: an unnamed lake, hereafter Lake A (82°06'N, 68°37'W) in 1996, Alexandra Lake (81°46'N, 65°32'W) in 1995, an unnamed lake, hereafter

Lake B (82°09'N, 68°29'W) in 1996, Craig Lake (81°52'N, 68°47'W) in 1992, Ekblaw Lake (81°40'N, 75°40'W) in 1998, Lake Hazen (81°50'N, 70°25'W) in 1981, 1990, 1992 and 1998, Heintzelman Lake (81°42'N, 66°56'W) in 1995, Kilbourne Lake (81°52'N, 68°25'W) in 1996, Lewis Lake (81°30'N, 74°35'W) in 1998, and Murray Lake (81°20'N, 69°34'W) in 1998 (Fig. 1). Table 1 shows the number of otoliths analysed from each lake.

OTOLITH PREPARATION

One of each pair of otoliths was prepared for micro-PIXE analysis. The otoliths were embedded in epoxy resin and a transverse cut was made such that a dorso-ventral cross section through the core of the otolith was created exposing all annuli (yearly growth increments) (Fig. 2a, b). The posterior half of each otolith was re-embedded in a standard 25 mm diameter leucite, geological probe-mount. The exposed otolith surfaces were sequentially ground (240, 320, 400, and 600 grit silicon carbide), then polished (5, 3, 1, and 0.3 µm aluminum oxide) and finally carbon-coated to prevent charging during microprobe analysis.

MICRO-PIXE ANALYSIS

The general arrangement and operation of the University of Guelph proton microprobe is shown in Figure 2c. Videomicrographs of the otolith sections were used as reference "maps" for the microprobe. A beam of protons, produced by an electrostatic accelerator, is focused on a spot, 5 - 10 µm wide, on the otolith. The excitation volume is a cylinder of approximately 5 - 10 µm in diameter and 30 µm in depth. Ninety percent of the observed Sr X-rays from the aragonite matrix of the otolith originate within a depth of 31 µm (Halden et al. 1996). The resultant X-rays emitted from the otolith were analysed using a silicon/lithium detector equipped with a compound filter of 125 µm Mylar and 106 µm aluminum. One-dimensional line-scans were obtained by rastering the proton beam along a transect from the core area to the dorsal edge of the otolith, incorporating all annuli (see Fig. 2b). Line-scanning of a typical Arctic char otolith took approximately 25 minutes at a proton beam current of 5 - 7 nA. The X-ray spectrum was used to identify trace elements present and the amount of a particular element (Sr and Zn, in this case) was measured by the height

of the X-ray peaks as counts. Three point (spot) analyses and, in some cases, six analyses in the core of each otolith were conducted (see Fig. 2b). Detection limits for Sr and Zn were 1 and 2 parts per million (ppm), respectively. Further details on proton microprobe theory and procedures are provided by Campbell et al. (1995).

DATA PRESENTATION

A summary of micro-PIXE point (spot) analysis data for Sr and Zn from Arctic char otoliths from lakes in Quttinirpaaq National Park is presented in Table 1. Line-scan data (as profiles) for Sr and Zn for Arctic char from these lakes are presented in Figures 3 - 138. For Lake Hazen Arctic char, the putative form of each fish is given when possible (1990, 1992 and 1998 samples). Point analysis data for Sr and Zn for individual Arctic char are presented as part of Figures 3 - 138 (insets). An electronic version of the raw data is available from the senior author c/o Fisheries and Oceans Canada, 501 University Crescent, Winnipeg, Manitoba, R3T 2N6, Canada.

ACKNOWLEDGMENTS

The work was done in part with the support of Parks Canada. Natural Resources Canada (Polar Continental Shelf Project) at Resolute, Nunavut provided logistical support. The authors thank D. Clark, E. Gyselman, J. Johnson, J. Maurice, V. Sahanatien, B. Troke, P. Wilkinson, R. Wissink, and G. Walker for field assistance and logistical support while at Quttinirpaaq National Park; K. Mathers, S. Mejia, and R. Wastle for otolith preparation; W. Teesdale for micro-PIXE analyses; P. Wilson for assistance with map preparation; and D. Laroque for assistance in preparing the manuscript. C. Read and T. Siferd provided constructive comments on an earlier version of the manuscript.

REFERENCES

BABALUK, J.A., J.L. CAMPBELL, C.L. EVANS, N.M. HALDEN, S.R. MEJIA, Z. NEJEDLY, J.D. REIST, and W.J. TEESDALE. 2002. Micro-PIXE analysis of strontium in Arctic char, *Salvelinus alpinus*, otoliths from

Quttinirpaaq National Park, Nunavut, Canada. Nucl. Instr. Meth. Phys. Res. B 189: 190-195.

BABALUK, J.A., N.M. HALDEN, J.D. REIST, A.H. KRISTOFFERSON, J.L. CAMPBELL, and W.J. TEESDALE. 1997. Evidence for non-anadromous behaviour of arctic charr (*Salvelinus alpinus*) from Lake Hazen, Ellesmere Island, Northwest Territories, Canada, based on scanning proton microprobe analysis of otolith strontium distribution. Arctic 50: 224-233.

BABALUK, J.A., L.G. HEURING, J.D. REIST, and B.N. BILLECK. 1999. Selected chemical analyses of water from lakes in Ellesmere Island National Park Reserve. Can. Data Rep. Fish. Aquat. Sci. 1050: iv + 7 p.

BABALUK, J.A., J.D. REIST, V.A. SAHANATIEN, N.M. HALDEN, J.L. CAMPBELL, and W.J. TEESDALE. 1998. Preliminary results of stock discrimination of chars in Ivavik National Park, Yukon Territory, Canada, using microchemistry of otolith strontium, p. 991-998. In N.W.P. Munro and J.H. Martin (eds.) Linking Protected Areas with Working Landscapes Conserving Biodiversity, Proceedings of the Third International Conference on Science and Management of Protected Areas, 12-16 May 1997. Wolfville, Canada: SAMPA.

BABALUK, J.A., H.R. WISSINK, B.G. TROKE, D.A. CLARK, and J.D. JOHNSON. 2001. Summer movements of radio-tagged arctic charr (*Salvelinus alpinus*) in Lake Hazen, Nunavut, Canada. Arctic 54: 418-424.

BROWN GLADDEN, J.G., L.D. POSTMA MAIERS, T.J. CARMICHAEL, and J.D. REIST. 1995. Mitochondrial DNA control region sequence variation in Arctic char (*Salvelinus alpinus* (L.)). Can. Data Rep. Fish. Aquat. Sci. 968: iv + 18 p.

CAMPBELL, J.L., W.J. TEESDALE, and N.M. HALDEN. 1995. Theory, practice and application of micro-PIXE analysis and element-distribution maps. Can. Mineral. 33: 279-292.

DEGENS, E.T., W.G. DEUSER, and R.L. HAEDRICH. 1969. Molecular structure and composition of fish otoliths. Mar. Biol. 2: 105-113.

- GUIGUER, K.R.R.A., J.D. REIST, M. POWER, and J.A. BABALUK. 2002. Using stable isotopes to confirm the trophic ecology of Arctic charr morphotypes from Lake Hazen, Nunavut, Canada. *J. Fish Biol.* 60: 348-362.
- HALDEN, N.M., J.A. BABALUK, A.H. KRISTOFFERSON, J.L. CAMPBELL, W.J. TEESDALE, J.A. MAXWELL, and J.D. REIST. 1996. Micro-PIXE studies of Sr zoning in Arctic charr otoliths: migratory behaviour and stock discrimination. *Nucl. Instr. Meth. Phys. Res. B* 109/110: 592-597.
- HALDEN, N.M., S.R. MEJIA, J.A. BABALUK, J.D. REIST, A.H. KRISTOFFERSON, J.L. CAMPBELL, and W.J. TEESDALE. 2000. Oscillatory zinc distribution in Arctic char (*Salvelinus alpinus*) otoliths: the result of biology or environment? *Fish. Res.* 46: 289-298.
- HUNTER, J.G. 1960. Hazen Lake, p. 18-20. *In* Annual report and investigators summaries (April 1, 1959 to March 31, 1960). *Fish. Res. Board Can. Arctic Unit*. Ottawa.
- ICHII, T., and Y. MUGIYA. 1983. Effects of dietary deficiency in calcium on growth and calcium uptake from the aquatic environment in the goldfish, *Carassius auratus*. *Comp. Biochem. Physiol.* 74(A): 259-262.
- JOHNSON, L. 1983. Homeostatic characteristics of single species fish stocks in Arctic lakes. *Can. J. Fish. Aquat. Sci.* 40: 987-1024.
- LALL, S.P. 1989. The minerals, p. 219-257. *In* J.E. Halver (ed.). *Fish nutrition*. Academic Press, Inc., Toronto.
- MUIR, D.C.G., N.P. GRIFFIT, W.L. LOCKHART, P. WILKINSON, B.N. BILLECK, and G.J. BRUNSKILL. 1995. Spatial trends and historical profiles of organochlorine pesticides in Arctic lake sediments. *Sci. Total Environ.* 160/161: 447-457.
- MUIR, D.C.G., and W.L. LOCKHART. 1992. Contaminant trends in freshwater biota, p. 121-125. *In* Synopsis of research conducted under the 1991/1992 Northern Contaminants Program. Indian and Northern Affairs Canada, Environmental Studies Report No. 68. Ottawa.
- MUIR, D.C.G., and W.L. LOCKHART. 1993. Contaminant trends in freshwater biota, p. 167-173. *In* Synopsis of research conducted under the 1992/1993 Northern Contaminants Program. Indian and Northern Affairs Canada, Environmental Studies Report No. 70. Ottawa.
- PARKS CANADA. 1994. Ellesmere Island National Park Reserve resource description and analysis. Department of Canadian Heritage, Winnipeg. Vol. 1 and 2.
- REIST, J.D., E. GYSELMAN, J.A. BABALUK, J.D. JOHNSON, and R. WISSINK. 1995. Evidence for two morphotypes of arctic char (*Salvelinus alpinus*) from Lake Hazen, Ellesmere Island, Northwest Territories, Canada. *Nord. J. Freshwater Res.* 71: 496-410.
- ROSENTHAL, H.L., M.M. EVES, and O.A. COCHRAN. 1970. Common strontium concentration of mineralized tissues from marine and sweet water animals. *Comp. Biochem. Physiol.* 32: 445-450.
- SCOTT, W.B., and E.J. CROSSMAN. 1973 (1979). *Freshwater fishes of Canada* (revised printing). *Bull. Fish. Res. Board Can.* 184: 966 p.
- SIMKISS, K. 1974. Calcium metabolism of fish in relation to ageing, p. 1-12. *In* T.B. Bagena (ed.). *Ageing of fish*. London: Unwin Brothers.

Table 1. Summary of micro-PIXE point analysis data from Arctic char otoliths from Quttinirpaaq National Park, Nunavut.

Lake	Year	Form	No. of otoliths/ no. of analyses	Mean Sr conc. (ppm \pm 1 standard deviation)	Mean Zn conc.
A	1996	all fish	10/30	426 \pm 52	64 \pm 21
Alexandra	1995	all fish	10/30	165 \pm 12	58 \pm 19
B	1996	all fish	10/30	490 \pm 64	61 \pm 18
Craig	1992	all fish	9/39	623 \pm 103	69 \pm 25
Ekblaw	1998	all fish	10/30	364 \pm 49	46 \pm 13
Hazen	1981	all fish	15/45	417 \pm 99	58 \pm 16
Hazen	1990	all fish	6/36	474 \pm 46	64 \pm 24
Hazen	1992	small	4/24	533 \pm 73	73 \pm 27
Hazen	1992	large	13/78	482 \pm 102	77 \pm 30
Hazen	1998	small	2/6	454 \pm 25	56 \pm 9
Hazen	1998	large	8/24	323 \pm 69	54 \pm 19
Hazen	all years	all fish	48/213	456 \pm 99	67 \pm 26
Heintzelman	1995	all fish	10/30	440 \pm 74	56 \pm 28
Kilbourne	1996	all fish	10/30	396 \pm 35	48 \pm 14
Lewis	1998	all fish	10/30	520 \pm 81	59 \pm 26
Murray	1998	all fish	10/30	179 \pm 55	63 \pm 16




Fig. 1. Map of the Lake Hazen area, Quttinirpaaq National Park, Nunavut showing collection locations of Arctic char used for micro-PIXE analysis and dates (year) fish were collected.




Fig. 2. (a) Lateral side of the external surface of an Arctic char otolith: broken line (---) indicates transverse cut to expose the internal surface, (b) diagram of the exposed surface of an otolith showing typical areas (dots) where point analyses and transect (---) along which line-scan were conducted and (c) diagram showing the general features and arrangement of the University of Guelph scanning proton microprobe.




Fig. 3. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (291 mm, 160 g, female, 13 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 4. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (588 mm, 1730 g, male, 28 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 5. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (539 mm, 1246 g, male, 20 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 6. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (541 mm, 1678 g, male, 17 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 7. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (582 mm, 1965 g, male, 17 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 8. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (524 mm, 1352 g, female, 20 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 9. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (289 mm, 210 g, female, 15 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 10. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (290 mm, 189 g, female, 18 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 11. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (259 mm, 163 g, female, 10 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 12. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (286 mm, 171 g, male, 16 yr) caught in Lake A, May 1996. Point analysis results are also indicated.




Fig. 13. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (366 mm, 869 g, female, 19 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 14. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (310 mm, 618 g, male, 21 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 15. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (519 mm, 3242 g, female, 14 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 16. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (201 mm, 156 g, male, 18 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 17. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (222 mm, 184 g, male, 19 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 18. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (177 mm, 101 g, female, 18 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 19. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (220 mm, 100 g, male, 21 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 20. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (219 mm, 87 g, female, 14 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 21. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (262 mm, 150 g, female, 13 yr) caught in Alexandra Lake, May 1995. Point analysis results are also indicated.




Fig. 22. Strontium (top) and zinc (bottom) profiles from two scanning proton microprobe line-scans of an otolith from an Arctic char (519 mm, 3242 g, female, 14 yr) caught in Alexandra Lake, May 1995. Point analysis was conducted in conjunction with one of the line-scans and the results are indicated.




Fig. 23. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (257 mm, 119 g, female, 14 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 24. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (245 mm, 131 g, female, 10 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 25. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (480 mm, 916 g, male, 26 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 26. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (376 mm, 348 g, male, 27 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 27. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (262 mm, 123 g, female, 19 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 28. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (257 mm, 137 g, female, 15 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 29. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (269 mm, 169 g, male, 14 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 30. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (244 mm, 121 g, female, 11 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 31. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (247 mm, 143 g, male, 11 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 32. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (272 mm, 150 g, male, 16 yr) caught in Lake B, May 1996. Point analysis results are also indicated.




Fig. 33. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (243 mm, 141 g, female, 12 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 34. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (244 mm, 124 g, male, 11 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 35. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (255 mm, 148 g, male, 11 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 36. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (238 mm, 123 g, male, 11 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 37. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (232 mm, 123 g, female, 10 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 38. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (223 mm, 95 g, male, 10 yr) caught in Craig Lake, June 1992. Point analysis results were also indicated.




Fig. 39. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (281 mm, 184 g, male, 16 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 40. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (274 mm, 184 g, male, 13 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 41. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (259 mm, 175 g, female, 13 yr) caught in Craig Lake, June 1992. Point analysis results are also indicated.




Fig. 42. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (173 mm, 44 g, female, 20 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 43. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (178 mm, 54 g, male, 14 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 44. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (180 mm, 58 g, female, 24 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 45. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (123 mm, 22 g, female, 15 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 46. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (131 mm, 21 g, male, 13 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 47. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (126 mm, 22 g, male, 13 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 48. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (218 mm, 90 g, female, 20 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated,




Fig. 49. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (216 mm, 103 g, female, 21 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 50. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (489 mm, 1540 g, male, 20 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 51. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (170 mm, 46 g, female, 21 yr) caught in Ekblaw Lake, July 1998. Point analysis results are also indicated.




Fig. 52. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (633 mm, 2450 g, male, 20 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 53. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (347 mm, 450 g, male, not aged) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 54. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (574 mm, 1475 g, male, 25 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 55. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (420 mm, 625 g, male, 18 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 56. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (717 mm, 4800 g, male, 22 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 57. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (610 mm, 2250 g, male, 19 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 58. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (390 mm, 600 g, female, 23 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 59. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (664 mm, 3250 g, male, 21 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 60. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (557 mm, 1700 g, male, 21 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 61. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (472 mm, 1000 g, female, 18 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 62. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (417 mm, 700 g, male, 10 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 63. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (470 mm, 1000 g, female, 14 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 64. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (365 mm, 450 g, male, 17 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 65. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (353 mm, 400 g, male, 21 yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 66. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (435 mm, 725 g, male, 18yr) caught in Lake Hazen, August 1981. Point analysis results are also indicated.




Fig. 67. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (357 mm, 333 g, male, 22 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated.




Fig. 68. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (348 mm, 416 g, female, 26 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated.




Fig. 69. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (326 mm, 365 g, male, 19 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated.




Fig. 70. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (328 mm, 347 g, female, 21 yr) Caught in Lake Hazen, May 1990. Point analysis results are also indicated.




Fig. 71. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (362 mm, 448 g, male, 22 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated.




Fig. 72. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (370 mm, 482 g, male, 22 yr) caught in Lake Hazen, May 1990. Point analysis results are also indicated.




Fig. 73. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (391 mm, 533 g, male, 23 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 74. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (362 mm, 483 g, male, 25 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 75. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a small-form Arctic char (366 mm, 472 g, male, 24 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 76. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (611 mm, 2170 g, male, 22 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 77. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (635 mm, 2216 g, male, 24 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 78. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (478 mm, 979 g, female, 22 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 79. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (578 mm, 1956 g, male, 21 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 80. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (618 mm, 2050 g, male, 24 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 81. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (570 mm, 1768 g, male, 27 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 82. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (560 mm, 1733 g, male, 27 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 83. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (452 mm, 836 g, male, 19 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 84. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (569 mm, 2037 g, male, 21 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 85. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (301 mm, 236 g, female, 16 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 86. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (565 mm, 1645 g, male, 21 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 87. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (672 mm, 1836 g, male, 25 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 88. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (518 mm, 1275 g, female, 19 yr) caught in Lake Hazen, June 1992. Point analysis results are also indicated.




Fig. 89. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (435 mm, 810 g, female, 16 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated.




Fig. 90. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (533 mm, 1270 g, female, 25 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated.




Fig. 91. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (630 mm, 1820 g, male, 21 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated.




Fig. 92. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (177 mm, 48 g, male, 11 yr) of unknown form caught in Lake Hazen, July 1998. Point analysis results are also indicated.




Fig. 93. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (644 mm, 2410 g, male, 21 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated.




Fig. 94. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (172 mm, 42 g, female, 13 yr) of unknown form caught in Lake Hazen, July 1998. Point analysis results are also indicated.




Fig. 95. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (524 mm, 1190 g, female, 22 yr) caught in Lake Hazen, August 1998. Point analysis results are also indicated.




Fig. 96. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (549 mm, 1805 g, female, 20 yr) caught in Lake Hazen, July 1998. Point analysis results are also indicated.




Fig. 97. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (531 mm, 1803 g, female, 18 yr) caught in Lake Hazen, July 1998. Point analysis results are also indicated.




Fig. 98. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from a large-form Arctic char (397 mm, 520 g, male, 21 yr) caught in Lake Hazen, July 1998. Point analysis results are also indicated.




Fig. 99. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (174 mm, 126 g, female, 9 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 100. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (243 mm, 261 g, male, 8 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 101. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (231 mm, 227 g, female, 17 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 102. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (187 mm, 146 g, male, 12 yr) caught in Heintzman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 103. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (198 mm, 164 g, female, 13 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 104. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (224 mm, 209 g, male, 24 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 105. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (246 mm, 292 g, female, 20 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 106. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (252 mm, 314 g, male, 11 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 107. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (285 mm, 511 g, female, 14 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 108. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (336 mm, 565 g, female, 11 yr) caught in Heintzelman Lake, May/June 1995. Point analysis results are also indicated.




Fig. 109. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (268 mm, 167 g, female, 17 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 110. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (292 mm, 165 g, female, 22 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 111. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (275 mm, 197 g, female, 15 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 112. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (308 mm, 283 g, male, 22 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 113. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (310 mm, 252 g, male, 19 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 114. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (277 mm, 200 g, female, 20 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 115. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (293 mm, 218 g, male, 19 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 116. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (283 mm, 223 g, male, 18 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 117. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (264 mm, 177 g, female, 16 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 118. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (260 mm, 171 g, male, 13 yr) caught in Kilbourne Lake, May 1996. Point analysis results are also indicated.




Fig. 119. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (158 mm, 54 g, male, 10 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 120. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (124 mm, 26 g, male, 6 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 121. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (134 mm, 33 g, female, 5 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 122. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (161 mm, 47 g, male, 7 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 123. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (145 mm, 42 g, female, 5 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 124. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (153 mm, 46 g, male, 11 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 125. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (539 mm, 1949 g, male, 18 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 126. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (168 mm, 47 g, female, 11 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 127. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (176 mm, 53 g, female, 7 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 128. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (153 mm, 46 g, female, 9 yr) caught in Lewis Lake, July 1998. Point analysis results are also indicated.




Fig. 129. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (590 mm, 1946 g, female, 24 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 130. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (630 mm, 2617 g, female, 20 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 131. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (196 mm, 63 g, female, 15 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 132. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (669 mm, 2639 g, male, 30 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 133. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (529 mm, 1522 g, female, 27 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 134. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (635 mm, 1928 g, male, 27 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 135. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (227 mm, 83 g, female, 17 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 136. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (182 mm, 59 g, male, 14 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 137. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (689 mm, 3316 g, male, 27 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.




Fig. 138. Strontium (top) and zinc (bottom) profiles from a scanning proton microprobe line-scan of an otolith from an Arctic char (168 mm, 46 g, male, 13 yr) caught in Murray Lake, July 1998. Point analysis results are also indicated.