



Agriculture and
Agri-Food Canada

Agriculture et
Agroalimentaire Canada



Algae Identification

FIELD GUIDE

Agriculture and Agri-Food Canada
Agri-Environment Services Branch

Canada 

This publication is cited as:
Huynh, M. and N. Serediak. 2006.
Algae Identification Field Guide.
Agriculture and Agri-Food Canada.
40 pages.

This publication is accompanied
by the Algae Identification Lab Guide
by Huynh and Serediak (2006).

© Her Majesty the Queen in
Right of Canada, 2011
Cat. No. A125-8/2-2011E-PDF
ISBN 978-1-100-18309-1
AAFC No. 11431E

Aussi offert en français sous le titre :
Identification des algues : Guide de terrain

For additional copies of this publication or to
request an alternate format, please contact:
Publications Services

Agriculture and Agri-Food Canada
Tower 4, Floor 0, Room 160
1341 Baseline Road
Ottawa (Ontario) K1A 0C5 Canada
Telephone: (613) 773-1444
Fax: (613) 773-1498
E-mail: publications@agr.gc.ca

Disclaimer

Responsibility for interpretation of the content
contained in this publication rests with the user.
Information in this publication is provided solely
for the user's information and, while thought to be
accurate, is provided strictly "as is" and without
warranty of any kind, either expressed or implied.
Agriculture and Agri-Food Canada will not be liable
to you for any damages, direct or indirect, or lost
profits arising out of your use of this publication.

Algae Identification **Field Guide**

AN ILLUSTRATIVE FIELD GUIDE ON IDENTIFYING
COMMON ALGAE FOUND IN THE CANADIAN PRAIRIES

Agriculture and Agri-Food Canada
Agri-Environment Services Branch

Nancy Serediak
Streamline Consulting
Edmonton, AB Canada



Mai-Linh Huynh
Agriculture and Agri-Food Canada
Regina, SK Canada

Acknowledgments

The authors sincerely thank the reviewers and supporters of this project: AAFC/AESB Water Quality Division; Glen Brandt, AAFC-AESB; Garth Mottershead, AAFC-AESB; Wayne Wark, AAFC Communications; Mark Graham, University of Alberta; Ron Zurawell, Government of Alberta; and Bob Klemmer, Government of Saskatchewan. Their comments and recommendations have proven invaluable for ensuring the relevance and scientific accuracy of the reference manual and field guide. The authors are also grateful to the following for allowing use of their photos: Charles Delwiche, David Patterson (micro*scope website), Morgan Vis, David Krogmann and Mark Schneegurt (Cyanosite website), Mark Graham, Steve Murrell, Ron Zurawell, David John, Brian Whitton, Peter York, Jane Jamieson, and Nick Stewart.

The authors would also like to acknowledge Steve Murrell, AAFC/AESB Water Quality Division, for coordinating and leading the quality assurance component of this project.

The authors would also like to sincerely thank Michael Parry, AAFC-AESB GIS Unit in Calgary Alberta, for his hard work and collaboration in developing the GIS-based program tool for this project.

Foreword

The algae identification field guide and accompanying lab reference manual were created for agricultural field personnel interested in algae, such as agricultural extension staff, watershed groups and individuals who work with agricultural surface water supplies. Algae listed in this guide and manual are not organized according to their evolutionary relationship (i.e. taxonomically) as with most traditional classification keys, but are presented based on common situations or issues that are significant to our target audience. Algae are also classified into four main groups (Palmer 1962): Blue-Greens, Greens, Diatoms and Flagellates. Information on these algal groups are found in the Algae Identification Lab Guide.

The objective of the guide and manual is to provide a starting point in identifying the most common or easily identifiable algal genera typically found in lakes, dugouts, wetlands and other freshwater lentic systems that are significant to our target audience. This starting point for algae identification is outlined as a two-step approach:

- A.** In the field, where depending on the situation, using the three senses of sight, smell and touch will help give the user a primary assessment of which algal group or genera the algae might belong to; and
- B.** In the lab, where the identification of algae is validated using a compound microscope. Since there exists many diverse species within each genus of algae and many looking very similar macroscopically, verification of algae must be done using a microscope.

Provided is a two part document on algae identification: the Algae Identification Field Guide (Part A) and Algae Identification Lab Reference Manual (Part B). As well, the field guide is available as a program tool compatible with ESRI's GIS program ArcPAD, and is found attached to this guide as a CD-ROM. This program tool improves the efficiency of algae identification and data collection than can otherwise be done with paper-based field guides and data collection forms, as well as provides a way in which each sample can be plotted spatially for future reference.

Table of Contents

Introduction	8
The starting point	10
Surface bloom-forming algae	12
Algae interfering with coagulation	26
Attached algae and algal mats found near shore	28
Potentially threatening freshwater algae	30
Glossary of Terms	33
Photo credits	34
References	36
Recommended Reading	37
Index	38

Introduction

What are algae?

They are organisms with a mixed evolutionary history that have four main commonalities:

1. They are very simple organisms with no vascular tissue. The only exception are brown algae where they have a higher degree of organ differentiation.
2. They have naked reproductive structures, i.e. there is no protective layer of cells around reproductive structures.
3. They are photoautotrophic, i.e. they produce their own food materials through photosynthesis by using sunlight, water and CO₂. There are some exceptions, such as some species of *Euglena*, where they ingest other organisms for food because they do not have any chloroplasts.
4. Similar to plants, they contain chlorophyll. However, not all algae are classified as plants.

Where are they found?

Algae commonly grow in any habitat where water or moisture is found. Habitats include fresh and salt water bodies, hot springs, ice, air, and in or on other organisms and substrate. In the Canadian Prairies, they are more common in summer than winter.

They are also found in communities, living with many other different species of algae, plankton and zooplankton. These communities can tell you a lot about the health of the ecosystem. For example, a community of *Euglena*, *Scenedesmus* and *Selenastrum* likely indicates a eutrophic water body.

Why study algae?

Algae are important indicators of ecosystem health and integrity because they form the base of most aquatic food chains. Virtually all aquatic animals are dependent on this primary producer. Algae are also an excellent indicator of water quality, as their abundance and community composition most often reflects (and has the capacity to affect) the chemical properties of water such as pH and nutrient levels.

Their ability to grow in large quantities can lead to dramatic changes in the appearance, taste and odour of water, and can negatively affect organisms in higher trophic levels (e.g. fish kills). Algae also can interfere with general water use and distribution, impairing water pumps, filters, pipes, animal troughs and misters, boilers and cooling equipment. For prairie water sources, algae can pose unique challenges for a variety of water uses because of the common and prolific growth cycles of algae during the open water season.

Knowledge about algae can provide valuable insights into water quality, indicating concerns such as pollution to waterways, reduction in water flow, or a health and safety risk, as some algal species are highly toxic to humans and livestock. Of significant concern for water users is Cyanobacteria (sometimes referred to as “blue-green algae”). Cyanobacteria is actually a bacteria that may produce lethal neurotoxins (brain toxin) or hepatotoxins (liver toxin), which can cause serious illness or death in humans and animals if the toxins are ingested.

It is important to note that while algae tend to be the most notable problem or issue in an aquatic ecosystem, other planktonic biota or organisms, including zooplankton and macroorganisms (both plants and animals), can also affect the physical, chemical and biotic properties of a water body. These factors should be taken into account when studying the aquatic ecosystem as a whole.

The starting point



1

Step One.

Which type of situation are you dealing with?

- A. A surface water bloom
Go to page 12
- B. Algae interfering with coagulation
Go to page 26
- C. Attached algae and algae mats found near shore
Go to page 28
- D. Potentially threatening freshwater algae
Go to page 30



2

Step Two.

Once you have matched your alga to a close description, there is a page number beside the algal name. This page number makes reference to the *Algae Identification Lab Guide* where you can find more information about the alga and its characteristics under the microscope. For example, “*Cladophora*, pg 28” indicates that your alga may be from the genus *Cladophora* and “pg 28” refers you to page 28 of the *Algae Identification Lab Guide* where you can find more information about this genus.

If you can not successfully match your alga to any of the algae descriptions, it is likely that your alga may not be listed in this guide.

3

Step Three.

To validate your algae identification, collect a sample that can be brought back to the lab and examined under a compound microscope.

Legend



Blue Green
(cyanobacteria)



Green Algae



Diatoms



Flagellates

Surface Bloom-forming Algae

Algae found in abundance near the surface of the water where light is present.

To ensure the best possible algal match or matches, read all questions under each category before selecting a genus or page number.

Is the algae:

1. Filamentous*?

- A. Does it have a septic odour and net-like appearance?
Hydrodictyon Pg 14
 - B. When squeezed, does it look like cotton?
Cladophora Pg 14
 - C. Does it feel silky or slippery?
Spirogyra Pg 15
 - D. Is it abundant near shore and leaves behind dried masses of algal paper?
Oedogonium Pg 15
 - E. Also see “Attached algae and algal mats” Pg 28
-

2. Planktonic*?

- A. Is it like pea-soup? And/or has a pigpen odour?
Anabaena, Microcystis, Aphanizomenon Pg 16-17
 - B. Is it spherical or globular?
Gloetrichia, Volvox, Nostoc Pg 18
 - C. Does it have a grassy odour?
Green algae, Ankistrodesmus Pg 19
 - D. Is the bloom throughout the water column? And deep green or green-black?
Oscillatoria, Pediastrum, Ankistrodesmus Pg 19-20
 - E. Does it look like grass clippings?
Aphanizomenon Pg 20
 - F. Does it look like bright green foam?
Euglena Pg 20
-

3. Jelly ball or aquatic plant-like?

A. Are they small, disc-like, bright green and have small roots on their undersides?

Duckweed Pg 21

B. Are they large, jelly-like balls?

Nostoc Pg 21

C. Is it plant-like, has whorled branches and smells of garlic or skunk?

Chara Pg 22

4. Too small to see individuals or colonies? And the water is cloudy or coloured?

A. Does it smell like geraniums or ripe cucumber? Musty or fishy?

Asterionella, *Synura*, *Dinobryon*, *Synedra* Pg 23

B. Also see Question #2, as the algae might be at different stage of development

C. No smell, but the water is coloured reddish brown? Pg 25

D. Common bloomers Pg 24

Bright green foam - *Euglena*

Fishy smell and green water - *Chlamydomonas*

"Clean water" and clogged filter - *Cyclotella*

E. Other possibilities (no distinguishable field cues)

Scenedesmus, *Closterium*, *Spirulina* Pg 25

Planktonic or Filamentous Finger Test

Wearing gloves, scoop a handful of whatever is blooming in the water with fingers spread slightly apart. Let the water drain and examine what remains. If long, stringy masses are left dangling from the fingers, it is a filamentous form. If mostly everything drains through the fingers and only a few bits stick to the glove, then it is a planktonic form.

FILAMENTOUS



PLANKTONIC




Filamentous Blooms


Stringy algal mats

Does it have a septic odour and net-like appearance?

► HYDRODICTYON pg 25 GR



In situ: D.M. John, York et al. (2002)



Microscale: M. Vis, Ohio University (2002)

The common name is “water net”. It can be found growing in hard water lakes and ponds. This genus is very susceptible to copper treatment.

When squeezed, does it look like cotton?

Colour also varies from green, yellow-brown to orange.

► CLADOPHORA pg 29 GR



In situ: W. Bourland, micro*scope (2006)



Microscale: W. Bourland, micro*scope (2006)

It is commonly found attached to rocks or substrates but can be detached by wave action. It is most often associated with streams, rivers and waterfalls. Septic odour can be present when abundant from the decomposition of the mat.

Filamentous Blooms


Stringy algal mats

Is it very silky or slippery?

► SPIROGYRA pg 30 GR



In situ: J. Jamieson, York et al. (2002)



Microscale: Australian Biological Resources Study, micro*scope (2006)


It is most likely a 'frog-spit' algae from the Order Zygnematales because of its silky/slippery qualities. A common genus from this Order is *Spirogyra*, also known as water silk or pond scum.

Is it abundant near the shore and leaves behind dried masses of 'algal paper'?

► OEDOGONIUM pg 28 GR



In situ: D. Patterson, micro*scope (2006)



Microscale: D. Patterson, micro*scope (2006)

When scooped up, it adheres to your hand rather than slipping through your fingers like *Spirogyra* (above). This genus is very susceptible to copper treatment.

Planktonic Blooms

Dispersed in water and unattached to solid objects

Is it like pea-soup? And/or has a pigpen odour? *Anabaena* and *Microcystis* require a microscope to view colonies, but are often distinguished by their pea-soup conditions. *Aphanizomenon* can often be identified macroscopically.

▶ ANABAENA pg 15 **BG**



In situ: S. Murrell



Microscale (M. Graham and S. Murrell)

This blue-green is capable of producing anatoxin, microcystin, or both. Blooms of some species of *Anabaena* can cause death in pets, livestock and wildlife. It is prevalent in phosphorus-rich water.

▶ APHANIZOMENON pg 18 **BG**



In situ: N. Serediak



In situ: E. Zoski



Microscale: W. Bourland, microscope (2006)

The most easily recognized blue-green. Colonies form a distinctive shape that resembles tiny grass or green fingernail clippings. They are roughly 0.5 to 1.0cm long.

Some species are capable of producing a neurotoxin called saxitoxin, similar to the paralytic shellfish poison that occur in marine environments. In North American freshwater environments, *Aphanizomenon* appears to be primarily non-toxic, although it can cause skin irritation to recreational water users.

Planktonic Blooms

Dispersed in water and unattached to solid objects

Is it like pea-soup? And/or has a pigpen odour? Continued....


► MICROCYSTIS pg 10 **BG**



In situ (R. Zurawell)



Microscale: D. Krogmann and M. Schneegurt, Cyanosite (2006)



In situ: P.V. York, York et al. (2002)

Blooms appear to be very similar to *Anabaena* blooms. *Microcystis* is found throughout the water column because they have gas vesicles that allow the colonies to regulate buoyancy.

Some species are capable of producing *microcystin*, a hepatotoxin (liver toxin) lethal to pets, livestock and humans. They may cause skin irritation for recreational water users.

Planktonic Blooms

Dispersed in water and unattached to solid objects

Is it spherical or globular?

And has a soft mucilage that is yellow-brown to light green, and slimy to touch?

► GLOEOTRICHIA pg 17 **BG**




Microscale : P York, York et al (2002).

Heavy blooms may cause serious skin irritations to recreational water users. When mature, colonies are visible to the naked eye and look like tiny fish eyes, tapioca beads or pom poms.

In situ photo not available.

► VOLVOX pg 20 **GR**



Microscale: D. Patterson and M. Farmer micro*scope (2006)

Blooms are infrequent but will occur over short periods during warm months, especially in waters contaminated with nitrogenous wastes or organically enriched water.

Spherical colonies at their largest are visible to the naked eye and can reach about 1mm in diameter. They are usually yellowish-brown to light green.

In situ photo not available.

► NOSTOC pg 16 **BG**



In situ (N. Serediak)

Nostoc is the most readily identified blue green in the field and sometimes called “freshwater grapes”. The colonies are in a jelly-like ball, which are usually surrounded by thick mucilage. Some species are capable of producing cyanotoxins.

It can be macroscopic in size and grow up to a few centimeters in diameter. Colour of the mucilage ranges from yellow-clear to blue-olive green.

Planktonic Blooms

Dispersed in water and unattached to solid objects

Does it have a grassy odour? (Like freshly cut grass)

▶ ANKISTRODESMUS pg 22 GR



Microscale: M Bahr and D Patterson, micro*scope (2006)

Grassy odour is the most common odour produced by green algae (not during decomposition). The smell is apparent only when present in large numbers. Common green algae are *Chlorella* (Lab Guide pg 23 and *Ankistrodesmus*).

Presence of *Ankistrodesmus* is a good indicator of clean water, since they tend to disappear from the algal community in very polluted systems.

In situ photo not available.

Is the bloom throughout the water column? And deep green or green-black?

▶ OSCILLATORIA pg 13 BG



In situ: D. Krogmann and M. Schneegurt, Cyanosite (2006)



B Anderson and D Patterson, micro*scope (2006)

It is a blue-green that can produce *microcystin* (hepatotoxin) and *anatoxin-a* (neurotoxin)

▶ PEDIASTRUM pg 26 GR



Microscale: D. Patterson and M. Farmer, micro*scope (2006)

Blooms are throughout the water column and thus do not result in surface scums. It is common in moderately nutrient-enriched water bodies. Also see *Ankistrodesmus* (above).

In situ photo not available.

Planktonic Blooms

Dispersed in water and unattached to solid objects

Does it look like grass clippings?

▶ APHANIZOMENON pg 18 **BG**



In situ: N Serediak



In situ, E Zoski



Microscale: W. Bourland, microscope (2006)

The most easily recognized blue-green. Colonies form a distinctive shape that resembles tiny grass or green fingernail clippings. They are roughly 0.5 to 1.0cm long.

Some species are capable of producing a neurotoxin called saxitoxin, similar to the paralytic shellfish poison that occur in marine environments. In North American freshwater environments, *Aphanizomenon* appears to be primarily non-toxic, although it can cause skin irritation to recreational water users.

Does it look like bright green foam or a thin film of scum?

▶ EUGLENA pg 36 **FL**



In situ (R. Zurawell)



In situ (S. Reedyk)



Microscale: D. Patterson and M. Farmer, microscope (2006)

Euglena is mostly found in freshwater habitats that are polluted by organic matter. *Euglena* blooms can also be red and exhibit a foam-like appearance in full bloom.

Other

Are they small, disc-like, bright green and have small roots on their undersides?

It is most likely not algae.

► **DUCKWEED** pg 39



In situ (N. Serediak)

Duckweed is often referred to as algae, but it is an aquatic plant. Individuals are typically 2-5cm long and by looking at the undersides, you can see short hair-like roots.

Are they large, jelly-like balls?

► **NOSTOC** pg 16 **BG**



In situ (N. Serediak)

Nostoc is the most readily identified blue green in the field and sometimes called “freshwater grapes”. The colonies are in a jelly-like ball, which are usually surrounded by thick mucilage. Some species are capable of producing cyanotoxins.

It can be macroscopic in size and grow up to a few centimeters in diameter. Colour of the mucilage ranges from yellow-clear to blue-olive green.

Also see:

Volvox and ***Gloeotrichia***
go to page 18

Other

Is it plant-like, has whorled branches and smells of garlic or skunk?

► **CHARA** pg 32 **GR**



In situ: N.F. Stewart, York et al. (2002)

Chara is also known as “stonewort”. It has a plant-like appearance and can grow several decimeters tall. It is easily recognized by their musky or garlicky odour.


Chara is abundant in hard, basic water ($\text{pH} > 7$), where calcium is abundant.

Too small for the naked eye

Water appears cloudy or coloured. Also see Planktonic Blooms on pages 16-20.

Does it smell like Geraniums? Musty? Ripe cucumber? Or fishy?

► **ASTERIONELLA** pg 33 **DM**



Microscale: M.Vis, Ohio University (2002)

Abundant in hard water lakes and known to clog filters. Geranium odour is present in low to moderate numbers; but changes to fishy in high numbers.

In situ photo not available.

► **SYNURA** pg 37 **FL**



Microscale: W. Bourland, micro*scope (2006)

Common in hard water lakes. Exhibits a ripe cucumber/musk melon smell in low to moderate numbers; but fishy in high abundance.

In situ photo not available.

► **DINOBRYON** pg 38 **FL**



Microscale: W. Bourland, micro*scope (2006)

Commonly bloom in summer in basic eutrophic bodies and exhibits a fishy odour. Also known to clog filters.

In situ photo not available.

► **SYNEDRA** pg 39 **DM**



Microscale: M.Vis, Ohio University (2002)

In moderate to high abundance, Synedra has an earthy to musty odour. Well-known to clog filters because of its rigid walls.

In situ photo not available.

Too small for the naked eye

Water appears cloudy or coloured. Also see Planktonic Blooms on pages 16-20.

Easily identified algae using a microscope. Common bloomers.

► **EUGLENA** pg 36 **FL**



In situ (R. Zurawell)



In situ (S. Reedyk)



Microscale:
D. Patterson
and M. Farmer,
micro*scope (2006)

Euglena is mostly found in freshwater habitats that are polluted by organic matter. *Euglena* blooms can also be red and exhibit a foam-like appearance in full bloom.

► **CHLAMYDOMONAS** pg 19




Microscale: D. Patterson and A. Laderman, micro*scope (2006)

Chlamydomonas is found in a variety of habitats, even snow in the high alpine! It exhibits a fishy smell in high numbers, and often colours the water green.

In situ photo not available.

► **CYCLOTELLA** pg 35



Microscale: M. Bahr and D. Patterson, micro*scope (2006)

Cyclotella is a common bloomer of the diatoms. It is a well known filter clogger and often found in 'clean' water..

In situ photo not available.

Too small for the naked eye

Water appears cloudy or coloured. Also see Planktonic Blooms on pages 16-20.

**Easily identified algae using a microscope.
Algae commonly found in the plankton community.**

► SCENEDESMUS pg 24 **GR**



Blooms can colour the water a lovely shade of green. It is common in freshwater aquaria and is a typical member of the open water plankton community.

In situ photo not available.

Microscale: L.A. Zettler and D. Patterson, micro*scope (2006)

► CLOSTERIUM pg 31 **GR**



Closterium is common in acid waters, while only a few species are found in nutrient rich hard waters. It is generally found mixed with other species of green algae.

In situ photo not available.

Microscale: D. Patterson and A. Laderman, micro*scope (2006)

► SPIRULINA pg 14 **BG**



Microscale: D. Patterson, L.A. Zettler and V. Edgecomb, micro*scope (2006)

Spirulina does not produce intense blooms or toxins. It is commonly marketed by health food producers as a general curative. In situ photo not available.

No smell? But the water is reddish-brown?



In situ: M. Huynh

It could likely be that the water is high in dissolved organic carbon- dissolved compounds from organic material. Leached tannins and lignins from decomposing plant matter can create tea-coloured water.

Another possibility is that the water may be high in iron, or polluted with other chemical compounds.

Algae interfering with coagulation

► EUGLENA pg 36 **FL**



In situ (R. Zurawell)



In situ (S. Reedyk)



Microscale:
D. Patterson
and M. Farmer,
microscope (2006)

Euglena is mostly found in freshwater habitats that are polluted by organic matter. *Euglena* blooms can also be red and exhibit a foam-like appearance in full bloom.

► ANABAENA pg 15 **BG**



In situ: S. Murrell




Microscale (M. Graham and S. Murrell)

This blue-green is capable of producing anatoxin, microcystin, or both. Blooms of some species of *Anabaena* can cause death in pets, livestock and wildlife. It is prevalent in phosphorus-rich water.

Algae interfering with coagulation

▶ ASTERIONELLA pg 33 **DM**



Microscale: M.Vis, Ohio University (2002)

Abundant in hard water lakes and known to clog filters. Geranium odour is present in low to moderate numbers; but changes to fishy in high numbers.

In situ photo not available.

▶ SYNEDRA pg 34 **DM**



Microscale: M.Vis, Ohio University (2002)

In moderate to high abundance, *Synedra* has an earthy to musty odour. Well-known to clog filters because of its rigid walls.

In situ photo not available.

FYI:
Relative
toxicity
to *copper
sulphate**

Susceptible

Anabaena, Microcystis, Aphanizomenon, Oscillatoria, Closterium, Hydrodictyon, Spirogyra, Cladophora, Asterionella, Synedra, Dinobryon, Synura, Volvox, Euglena

Resistant

Nostoc, Chlorella, Chara, Stigeoclonium, Ankistrodesmus, Scenedesmus, Chlamydomonas

*See Recommended Reading Material "Algal treatment and removal" on page 37 at the back of this guide.

Attached algae

and algal mats commonly found near shore

Mucilaginous and slimy film or tube on the substrate?

▶ CHROOCOCCUS pg 11 **BG**




Microscale: B.A. Whitton, York et al. (2002)

Chroococcus is found attached to rocks or soil in or near water and has a colourless mucilaginous sheath.

In situ photo not available.

▶ GLOEOCAPSA pg 12 **BG**



Microscale: M. Vis, Ohio University (2002)

A subaerial genus that forms red, brown or orange jelly globs on substrates in or near water.

In situ photo not available.

▶ TETRASPORA pg 21 **GR**



In situ- collapsed tube: P.V. York, York et al. (2002)

Colonies are in a soft, fragile, mucilaginous common tube or sac, which can be found attached to aquatic plants.

In situ photo not available.

Attached algae

and algal mats commonly found near shore

Floating mats or filamentous?

► STIGEOCLONIUM pg 27 **GR**



In situ: P.V. York, York et al. (2002)



Microscale: P.V. York, York et al. (2002)

It is a distinctly bright green alga often attached to rocks in fast-flowing water. It is also a filamentous mat that will not pass through open fingers of a cupped hand.

Also see:

Cladophora

go to page 14

Chara

go to page 22

Oedogonium

go to page 15

Spirogyra

go to page 15



Common blue-greens not listed in this manual that also form dense mats include: *Phormidium*, *Lyngbya*, *Tolypothrix* and *Stigonema*.

Potentially threatening freshwater algae

Listed below are common blue-greens known to cause low to serious risk to human health.


Only certain species of the genera below are toxic. Microscopic work or lab testing is required.

Is it like pea soup? And/or has a pigpen odour? A pigpen odour is usually associated with *Anabaena*, *Microcystis* and *Aphanizomenon*.

► **ANABAENA** pg 15 **BG**



In situ: S. Murrell



Microscale (M. Graham and S. Murrell)

This blue-green is capable of producing *anatoxin*, *microcystin*, or both. Blooms of some species of *Anabaena* can cause death in pets, livestock and wildlife. It is prevalent in phosphorus-rich water.

► **APHANIZOMENON** pg 18 **BG**



In situ: N. Serediak



In situ: E. Zoski.



Microscale: W. Bourland, micro*scope (2006)

The most easily recognized blue-green. Colonies form a distinctive shape that resembles tiny grass or green fingernail clippings. They are roughly 0.5 to 1.0cm long.

Some species are capable of producing a neurotoxin called saxitoxin, similar to the paralytic shellfish poison that occur in marine environments. In North American freshwater environments, *Aphanizomenon* appears to be primarily non-toxic, although it can cause skin irritation to recreational water users.

Potentially threatening freshwater algae


► MICROCYSTIS pg 10



In situ (R. Zurawell)



In situ: P.V. York, York et al. (2002)



Microscale: D. Krogmann and M. Schneegurt, Cyanosite (2006)

Blooms appear to be very similar to *Anabaena* blooms. *Microcystis* is found throughout the water column because they have gas vesicles that allow the colonies to regulate buoyancy.

Some species are capable of producing *microcystin*, a hepatotoxin (liver toxin) lethal to pets, livestock and humans. They may cause skin irritation for recreational water users.

Glossary of Terms

Algae - a group of organisms with mixed evolutionary history that share common characteristics. The most notable being that they have no vascular tissue (with some exceptions), have naked reproductive structures, are photoautotrophic, and contain chlorophyll (with some exceptions).

Chloroplast - pigmented structure (organelle) within the cell that conducts photosynthesis.

Coagulation - a chemical process that can reduce turbidity, dissolved organic compounds, and colour (AAFRD 2002).

Epiphytic/epiphyte - an organism attached to substrate or another organism.

Filamentous algae - long, multicellular threadlike filaments or masses of algae.

Hepatotoxin - a toxin that affects liver function, such as microcystin.

Lentic - standing water such as wetlands, dugouts, reservoirs and lakes. Antonym: lotic.

Neurotoxin - a toxin that affects brain function, such as anatoxin and saxitoxin.

Photoautotrophic - the ability to make food materials through photosynthesis by using sunlight, water and CO₂. Synonym: photosynthetic.

Planktonic algae - single-celled organisms or aggregations of single-celled algae that live near the water surface.

Surface bloom-forming algae - algae found in abundance near the surface of the water where light is present.

Photo credits

Planctonic/Filamentous (page 7): N. Serediak.

Microcystis: (in situ) R. Zurawell; (microscopic photo) D. Krogmann and M. Schneegurt. 2006. Cyanosite. <<http://www-cyanosite.bio.purdue.edu>>; (Algae in container) P.V. York in York et al. (2002).

Anabaena: (in situ) S. Murrell; (microscopic photo) M. Graham and S. Murrell.

Aphanizomenon: (microscopic photo) W. Bourland. 2006. Micro*scope. <<http://starcentral.mbl.edu/microscope/portal.php>>; (In situ) N.Serediak.

Oscillatoria: (In situ) D. Krogmann and M. Schneegurt. 2006. Cyanosite. <<http://www-cyanosite.bio.purdue.edu>>; (microscopic photo) B. Anderson and D. Patterson. 2006. Micro*scope. <<http://starcentral.mbl.edu/microscope/portal.php>>.

Gloeotrichia: P. York in York et al. (2002).

Nostoc: N. Serediak.

Gloeocapsa: M. Vis. 2002. Ohio University. Algae homepage. <<http://vis-pc.plantbio.ohiou.edu/algaeindex.htm>>.

Chroococcus: B. Whitton in York et al. (2002).

Tetraspora: P.V. York in York et al. (2002).

Ankistrodesmus: M. Bahr and D. Patterson. 2006. Micro*scope. <<http://starcentral.mbl.edu/microscope/portal.php>>.

Volvox: D. Patterson and M. Farmer. 2006. Micro*scope. <<http://starcentral.mbl.edu/microscope/portal.php>>.

Closterium: D. Patterson and A. Laderman. 2006. Micro*scope. <<http://starcentral.mbl.edu/microscope/portal.php>>.

Spirogyra: (in situ) J. Jamieson in York et al. (2002); (microscopic photo) Australian Biological Resources Study. 2006. Micro*scope. <<http://starcentral.mbl.edu/microscope/portal.php>>.

Oedogonium: (in situ and microscopic photo) D. Patterson. 2006. Micro*scope. <<http://starcentral.mbl.edu/microscope/portal.php>>.

Hydrodictyon: (in situ) D.M. John in York et al. (2002); (microscopic photo) M. Vis. 2002. Ohio University. Algae homepage. <<http://vis-pc.plantbio.ohiou.edu/algaeindex.htm>>.

Stigeoclonium: P.V. York in York et al. (2002).

Cladophora: (in situ and microscopic photo) W. Bourland. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Chara: N.F. Stewart in York et al. (2002).

Chlamydomonas: D. Patterson and A. Laderman. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Synedra: M. Vis. 2002. Ohio University. Algae homepage.
<<http://vis-pc.plantbio.ohiou.edu/algaeindex.htm>>.

Cyclotella: M. Bahr and D. Patterson. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Pediastrum: D. Patterson and M. Farmer. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Scenedesmus: L.A. Zettler and D. Patterson. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Dinobryon: W. Bourland. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Synura: W. Bourland. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Euglena: (in situ) R. Zurawell; (microscopic photo) D. Patterson and M. Farmer. 2006.
Micro*scope.<<http://starcentral.mbl.edu/microscope/portal.php>>.

Duckweed: N. Serediak.

Asterionella: M. Vis. 2002. Ohio University. Algae Homepage.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Spirulina: D. Patterson, L.A. Zettler and V. Edgecomb. 2006. Micro*scope.
<<http://starcentral.mbl.edu/microscope/portal.php>>.

Colored water (page 19): M. Huynh

References

- Bold, H. and M. Wynne. 1978. Introduction to the algae structure and reproduction, 2ed. Prentice-Hall, INC., Englewood Cliffs, N.J. 662pp.
- Fritsch, H.W., Graham, O.P., Iyengar, D.A., Johansen, H.H., Johnson, B.H., Kethcum, G.F., Papenfuss, G.W., Prescott, E.G., Pringshsheim, H.H Strain, and L.H., Tiffany. 1951. Manual of Phycology: An introduction to the algae and their biology. Chronica Botanica Co. (Pub). Waltham, Mass., USA.
- Hickman, M. 2000. A brief introduction to algae- Botany 333, Laboratory Manual. University of Alberta, Edm. Canada.
- John, D.M., B.A. Whitton and A.J. Brook. 2002. The Freshwater Algal Flora of the British Isles: An Identification Guide to Freshwater and Terrestrial Algae. The Natural History Museum. Cambridge.
- Palmer, Mervin. 1962. Algae in water supplies: An illustrated manual on the identification, significance, and control of algae in water supplies. U.S. Department of Health, Education and Welfare. Public Health Service. Division of water supply and pollution control. #Pub.-216489.
- Palmer, Mervin. 1977. Algae and water pollution. EPA-600/9-77-036. Environmental Protection Agency, United States of America.
- Prescott, G.W. 1964. How to know the fresh-water algae- an illustrated key for identifying the more common freshwater algae to genus, with hundreds of species named and pictured and with numerous aids for the study. Dubuque, Iowa, W.C. Brown Co. 293pp.
- P.V. York, D.M. John, and L.R. Johnson. 2002. A Photo Catalogue of Images of Algae and Algal Habitats. A CD-ROM accompanying 'The Freshwater Algal Flora of the British Isles' (eds. D.M. John, B.A. Whitton and A.J. Brook). Cambridge University Press.
- Smith, G.M. (ed.), L.R., Blinks, H.C., Bold, K.M., Drew, F., Drouet, J., Feldmann, F.E.,

Recommended Reading

Algal treatment and removal, and management practices

Alberta Agriculture, Food and Rural Development, Agriculture Education and Training Branch. 2002. Quality Farm Dugouts. Prairie Water News. Alberta, Canada.

Palmer, Mervin. 1962. Algae in water supplies: An illustrated manual on the identification, significance, and control of algae in water supplies. U.S. Department of Health, Education and Welfare. Public Health Service. Division of water supply and pollution control. #Pub.-216489.

Field sampling methods

John, D.M., B.A. Whitton and A.J. Brook. 2002. The Freshwater Algal Flora of the British Isles: An Identification Guide to Freshwater and Terrestrial Algae. The Natural History Museum. Cambridge.

Preserving and preparing mounts

Prescott, G.W. 1964. How to know the fresh-water algae- an illustrated key for the identifying the more common freshwater algae to genus, with hundreds of species named and pictured and with numerous aids for the study. Dubuque, Iowa, W.C. Brown Co. 293pp.

Index

Blue Greens

Anabaena	pg 16, 26, 30
Aphanizomenon	pg 16, 20, 30
Chroococcus	pg 28
Gloeocapsa	pg 28
Gloeotrichia	pg 18
Oscillatoria	pg 19
Microcystis/Anacystis	pg 17, 31
Nostoc	pg 18, 21
Spirulina	pg 25

Diatoms

Asterionella	pg 23, 27
Cyclotella	pg 24
Synedra	pg 23, 27

Flagellates

Dinobryon	pg 23
Euglena	pg 20, 24, 26
Synura	pg 23

Green Algae

Ankistrodesmus	pg 19
Chara	pg 31
Chlamydomonas	pg 24
Chlorella	pg 28
Cladophora	pg 14
Closterium	pg 25
Hydrodictyon	pg 14
Oeodogonium	pg 15
Pediastrum	pg 19
Scenedesmus	pg 25
Spirogyra	pg 15
Stigeoclonium	pg 29
Tetraspora	pg 28
Volvox	pg 18

Other

Duckweed	pg 21
----------	-------

