

Figure 1. East side of Pangnirtung Fiord, first side valley north of the Hamlet of Pangnirtung. A prominent ridge of the Quaternary lateral moraine system trends across the middle stage. 2011-056





Figure 2. A line of 5 tons developed in coarse grained granitic rock of the Cumberland Batholith east of Pangnirtung (65°11'N, 65°54'W). The tors are separated by weathering along near vertical joints. Note large weathering pits (gneiss) on top. 2011-052

Abstract
 In 2009, as part of the CGM Program of the Geological Survey of Canada, Cumberland Peninsula east and southeast of the National Park was mapped. Mapping included several aspects of the regional Precambrian geology as well as the Quaternary geology. Regional geology was mapped for sedimentological and geomorphological purposes, and moraines and other deposits were mapped for geomorphological purposes. The eastern part of the peninsula is today an area of intensive glacial and evidence has experienced this style of glaciation throughout the Quaternary. During the last glacial maximum (c. 120ka) the alpine glaciers thickened to form regional ice divides over the mountains. Ice flow from these divides covered most, possibly all, of the region and supplied ice sheets along the coast. This local ice divide was the Laurentide Ice Sheet in Cumberland Sound. Despite possibly repeated glaciation, substantial areas show little or no sign of glacial erosion and retain tertiary surface marked with block fields and tors.


Résumé
 En 2009, dans le cadre du programme GEM, la Commission géologique du Canada a cartographié les régions de la péninsule de Cumberland à l'est et au sud-est du parc national. La cartographie inclut plusieurs aspects de la géologie régionale, y compris la géologie sédimentologique et géomorphologique, et des dépôts de moraines et d'autres dépôts cartographiés à des fins de cartographie géomorphologique. La partie est de la péninsule est aujourd'hui une zone de glaciation intensive et il existe des preuves de ce style de glaciation tout au long du Quaternaire. Pendant le dernier maximum glaciaire (c. 120 ka), les glaciers alpins se sont épaissis pour former des lignes de partage glaciaire couvrant une grande partie sinon toute la région et formant des cônes glaciaires en bordure de plusieurs fjords. Cette glace locale entrait en collision avec l'iceberg laurentien dans la baie Cumberland. Malgré plusieurs glaciations, des régions importantes ne montrent que très peu ou même aucun signe d'érosion glaciaire et retiennent les surfaces du tertiaire recouvertes de champs de blocs et de tors.



Cover illustration: Lateral Glacier, Cumberland Peninsula. Photograph by Art Dyke. 2002-235
 Printed map: Catalogue No. M183-118-2011E ISBN 978-0-600-20085-9
 Digital map: Catalogue No. M183-118-2011E-PCF ISBN 978-1-100-10628-0
 © Her Majesty the Queen in Right of Canada 2011

Natural Resources Canada / Ressources naturelles Canada

CANADIAN GEOSCIENCE MAP 18
 (preliminary version)
SURFICIAL GEOLOGY
CLEARWATER FIORD-PANGNIRTUNG SOUTH
 Baffin Island, Nunavut
 1:100 000



SURFICIAL DEPOSITS

QUATERNARY

HOLOCENE

- ngF** Proglacial outwash: gravel and sand, 1–10 m thick, forming fans.
- ngFp** Proglacial outwash: gravel and sand 1–10 m thick, forming active terraced fans.
- ntm** Till: nonsorted glacial debris commonly very bouldery with a silty sand matrix.
- mf** Lateral moraine: 5–10 m high moraine ridges with over-stepped, fan-like ridges on shallowly bedded glacial ice cores and associated ground moraine with minor glacioluvial sediments, distinguished from older moraines by lack of moraine lobes and past cover.
- AF** Alluvial fans: gravel and sand commonly bouldery, with detrital organic layers and buried soils, 1–20 m thick, formed by steep-gradient streams and debris flows forming channels and levees.
- Ap** Alluvial plains: gravel and sand, 1–10 m thick.
- At** Alluvial terraces: gravel and sand, 1–10 m thick, above limit of modern flooding.
- Cz** Landslide deposits: rock avalanche debris of coarse blocks, 10 or more metres thick, derived from cliff failure.
- Cf** Colluvial fan deposits: blocky to gravely debris-flow accumulations mixed with silt, 1–20 m thick, traversed by narrow channels and debris levees, typically intersect steep slopes (Ca) at the ends of prominent debris hops on cliff surface slopes less than angle of repose but steeper than those of alluvial fans.
- Ca** Talus: generally active accumulations of blocks and rubble, as much as 50 m thick forming talus (snow) aprons at angle of repose below cliffs derived from rock falls.
- Cr** Rock glacier debris: talus, generally 10–50 m thick, deformed by interstitial flow of buried ice to form talus glaciers; irregular terraces on talus slopes with transverse ridges or trails above steep frontal faces, some faces stable and well-vegetated; most rivers unstable, unvegetated, and at angle of repose.
- MR** Beach sediments: gravel and sand, commonly bouldery, 1–5 m thick, forming raised beach ridges and swales and the modern, transgressive beach, deposited in beach, deltaic, and offshore environments during regression of postglacial sea.
- Ms** Deltaic sediments: sand and gravel, typically overlying fine sand and silt bottomset beds, 5–20 m thick, forming raised terraces, terraces at marine limit formed on or near the ice margin.
- Mv** Nearshore to offshore veneer: sand, silt, and minor clay, with dropstones, 1–2 m thick.
- Mb** Nearshore to offshore blanket: sand, silt, and minor clay, with dropstones, 2–20 m thick.

EARLY HOLOCENE AND WISCONSINAN

- GLr** Beach or deltaic sediments: sandy gravel, 1–10 m thick, typically formed at the maximum lake extent.
- GLs** Nearshore proglacial sediment: sand, silt, and minor clay, 1–2 m thick.

GLACIOLUVIAL SEDIMENTS: gravel and sand, 1–10 m thick, deposited behind, at, and in front of the ice margin

- GFf** Proglacial outwash: gravel and sand, 1–10 m thick, forming fan-shaped deposits.
- GFp** Proglacial outwash: gravel and sand, 1–10 m thick, forming inactive terraced fans.
- GFt** Proglacial outwash: gravel and sand, 1–10 m thick, forming terraced deposits.
- GFh** Ice-contact sediments: gravel and sand, 1–10 m thick, forming hummocky deposits.

TILL, nonsorted bouldery diamictics, 1–40 m thick, deposited in subglacial and ice marginal environments, lithic composition generally reflecting underlying bedrock

- Tm** Lateral moraine: 5–40 m high ridges and hummocks comprised mainly of till probably overlying debris on glacial ice cores, forming lateral and end moraine ridges and less organized, hummocky accumulations formed during ice-marginal recession; moraine crests muted due to colluviation during partial deglaciation of ice cores; matrix somewhat more sandy and less silty than till forming ground moraine; locally contains ice-contact stratified drift and outwash.
- Tr** Rock-glacierized moraine: ice-cored end or lateral moraine, 5–40 m high, displaced from original site of glacial deposition by downslope flow of ice-debris mixture. It moved in places with silt; commonly act as local base level for silt accumulation; hence difficult to distinguish from C2 in places; many stable rivers.
- Tv** Till veneer: variably bouldery (10–60% cover; typically 20–40% diameter) with silty sand matrix, 0.2–2 m thick and discontinuous; insufficiently thick to obscure relief of underlying bedrock.
- Tb** Till blanket: variably bouldery (10–60% cover; typically 20–40% diameter) with silty sand matrix, 2–10 m thick, sufficiently thick to obscure relief of underlying bedrock.

PRE-WISCONSINAN

- Wv** Residual: talus, rock rubble, and gravel, 1–2 m thick, mantling bedrock, formed by disintegration of Precambrian bedrock prior to Last Glacial Maximum but including sparse erratics and possibly morphologically degraded old till; matrix leucocratic comprised of more-scale blocks in silted matrix with typically 60–80% block cover; finer rubble and gravels on coarse microtopography of Cumberland Batholith, interspersed by less 1–4 m high bedrock surface shows little or no sign of glacial scouring; veneer retains its pre-Quaternary form, mainly on flat or gently graded, cliff-bordered, upland plateaus, but on moderately steep slopes in places, most surfaces probably covered by cold-based ice during one or more glaciations.

BEDROCK

PRE-QUATERNARY

ROCK: rock of various compositions and Precambrian ages; Cumberland Batholith comprised largely of monzogranite dominant west of Kingiall Fiord, tonalite and metasediments further east.

R1 CMB: major escarpments, typically hundreds of metres high, forming serrated faces with multiple debris hoppers, glacially scoured surfaces removed by postglacial scarp retreat producing basal (or subbasal) talus accumulations.

R2 Scoured rock: hills and hummocky surfaces with lake basins and ice-moulded embayments resulting from glacial scouring and with patchy veneers of till; commonly depleted of matrix material; probably covered by warm-based ice during distal intervals of Wisconsinian Glaciation, including Last Glacial Maximum.

Legend:

- Glacier ice margin for 2009 from SPOT5 satellite imagery
- Geological boundary (defined)
- Geological boundary (gradational)
- Fluted bedrock (direction known)
- Esker
- Lateral moraine
- Lateral meltwater channel; barb on upslope side
- Glacial lake spillway
- Marine limit
- Clippe
- Cliff
- Glacial lake limit
- Dyke
- Field observation (point colour relative to geological units)
- Rock blower
- Strain flow (direction known)
- Station from literature (Kjelson et al., 2001)
- Station, marine elevation elevation in metres
- Station, marine elevation elevation in metres from literature (Kjelson, 2001)

Recommended citation:
 Dyke, A.S., 2011. Surficial geology, Clearwater Fiord-Pangnirtung south, Baffin Island, Nunavut. Geological Survey of Canada, Canadian Geoscience Map 18 (preliminary version), scale 1:100 000. doi:10.4095/28963

ess.nrcan.gc.ca

Canadian Geoscience Maps
 Cartes géoscientifiques du Canada

Canada

CANADIAN GEOSCIENCE MAP 18
 (preliminary version)
SURFICIAL GEOLOGY
CLEARWATER FIORD-PANGNIRTUNG SOUTH
 Baffin Island, Nunavut
 1:100 000

Author: A.S. Dyke
 Geology by A.S. Dyke, 2009
 Geological compilation by A.S. Dyke, 2009
 Cartography by L. Robertson

Initiative of the Geological Survey of Canada, conducted under the auspices of the Multiple Metals Cumberland Peninsula (Nunavut) project, as part of Natural Resources Canada's Geomapping for Energy and Minerals (GEM) program.

Logistical support provided by the Polar Continental Shelf Project as part of its mandate to promote scientific research in the Canadian North. FCSP 002-09

Map projection: Universal Transverse Mercator, zone 20, North American Datum 1980
 Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications. Elevations in feet above mean sea level.
 Mean magnetic declination 2011, 3°39'W, decreasing 29.7' annually. Reading vary from 3°31'W in the SW corner to 3°47'W in the NE corner of the map.

The Geological Survey of Canada welcomes corrections or additional information from users. This map conforms to the ISO 9001:2008 standard used by the Scientific and Technical Publishing Services Quality Management System.
 This publication, including digital data, can be downloaded free of charge from GeoPlus (http://geoplus.nrcan.gc.ca/). It is also available from the Geological Survey of Canada Bookstore (http://geoscan.nrcan.gc.ca/bookstore).

Recommended citation:
 Dyke, A.S., 2011. Surficial geology, Clearwater Fiord-Pangnirtung south, Baffin Island, Nunavut. Geological Survey of Canada, Canadian Geoscience Map 18 (preliminary version), scale 1:100 000. doi:10.4095/28963

CANADIAN GEOSCIENCE MAP 18
 (preliminary version)
SURFICIAL GEOLOGY
CLEARWATER FIORD-PANGNIRTUNG SOUTH
 Baffin Island, Nunavut