



## Risk Evaluation Determining Whether Environmental Emergency Planning is Required Under the **Environmental Emergency Regulations set under the Canadian Environmental Protection Act, 1999 (CEPA 1999)**

This report is a consolidation of two substances listed under Schedule 1 of the *Environmental Emergency Regulations* set under CEPA 1999. The substances inherent hazardous properties are virtually the same.

**Nickel Nitrate (CAS #: 13138-45-9)**

*(Nickel(II) Nitrate)*

**AND Nickel(II) Nitrate, Hexahydrate (CAS #: 13478-00-7)**

*(Oxidic, Sulphidic, and Soluble Inorganic Nickel Compounds)*

**(CEPA, 1999 – Schedule 1, #42)**

### Summary of Risk Evaluation Analysis

The Risk Evaluation Framework (REF), developed by Environment Canada, is applied to chemical substances in order to determine whether or not an environmental emergency (E2) plan should be required. For those substances that are evaluated as requiring an E2 plan, the next step is to determine a threshold quantity for the total quantity of the substance and the maximum storage container size on site. Within the REF, three categories are used to evaluate chemicals and assign threshold quantities. Shown below are the details within the three categories for *nickel nitrate*:

#### 1. Environmental Hazard

##### **(Persistence, Bioaccumulation and Aquatic Toxicity Analysis)**

- According to Mackay *et al.* (2003), nickel, chromium, arsenic, and cadmium are classified as indefinitely persistent. As a result, *nickel nitrate* is also classified as indefinitely persistent in the environment.
- No data are available to determine the bioaccumulation potential.

- For aquatic toxicity, it was determined that *nickel nitrate* is slightly toxic (LC<sub>50</sub> at 96 hrs between 10 and 100 mg/L) based on the most sensitive species, Atlantic salmon (Grande *et al.*, 1983).

## 2. Human Hazard

### (Inhalation Toxicity and Carcinogenicity Analysis)

- Since *nickel nitrate* has a vapour pressure below 1.33 kPa at 20°C (ATSDR, 2003; HSDB, 2004), there is insufficient vapour from the substance to cause an inhalation problem.
- Nickel nitrate* is categorized as an International Agency for Research on Cancer (IARC) carcinogen with a classification rating of '1' (ATSDR, 2003; HSBD, 2004; IARC, 1990; Genium, 2004).

## 3. Physical Hazard

### (Flammability and Combustibility Analysis)

- Nickel nitrate* is a substance normally stable, even under fire exposure conditions, and is not reactive to water (Genium, 2004; CSST, 2000).

### Flammability and Combustibility Analysis Table

	Value	Vapour Cloud Explosion (Yes/No)	Combustible (Yes/No)	References
Boiling point (°C)	136.7	No	No	(Lewis, 2000; ATSDR, 2003)
Flash point (°C)	--			(HSDB, 2004; ATSDR, 2003)

## Threshold

The data from the above mentioned categories were used to determine the various environmental/human thresholds. The threshold quantities relate to: carcinogenicity, aquatic toxicity, inhalation toxicity, vapour cloud explosion, combustibility, and reactivity.

The threshold quantity for carcinogenicity is defaulted to 0.22 tonnes, if it meets the following criteria:

- 1) have an IARC rating of 1, 2A or 2B, or have a US EPA rating of A, B1, or B2;
- 2) be persistent in any media for greater than 5 years.

Threshold quantities assigned as a result of aquatic toxicity are based on the following table:

### Threshold Quantities Assigned as a Result of Aquatic Toxicity

Criteria	Extremely Toxic	Highly Toxic	Moderately Toxic	Slightly Toxic
Persistence (water)	≥ 6 months	≥ 2 months to < 6 months	N/A	N/A
Bioaccumulation	BCF ≥ 5000 or Log Kow ≥ 5	BCF ≥ 500 to < 5000 or Log Kow ≥ 4 to < 5 (unless BCF < 500)	N/A	N/A
Acute Aquatic Toxicity (96 hrs LC <sub>50</sub> – mg/L)	≤ 0.1	> 0.1 to ≤ 1	> 1 to ≤ 10	> 10 to ≤ 100
Threshold Quantity Tonnes (lbs)	0.22 (500 lbs)	1.13 (2 500 lbs)	4.50 (10 000 lbs)	9.10 (20 000 lbs)

A substance is a candidate for a vapour cloud explosion if its flash point is < 23°C and its boiling point is < 35°C. A substance is considered combustible if its flash point is < 23°C or its boiling point is < 35°C. In accordance with the precautionary principle, the category with the lowest threshold will be used. For further explanation, please refer to the *Implementation Guidelines for Part 8 of the Canadian Environmental Protection Act, 1999 – Environmental Emergency Plans* (<http://www.ec.gc.ca/ee-ue/>).

### Conclusion

For *nickel nitrate* # 42, the E2 hazard threshold was triggered by:

- 1) Aquatic toxicity at 0.22 tonnes;
- 2) Carcinogenicity at 0.22 tonnes.

Therefore, *nickel nitrate* # 42 is recommended for addition to Schedule 1 of the *Environmental Emergency Regulations* under Part 3 with a threshold quantity of 0.22 tonnes.

## References

ATSDR (Agency of Toxic Substances for Disease Registry). *Toxicological Profile for Nickel CAS # 7440-02-0*. PB/98/101199/AS. 2003. World Wide Web accessed October 2004.

<http://www.atsdr.cdc.gov/toxprofiles/tp.asp?id=245&tid=44>

CSST (Commission de la santé et de la sécurité au travail). *Service du répertoire toxicologique: Dinitrate de nickel anhydre*. Gouvernement du Québec, 2000. World Wide Web accessed October 2004.

<http://www.reptox.csst.qc.ca/RechercheProduits.asp>

Genium. "Material Safety Data Sheet – Nickel Nitrate." Release 58. CD-ROM. Genium Publishing Corp. Schenectady, NY. 2004.

Grande, M., and S. Andersen. "Lethal Effects of Hexavalent Chromium, Lead and Nickel on Young Stages of Atlantic Salmon (*Salmo salar* L.) in Soft Water". 1983. *Vatten*. 39(4): 405-416.

HSDB (Hazardous Substances Data Bank). *Nickel Nitrate*. Canadian Centre for Occupational Health and Safety. U.S. National Library of Medicine. 2004. World Wide Web accessed October 2004.

<http://toxnet.nlm.nih.gov/cgi-bin/sis/htmlgen?HSDB>

IARC (International Agency for Research on Cancer). *Nickel and Nickel Compounds*. World Health Organization. 1990. World Wide Web accessed September 2004.

<http://monographs.iarc.fr/ENG/Monographs/vol49/mono49-7.pdf>

Lewis Sr., R.J. *Sax's Dangerous Properties of Industrial Materials*. 10<sup>th</sup> Edition. Wiley-Interscience Publication. 2000. Volume 3. p. 2632.

Mackay, D., E. Webster, D. Woodfine, T.M. Cahill, P. Doyle, Y. Couillard, and D. Gutzman. "Contributed Articles Towards Consistent Evaluation of the Persistence of Organic, Inorganic and Metallic Substances." 2003. ASP. Canadian Environmental Modelling Centre and Environment Canada. *Human and Ecological Risk Assessment*. 9: 1445-1474. ISSN: 1080-7039.

Cat. No.: En14-55/19-2011E-PDF

ISBN: 978-1-100-19708-1

For information regarding reproduction rights, please contact Public Works and Government Services Canada at 613-996-6886 or at [droitdauteur.copyright@tpsgc-pwgsc.gc.ca](mailto:droitdauteur.copyright@tpsgc-pwgsc.gc.ca)

© Her Majesty the Queen in Right of Canada, represented by the Minister of the Environment, 2011

Aussi disponible en français