


13736 12

Catalogue of Salmon Streams
and Spawning Escapements
of Statistical Area 28
Howe Sound–Burrard Inlet



M.J. Hancock and D.E. Marshall

Department of Fisheries and Oceans
Salmonid Enhancement Program
1090 West Pender Street
Vancouver, British Columbia V6E 2P1

January, 1986

Canadian Data Report of
Fisheries and Aquatic Sciences
No. 557



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Data Report of Fisheries and Aquatic Sciences

Data reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of data reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was introduced with the publication of report number 161.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport statistique canadien des sciences halieutiques et aquatiques

Les rapports statistiques servent à classer et à archiver les compilations de données pour lesquelles il y a peu ou point d'analyse. Ces compilations auront d'ordinaire été préparées à l'appui d'autres publications ou rapports. Les sujets des rapports statistiques reflètent la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports statistiques ne sont pas destinés à une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans autorisation écrite préalable de l'établissement auteur. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 25 de cette série ont été publiés à titre de relevés statistiques, Services des pêches et de la mer. Les numéros 26 à 160 ont été publiés à titre de rapports statistiques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 161.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Data Report of
Fisheries and Aquatic Sciences

No. 557

January 1986



CATALOGUE OF SALMON STREAMS AND SPAWNING ESCAPEMENTS

STATISTICAL AREA 28

HOWE SOUND - BURRARD INLET

by

M.J.Hancock and D.E.Marshall

Department of Fisheries and Oceans
Salmonid Enhancement Program
1090 West Pender Street
Vancouver, B.C. V6E 2P1

Minister of Supply and Services Canada 1986
Cat. No. Fs 97-13/557 ISSN 0706-6465

Correct citation for this publication:

Hancock, M.J. and D.E.Marshall. 1986 Catalogue of Salmon Streams
and Spawning Escapements of Statistical Area 28 Howe Sound - Burrard Inlet
Can. Data Rep. Fish and Aquat. Sci. 557: xiv + 190 p.

ABSTRACT

Hancock, M.J. and D.E.Marshall, 1986, Catalogue of Salmon Streams and Spawning Escapements of Statistical Area 28, Howe Sound - Burrard Inlet. Can. Data Rep. Fish and Aquat. Sci. 557: xiv + 190 p.

Catalogue containing each stream's location, spawning distribution, barriers and points of difficult ascent, escapement records and other general data pertaining to the stream. The catalogue also includes a topographical map of the stream and in some cases a sketch map which further describes the surrounding area.

Keywords: British Columbia, Statistical Area 28, Howe Sound - Burrard Inlet, salmon streams, spawning escapements.

RÉSUMÉ

Hancock, M.J. and D.E.Marshall, 1986. Catalogue of Salmon Streams and Spawning Escapements of Statistical Area 28, Howe Sound - Burrard Inlet. Can. Data Rep. Fish and Aquat. Sci. 557: xiv + 190 p.

Le présent répertoire donne l'emplacement de chaque cours d'eau, la répartition de fraie, les points de remonte difficile, les données sur les saumons de remonte et d'autres information générales concernant le cours d'eau. On y trouve aussi une carte topographique de l'emplacement du cours d'eau et, dans quelque cas, un croquis décrivant la zone environne.

Mots-cles: Colombie-Britannique, zone statistique 28, Howe Sound - Burrard Inlet, cours d'eau a saumons, remonte.

CONTENTS

Abstract/Resume.....	iii
Table of Contents	iv
Statistical Maps	v,vi
Management Divisions	vii
Districts and Subdistricts	viii
Standards Used	ix,x
Federal Fisheries District Addresses	xi
Map of Salmon Spawning Streams of Statistical Area 28.....	xii
Index to Map of Salmon Spawning Streams of Statistical Area 28	xiii
Escapement Summary for Statistical Area 28, Howe Sound - Burrard Inlet	xiv

STREAM DATA

ASHLU CREEK.....	1,2	MCCARTNEY CREEK	101
(Avalon Creek).....	7	MCNAB CREEK	104
(Branch 100 Creek	9	MCNAIR CREEK (Hastings Creek).....	107
BROHM RIVER.....	12	(Meighan Creek)	110
BROTHERS RIVER	15	MOSQUITO CREEK	113
CAPILANO RIVER	18,19	NELSON CREEK	117
CHAPMAN CREEK (Mission Creek).....	23	OUILLET CREEK (Jap Creek, Oulette Cr.)	121
CHASTER CREEK (Gower Creek).....	26	PILLCHUCK CREEK (Pilchuck Creek).....	125
CHEAKAMUS RIVER.....	29	POTLATCH CREEK	128
(Chuck-Chuck Creek).....	34	RAINY RIVER	131
CYPRESS CREEK	37	(Richards Creek).....	135
DAKOTA CREEK	40	ROBERTS CREEK	138
DRYDEN CREEK	43	SEYMOUR RIVER	141,142
EAGLE CREEK (Eagle Harbour Creek)...	46	(Shop #3 Creek).....	145
FLUME CREEK.....	49	SHOVELNOSE CREEK	148
HASTINGS CREEK.....	52	(Spring Creek)	151
HIGH FALLS CREEK	55	SQUAMISH RIVER	154,155
HOP RANCH CREEK (Hop Ranch	58	STAWAMUS RIVER (Little Stawamus River)	159
Creek System (4))		(Tenderfoot Creek).....	164
INDIAN RIVER (Burrard River).....	61,62	(Thirty Seven Mile Creek).....	167
JUDD SLOUGH.....	66	(Twenty Eight Mile Creek).....	170
LANGDALE CREEK.....	70	(Twin Creek, Archies Creek).....	173
(Loggers Lane Creek).....	73	(Upper Paradise Channel).....	176
(Long Bay Creek).....	76	(Lower Paradise Channel).....	-
LYNN CREEK.....	79	WAKEFIELD CREEK	179
MACKAY CREEK (McKay Creek)	83	(West Bay Creek).....	182
MAMQUAM RIVER	86,87	(Williamson Creek).....	185
(Mamquam Spawning Channell)	92	WILSON CREEK.....	188
MANNION CREEK (Cotton Creek).....	95		
MASHITER CREEK	98		


- STATISTICAL AREAS ARE DIVIDED BY RED LINES.
- SALMON FISHING WITH NETS OF ANY KIND IS NOT PERMITTED OUTSIDE OF — THAT IS SEAWARD OF — THE HEAVY BLACK LINE.

NOTE: AREAS 5, 6, 7, 8 AND 9 REVISED JAN 1974.

DEPARTMENT OF THE ENVIRONMENT FISHERIES OPERATIONS

STATISTICAL MAP

SHOWING AREAS OF CATCH FOR
BRITISH COLUMBIA WATERS
(NORTHERN HALF)



READ CAREFULLY

1. PIN UP IN WHEELHOUSE
2. WHEN DELIVERING YOUR CATCH, GIVE TALLY MAN THE MAP NUMBER, OR NUMBERS SHOWING THE AREA IN WHICH YOUR FISH WERE CAUGHT
3. ACCURATE CATCH REPORTS WILL HELP PRESERVE YOUR FISHERIES.

READ CAREFULLY

1. PIN UP IN WHEELHOUSE
2. WHEN DELIVERING YOUR CATCH, GIVE TALLY MAN THE MAP NUMBER, OR NUMBERS SHOWING THE AREA IN WHICH YOUR FISH WERE CAUGHT.
3. ACCURATE CATCH REPORTS WILL HELP PRESERVE YOUR FISHERIES
4. FOR COMPLETE DETAILS, CONSULT BRITISH COLUMBIA FISHERIES REGULATIONS.

— STATISTICAL AREAS ARE DIVIDED BY RED LINES


— SALMON FISHING WITH NETS OF ANY KIND IS NOT PERMITTED OUTSIDE OF — THAT IS SEAWARD OF — THE HEAVY BLACK LINE.

● FISHERIES SERVICES OFFICES

DEPARTMENT OF THE ENVIRONMENT FISHERIES SERVICE


STATISTICAL MAP SHOWING AREAS OF CATCH FOR BRITISH COLUMBIA WATERS (SOUTHERN HALF)

0 10 20 30 40 50
SCALE IN MILES



FOR ALL FISH CAUGHT
OFF THE COAST OF THE
STATE OF WASHINGTON
SOUTH OF CAPE FLATTERY
REPORT AS AREA "C"

MANAGEMENT DIVISIONS
FISHERIES & OCEANS
PACIFIC REGION





BRITISH COLUMBIA

FISHERIES & OCEANS - Pacific Region

WHITEHORSE DISTRICT

YUKON-ARCTIC

ALSEK-TAKU

YUKON SOUTH

NORTHERN B.C.



LEGEND

DISTRICT BOUNDARY ———
SUBDISTRICT BOUNDARY ———

VANCOUVER

COQUITLAM

MISSION

STEVESTON

SURREY

CHILLIWACK

QUATSINO SOUND

KYUQUOT

TAHSIS

TOFINO

PORT ALBERNI

SOOKE

VICTORIA

SAANICH

COMOX

PENDER HARBOUR

NANAIMO

DUNCAN

CHILLIWACK

COQUITLAM

SURREY

STEVESTON

MISSION

CHILLIWACK

COQUITLAM

SURREY

STEVESTON

CHILLIWACK

COQUITLAM

SURREY

STEVESTON

CHILLIWACK

STANDARDS USED ON STREAM DATA PAGE

Name of Stream: Name given in Gazetteer of Canada, British Columbia, 1985, third edition.

Local names are in lower case and enclosed in brackets.

Statistical Area: As defined by D.F.O., showing areas of catch for B.C. Waters (Map dated Jan. 1974)

Districts and Subdistricts: As defined by D.F.O. (Map 1985)

RAB Numbers: The Aquatics Unit of the Resource Analysis Branch, Ministry of the Environment have assigned a hierarchical coding system (RAB number) to drainage basins of British Columbia. RAB numbers classify catchment areas and river channels. Further information on RAB coding system can be found in "A Hierarchical Watershed Coding System for British Columbia", RAB Technical Paper #3, Ministry of the Environment, Victoria, B.C. June 1980.

Geographic Co-ordinate Reference: The geographic co-ordinates given indicate the centre of each feature, except for streams, where the position of the mouth is given. The geographic co-ordinates are to the nearest minute.

Length: The portion of the stream accessible to spawning salmon.

Drainage: Area in square kilometers of the entire drainage basin feeding the stream.

Discharge: Extremes of maximum and minimum daily discharge for the period of the last 30 years. Discharge date is taken from "Historical Stream Flow Summary", British Columbia, Water Survey of Canada.

Temperature: As described ($^{\circ}\text{C}$)

Barriers and Points of difficult ascent: Complete and partial barriers to salmon and their distance from the stream mouth. Species likely to be affected may be listed. Both natural and man-made obstructions are defined.

Spawning distribution:

Portion of the stream utilized by each species. Distribution is indicated by brief comments opposite the species.

General remarks: Emphasizes features of stream and spawning populations. Also includes industrial activity, routes of accessibility, etc. The comments and dates are taken from "Annual Reports of Salmon Streams and Spawning Grounds". In some cases, references to additional information not included in the General remarks may be given.

Escapement Records: The escapement represents the mid point of the coded range of escapement for each species. For example: the letter "H" representing 5000-10000 fish would be entered as 7500. Where absolute numbers are provided by Fisheries Personnel, these numbers are entered. N/O means the stream was inspected, but no fish were observed; UNK means there was evidence of fish present, but no estimates were made; NO RECORDS means no escapement records for the applicable years could be found in the escapement files.

Timing: Dates which salmon arrive in the stream, begin to spawn, reach peak spawning period and finish spawning.

E = early (1st to 10th of the month)

M = mid (11th to 20th of the month)

L = late (21st to end of the month)


NB: Distance references are from the mouth of the stream unless otherwise stated.

FISHERIES & OCEANS - Pacific Region

DISTRICT/SUB-DISTRICT OFFICES

<u>DISTRICT/SUB-DISTRICT</u>	<u>ADDRESS</u>	<u>TELEPHONE</u>	<u>SUB-DISTRICT NUMBER</u>
<u>DISTRICT #1 - Kamloops</u>	202 - 317 Seymour St., Kamloops, V2C 2E9	374-4322	
Salmon Arm	Box 1160, 461 Beatty Ave. NW, Salmon Arm V0E 2T0	832-8037	29K
Prince George	2392 Ospika Blvd., Prince George, V2N 3N5	564-7030	29J
Clearwater	Box 610, Clearwater, V0E 1N0	674-2633	29J
Lillooet	Box 315, Lillooet, V0K 1V0	256-4525	29F
Quesnel	Box 4340, Quesnel, V2J 3J3	992-2434	29H
Williams Lake	540 Borland St., Williams Lake, V2G 1R9	398-6544	29G
<u>DISTRICT #2 - New Westminster</u>	309 - 549 Columbia St., New West., V3L 1B3	524-7181	
Vancouver	309 - 549 Columbia St., New West., V3L 1B3	524-7306	28A
Surrey	309 - 549 Columbia St., New West., V3L 1B3	524-7171	29B
Coquitlam	309 - 549 Columbia St., New West., V3L 1B3	524-7169	29C
Steveston	1255 No. 1 Road, Richmond, V7E 1T7	274-7217	29A
Squamish	Box 85, Squamish, V0N 3G0	892-3230	28B
Mission	Box 3308, Mission, V2V 4J5	826-3664	29D
Chilliwack	Suite 5, 9375 Mary St., Chilliwack, V2P 4G9	792-6011	29E
<u>DISTRICT #3 - Nanaimo</u>	60 Front St., Nanaimo, V9R 5H7	754-3257	
Nanaimo/Ladysmith	60 Front St., Nanaimo, V9R 5H7	754-3257	17
Qualicum Beach	Box 1270, Qualicum Beach, V0R 2T0	752-9712	14S
Comox	Box 1328, Comox, V9N 3Z0	339-2031	14N
Duncan	Box 241, 191 Ingram St., Duncan, V0L 3X3	746-6221	18
Powell River	4488 Marine Avenue, Powell River, V8A 2K2	485-9621	15
Pender Harbour	Box 10, Madeira Park, V0N 2H0	883-2313	16
<u>DISTRICT #4 - Port Alberni</u>	Box 280, Federal Building, Port Alberni, V9Y 7M7	724-0195	
Port Alberni	Box 280, Federal Building, Port Alberni, V9Y 7M7	724-0195	23
Quatsino Sound	Box 10, Port Hardy, V0N 2P0	949-6422	27
Kyuquot	Box 549, Tahsis, V0P 1X0	934-6606	26
Tahsis	Box 549, Tahsis, V0P 1X0	934-6606	25
Tofino	Box 48, Tofino, V0R 2Z0	725-3468	24
<u>DISTRICT #5 - Campbell River</u>	215 - 950 Alder St., Campbell River, V0W 2P8	287-2102	
Campbell River	215 - 950 Alder St., Campbell River, V0W 2P8	287-2102	13
Seymour Inlet	Box 10, Port Hardy, V0N 2P0	949-6422	11
Alert Bay	Box 10, Alert Bay, V0N 1A0	974-5216	12
<u>DISTRICT #6 - Victoria</u>	116 - 816 Government St., Victoria, V8W 1W9	566-3252	
Victoria/Saanich	116 - 816 Government St., Victoria, V8W 1W9	566-3252	19
Sooke	Box 460, Sooke, V0Z 1N0	642-5322	20
<u>DISTRICT #7 - Kitimat</u>	315 - 450 Federal Building, Kitimat, V8C 1T6	632-4884	
Butedale	315 - 450 Federal Building, Kitimat, V8C 1T6	632-4884	6
Bella Bella	Box 38, Bella Bella, V0T 1B0	957-2363	7
Bella Coola	Box 130, Bella Coola, V0T 1C0	799-5345	8
Rivers Inlet	Dawson Landing P.O., Rivers Inlet, V0N 1M0		9
Smith Inlet	Dawson Landing P.O., Rivers Inlet, V0N 1M0		10
<u>DISTRICT #8 - Prince Rupert</u>	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	
Waterfront	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	
Skeena	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	4A
Grenville - Principe	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	5
Lower Nass	109 - 417 2nd Ave. West, Prince Rupert, V8J 1G8	624-9137	3A
Upper Nass	Box 29, Nass Camp, V0J 3J0	633-2408	3B
Hazelton	Box 327, Field Street, Hazelton, V0J 1Y0	842-6327	4C
Smithers	Box 578, Smithers, V0J 2N0	847-2312	4D
Terrace	4721-B Lazelle Ave., Terrace, V8G 1R6	635-2206	4B
<u>DISTRICT #9 - Queen Charlotte Is.</u>	Box 99, Queen Charlotte City, V0T 1S0	559-4413	
West Coast Q.C.I.	Box 99, Queen Charlotte City, V0T 1S0	559-4413	2W
Masset	Box 99, Masset, V0T 1M0	626-3316	1
Sandspit	Box 222, Sandspit, V0T 1T0	637-5340	2E
<u>DISTRICT #10 - Whitehorse</u>	122 Industrial Road, Whitehorse, Y.T., Y1A 2T9	667-2235	
Yukon South/Northern B.C.	122 Industrial Road, Whitehorse, Y.T., Y1A 2T9	667-2235	120
Yukon-Arctic	122 Industrial Road, Whitehorse, Y.T., Y1A 2T9	667-2235	110
Alsek-Taku	Box 5341, Haines Junction, Y.T., Y0B 1L0	634-2235	130

Salmon Spawning Streams of
Howe Sound - Burrard Inlet



INDEX TO MAP OF SPAWNING STREAMS OF STATISTICAL AREA 29

HOWE SOUND - BURRARD INLET

1. ASHLU CREEK
2. (Avalon Creek)
3. (Branch 100 Creek)
4. BROHM RIVER
5. BROTHERS RIVER
6. CAPILANO RIVER
7. CHAPMAN CREEK (Mission Creek)
8. CHASTER CREEK (Gower Creek)
9. CHEAKAMUS RIVER
10. (Chuck-Chuck Creek)
11. CYPRESS CREEK
12. DAKOTA CREEK
13. DRYDEN CREEK
14. EAGLE CREEK (Eagle Harbour Creek)
15. FLUME CREEK
16. HASTINGS CREEK
17. HIGH FALLS CREEK
18. HOP RANCH CREEK (Hop Ranch
Creek System (4))
19. INDIAN RIVER (Burrard River)
20. JUDD SLOUGH
21. LANGDALE CREEK
22. (Loggers Lane Creek)
23. (Long Bay Creek)
24. LYNN CREEK
25. MACKAY CREEK (McKay Creek)
26. MAMQUAM RIVER
27. (Mamquam Spawning Channel)
28. MANNION CREEK (Cotton Creek)
29. MASHITER CREEK
30. MCCARTNEY CREEK
31. MCNAB CREEK
32. MCNAIR CREEK (Hastings Creek)
33. (Meighan Creek)
34. MOSQUITO CREEK
35. NELSON CREEK
36. OUILLET CREEK (Jap Creek, Oulette Cr.)
37. PILLCHUCK CREEK (Pilchuck Creek)
38. POTLATCH CREEK
39. RAINY RIVER
40. (Richards Creek)
41. ROBERTS CREEK
42. SEYMOUR RIVER
43. (Shop # 3 Creek)
44. SHOVELNOSE CREEK
45. (Spring Creek)
46. SQUAMISH RIVER
47. STAWAMUS RIVER (Little Stawamus River)
48. (Tenderfoot Creek)
49. (Thirty Seven Mile Creek)
50. (Twenty Eight Mile Creek)
51. (Twin Creek, Archies Creek)
52. (Upper Paradise Channel)
53. (Lower Paradise Channel)
54. WAKEFIELD CREEK
55. (West Bay Creek)
56. (Williamson Creek)
57. WILSON CREEK

SUMMARY
ESCAPEMENT RECORD FOR STATISTICAL AREA 28 HOWE SOUND - BURRARD INLET

YEAR	SOCKE YE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
47	25	15625	11025	62275	519575	17875
48		15500	93000	102550	25	17500
49	25	15500	17300	52500	299850	37400
50		15625	13125	52725	25	18550
51		15975	97575	98775	304625	18225
52		15500	103450	240875	150	41750
53		18900	13250	23250	201725	10175
54		16700	45700	67150	300	18900
55	4	15975	31998	30510	192550	18045
56		15550	8540	14500		9165
57		19100	51300	78875	237175	17445
58		18900	25820	87950		17275
59		17100	6175	158925	157125	8200
60		15550	17514	30450		10101
61		19100	50464	23700	452500	17661
62		18900	13686	54125		16622
63		9100	14721	44375	832625	8947
64		7575	57097	39225		17836
65		35750	18750	13350	110800	16350
66		18900	15750	26750	75	9450
67		6900	20425	46675	47000	12325
68		10600	18025	125650		14300
69	25	24800	14550	70000	32100	10275
70	25	31000	37825	132025		8675
71	25	11279	38300	45990	59325	5081
72		9488	11915	364745	757	23341
73	12	14015	30850	277247	173300	19981
74		9343	146375	155200		22181
75	75	4817	64997	71000	79000	17710
76	2	6008	35503	134644	300	10112
77	300	4170	14963	138675	25255	5750
78	297	1563	15754	123063	512	2980
79	12	5303	56804	37856	27226	5231
80	25	5794	43321	193522	400	7434
81	8	6126	43434	149072	55480	5746
82	30	4717	48057	158158		3320
83	419	4516	47007	113577	26574	1865
84	62	5191	45602	187118		1198
85						

TIMING

ARRIVE						
START						
PEAK						
END						

REMARK



NAME OF STREAM ASHLU CREEK RAB NO. 90-1300-140
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 54' 123° 17'
 LOCATION OF MOUTH Flows S.E. into Squamish River., North of mouth of Cheakamus River,
New Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 3.2km

SPAWNING DISTRIBUTION

Species


Section of Stream Used

chinook	evenly distributed throughout
coho	" " "
steelhead	" " "
chum	" " "
1984 reported all species observed downstream from bridge.	

GENERAL REMARKS

- 1973 The watershed of this stream has been almost completely denuded by logging over the last 30 years, consequently, the stream is subject to rapid rise and fall in water levels. In 1973 flooding caused severe bank erosion on the south shore and extensive silting in the lower spawning area of the river. Approx 25% of the streambed was affected by silting. The stream also shifted 91m to the south. Extensive scouring also affected approx. 25 - 35% of the pink and chinook spawning area. Steelhead, coho and chinook are subject to heavy sport fishing. This stream is also prone to heavy debris buildups, especially along the bars.
- 1978 This creek supports a heavy sports fishery for both chinook and coho.
- 1980 Severe flood late Dec. Estimate 75 - 90% loss of spawn.
- 1982 Very high w/l in Dec. some scouring and debris.
- 1983 Stream is very fast flowing and subject to many freshets, resulting in a substrate of large gravel. Only isolated pockets of gravel suitable for spawning. Chum run has declined dramatically over the past 9 years.
- Light predation by bears and birds. 1974 reported grizzly bear present.
- Seasonal fluctuations in water levels.

Sketch of lower Ashlu Creek, 1968



ESCAPEMENT RECORD FOR ASHLU CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947		25	200	75	1500	200
48		25	400	750	N/O	200
49		25	25	400	1500	75
50		25	200	400		75
51		25	200	75	1500	200
52		25	1500	3500		400
53		200	400	400	1500	75
54		200	75	400		200
55		25	750	75	750	75
56		75	200	200		75
57		200	75	750	750	200
58		200	750	400		75
59		400	75	400	3500	75
60		75	200	75		200
61		200	UNK	750	2500	UNK
62		200	750	75		75
63		400	200	400	7000	50
64		100	700	400		100
65		1500	700	400	7000	75
66		200	200	100		50
67		200	200	400	3500	100
68		200	2000	700		100
69		2000	300	700	2000	50
70		1500	1500	400	N/O	150
71		300	700	400	5000	400
72		700	400	1500	N/O	400
73		600	600	7500	5500	150
74		750	3500	8000		1200
75		200	3500	1500	1500	400
76		400	750	3500		400
77		200	200	75	75	75
78	5	250	200	25		100
79		200	200	25	25	200
80		120	400	200		120
81		400	100	100	NO REC	150
82		200	75	25		100
83	25	150	100	N/O	50	50
84		250	400	400		
85						

TIMING

ARRIVE		E JUL-E AUG	AUG - L OCT	E OCT - NOV	AUG	M FEB
START		JUL - L AUG	AUG - M NOV	L OCT - NOV	AUG - SEPT	
PEAK		JUL - SEPT	SEPT - M DEC	NOV - E DEC	SEPT	
END		L SEPT-E OCT	OCT - E JAN	L NOV-L DEC	SEPT - M OCT	JUNE

REMARK

Avalon Creek see Dakota Creek P 40

NAME OF STREAM (Avalon Creek) RAB NO. -LOCAL NAME DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 31' 123° 30'LOCATION OF MOUTH Flows into Thornbrough Channel, S. of Dakota Cr.LENGTH 1.6 km WIDTH m DRAINAGE km²DISCHARGE (m³/s) MAX MIN Temperature (°C) COMPOSITION: Bedrock Boulder Coarse Fine Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

gravel bar at mouth -- passable at high flows

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho

- unknown

chum

- evenly distributed up to 200 yds

GENERAL REMARKS

1971 This stream flows into Thornbrough Channel near the site of a large log dumping and sorting area. It is short and the water disappears underground about 1.6km from mouth. It has several short stretches where very good spawning conditions exist. Could support up to 1000+ chum.

1976 This minor stream has been subjected to severe abuse by the Rivtow development at the mouth.

1980 Low water levels in early Oct. resulted in stream going underground near mouth. High flows in December caused scouring and some erosion.

1981/82 If an enhancement box were to be put on the stream, I would recommend that a cat be brought in to move the gravel bar so that fish can get up creek at all flows.


ESCAPEMENT RECORD FOR (Avalon Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71				75		
72				200		
73				200		
74				75		
75				25		
76				50		
77				-		
78				-		
79				N.O.		
80				N.O.		
81				N.O.		
82				N.O.		
83				N.I.		
84				N.O.		
85						

TIMING

ARRIVE				E - M OCT		
START				M - L OCT		
PEAK				L OCT-M NOV		
END				E NOV-M DEC		

REMARK



NAME OF STREAM (Branch 100) RAB NO. -
 LOCAL NAME
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 56' 123° 18'
 LOCATION OF MOUTH Flows W. into Squamish River near Skaweskin Indian Reserve 7

LENGTH km WIDTH m DRAINAGE km²
 DISCHARGE (m³/s) MAX MIN
 Temperature (°C)
 COMPOSITION: Bedrock Boulder Coarse Fine
 Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

Impassable rock falls at 250 - 300 yds from mouth

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- in upper and lower reaches
------	------------------------------

GENERAL REMARKS

- 1979 Previously included with Squamish River salmon counts
 1980 Flood conditions late December -- estimated losses of 50% spawners.
 Erosion and silting during flood.
 1983 Will always be a low production stream given its general characteristics.

Water levels normal throughout most of year -- frequent high water
 early October to late December


ESCAPEMENT RECORD FOR (Branch 100 Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25			
80		25				
81			25			
82			25			
83			10			
84			30			
85						

TIMING

ARRIVE		OCT	OCT			
START		NOV	NOV - DEC			
PEAK		L NOV	L NOV-E JAN			
END		M DEC	M DEC-M JAN			

REMARK



NAME OF STREAM BROHM RIVER RAB NO. 90-1300-050-010-010

LOCAL NAME _____

DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 48' 123° 08'LOCATION OF MOUTH Flows S. into Cheekye River, E. of Cheakamus River, New Westminster
Dist. _____LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²DISCHARGE (m³/s) MAX _____ MIN _____

Temperature (°C) _____

COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Rock falls at North end of Brohm Lake at 4.8km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- scattered throughout
steelhead	- scattered throughout

GENERAL REMARKS

- 1979 This streams population formerly included in Cheakamus River counts.
 1981 Some scouring, erosion and silting during October high water.
 1983 Some minor alterations in stream course by Highway Dept. near lake.
 Stock movements upstream may be hindered by harsh stream conditions
 between Cheekye and Brohm Creeks. Access to the lake is questionable
 due to conditions at Hwy. 99 culvert.
 1984 All fish observed within 200 yds of mouth.


ESCAPEMENT RECORD FOR BROHM RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			75			25
80			150			50
81			50			35
82			50			30
83			20			40
84			25			
85						

TIMING

ARRIVE		OCT - E NOV		MAR
START		NOV - M DEC		APR
PEAK		NOV - DEC		MAY
END		DEC - E JAN		JUN

REMARK Figures formerly included in Cheakamus reports.



NAME OF STREAM BROTHERS CREEK RAB NO. 90-0900-010
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 20' 123° 08'
 LOCATION OF MOUTH Flows S. into Capilano River, New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable at .80km-- stream narrow with large boulders

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- scattered
chum	- upper area from Keith Rd. to Hwy. 401

GENERAL REMARKS

1977 First report. Some problems encountered in protecting stream from fish molesting. Illegal gaffing.
 1978 Water levels extremely low during the summer months.
 1979 Erosion and silting 12 - 15%. Water levels extremely low during summer and fall. One half of normal yearly precipitation -- flood mid.Dec.
 1980 Water levels extremely low during Jun-Jul-Aug -- high last week Oct.
 1984 Low water during June to Oct. Predation by humans still a problem.


ESCAPEMENT RECORD FOR BROTHERS RIVER


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77			12	55		
78			66	48		
79		13	54	38		
80		20	85	65		
81		12	110	160	35	10
82		18	194	56		6
83		27	186	124	64	
84		17	183	97		11
85						

TIMING

ARRIVE		E OCT	E OCT	L OCT	-	M DEC
START		M OCT	SEPT - L OCT	SEPT - L OCT	-	MAR
PEAK		-	SEPT - M NOV	OCT - M NOV	-	-
END		L OCT	DEC - E JAN	NOV - M DEC	-	A?L

REMARK





NAME OF STREAM CAPILANO RIVER RAB NO. 90-0900
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 19' 123° 08'
 LOCATION OF MOUTH Flows S. into First Narrows, Burrard Inlet, New Westminster
Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 360 Jan 15, 1961 MIN 1.10 Oct. 3, 1950
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Cleveland Dam, Greater Vancouver Water Board at 4.8km

SPAWNING DISTRIBUTION

Species	Section of Stream Used
chinook	- to hatchery
coho	- to hatchery, some trucked above the dam
chum	- lower reaches and Brothers Creek confluence
pink	- lower reaches
steelhead	- to hatchery, some trucked above the dam

GENERAL REMARKS

This stream is connected with the water supply system of greater Vancouver. In 1954, the construction of Cleveland Dam and the adjoining reservoir was completed. This impassable dam was overcome by constructing a fishway and facilities for trucking salmon above the dam. This equipment is still in operation. The Capilano Hatchery was completed in 1972 and was constructed as part of a program to increase stocks of coho and chinook salmon and steelhead trout. Its yearly designed operating capacity is 1,000,000 coho smolts, 2 - 3,000,000 chinook smolts and 20-30,000 steelhead smolts.

1955/63 Fishway and facilities for trucking salmon above dam in operation.

1966 A steel mesh fence covering was put over the top of the holding pool at Dam site to stop the public from jigging for fish.

1968 Some rock work done above trapping facilities and below the dam. This should have the effect of permitting fry and yearlings a better chance to survive plunge over spillway of dam.

1969 Removal of gravel bars by G.V.W.B. for use as road gravel 3.2km above West Fork Bridge -- no damage to spawning areas.

1971 Hatchery in partial operation -- this stream subject to much abuse by general public and the salmon runs are often molested.

1973 Heavy poaching.

1976 120 chum lost to human predators.

1977 A good return of hatchery jacks.

continuation

CAPILANO RIVER

- 1979 Flooding in Dec. has probably decimated 50% of wild stocks.
- 1981 Squamish Indian Band By-law #10 is in effect on this system.
- 1982 50 chum lost to human molestation.

Predation: Merganser and bear in upper reaches. Very heavy poaching in lower river and fish are molested by dogs and juveniles.

Seasonal fluctuations in water levels.


ESCAPEMENT RECORD FOR CAPILANO RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			3500	3500	7500	750
48			7500	1500	N/O	750
49			3500	1500	3500	750
50			3500	1500	N/O	1500
51			3500	3500	750	750
52			7500	1500	25	1500
53			3500	750	1500	750
54			3500	3500	75	1500
55	4		4998	400	400	95
56			1840	25		65
57			5100	200	75	95
58			3745	400	N/O	75
59			NO RECORD			
60			3614	25		251
61			2114	25	25	86
62			2636	25		97
63			2071	75	100	97
64			2622	25		161
65			750	25	25	25
66			3500	25		75
67			1500	25		200
68			1500	200	N/O	25
69			1500	200	25	75
70			3500	75	N/O	75
71		44	4000	75	25	91
72		38	1200	700	7	91
73		165	1100	1100	150	56
74		93	40200	1500		31
75		767	6391	400	200	35
76	2	1102	25248	40	-	12
77		-	NO REC	120	30	150
78			500	250	-	35
79		3000	43000	280	200	100
80		2839	25434	200	-	250
81		1330	24100	400	450	200
82		463	27500	100	-	120
83	3	1133	20186	500	70	237
84		1694	16859	205	-	380
85						

TIMING

ARRIVE	E OCT	E JUN	E AUG-M OCT	L JUL	DEC
START	AUG - E OCT	JUN - JUL	SEPT - M OCT	JUL - E OCT	-
PEAK	OCT	AUG - M OCT	OCT	L AUG-M OCT	-
END	NOV	SEPT - DEC	E NOV - DEC	SEPT - E NOV	AUG

REMARK 1981 coho includes 2804 jacks, chinook includes 420 jacks.



NAME OF STREAM CHAPMAN CREEK RAB NO. 90-1600
 LOCAL NAME (Mission Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°26' 123°43'
 LOCATION OF MOUTH Flows S.W. into Str. of Georgia at Wilson Creek P.O., New
Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 193 Oct. 13, 1962 MIN 0.034 Jul. 22, 1962
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 4.8km

SPAWNING DISTRIBUTION

Species	Section of Stream Used
chum	concentrated - in lower 1.2km, then scattered throughout
coho	- scattered

GENERAL REMARKS

This stream is the main water source for the Sunshine Coast Regional Water Board. Consumption of this water began in 1970 and will probably increase rapidly in the near future (1969-70)

1970 The Local Water District constructed a small dam near the falls.

1972 Heavy rains in July washed out most of the steelhead spawning site. Up to 50% of the chum spawn was lost during December because of a severe freshet.

1979 This creek is subject to flash flooding from the extensive logging operations in the upper watershed. There was no fishery in District #1 on chum this year resulting in four times the return from the '75 brood year.

1980 Flash flooding is becoming a problem to private property near the mouth and extensive rip-rap will have to be installed. Extensive scouring due to high Dec. floods.

1982 Poor assessment due to high water levels

Predation: mainly light, some jigging problems and fish molestation -- juveniles trampling through reeds near hwy. bridge (1980)


ESCAPEMENT RECORD FOR CHAPMAN CREEK (Mission Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75	1500	750	75
48			25	750	25	25
49			200	1500	25	25
50			25	750	25	
51			75	1500	200	75
52			75	750	25	25
53						
54			NO	RECORDS FOR	1953 - 1964	
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65				50		
66				50		
67				100	50	
68				50		
69				200		
70				1500		
71				2200		40
72				3300		75
73				3500	100	200
74			75	3500		400
75			75	200	25	200
76			100	100		
77			UNK	2500		
78			75	400	12	
79			50	1200		
80			50	500		
81			25	600		
82			N.O.	20		
83			50	400		
84			50	200		
85						

TIMING

ARRIVE			E - M SEPT	E - M OCT	JUL	DEC
START			M - L SEPT	M OCT	JUL - L AUG	JAN
PEAK			E - L OCT	E - L NOV	AUG - SEPT	MAR
END			M NOV-E DEC	F - M DEC	SEPT	MAY

REMARK



NAME OF STREAM CHASTER CREEK RAB NO. 90-1500
 LOCAL NAME (Gower Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 23' 123° 33'
 LOCATION OF MOUTH Flows S.W. into Straits of Georgia, N.W. of Gower Pt.,
New Westminster Dist.
 LENGTH km WIDTH m DRAINAGE km²
 DISCHARGE (m³/s) MAX 0.351 May 16, 1965 MIN 0 Aug. 2, 1965
 Temperature (°C)
 COMPOSITION: Bedrock Boulder Coarse Fine
Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

Impassable falls at 4.8km and steep terrain with large boulders and rapids at 2.5km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- up to 1.6km
coho	- upstream to hwy.

GENERAL REMARKS

- 1972 This stream has good spawning areas scattered throughout the lower 2.4km and probably has spawning capacity for 1500 chum. Heavy rains in Dec. affected 70% of the stream bed -- extensive erosion, scouring and heavy siltation in the lower spawning areas. An estimated 60-80% of the spawn was lost.
- 1973 Water Licences on this stream reduce water supply in summer low flows to almost nothing.
- 1975 High water damaged about 30% of spawning area.
- 1979 There was virtually no commercial fishery on chum in District #1 this year which may have contributed greatly to good escapements along the Sechelt Peninsula. Some scoring during latter part of December due to extremely high water levels.
- 1980 A Gibson's school put a S.E.P. box in with chum eggs, but it filled with silt from a natural mud slide that occurred upstream. Severe erosion due to flooding in December.

Light predation by dogs and public.


ESCAPEMENT RECORD FOR CHASTER CREEK (Gower Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947				400		
48				400		
49				750		
50			NO	RECORDS		
51			NO	RECORDS		
52				200		
53						
54			NO	RECORDS FOR	1953 - 1969	
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70				200		
71				20		
72				75		
73			NO	RECORDS		
74				75		
75			25	75		
76			N.O.	N.O.		
77				150		
78			10	50		
79			N.O.	230		
80				75		
81			30	75		
82			N.O.	6		
83			30	200		
84			25	500		
85						

TIMING

ARRIVE			SEPT - E NOV	M OCT-E NOV		
START			OCT - E NOV	L OCT-E NOV		
PEAK			NOV	E - L NOV		
END			M DEC	L NOV-M DEC		

REMARK



NAME OF STREAM CHEAKAMUS RIVER RAB NO. 90-1300-050
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 47' 123° 10'
 LOCATION OF MOUTH Flows S. and S.W. into Squamish River, New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls 9m high at 14km

SPAWNING DISTRIBUTION

Species

Section of Stream Used


chinook	- above Culliton Creek
chum	- from mouth to Culliton
coho	- throughout

GENERAL REMARKS

Cheakamus Lake elevation is 830 meters. Peak run-off occurs between May and August. Since 1957, flows have been affected by storage and diversion from Daisy Lake via a penstock discharging through a powerhouse into the Squamish River. The river is continually being encroached upon by stream-side land development, i.e. dyking, channelizing, diversions etc. It is also subject to an extremely heavy sports fishery for steelhead, dolly varden, chinook and coho. Migrating fish are also being subjected to an Indian net fishery on the Squamish River before they enter the Cheakamus River.

- 1975 35% of stream bed eroded and silted during late Oct. floods -- several major channel changes. Estimated loss of 75 - 90% pink and chinook spawn.
- 1977 Unusually high numbers of sockeye seen. The sports fishery is very heavy and the catch of coho, chinook and steelhead represents a fair percentage of the runs.
- 1989 Extensive highway relocation planned for near future. Work will be in canyon above spawning area, but F.O. concerned about possible heavy silt.
- 1981 Some damage to chinook eggs during October flood -- gravel movement.
- 1982 This is the first year of use for the enhanced Upper Paradise Channel which received approx 6,000 chum. Possible damage to chinook salmon spawn during flood in late October.
- 1983 B.C. dam test in Aug. showed substantial bedload movements began when flows exceeded 8000 cfs. Tests may result in loss of chinook spawn.
- Dyking work took place just above B.C. Hydro Bridge and near canyon area.

Sketch of chum spawning area
on Cheakamus River, 1968



ESCAPEMENT RECORD FOR CHEAKAMUS RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947		400	400	3500	75000	400
48		400	1500	3500		400
49		400	750	3500	35000	400
50		400	1500	3500		400
51		750	3500	15000	15000	400
52		400	3500	15000		3500
53		3500	1500	3500	15000	400
54		750	1500	5000		750
55		750	1500	7500	15000	1500
56		400	1500	3500		400
57		3500	3500	35000	35000	750
58		3500	3500	35000		1500
59		1500	3500	75000	75000	400
60		400	1500	15000	UNK	750
61		3500	7500	9300	351700	1500
62		3500	3500	35000		400
63		1500	1500	30000	555000	750
64		400	15000	15000		1500
65		3500	7500	3500	35000	400
66		3500	1500	15000		750
67		1500	5000	20000	20000	1500
68		300	4000	30000		3000
69		2000	2000	20000	12000	1000
70		3000	6000	25000	N/O	1500
71		2200	4500	4500	2200	1200
72		400	1500	60000		2500
73		300	4000	50000	25000	3000
74		400	15000	35000		3500
75		200	15000	7500	3500	1500
76		400	1500	15000	-	1500
77	200	200	1500	75000	750	1500
78	100	150	1500	60000	500	1000
79		200	200	7500	400	750
80		200		7500	400	750
81		800	2500	40000	3000	1200
82	30	200	1500	20000		1200
83	25	100	1500	20000	75	250
84	N.O.	400	1500	50000	-	UNK
85						

TIMING

ARRIVE		JUN - JUL	AUG - L NOV	M SEPT-M OCT	E AUG	DEC
START		JUN - AUG	AUG - E DEC	OCT - NOV	M AUG	
PEAK		JUL - SEPT	SEPT - L DEC	M NOV - DEC	E - L SEPT	
END		AUG - M OCT	OCT - L JAN	L DEC-E JAN	OCT	JUN

REMARK

Chuck-Chuck Creek

see

Branch 100 Creek p.9

NAME OF STREAM (Chuk-Chuk Creek) RAB NO. 90-1300-160
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 56' 123° 19'
 LOCATION OF MOUTH Flows S. into Squamish River, N. of High Falls Creek

LENGTH 2.4 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chinook	- over lower 100 yds
coho	- throughout

GENERAL REMARKS

- 1979 Figures previously included in Squamish River populations.
 1980 Flood conditions in late December -- coho and chinook spawning areas probably alright, but the chum spawning area is badly silted.
 1981 This creek breaks up into several small streams at 2.4km.
 1983 No indication of chum spawning at all this year. Also no evidence of pinks. Heavy silt load in bottom portion of stream may be result of relative closeness to Weldwood T.F.L. mainline haul road.
 1984 Enumeration estimate for chinook is from S.E.P. personnel who surveyed the area for broodstock.

Physical condition: 1983 Bottom portion of stream has heavy silt load over gravel substrate which is common in this stream.


ESCAPEMENT RECORD FOR (Chuk-Chuk Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25		25	25
80		15	40	25		25
81		10	40	20		25
82		20	30	25		
83		5	50			
84						
85						

TIMING

ARRIVE		M AUG	SEPT	SEPT		APL
START		L AUG-E SEPT	E - M OCT	OCT		MAY
PEAK		M - L SEPT	OCT - NOV	NOV		MAY - JUN
END		E - M OCT	L - M DEC	DEC		JUN

REMARK Figures previously included in Squamish River Report



NAME OF STREAM CYPRESS CREEK RAB NO. 90-0970
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°20' 123°15'
 LOCATION OF MOUTH Flows S. into Burrard Inlet, N.E. of Point Atkinson, New West.Dist.

LENGTH .40 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- below Keith Road
------	--------------------

GENERAL REMARKS

First report 1981.

- 1981 10% erosion and silting -- low w/l in September, flood in Oct.
- 1983 Early November floods
- 1984 Water levels low June - October, 1981/82 floods and continuous maintainance on lower portion of spawning area affected this stream.


ESCAPEMENT RECORD FOR CYPRESS CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81			12			
82			N.O.			
83			2	N/O		
84			N.O.			
85						

TIMING

ARRIVE						
START						
PEAK						
END						

REMARK



NAME OF STREAM DAKOTA CREEK RAB NO. 90-1430-010
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 31' 123° 30'
 LOCATION OF MOUTH Flows E. into Thornbrough Channel, New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN 0 Aug. 17, 1951
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock rapids and very steep at 3.2km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- at mouth of Dakota Creek
chum	- " "

GENERAL REMARKS

- 1970 First report. This stream is very similar to McNair Creek -- small spawning area near tide water -- very steep and bouldery a short distance from mouth.
- 1971 This stream had salmon runs some years back according to long time residents. A short area suitable for spawning salmon exists near the mouth of the stream.
- 1974 First report of salmon seen by F.O.
- 1978 Water level up for time of year (Oct. and Nov.)
- 1979 Extreme siltation from Construction Aggregates gravel pit. The company has taken measures to prevent further siltation.


ESCAPEMENT RECORD FOR DAKOTA CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70				N.O.		
71				N.O.		
72				N.O.		
73			N.O.			N.O.
74			25	25		
75			6	75		
76				N.O.		
77						
78				N.O.		
79			N.O.	2		
80			N.O.	N.O.		
81			N.O.	N.O.		
82			NOT INSPECTED			
83			N.O.	6		
84			N.O.	40		
85						

TIMING

ARRIVE		SEPT	M OCT	
START		E - M OCT	M - L OCT	
PEAK		L OCT	M NOV	
END		L - M NOV	E - M DEC	

REMARK First report 1970



NAME OF STREAM DRYDEN CREEK RAB NO. -LOCAL NAME DISTRICT 2 STATISTICAL AREA 28 POSITION 49°45' 123°08'LOCATION OF MOUTH Flows S. into Hop Ranch Creek, New West. Dist.LENGTH 1.6 km WIDTH m DRAINAGE km²DISCHARGE (m³/s) MAX MIN Temperature (°C) COMPOSITION: Bedrock Boulder Coarse Fine Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	unknown
------	---------

GENERAL REMARKS

1984 Most of the substrate in this system is made up of fines and survival is thought to be low. Normal water levels.

ESCAPEMENT RECORD FOR (Dryden Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84			50			
85						

TIMING

ARRIVE			L OCT			
START			M NOV			
PEAK			E DEC			
END			M DEC			

REMARK



NAME OF STREAM EAGLE CREEK RAB NO. 90-0988
 LOCAL NAME (Eagle Harbour Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 21' 123° 16'
 LOCATION OF MOUTH Flows S.E. into Queen Charlotte Channel, Burrard Inlet, N. of
Point Atkinson, New Westminster Dist.
 LENGTH .80 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 1.2km culvert 3 meters from mouth

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- scattered in tidal flats at mouth of stream during years when the culvert is impassable
------	---

GENERAL REMARKS

- 1971 This small stream is located in the middle of a residential area in North Vancouver. Fish molestation is a problem as the stream flows through many back yards.
 1979 Recommend measures be taken to allow fish access past culvert -- approx 274 m of usable spawning grounds.
 1980 No work yet done on improving access.
 1984 W/L low June to October.


ESCAPEMENT RECORD FOR EAGLE CREEK (Eagle Harbour Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69		NO RECORDS PRIOR TO 1970				
70				25		
71				15		
72				25		
73				6		
74				25		
75				25		
76				N.O.		
77				N.O.		
78				-		
79				6		
80				N.O.		
81			6			
82				-		
83				N/O		
84			7	3		
85						

TIMING

ARRIVE			E NOV	NOV		
START			M NOV	NOV		
PEAK			E DEC	-		
END			-	NOV		

REMARK



NAME OF STREAM FLUME CREEK RAB NO. 90-1580
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 26' 123° 40'
 LOCATION OF MOUTH Flows S.W. into Straits of Georgia, W. of Roberts Cr. P.O.
New Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 1.6km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- evenly distributed in lower reaches
------	---------------------------------------

GENERAL REMARKS

- 1972 First report. This stream has very good potential - slow moving, good steady flow conditions plus good spawning gravel. The stream has been hard hit by sub-division development and the road construction around the area.
- 1979 This stream has potential if passage was available past the falls at the mouth of the stream. The Sechelt Fish and Game Club have undertaken to install an incubation box (S.E.P. Program). High flooding in Dec. this year may have destroyed the egg take.
- 1980 Sechelt Fish and Game Club intend to blast out the falls at the mouth of the creek and intend to plant chum in the system.
- 1981 Falls blasted to allow fish passage upstream.

ESCAPEMENT RECORD FOR FLUME CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70			NO RECORDS PRIOR TO 1972			
71						
72				25		
73				75		
74				25		
75						
76			UNK	N.O.		
77			UNK	N.O.		
78			UNK	N.O.		
79			UNK	N.O.		
80			N.O.	UNK		
81			N.O.	UNK		
82			N.O.	2		
83			UNK	6		
84			N.O.	N.O.		
85						

TIMING

ARRIVE				M OCT		
START				L OCT		
PEAK				E - L NOV		
END				L NOV-M DEC		

REMARK

Hastings Creek

see

Lynn Creek p.79

NAME OF STREAM HASTINGS CREEK RAB NO. 90-0800-020
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 19' 123° 01'
 LOCATION OF MOUTH Flows SE into Lynn Cr. E. of North Vancouver, New West. Dist.

 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable concrete culvert at .40km
 Impassable culvert at Lynn Valley Rd. 2km upstream.

SPAWNING DISTRIBUTION

Species Section of Stream Used

coho	from Arbourlynn Dr. to Ross Rd. (83) above fish ladder to Lynn Valley Rd. (1981)
------	---

GENERAL REMARKS

- 1977 Recommend that part of the culvert at Arbourlynn Rd. be removed and a small fish ladder be installed. Hastings Creek had more coho and steelhead on the spawning grounds than Lynn Cr. prior to the installation of the culvert in the early 60's
- 1978 Fish ladder was installed by the Squaretailers Fish and Game Club in Aug. of this year. No coho were observed upstream. By mid December several steelhead were observed above the ladder.
- 1979 Recommend that baffles are installed at the Lynn Valley Rd. culvert. An incubation box was installed at Lynn Valley Road and 30,000 Capilano eggs were transported and transplanted.
- 1981 Approx. 30,000 Capilano Hatchery eggs were placed in Hastings Cr.
- Most years reported predation by humans.
 Seasonal fluctuations in water levels.


ESCAPEMENT RECORD FOR HASTINGS CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77			16			
78			12			
79			50			6
80			175			18
81			85			15
82			78			2
83			16			
84			32			3
85						

TIMING

ARRIVE			OCT			
START			NOV			DEC
PEAK			NOV - E DEC			MAR
END			E - L DEC			APL

REMARK



NAME OF STREAM HIGH FALLS CREEK RAB NO. 90-1300-150
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 56' 123° 19'
 LOCATION OF MOUTH Flows S.W. into Squamish River, New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 3.2km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	<ul style="list-style-type: none"> - random distribution from Squamish confluence to just above logging road bridge (1982) - main concentration between confluence of B.C.Hydro powerhouse channel and T.F.L.main road (1983)
------	---

GENERAL REMARKS

1979 First Report -- this streams populations previously included in Squamish River figures.

1981 High w/l on October 31.

1980 Flood conditions in late December. Losses to coho in the 40% range. Some erosion silting and scouring.

1982 Stream is very ill defined with many small branches within the lower reaches.

1983 Channel flows into B.C.Hydro powerhouse channel which is considered part of the Squamish River. Chinook primarily spawn in the channel and seem to migrate into High Falls Creek on only an incidental basis.

Water levels normal throughout most of the year -- high flows over fall and early winter months.

Light predation by bears and eagles.

ESCAPEMENT RECORD FOR HIGH FALLS CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			75			25
80			200			
81			400			
82		10	50			
83			50			
84			50			
85						

TIMING

ARRIVE			OCT			APL
START			OCT - E NOV			APL
PEAK			DEC - E JAN			MAY
END			DEC - M JAN			JUN

REMARK Figures for this stream previously included in Squamish River population.

Hop Ranch Creek

see

Dryden Creek p.43

NAME OF STREAM HOP RANCH CREEK RAB NO. 90-1300-030
 LOCAL NAME (Hop Ranch Creek System (4))
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 45' 123° 08'
 LOCATION OF MOUTH Flows S.W. into Squamish River, New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 1.00 Jan. 20, 1968 MIN -
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 3.2km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- even distribution between highway 99 and Squamish confluence
chum	- " " "

GENERAL REMARKS

- 1979 This streams population previously included with Squamish River figs.
 1980 Flood conditions in late December. Estimate 25% loss of coho spawn.
 Some silting erosion and scouring.
 1982 Slight change in stream course in lower reach due to flood control.
 Water levels normal except during late Oct. flood when water levels
 very high.
 Pumping station for flood control put in place at Squamish River con-
 fluence. During flood periods control gates to culverts are closed
 and runoff is either allowed to back up in creek or is pumped over
 dyke.
 1983 This stream is very susceptible to public harassment -- flows through
 major residential area.


ESCAPEMENT RECORD FOR HOP RANCH CREEK (Hop Ranch Creek System (4))


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25			
80			200			
81			50	50		
82			30	50		
83			30			
84			30			
85						

TIMING

ARRIVE			OCT			
START			E - M NOV			
PEAK			L NOV - DEC			
END			M DEC - JAN			

REMARK This stream previously included with Squamish River counts.





NAME OF STREAM INDIAN RIVER RAB NO. 90-0500
 LOCAL NAME (Burrard River)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 28' 122° 53'
 LOCATION OF MOUTH Flows S. into Indian Arm, New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

falls at 9.6km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum
pinks
chinook
coho

- in first 6.4km heavy concentration in sloughs
 - in lower 5.6km concentrated in main river
 - above Twin Bridges and below falls
 - in sloughs and lower end of tributary streams

GENERAL REMARKS

- 1959 Canadian Colliery Resources constructed a logging road up the right bank of the river to the falls. In 1969 they started logging and shake manufacturing at the river mouth.
- 1961 The Department erected a counting weir and holding facilities at 1.6km. Pink salmon eggs were collected for transplant purposes.
- 1966 A transplant of chinook fry from the Qualicum River was undertaken in an effort to establish a run of this species. Results were negative.
- 1968 When B.C.Hydro cleared their transmission line right of way, they left areas in the bottom of the valley which were subject to breakthrough.
- 1972 Flood water cut through the B.C.Hydro right of way and the existing channel was cut off when the river reverted to a channel cut in 1968. The length of the changed portion of the stream is approx. 4km.
- 1973 Most of the egg taking facilities installed by the R.D.Branch were removed by Weldwood for the Department.
- 1974 B.C.Hydro did usual maintainance work on towers. Weldwood curtailed logging operations and will resume in 1975. Minor repairs on bridge will be carried out, but no Fisheries problems expected.
- 1975 Pinks arrived at mouth last week in August and unseasonal heavy rains resulted in brief period of stay in estuary.
- 1976 Approx 12 redds were observed high and dry.
- 1977 Any future herbicide spraying must be monitored by Fisheries personnel.
- 1978 Historically chinook have not utilized this river, but a small run has established itself.

continued

INDIAN RIVER

1984 Some over spawning .

Physical Conditions:

- 1968 30% silting during heavy freshets — heavy scouring and stream changes in 3 locations.
- 1969 Some changes of location of main channel in Hixon Cr. area.
- 1978 Minor silting, 15 - 20%, and slight scouring.
- 1981 50-60% erosion and silting, extensive changes in course after high water. 60% loss of pink spawn due to scouring.
- 1983 20% erosion and silting — 25% of pink and chum spawning area affected by scouring. Water levels low during summer and fall — high in first week of November floods.
- 1984 Side channel has changed in some areas. Erosion above falls by road side. Flooding and freshetts cause river to change course each year.

Predation: bears, merganzers, humans. 1984 reported 100 bald eagles in vicinity.


ESCAPEMENT RECORD FOR INDIAN RIVER (Burrard River)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947	25		750	15000	75000	75
48			1500	15000	N/O	75
49	25		1500	3500	100000+	75
50			1500	3500	N/O	400
51			3500	35000	75000	750
52			3500	35000	75	400
53			1500	1500	100000+	400
54			1500	35000	200	400
55			3500	3500	75000	400
56			1500	1500		400
57			1500	3500	125000	400
58			750	15000	N/O	200
59			UNK	UNK	UNK	UNK
60			1500	4000		400
61			3500	2500	67800	200
62			400	3500		400
63			1500	3000	200000	400
64			3500	5000		200
65			400	3500	35000	400
66			1500	3500	75	200
67			1500	3500	7500	200
68			750	15000	N/O	400
69	25		400	15000	7500	200
70	25		750	15000	N/O	200
71	25		750	7500	35000	200
72	N/O		400	35000	N/O	200
73	N/O		750	35000	35000	200
74	N/O		750	7500	N/O	200
75	75		200	15000	35000	200
76			200	20000	-	UNK
77	25	25	500	14000	22000	25
78	7	6	150	7000	-	25
79	12	180	280	7500	22000	50
80	25	50	300	15000	-	150
81	8	20	800	17500	40000	80
82		50	450	24000	-	75
83	16	70	700	26000	24000	55
84	12	100	600	30000		180
85						

TIMING

ARRIVE	E SEPT	E SEPT	E SEPT	E - M OCT		DEC
START	M AUG	E SEPT	JUL -L SEPT	E - M OCT	JUL - SEPT	
PEAK	SEPT - E OCT	L OCT	OCT - NOV	E OCT - NOV	SEPT	
END	L OCT	E DEC	NOV - L JAN	NOV - L DEC	L SEPT-OCT	MAY

REMARK



NAME OF STREAM JUDD SLOUGH RAB NO. 90-1300-040
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 46' 123° 10'
 LOCATION OF MOUTH Flows S. into Squamish River through Brackendale Community.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- throughout channel
chum	- throughout channel

GENERAL REMARKS

Formerly an active flood channel of the Squamish River and one of the principal chum spawning areas, Judd Slough has been affected by a number of engineering activities since 1967 as follows:

- 1967 An extensive log jam which formed at the upper end of the slough was partially cleared and two culverts were installed to restore flow to spawning beds. At the same time, a rip-rap armoured dyke was constructed at the inlet to prevent a major breakthrough of the Squamish River into the slough.
- 1968 Culverts were cleared of debris.
- 1969 The log jam was completely removed, the dyke strengthened and the twin culverts replaced with a single 5 foot diameter culvert.
- 1975 As part of a joint program to upgrade and extend flood control works in the Squamish - Brackendale area, the Province and Municipality of Squamish constructed a dyke parallel to the mainstem Squamish River which completely encompassed Judd Slough. The old Fisheries-built dyke was replaced and a new culvert was installed along with a manually-operated intake valve and intake trash rack to provide controlled inflow to the slough. At a point where the dyke crosses the lower end of the slough, a twin-culvert concrete outlet structure with drop gates was also installed. In the event of a severe freshet, both intake and outlet gates could be closed, isolating the slough and preventing flooding of

Judd Slough

residential property on lowlands adjacent to the slough. Internal drainage was provided by a large-capacity pumping station.

1978 and 1979 Improvement work was carried out on the main slough and two minor tributary channels by channelizing with a bulldozer. Channelization created a more uniform width and gradient, increasing the area suitable for spawning. Coarse materials were excavated from the stream bed and placed on banks to increase their stability. At the same time, two experimental pond-type spawning areas were created by excavating adjacent to the channel. Graded gravel was added to one of the two ponds.

1979-84 Siltation resulting from the Squamish River inflow, and the rapid spread of grass-like rooted aquatic weed caused deterioration of spawning beds. At the request of the Small Projects Unit of SEP the Province removed the intake structure during the course of other flood control work in the Squamish area. Flows in Judd Slough were then provided only by silt-free ground water and an opportunity was created for further development aimed at a purely groundwater-fed spawning channel.

1984 The Small Projects Unit rechannelized Judd Slough, excavating the old channel to a greater depth to increase the flow of groundwater. Rip-rap armouring was added to the banks to prevent undermining by spawning fish and to create cover for juvenile coho salmon. Size graded spawning gravel was not added since the native material was of good quality. The area of improved spawning/rearing habitat created in the developed portion of Judd Slough is 5360 m².


ESCAPEMENT RECORD FOR (Judd Slough)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84			50	7000		
85						

TIMING

ARRIVE			E DEC	L OCT		
START			M DEC	E NOV		
PEAK			E JAN	E DEC		
END			L JAN	L DEC		

REMARK



NAME OF STREAM LANGDALE CREEK RAB NO. 90-1525
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 26' 123° 28'
 LOCATION OF MOUTH Flows E. into Thornbrough Channel, N. of Hopkins Landing,
New Westminster Dist.
 LENGTH 1.6 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 3.74 Sept. 18, 1968 MIN 0.001 Sept. 19, 1968
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 1.6km, culvert at 1.2km -- passable at suitable water levels

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- coho fry observed above culvert during inspection Oct. 1974
chum	- evenly distributed below culvert

GENERAL REMARKS

- 1971 A second growth is becoming well established on the watershed. This should help to stabilize the stream. Fish molestation by juveniles is a problem.
- 1972 Heavy rains in Dec. caused silting and erosion which affected 20% of the stream and scouring which affected the entire spawning area. Estimate 60 - 70% of spawn was lost.
- 1977 Gravel scoured out.
- 1979 Some scouring due to high water levels in December. Some fish molesting. This stream fluctuates considerably. Spawning gravel is concentrated in the lower reaches with large rocks and fast waters further upstream.
- 1982 High water in October from heavy rains.


ESCAPEMENT RECORD FOR LANGDALE CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69		NO RECORDS PRIOR TO 1970				
70				200		
71				50		
72				150		
73				75		
74				25		
75				25		
76				N.O.		
77				125		
78				50		
79				25		
80				12		
81				N.O.		
82				24		
83				6		
84				6		
85						

TIMING

ARRIVE				E OCT-E NOV		
START				E OCT-M NOV		
PEAK				E-M NOV		
END				E NOV-M DEC		

REMARK



NAME OF STREAM (Loggers Lane Creek) RAB NO. -
 LOCAL NAME
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°43' 123°09'
 LOCATION OF MOUTH Flows into Mamquam Blind Channel, near Dentville Community.

LENGTH 4.8 km WIDTH m DRAINAGE km²
 DISCHARGE (m³/s) MAX MIN
 Temperature (°C)
 COMPOSITION: Bedrock Boulder Coarse Fine
 Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- in pond at mouth of stream -- upstream side of Loggers Lane Rd.
------	---

GENERAL REMARKS

1981 Reported spawning in top end of creek.


ESCAPEMENT RECORD FOR (Loggers Lane Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25			
80			25			
81			20			
82			25			
83			5			
84			10			
85						

TIMING

ARRIVE			NOV			
START			DEC			
PEAK			DEC-JAN			
END			JAN-FEB			

REMARK



NAME OF STREAM (Long Bay Creek) RAB NO. 90-1500-027
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 30' 123° 23'
 LOCATION OF MOUTH Flows S.E. into Port Graves Bay, Gambier Island, New Westminster
 Dist. _____
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable at 1.2km where stream narrows and gradient rises rapidly
 some storm debris (leaves and branches)

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- evenly distributed -- some overspawning in heavy run years.
------	---

GENERAL REMARKS

- 1969/70 This small stream has excellent spawning beds, good gravel, well defined banks and a stable water flow.
 1976/77 Considerable overspawning and egg-digging in shallow gravel areas. Addition of gravel would be a good project on this stream. The stream is potentially a good producer. It is away from civilization and there are no predators of any consequence.
 1978 Approx 30% of stream affected by erosion -- flash floods and high water cause scouring and shifting gravel areas. Debris removed by hand after storms.
 1979 25% of lower stream below culvert affected by siltation. Heavy rains in mid Dec. scoured out spawning bed and destroyed 80% of spawn. Sub-division development in area may lead to frequent flash flooding in future if a substantial area of land is cleared
 1981/82 Scouring through right fork of stream.


ESCAPEMENT RECORD FOR (Long Bay Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64		NO RECORDS PRIOR TO 1965				
65				100		
66				1500		
67				300		
68				900		
69				2000		
70				3000		
71				1100		
72				2200		
73				3000		
74				200		
75				200		
76				1500		
77				3600		
78				1850		
79				1700		
80				1500		
81				2000		
82				520		
83				2100		
84				3900		
85						

TIMING

ARRIVE				E-L OCT		
START				E OCT-M NOV		
PEAK				M OCT-M NOV		
END				E-L DEC		

REMARK



NAME OF STREAM LYNN CREEK RAB NO. 90-0800
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°19' 123°02'
 LOCATION OF MOUTH Flows S. into Burrard Inlet, W. of Seymour R. New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls in canyon at 5 - 6 km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- in canyon
chum	- in lower reaches

GENERAL REMARKS

- 1968 The tidal portion of this stream was deepened in 1968 by the Vancouver Cruising Club.
- 1969 Steelhead and coho in this stream are subject to heavy sport fishing pressure during the open season.
- 1970, 71, 73, Industrial pollution, leachate from Lynn Creek Garbage Dump and gravel removal by North Shore Municipalities over the past 10-15 years may account for the lack of chum and pink in this area. The municipalities are presently observing gravel removal restrictions.
- 1971 Because the city of North Vancouver takes its water supply from this creek, the water flow is critically low during dry periods. In 1971, the guaranteed flow was 2 cfs.
- 1974 A partial obstruction was eliminated when the Dist. of North Vancouver replaced and buried a water main at a crossing site below the canyon.
- 1975 Estuary of the creek channelized and filled for large deep-sea lumber terminal.
- 1977 A renewed application to remove gravel from lower portion of creek will have an adverse effect on salmon enhancement. The continued use of the garbage dump site adjacent to the creek contributes to the degradation of spawning area used by chum and pinks.
- 1979 Heavy December rains probably destroyed the majority of this years spawn.
- 1981 Heavy October rains have probably destroyed the majority of this years early spawn. Toxic Leachates from Premier Landfill site were found in the creek.

LYNN CREEK

Physical conditions:

1979 10% erosion and silting -- heavy scouring of bed during heavy Dec. rain.

1981 75% erosion and silting -- scouring from canyon to 3rd St. Bridge.

1982 Water levels low during June to September.

1979/84 Reported predation by humans.


ESCAPEMENT RECORD FOR LYNN CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75	25	75	25
48			400	200	N/O	25
49			75	200	75	25
50			200	75	N/O	25
51			200	25	25	75
52			200	200	N/O	25
53			75	25	25	25
54			75	200	N/O	75
55			200	25	25	75
56			75	UNK	N/O	75
57			200	25	25	25
58			75	75	N/O	75
59			UNK	UNK	UNK	UNK
60			25	25		25
61			25	25	25	25
62			25	25		25
63			25	25	25	25
64			25	25		25
65			25	25	25	25
66			25	25		750
67			25	25		25
68			25	25		25
69			200	N/O	N/O	200
70			75	N/O	N/O	25
71			75	N/O	N/O	25
72			400	400	N/O	200
73			75	25	N/O	25
74			75	25	N/O	25
75			75			25
76			30	N/O		UNK
77			60	N/O		25
78			85	6		25
79		4	42	10	26	
80			250	10		60
81		4	120	10		35
82		6	121			42
83		6	175	N/O	N/O	18
84			185	16		44
85						

TIMING

ARRIVE		JUN-M NOV	E-M OCT		JAN
START		M OCT	JUN-M NOV	E OCT-M NOV	E SEPT-L OCT
PEAK			SEPT-L NOV	M OCT-L NOV	SEPT
END			NOV-E JAN	L OCT-L NOV	L OCT DEC

REMARK



NAME OF STREAM MACKAY CREEK RAB NO. 90-0860
 LOCAL NAME (McKay Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 13' 123° 06'
 LOCATION OF MOUTH Flows S. into Burrard Inlet, E. of Capilano R. New Westminster
Dist.
 LENGTH km WIDTH m DRAINAGE km²
 DISCHARGE (m³/s) MAX 6.14 Dec. 17, 1979 MIN 0.005 Aug. 22, 1977
 Temperature (°C)
 COMPOSITION: Bedrock Boulder Coarse Fine
 Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

Impassable small dam at 3.2 km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- even distribution from Marine Drive to Ridgewood Avenue
------	---

GENERAL REMARKS

- 1958 Water conditions are seriously affected by the extensive land clearing and residential development of the watershed areas of the stream. During the past dry season the stream bed was practically dry and it is to be expected that the resident coho fry population suffered heavily.
 NO RECORDS UNTIL 1978
 1978 Reports predation by public.
 Seasonal fluctuations in water levels.


ESCAPEMENT RECORD FOR MACKAY CREEK (McKay Creek)


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			25			
48			75			
49			25			
50			25			
51			75			
52			25			
53			25			
54			25			
55			25			
56			25			
57			25			
58			25			
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75		NO RECORDS SINCE 1958				
76						
77						
78			8			
79			18	4		
80			12	4		6
81			25	6		6
82			16			
83			16			
84			32			
85						

TIMING

ARRIVE			M OCT			
START			JUL-L OCT			
PEAK			SEPT-L NOV	M NOV		MAR
END			L OCT-L DEC			

REMARK





NAME OF STREAM MAMQUAM RIVER RAB NO. 90-1300-020
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°44'123°09'
 LOCATION OF MOUTH Flows W. and S. into mouth of Squamish River, New Westminster
Dist. _____
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 248 July 12, 1972 MIN 2.13 Sept. 15, 1970
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 4.8km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chinook pink chum sockeye	- scattered mainly above Mashiter Cr. confluence. - scattered primarily below Hwy. 99 bridge - in mainstream and in new spawning channel - scattered
------------------------------------	---

GENERAL REMARKS

- 1963 The Provincial Government built a rock groin on the left bank of the river in the lower reaches to help stabilize the bank.
- 1970 An estimated 1000-1500 chum were destroyed by uninformed citizens.
- 1971 Very heavy sports fishery -- large numbers of fish killed by juveniles.
- 1972 25 - 30% loss of spawn due to very high water in Dec. Since the watershed in the upper reaches has been logged off, the stream is subject to rapid fluctuations in water levels.
- 1973 Fish molestation is a problem as the main part of the stream runs through the municipality.
- 1974 Flash flooding causes up to 35% of the gravel bars in the spawning grounds to shift after spawning. Most of the side channels are stable, but the main channel moves several times a year. Stabilization of flows will occur when the second growth is well established. This should not be too far in the future as some improvement has been noted in the last four years.
- 1975 Heavy floods in Oct. caused estimated loss of up to 80% in pink and chinook spawn.
- 1978 Stream subject to a very heavy sports fishery both spring and fall. Chinook return very poor. Coho did not arrive in any numbers until late December.

continued.....

continuation

MAMQUAM RIVER

- 1981 Chinook and pink spawn would have been damaged during the flood conditions on Oct. 31.
- 1982 Up to 700,000 cu. yds. of gravel removed from lower reaches to build up gravel bars. This amount may change the stream dynamics of the lower reaches.
- 1983 20% chum spawned in mainstream while 80% spawned in newly created ground water fed channel dug in conjunction with construction of dykes. Most sockeye and coho seen in this channel, which offers far more favorable spawning habitat as compared to the fast flowing Mamquam River. Extensive gravel removal and dyking operations took place from July to October. Dyking program for Mamquam now almost complete.

Physical conditions:


- 1974 Much gravel shifting due to flash flooding -- some 35% area shifted after spawning took place.
- 1975 60-75% of spawning beds affected during late October.
- 1980 Heavy siltation late Dec. -- much gravel movement. Extremely high W/L in late October.
- 1982 Some movement and gravel shifting -- mainstream relocated 36m closer to South bank just below Mashiter Creek.
- 1983 Major change in stream course below B.C. Railway Bridge as a result of gravel removal operations.

Biological conditions:

- 1978 reported some egg-digging due to late fall low water conditions.
- 1983 Some egg digging, 10%.

Predation: Birds, bears and public harassment.

Sketch of Lower Portion of Mamquam River, 1968



ESCAPEMENT RECORD FOR MAMQUAM RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947		200	400	750	7500	400
48		75	750	1500	N/O	400
49		75	200	1500	7500	75
50		200	200	3500		200
51		200	7500	3500	7500	200
52		75	3500	7500		75
53		200	200	750	3500	200
54		750	1500	3500		200
55		200	1500	3500	750	75
56		75	200	1500		200
57		400	1500	3500	750	200
58		200	750	1500		75
59		200	200	7500	3500	200
60		75	1500	3500		200
61		400	1500	3500	15000	75
62		200	400	400		200
63		200	750	3500	100000+	200
64		75	3500	3500		75
65		750	750	400	3500	200
66		200	400	200		200
67		200	1500	300	500	50
68		100	500	2500		300
69		800	500	10000	500	300
70		1500	8000	25000	N/O	500
71		700	6000	3000	1500	400
72		350	1500	45000		1300
73	12	750	2500	45000	35000	4000
74		400	3000	25000		1500
75		75	750	7500	3500	75
76		400	750	15000	200	400
77	75	200	200	750	750	200
78	80	60	400	30000		150
79		25	200	400	750	200
80		150	500	12000		100
81		150	250	5000	1500	125
82		150	300	6000		125
83	30	50	300	5000	300	75
84		200	400	15000		
85						

TIMING

ARRIVE		M JUL-AUG	M SEPT-E NOV	SEPT-L OCT	E AUG	M FEB
START		JUN-M AUG	SEPT-L NOV	L OCT-NOV	AUG	
PEAK		JUL-E SEPT	SEPT-M DEC	NOV-DEC	SEPT	
END		JUL-L SEPT	OCT-M JAN	DEC-M JAN	OCT	MAY

REMARK

Mamquam Spawning Channel

see

Mamquam River p.87

NAME OF STREAM (Mamquam Spawning Channel) RAB NO. -LOCAL NAME DISTRICT STATISTICAL AREA POSITION 49° 44' 123° 09'LOCATION OF MOUTH Flows into Mamquam River near Squamish Golf and Country ClubLENGTH .80 km WIDTH m DRAINAGE km²DISCHARGE (m³/s) MAX MIN Temperature (°C) COMPOSITION: Bedrock Boulder Coarse Fine Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

sockeye	- throughout channel
coho	- " "
chum	- " "

GENERAL REMARKS

1984 SEP personnel conducted a dead pitch on this system during late 1984 and early 1985

This Channel, developed in 1983, is a cooperative enhancement project between the Small Projects Unit of SEP and the agencies involved in local dyke construction: the Provincial Water Management Branch and the Municipality of Squamish. Material excavated in developing the channel was used to construct an adjacent dyke. The new channel, fed by groundwater, provides 2000 m² (about 400 m long by 5 m wide) of new spawning habitat for chum and coho and rearing/overwintering habitat for juvenile coho.


ESCAPEMENT RECORD FOR (Mamquam Spawning Channel)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84	50		100	2000		
85						

TIMING

ARRIVE	M AUG		E DEC	E OCT		M FEB
START	L AUG		L DEC	E NOV		TO
PEAK	E SEPT		E JAN	E DEC		
END	L SEPT		L JAN	E JAN		MAY

REMARK



NAME OF STREAM MANNION CREEK RAB NO. 90-1500-040
 LOCAL NAME (Cotton Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°28' 123°27'
 LOCATION OF MOUTH Flows S.W. into Thornbrough Channel, S.W. side of Gambier Island,
New Westminster Dist.
 LENGTH km WIDTH m DRAINAGE km²
 DISCHARGE (m³/s) MAX MIN
 Temperature (°C)
 COMPOSITION: Bedrock Boulder Coarse Fine
 Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

Impassable rock falls at .80km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- evenly distributed throughout
------	---------------------------------

GENERAL REMARKS

- 1969 A small stream with .80km of suitable spawning area -- remainder is short, steep and unsuitable.
 1978 Very poor escapement this year for unknown reasons. Possibly cleaned out by commercial fishery last week in October. Water conditions were ideal.
 1980 Heavy scouring of stream bed during December rains. W/L low in Sept. Oct. delayed migration. Water levels abnormally high in December -- gravel composition too coarse for spawning in many areas.
 1984 Water levels low Jun. to Oct. -- human predation.

Seasonal fluctuations in water levels.

Light predation.


ESCAPEMENT RECORD FOR MANNION CREEK (Cotton Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68		NO RECORDS PRIOR TO 1969				
69				500		
70				200		
71				100		
72				250		
73				750		
74				75		
75				400		
76				125		
77				750		
78				90		
79				175		
80				150		
81				300		
82			65	50		
83				400		
84				350		
85						

TIMING

ARRIVE				M OCT-M NOV		
START				E-M NOV		
PEAK				M NOV-M DEC		
END				M-L DEC		

REMARK



NAME OF STREAM MASHITER CREEK RAB NO. 90-1300-020-010
 LOCAL NAME (Little Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 44' 123° 04'
 LOCATION OF MOUTH Flows S.W. into Mamquam River, E. of Squamish River, New West.
District.
 LENGTH km WIDTH m DRAINAGE km²
 DISCHARGE (m³/s) MAX 44.2 Nov. 3, 1975 MIN 0.289 Sept. 16, 1973
 Temperature (°C)
 COMPOSITION: Bedrock Boulder Coarse Fine
 Silt & Sand Unclassified

Barriers or Points of Difficult Ascent:

Impassable falls at 1.20

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- scattered in the suitable gravel beds lying upstream
pink	- of logs and boulders.
	- " " " " "

GENERAL REMARKS

- 1979 Approx 2.4km of excellent spawning gravel was lost when this creek cut a new channel into the Mamquam River.
- 1980 Erosion and silting heavy during December flood -- some scouring -- estimated 90-100% loss of spawn.
- 1982 Some bank erosion -- approx 30% of stream bed load moved during late October flood. Fast flowing stream with steep gradient, can only support light runs.
- 1983 Major changes in physical condition of this stream has resulted in a sharp decline in its productivity which will most likely not be re-established.


ESCAPEMENT RECORD FOR MASHITER CREEK (Little Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25	25	25	
80			25	N/O		
81			25		75	
82			20	50		
83			20			
84			100			
85						

TIMING

ARRIVE			OCT-E NOV		JUL	
START			OCT-L NOV		AUG	
PEAK			M NOV-M DEC		M SEPT	
END			JAN		OCT	

REMARK



NAME OF STREAM McCARTNEY CREEK RAB NO. 90-0690
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 19' 123° 00'
 LOCATION OF MOUTH Flows S. into Vancouver Harbour

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable steep incline at .40m

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- unknown
------	-----------

GENERAL REMARKS

First Report: 1982 Water levels low June - September.
 1984 Water levels low June - October, predation by humans.


ESCAPEMENT RECORD FOR MCCARTNEY CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82			22			
83			13			
84			39			
85						

TIMING

ARRIVE			SEPT-M OCT			
START			SEPT-E NOV			
PEAK			OCT-L NOV			
END			NOV-L DEC			

REMARK



NAME OF STREAM McNAB CREEK RAB NO. 90-1370
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 33' 123° 24'
 LOCATION OF MOUTH Flows S. into Thornbrough Channel, N. of Gambier Isl., New West.
Dist.
 LENGTH 4 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 6.3km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- concentrated in small tributary .80 km from mouth
pink	- in lower .80km

GENERAL REMARKS

- This stream is subject to flooding and has changed course several times over the years.
- 1969 Some sections of the stream have excellent gravel. Gravel bars and minor debris in the lower reaches. As one point, where the road parallels a canyon, road material spills close to the creek and could become a problem during periods of heavy rain. Recommend that close attention be given to this potential slide area.
- 1973 In October, dredging operations on this stream were completed. A channel approx 18m wide and 6lm in length was dredged out of the mud flats immediately east of the mouth of the creek. The channel will be used for mooring small boats.
- 1979 More exploratory work is required on this creek to determine arrival dates and peak spawning periods. Some logging in the upper watershed should be watched.
- 1980 Majority of spawn occurs in a small unnamed creek that flows into McNab near the power lines. High December flooding will have caused severe scouring and erosion. SEP in looking at this creek for possible enhancement.
- 1982 SEP put an incubation box on this creek and took 40,000 eggs.


ESCAPEMENT RECORD FOR McNAB CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50				75		
51				1500	3500	
52				75		
53				25	3500	
54				25		
55					200	
56						
57				400	400	
58				200		
59				75	75	
60				200		
61				25	25	
62				25		
63				200	25	
64				200		
65				25	25	
66				75		
67				100	50	
68				450		
69				200	50	
70			200	600		
71			150	150		
72						
73			350	350		
74			75	400		
75			25	75	N/O	
76			25	75	N/O	
77			200	1500	25	
78			175	1500		
79		6	150	600		
80			150	1200		
81				1500		
82			N/O	500		
83			100	300		
84			100	2000		
85						

TIMING

ARRIVE		E NOV	E-M OCT		JAN
START		E NOV	SEPT-L OCT	AUG	
PEAK		L NOV-E DEC	L OCT-M NOV	SEPT	
END		E-M DEC	NOV-L DEC	OCT	APL

REMARK



NAME OF STREAM McNAIR CREEK RAB NO. 90-1430
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 31' 123° 30'
 LOCATION OF MOUTH Flows S.E. into Thornbrough Channel, W. of Gambier Isl.
New Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Rock falls at 4.8km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- in lower reaches
chum	- in lower reaches

GENERAL REMARKS

- 1972 First report of any salmon for some years.
 1973 Spawning areas well separated by steep and rocky sections.
 1974 Series of old dams hinder the recruitment of gravel for spawning beds in lower reaches. In Jan, silt from the Dept. of Highways excavations was released into the stream.
 1979 Some logging in the upper watershed. High water in December may have caused some damage.
 1984 New spawning area developed by high water forming side channel below highway.

ESCAPEMENT RECORD FOR McNAIR CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69		NO RECORDS PRIOR TO 1970				
70			N/O	N/O		
71			N/O	N/O		
72				10		
73				6		
74			25	25		
75			UNK	UNK		
76				N/O		
77				N/O		
78			N/O	N/O		
79			N/O	20		
80			N/O	N/O		
81			N/O			
82			N/O	N/O		
83			UNK	N/O		
84			UNK	50		
85						

TIMING

ARRIVE		L SEPT	E-M OCT		
START		E OCT	L OCT-E NOV		
PEAK		L OCT	E-M NOV		
END		E DEC	L NOV-M DEC		

REMARK

Meighan Creek

see

Loggers Lane Creek p.73

NAME OF STREAM (Meighan Creek) RAB NO. 90-1300-027
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°44' 123°09'
 LOCATION OF MOUTH Flows W. near Trailer Park at mouth of Manquam River.

LENGTH 4 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- all spawning observed in the vicinity of coho park
chum	- " " " "

GENERAL REMARKS

1979 This stream was formerly included with Squamish River runs.
 Water levels normal -- fish harassment by people and dogs.

ESCAPEMENT RECORD FOR (Meighan Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25			
80			50			
81			25			
82			-	-		
83			-	-		
84			25	200		
85						

TIMING

ARRIVE			OCT-L NOV	E NOV		
START			OCT-L NOV	M NOV		
PEAK			E NOV-M DEC	E DEC		
END			L NOV-E JAN	M DEC		

REMARK

NAME OF STREAM MOSQUITO CREEK RAB NO. 90-0850
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°19' 123°05'
 LOCATION OF MOUTH Flows S. through North Vancouver into Burrard Inlet, New Westminster
Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 11.8 Jan. 19, 1968 MIN 0.150 Aug. 11, 1968
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

passable to Hwy. 401 -- culvert at 1.6km

SPAWNING DISTRIBUTION

Species

Section of Stream Used


coho	- evenly distributed
------	----------------------

GENERAL REMARKS

- 1956 This stream is being continuously cleared as a flood control measure and very little area is left for spawning.
 1958 Water conditions are seriously affected by extensive land clearing and residential development. During the past dry season the stream beds were practically dry and a serious loss in resident coho fry is to be expected.
 No records until 1981 -- flow affected by diversion.
 1982 Human predation.
 1983 W/L abnormally high Sept.-Nov.
 1984 W/L low during June - October.

Sketch of Mosquito Creek, 1948

□ - coho spawning grounds




ESCAPEMENT RECORD FOR MOSQUITO CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			25			
48			25			
49			25			
50			25			
51			75			
52			25			
53			25			
54			25			
55			25			
56			25			
57			25			
58			25			
59						
60		NO RECORDS AFTER 1958				
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81			4			
82			4			
83			3			
84			2			
85						

TIMING

ARRIVE						
START			M SEPT-E OCT			
PEAK			OCT			
END			L OCT-NOV			

REMARK



NAME OF STREAM NELSON CREEK RAB NO. 90-0990
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 21' 123° 16'
 LOCATION OF MOUTH Flows S.W. into Fisherman's Cove, N. of Pt. Atkinson, New
Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Rock falls at 1.6km

Impassable sluiceway at Marine Drive during low flows

Impassable road culvert on Cranley Dr. at 609m

Impassable sluiceway under P.G.E. trestle in Nelson Canyon at 762m

Impassable bridge at 68m

SPAWNING DISTRIBUTION

Species


Section of Stream Used

chum	- scattered on available gravel
------	---------------------------------

GENERAL REMARKS

- 1958 This stream is subject to very low flows each year until the late fall rains commence. Water from Eagle Lake and the mainstem above the upper limit of migration is dammed and used for the municipality of West Vancouver's domestic water supply.
- 1959 Fisheries Officers reported that this stream was unsuitable for chum salmon propagation because of deforestation, water draw-off, housing, commercial development and poor passage at culverts. Logging and subdivision development has been responsible for denuding the forest cover resulting in extremes of discharge, gravel scouring and water temperature problems.
- 1967 Thunderbird Marine operators eliminated the lower 76m of the stream by dredging to enlarge the bay. As a result, fish are only able to enter the creek when the tides are above 14' (4.3m). A biological study conducted this year concluded that rehabilitation of this stream would not be practical.
- 1979 Slight scouring — approx 25% of spawn lost. Juveniles have killed a few spawners.
- 1981 High water late Oct. 1984 Low water Jun. - Oct.

Sketch of Nelson Creek




ESCAPEMENT RECORD FOR NELSON CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947				400		
48				750		
49				750		
50				1500		
51				750		
52				25		
53				400		
54				750		
55				35		
56				25		25
57				25		
58				25		
59						
60			NO RECORDS FROM 1959 - 1969			
61						
62						
63						
64						
65						
66						
67						
68						
69						
70				75		
71				30		
72				75		
73				35		
74				25		
75				N/O		
76						
77				N/O		
78						
79				50		
80				6		
81				6		
82				N/O		
83				-		
84				3		
85						

TIMING

ARRIVE						
START				E-M OCT		
PEAK				E OCT-M NOV		
END				L OCT-L NOV		

REMARK



NAME OF STREAM OUILLET CREEK RAB NO. 90-1490
 LOCAL NAME (Oulette Creek, Jap Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°28'123°29'
 LOCATION OF MOUTH Flows into Thornbrough Channel, N. of Williamsons Landing,
New Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species


Section of Stream Used

chum	- spawning to the lower hwy. bridge approx .40km from mouth
coho	- unknown

GENERAL REMARKS

- 1968 A 20% loss of spawn occurred this year due to over spawning.
 1969 The mouth of this stream was cleared by a lumber company for a sawmill.
 1972 The estimated loss of spawn due to Dec. floods was approx 40%.
 1973 10% erosion and silting in lower reaches -- slight scouring approx 1.2km from mouth.
 1977 L. and K. Lumber developing this property and #1 stream will be diverted.
 1978 Extensive erosion and silting due to Highways Construction of new culvert -- stream course changed, very low water levels.
 1979 Severe scouring due to instability and high water in December. Survival will be low.
 1980 B.C.I.T. has an incubation box on this creek; but no eggs were put in this year. High flood conditions in December caused some scouring.
 1984 Extreme silting in upper watershed may affect egg mortality. Extra eggs were incubated to compensate for any loss.

Sketch of Oulette Creek, 1969




ESCAPEMENT RECORD FOR (OUILLET CREEK) (Oulette Creek) (Jap Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63			NO RECORDS PRIOR TO 1965			
64						
65				450		
66				800		
67				100		
68				2400		
69				500		
70				4000		
71				3000		
72				3000		
73				1500		
74				25		
75				200		
76				N/O		
77				400		
78			6	50		
79			10	60		
80			N/O	150		
81			20	130		
82			N/O	45		
83			20	100		
84			50	750		
85						

TIMING

ARRIVE			E NOV	E-M OCT		
START			E NOV	E OCT-L NOV		
PEAK			M NOV	E-M NOV		
END			M DEC	L NOV-M DEC		

REMARK



NAME OF STREAM PILLCHUCK CREEK RAB NO. 90-1300-100
 LOCAL NAME (Pilchuck Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 52' 123° 14'
 LOCATION OF MOUTH Flows S.W. into Squamish River, N.W. of mouth of Cheakamus R.
New Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 3.2km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- scattered mainly in the area of Cloudburst Mountain
chum	- scattered mainly in the area of Elizabeth Creek.

GENERAL REMARKS

- 1955 30% of the stream bed was scoured during flooding in early Nov. and an estimated 40% of the coho spawn was lost. Most of the chum entered after the flood.
- 1971 This stream flows slowly along the flats up to 4.8km. Heavy deposits of mud and silt can be found between the scattered gravel beds.
- 1977 There are 3 small streams and one large stream on this system. The largest stream is Cloudburst Creek and is about 6.4km long. It is in this stream that the majority of coho spawn. The chum are mostly in the lower reaches of the Pillchuck itself.
- Seasonal fluctuation in water levels.
- Some predation by bears and dogs.


ESCAPEMENT RECORD FOR PILLCHUCK CREEK (Pillchuck Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			25	25		
48			400	75		
49			400	75		
50			75	75		
51			750	75		
52			750	75		
53			750	75		
54			400	75		
55			400	75		
56			25	25		
57			400	200		
58			200	75		
59			750	750		
60			25	25		
61			200	25		
62			750	25		
63			25	25		
64			400	25		
65			750	25		
66			25	25		
67			50	50		
68			500	100		
69			50	50		
70			200	300		
71			200	200		
72			200	700		
73			300	350		
74			750	200		
75			1500	200		
76			1500	200		
77			1500	200		
78			75			
79			75	25		
80			1000	600		
81			250	300		
82			50	100		
83			50	-		
84			100	N/O		
85						

TIMING

ARRIVE			E OCT	M-L OCT		
START			OCT-M NOV	L OCT-NOV		
PEAK			OCT-M DEC	NOV-E DEC		
END			NOV-E JAN	L DEC-M JAN		

REMARK



NAME OF STREAM POTLATCH RIVER RAB NO. 90-1350
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 35' 123° 19'
 LOCATION OF MOUTH Flows S. into Howe Sound, N. of Anvil Island, New Westminster
Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at approx .40 km from mouth

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- spawning up to .40km
------	------------------------

GENERAL REMARKS

1977 Some erosion along bank during flood -- this occurs near foot bridge approx .40 upstream from mouth. Considerable scouring during flash floods. Reports from watchman indicate that there was a run of chum in the fall of 1976.


ESCAPEMENT RECORD FOR POTLATCH CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77				50		
78				-		
79				-		
80				-		
81				-		
82				-		
83				75		
84						
85						

TIMING

ARRIVE				L OCT		
START				L OCT		
PEAK				L OCT		
END				NOV		

REMARK



NAME OF STREAM RAINY RIVER RAB NO. 90-1400
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 31' 123° 29'
 LOCATION OF MOUTH Flows S.E. into Thornbrough Channel at Port Mellon, New Westminster,
B.C.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 428 Dec. 1, 1958 MIN 0 Sept. 30, 1963
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 4.8km
 dam with fish ladder at 1.20km

SPAWNING DISTRIBUTION

Species

Section of Stream Used


chum	- below dam
coho	- evenly distributed, some above dam

GENERAL REMARKS

A diversion dam was built approx 1.2km from mouth to facilitate the water requirements of a pulp mill constructed at the mouth prior to 1930.
 A Denile-type fishway was constructed in 1954 to assist fish passage.
 No records between 1946 and 1971.
 1971 Coho observed above the dam, chum salmon in stream just above tidal influence. Large stockpile of gravel behind dam — lower reaches below dam very poor spawning ground — heavy boulders with a few patches of suitable gravel in between boulders. Ladder should be inspected to see if an improved design would facilitate passage. River above dam should be checked to see if improvements to the ladder are warranted.
 1978 Several sheets of plywood removed from ladder intake at dam location. Ladder impassable due to screen not having been cleaned out.
 1979 Water levels are controlled by Canfors Lake in the upper drainage area of Rainy River. Minor scouring. Dredging was carried out behind the dam to provide water for the mill.
 1980/81/82 Canfors water supply dam filled up with gravel twice this year. The gravel is then dredged out and hauled away. Some method of getting the gravel downstream should be devised.
 1984 Commencing in the summer of '84 Canfor will be allowed to push gravel directly over the dam to allow for gravel recruitment.
 Seasonal fluctuations in water levels.

Sketch of Rainy River, 1971
(lower spawning grounds)

- ▲ - chum
- - coho
- s - steelhead
- - pink




ESCAPEMENT RECORD FOR RAINY RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69			NO RECORDS PRIOR TO 1971			
70						
71			25	25		
72			25	25		25
73			25	25	25	25
74			75	25		25
75			75	25		25
76			N/O	N/O		
77			N/O	N/O		
78			60	4		
79			N/O	12		
80			N/O	N/O		
81			N/O	N/O		
82				12		
83			UNK	N/O		
84			13	70		
85						

TIMING

ARRIVE			M SEPT-L OCT	E OCT-E NOV		DEC
START			M OCT-M NOV	M OCT-M NOV		JAN
PEAK			L NOV	M OCT-M NOV		FEB
END			M-L DEC	M NOV-M DEC		APL

REMARK



NAME OF STREAM (Richards Creek) RAB NO. -
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA _____ POSITION 49° 23' 122° 52'
 LOCATION OF MOUTH Flows S.W. into Buntzen Bay, Indian Arm.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Steep at .40km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

	unknown
--	---------

GENERAL REMARKS

1981 First report. 10% erosion and silting -- low water levels during summer, flood in late October.
 1982 Low water June - Sept.
 1984 Low water June to Oct.


ESCAPEMENT RECORD FOR (Richards Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81			120		20	
82			27			
83			-			
84			30			
85						

TIMING

ARRIVE			E OCT			
START			-			
PEAK			OCT-L NOV			
END			L NOV-M DEC			

REMARK



NAME OF STREAM ROBERTS CREEK RAB NO. 90-1575

LOCAL NAME _____

DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 25' 123° 38'LOCATION OF MOUTH Flows S. into Str. of Georgia, S.E. of Trail Bay, New Westminster District.LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²DISCHARGE (m³/s) MAX 31.1 Jan. 29, 1965 MIN 0.024 Jul. 9, 1968

Temperature (°C) _____

COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 1.20km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum

- evenly distributed up to 1.20km

GENERAL REMARKS

- 1969 In March an eight foot wide trench was cut across the stream bed, for the purpose of installing a waterline, and the spawning grounds in the area were destroyed. It is recommended that the log jam below the falls be removed as it interferes with the natural movement of the gravel.
- 1972 Silting and erosion caused by high water levels affected about 20% of the spawning beds. An estimated 60-70% of the spawn was lost.


ESCAPEMENT RECORD FOR ROBERTS CREEK


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			25	750		
48			25	400		
49			25	1500		
50			25	750		
51			75	750		
52			25	400		
53						
54						
55			NO RECORDS FROM 1953 - 1964			
56						
57						
58						
59						
60						
61						
62						
63						
64						
65				1200		
66				1000		
67				1500		
68				1200		
69				1500		
70				3000		
71				2100		
72				2500		
73				1300		
74				75		
75				400		
76				100		
77				1000		
78				250		
79				1500		
80				1200		
81			N/O	200		
82				115		
83				1500		
84				6500		
85						

TIMING

ARRIVE				M OCT-E NOV		
START			M OCT	M OCT-M NOV		
PEAK			M NOV	M-L NOV		
END			NOV	L NOV-L DEC		

REMARK





NAME OF STREAM SEYMOUR RIVER RAB NO. 90-8500
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°18' 123°01'
 LOCATION OF MOUTH Flows S. into Second Narrows, Burrard Inlet, New Westminster Dist.

LENGTH 19 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 425 Jan. 15, 1961 MIN 0.314 Sept. 30, 1947
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable G.V.W.B. Dam at 19km

Large boulders and narrow canyon at 4km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- small numbers of coho in tributaries, the balance in main R.
chum	- below G.V.W.B. gate
pink	- " "

GENERAL REMARKS

- 1969 Dredging and dyking with dragline machines to deepen the river to a depth of 8' at zero tide, was carried out in the area from below the railroad crossing to the mouth on the east side of the river. In the past the depth of the river was 8'. However, the depth decreased due to freshets which caused material to move downstream. Because of its plentiful water supply, Seymour River is looked upon as a reservoir that could supplement other systems, such as Capilano Lake and Coquitlam Lake in times of high demand. Precipitation in this area is relatively heavy. The average annual rainfall at the southern end of Seymour Lake is 147" while Vancouver's annual average is 59".
- 1974 This river is subject to a heavy sports fishery because of its proximity to Vancouver. Stray coho from the return to Capilano Hatchery were observed this year.
- 1975 An additional closure was necessary to conserve coho stocks in canyon pools during low water period (15 days). Some pinks observed as far as the pipe line pools. Estimated 1 - 3% Capilano coho strayed into Seymour R.
- 1976 Approx 11% Capilano coho strayed into this river.
- 1981 Erosion and silting 60% below canyon to Dollarton Bridge.
- 1983 15% erosion and silting 15%, scouring in lower spawning area
- Predation by birds, bears and humans.


ESCAPEMENT RECORD FOR SEYMOUR RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			1500	750	1500	750
48			3500	1500	N/O	400
49			1500	1500	1500	750
50			1500	1500	N/O	750
51			1500	1500	750	750
52			3500	1500	25	750
53			1500	750	1500	750
54			1500	3500	25	750
55			3500	200	400	750
56			1500	200		400
57			3500	200	75	750
58			400	200	N/O	200
59			UNK	UNK	UNK	UNK
60			1500	25		750
61			400	25	400	750
62			1500	25		400
63			1500	75	400	400
64			750	25		750
65			400	25	200	200
66			1500	25		400
67			400	25	400	200
68			1500	25	N/O	400
69			1500	N/O	25	400
70			1500		N/O	200
71			3500	25	200	200
72			1500	750	N/O	200
73			1500	25	25	200
74			3500	N/O	N/O	200
75			1500	25	200	200
76		6	1000	N/O	100	75
77		20	3000	150	100	250
78		17	5000	220	-	350
79		150	4500	300	250	300
80		250	9000	250	-	300
81		50	8500	800	1000	600
82		250	8500	200	-	550
83		300	14000	500	1000	550
84		280	13000	600	-	580
85						

TIMING

ARRIVE			JUN - JUL			DEC
START		SEPT	JUN - JUL	OCT	JUL - AUG	TO
PEAK		OCT	M AUG-L SEPT	OCT	AUG - OCT	
END		NOV	SEPT - DEC	NOV	M SEPT-OCT	JUN

REMARK



NAME OF STREAM (Shop #3 Creek) RAB NO. -
 LOCAL NAME _____
 DISTRICT _____ STATISTICAL AREA _____ POSITION $50^{\circ}07'123^{\circ}24'$
 LOCATION OF MOUTH Flows into Elaho River near confluence of Elaho River and
 Squamish River
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 3.2km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- evenly distributed between forestry logging road and 3.2km upstream.
------	--

GENERAL REMARKS

- 1979 First report -- formerly included with Squamish River escapements.
 1980 10% erosion and silting -- minor scouring. High water levels during flood, but no evidence of damage.
 1982 Some bed load movement may have taken place during late Oct. flood.


ESCAPEMENT RECORD FOR (Shop 3 Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25			
80			100			
81			30			
82			200			
83			75			
84			100			
85						

TIMING

ARRIVE			OCT			
START			NOV			
PEAK			L NOV-DEC			
END			L DEC-JAN			

REMARK



NAME OF STREAM SHOVELNOSE CREEK RAB NO. 90-1300-180
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 50° 04' 123° 21'
 LOCATION OF MOUTH Flows S.W. into Squamish River, S. of Elaho River, New Westminster
Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 4.8m

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho chinook chum	- throughout on available gravel - concentrated in lower .40km - primarily in lower reaches.
-------------------------	--

GENERAL REMARKS

1973

An ideal spawning ground with consistent water levels, good gravel and plenty of shaded areas. This stream is groundwater fed — back up water is forced into Shovelnose Creek when Squamish River floods.

1978 Vegetation along this stream bank is coming back well and affords good cover for adults and fingerlings.

1980 Severe flood conditions in late December. The Squamish River cut through its bank and entered Shovelnose at just about the upstream limit of fish migration. The river then utilized the stream channel and scoured out all the spawning area. Estimated 100% loss of spawn.

1982 Outflow of the Squamish River extended 137m due to change in side channel during flood. Tenderfoot hatchery staff concentrated on getting brood stock from this stream. Eagle and bear predation very evident.

1983 The change in physical features of the lower portion of creek has resulted in fish having difficulties in migrating to the upper reaches of stream. Stream course lengthened by 2.4km.

1984 The significant increase in flows since the flood has resulted in a notable decrease in spawning stocks. It will be difficult to enumerate spawners due to increased flows and turbidity.

ESCAPEMENT RECORD FOR SHOVELNOSE CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			25			
48			400			
49			75			
50			25			
51			400			
52			3500			
53			75			
54			200			
55			200			
56			75			
57			75		75	
58			200			
59			75		25	
60			75			
61			200			
62			200			
63			75			
64			400			
65			400			
66			75			
67			200			
68			200			
69			50			
70			200			
71		35	200	75	400	25
72		N/O	250	700		350
73		200	1200	2500	2500	100
74		200	3500	1500		75
75		75	400	400	75	25
76		200	400	3500	-	200
77		25	200	1500	25	N/O
78	5	80	175	75	-	70
79		25	75	75	25	25
80		125	750	1500		80
81		150	600	1200	400	40
82		150	350	400		50
83	10	100	50	50	N/O	40
84		150	N/O	N/O		
85						

TIMING

ARRIVE		M JUN-E AUG	SEPT-E NOV	L SEPT-E NOV	E JUL	FEB
START		E JUL-L AUG	SEPT-L NOV	M OCT-E NOV	E AUG	TO
PEAK		M AUG-E SEPT	OCT-M DEC	M NOV	M SEPT	
END		L SEPT-M OCT	OCT-E JAN	L NOV-DEC	M OCT	JUN

REMARK

Spring Creek

see

Branch 100 Creek p.9

NAME OF STREAM (Spring Creek) RAB NO. -
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°56' 123°19'
 LOCATION OF MOUTH Flows into Squamish River, near 48 km point on W. side of
river.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 2.4km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chinook	-distributed evenly throughout first 1.6km
coho	- " " "
chum	- " " "

GENERAL REMARKS

1982

First report. This creek was previously included with Squamish River escapement. It is very productive for the accessible length, but as it is situated near the 48km point on the west side of the Squamish, it is very difficult to visit on a regular basis. Water levels are greatly influenced by extended periods of precipitation. Bird and bear predation evident. Some egg digging by chum. 1983 Frequent high water over fall and early winter months. Some egg digging, but not extensive. Escapement figures based on observation by hatchery staff and during a November 21 helicopter flight.


ESCAPEMENT RECORD FOR (Spring Creek)


YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82		200	350	4500		
83	10	75	200	4000	15	50
84		100	200	1000		
85						

TIMING

ARRIVE	E SEPT	E AUG	L SEPT-L NOV	L SEPT-E NOV	L AUG	APL
START	M SEPT	M AUG-E DEC	M NOV-E DEC	E OCT-L NOV	E SEPT	APL
PEAK	L SEPT	E-L SEPT	L NOV-L DEC	E NOV-M DEC	M SEPT	MAY
END	E OCT	E-L OCT	L DEC-M JAN	L NOV-L DEC	E OCT	JUN

REMARK





NAME OF STREAM SQUAMISH RIVER RAB NO. 90-1300
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°41' 123°10'
 LOCATION OF MOUTH Flows S. into head of Howe Sound, New Westminster Dist.
 LENGTH 69 km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 2230 Sept.6, 1957 MIN 21.2 Jan. 12, 1979
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 69km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chinook	- majority of chinook spawned above Ashlu Cr.
chum	- 65% chum below Cheakamus R. 35% above Ashlu confluence.
coho	- " " " "

(comments from 1982/83 reports)

GENERAL REMARKS

Accessible tributaries: Elaho R. Ashlu Cr.; Cheakamus R.; Shovelnose Cr.; Meighan Cr.; and Mamquam R.
 Precipitation on the Squamish River watershed ranges from 60" to over 150". Short duration floods occur as a result of heavy rains enhanced by rapid snow melt. Heavy siltation occurs during these freshets. In 1963, Canadian Collieries extended the logging road to the mouth of the Elaho River. Thus, the upper reaches of the Squamish River are now accessible to the public. Due to past development, a large part of the estuary in the area of Mamquam channel has been altered or eliminated by dredge and land fill operations and log storage. An additional port oriented land fill river training dyke and dredge spoil have recently eliminated nearly 100 acres of inter-tidal marsh and mud flat in the central portion of the estuary. B.C. Railway has also proposed expansion of existing port facilities to include a bulk and unit load port development which will occupy a major portion of the inner estuarine environment. In addition to direct habitat loss, this industrial development also causes environmental degradation. The remaining inner area constitutes the only source of inner estuarine food available to juvenile salmon, herring and other fish. Port development will permanently displace the fish which use the inner estuary into an exclusively outer estuarine environment. This could cause a huge decrease in fish populations.

continued.....

SQUAMISH RIVER

- 1963 The logging practiced on this watershed will have an adverse affect on run-off for the next 20 years.
Remedial work was carried out on Judd Slough in Sept. and Nov. 1969 to help alleviate problems related to the extremely low water levels caused by lack of precipitation. The rock dyke at the head of Judd Slough was extended and the two thirty-six inch culverts were replaced with one sixty inch culvert. Access to the main channel of the Squamish River and fresh water exchange between the main channel and inner delta is now possible.
- 1974 This stream supports a very heavy sports fishery and a large Indian food fishery. The course of the stream changes every year.
- 1975 The late October floods caused extensive damage to the egg depositions of the pink and chinook salmon -- possible loss as high as 80% of deposited eggs.
- 1978 Subject to a heavy sports fishery all year round and to a heavy Indian food fishery that probably accounts for 50% take of returning coho and chinook.
- 1980 Severe flood conditions existed at the end of December. Extensive damage was done to spawn in the entire Squamish system. Upper Squamish - estimate 30% chum survival. Coho and chinook will be lost. Mid Squamish - 30% of coho and chinook should survive. Lower Squamish, most chum lost except for Judd Slough.
- 1982 Flood conditions late October may have caused mortality to chinook. Some egg digging by chum.
- 1983 Sockeye appear to utilize confluence areas of west side streams which flow into Squamish River. Minimal egg digging by chum.
- 1984 Specific escapements: Eagle Run Slough 7,000 chum, Baynes Isl. 8000 chum, Three falls 3,000 chum.

Physical characteristics:

- 1975 Serious erosion and silting -- 30-50% of bed affected, heavy scouring throughout.
- 1978 Some stream-bed course changes between 24 and 29 miles.
- 1980 Mainstem affected by gravel movement. Much scouring occurred along the whole length of the river. Debris scattered everywhere. Extremely high water levels on December 26, 27, and 28.
- 1981 Approx 20% of the run was lost due to flooding Oct. and Nov 1 -- estimate about 15% of stream bed adversely affected. Some scouring and change of stream beds, damage moderate.
- 1982 Much scouring due to late Oct. flood conditions, 20% of stream beds may have been directly affected. Extensive scouring and siltation due to movement of tree debris downstream.
- 1983 Major erosion problem noted at clay banks just downstream from Cheakamus River confluence. Many changes in stream flow characteristic of Upper Squamish. Periodically high flows over fall and early winter months -- normal flow patterns over rest of year.
- 1984 Water levels abnormally high during Thanksgiving weekend.

Predation: eagles, seagulls, bears, wolves and coyotes.


ESCAPEMENT RECORD FOR SQUAMISH RIVER

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947		15000	3500	35000	350000	15000
48		15000	75000	75000	N/O	15000
49		15000	7500	35000	250000	35000
50		15000	3500	35000		15000
51		15000	75000	35000	200000	15000
52		15000	75000	175000		35000
53		15000	3500	15000	175000	7500
54		15000	35000	15000		15000
55		15000	15000	15000	100000	15000
56		15000	1500	7500		7500
57		15000	35000	35000	75000	15000
58		15000	15000	35000		15000
59		15000	1500	75000	75000	7500
60		15000	7500	7500		7500
61		15000	35000	7500	15000	15000
62		15000	3500	15000		15000
63		7000	7000	7000	70000	7000
64		7000	30000	15000		15000
65		30000	7000	3000	30000	15000
66		15000	7000	3000		7000
67		5000	10000	20000	15000	10000
68		10000	7000	70000		10000
69		20000	8000	18000	10000	8000
70		25000	15000	50000	N/O	6000
71		8000	18000	15000	15000	2500
72		8000	4500	200000	750	18000
73		12000	18000	120000	70000	12000
74		7500	75000	70000	N/O	15000
75		3500	35000	35000	35000	15000
76		3500	3500	75000	-	7500
77		3500	7500	35000	1500	3500
78	100	1000	7000	20000	-	1200
79		1500	7500	15000	3500	3500
80		2000	4000	150000	-	5500
81		3200	4500	75000	9000	3200
82		3000	7500	100000	-	1000
83	300	2500	5000	50000	1000	500
84	N/O	2000	10000	50000	UNK	
85						

TIMING

ARRIVE		JUN-E JUL	M AUG-E OCT	SEPT-OCT	E AUG	MAR
START	-	JUN-E JUL	SEPT-E NOV	E OCT-NOV	L AUG	TO
PEAK	-	JUL-L SEPT	SEPT-M DEC	M NOV-E DEC	SEPT	
END	-	JUL-M OCT	OCT-L JAN	DEC-M JAN	OCT	MAY

REMARK



NAME OF STREAM STAWAMUS RIVER RAB NO. 90-1200
 LOCAL NAME (Little Stawamus River)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°41' 123°09'
 LOCATION OF MOUTH Flows N.W. and S.W. into head of Howe Sound, New Westminster Dist.

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX 61.4 Oct. 25, 1979 MIN 0.170 Feb. 12, 1975
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Increased gradients creating rock falls
 Impassable rock falls at 3.2km

SPAWNING DISTRIBUTION

Species	Section of Stream Used
coho	- majority of fish located in upper section of Little Stawamas.
chum	- evenly scattered

GENERAL REMARKS

In the first .80km downstream of the obstruction, the stream has a fairly steep gradient and many large boulders. Although the lower 2.4km has a tendency to meander, the stream bed has finer gravel and is better suited to spawning.

This non-glacial stream has clear water and maintains a fairly constant flow throughout the year.

Fish molestation and interference are a problem as 700-800 people live along the water course (1974)

The watershed was previously logged off, but the second growth is now well established. (1973)

This 3.2km spawning area is capable of supporting a large number of chum spawners. Some artificial means or re-establishing chum production on the stream should be given serious consideration. (1973)

Nearly all the coho in this system are in the Little Stawamus River. (1973)

The Little Stawamus main tributary is 4km long, and is groundwater swamp fed. Coho escapement ranges from 150-1500 per year. (1973)

Bulldozer work and land clearing affected approx. 640m of the stream bed of the Little Stawamus River. Trees and brush from both banks, in the upstream end of the housing development project, have been removed.

Consequently, the banks have eroded in several places and silt has been deposited into the stream. There were pools in these affected areas before the land was cleared. The stream now has a man-made channelled

continued

STAWAMUS RIVER

appearance. The stream above and below the cleared area appears to provide an excellent habitat for rearing juvenile coho. There is an abundance of low brush and trees on the margins, and small pools are distributed throughout the stream. Fry were seen below the cleared section and near the stream source. None were seen in the cleared section.


- 1977 About 3.2km of Little Stawamus River is used for spawning, but occasionally fish use the marsh when a big run is in the stream.
- 1978 Fish numbers were fairly evenly split between Little Stawamus main tributary and the main Stawamus.
- 1980 Survival should be in the 70-80% range.
- 1981 Spawning occurred after Oct. 31 flood, so there was no damage done.
- 1982 A small bank stabilization program took place behind the residences of Valleycliff along the Little Stawamus Creek.
- 1983 Stawamus River supports very little spawning and is primarily a transition zone for stocks migrating to Little Stawamus Creek. This creek is subject to damage due to its relationship with the maintenance and development of the Valleycliff subdivision.

Physical characteristics:

- 1980 Severe erosion and silting in main Stawamus -- some in Little Stawamus. Severe scouring in main Stawamus.
- 1981 Gravel movement during flood.
- 1982 Some erosion to lower sections of Stawamus River. Movement of bedload very evident during late October flood.

Predation: Light bird predation, fish very vulnerable to dogs and human molestation.

Sketch of Stawamus River, 1968



ESCAPEMENT RECORD FOR STAWAMUS RIVER (Little Stawmus River)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			400	400	750	200
48			750	400		200
49			750	750	750	200
50			400	400		200
51			750	200	400	25
52			750	75		75
53			200	75	200	75
54			400	200		25
55			400	200	25	75
56			75	25		25
57			400	75	25	25
58			400	75		75
59			75	200	25	25
60			75	75		25
61			25	25	25	25
62			25	25		25
63			75	75	75	25
64			200	25		25
65			75	25	25	25
66			25	25		25
67			50	50	N/O	50
68			50	50		50
69			50	50	N/O	50
70			900	N/O	N/O	25
71			200	N/O	N/O	N/O
72			40	N/O		
73			450	25		25
74			750	N/O		
75			400	N/O		
76			400			25
77			25			25
78			150	UNK		25
79			25	N/O		25
80			N/O			25
81			75			25
82			100	25		20
83			75			
84			200	200		
85						

TIMING

ARRIVE		SEPT-L OCT	L SEPT-M OCT		
START		SEPT-M NOV	OCT-E NOV	SEPT	MAR
PEAK		SEPT-E DEC	OCT-M NOV	SEPT	APL
END		OCT-L DEC	NOV-M DEC	OCT	MAY

REMARK

Tenderfoot Creek

see

Cheakamus Riverp.30

NAME OF STREAM (Tenderfoot Creek) RAB NO. 90-1300-050-013

LOCAL NAME _____

DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 50' 123° 09'LOCATION OF MOUTH Flows S. into Cheakamus RiverLENGTH _____ km WIDTH _____ m DRAINAGE _____ km²DISCHARGE (m³/s) MAX _____ MIN _____

Temperature (°C) _____

COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____

Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Passable to lake, 1.6km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho chum	- on all available gravel and in the lake - " " "
--------------	--

GENERAL REMARKS

- 1981 The Salmon Enhancement Program built a hatchery by the lake this year.
- 1982 This stream is ground water fed from Tenderfoot Lake which eliminates potential for high water changes. Stream is primarily a transitional zone with most spawning taking place in the lake. This system is now greatly influenced by operations at Tenderfoot Hatchery. Coho brood stock is removed from a broom stick fence permanently installed between the Cheakamus confluence and the lake. 1500 cu.yd. of poor gravel was replaced in the lake with 1500 cu. yds. of clean washed gravel.
- 1983 Some bird predation -- a limited amount of egg digging. Coho escapements to system were very late but extremely high when they finally showed up. Access to stream through culvert under B.C.Rail tracks may have hindered adequate chum escapements.
- 1984 Counting fence at the mouth of stream manned by SEP personnel.


ESCAPEMENT RECORD FOR (Tenderfoot Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81		N/O	500	2000		
82			250	800		
83			3500	100		
84			700	10000		
85						

TIMING

ARRIVE			L SEPT-M NOV	L SEPT-E OCT		
START			L OCT-E DEC	E-L OCT		
PEAK			L NOV-E JAN	M NOV		
END			M FEB-L DEC	M DEC		

REMARK



NAME OF STREAM (Thirty Seven Mile Creek) RAB NO. _____

LOCAL NAME _____

DISTRICT 2 STATISTICAL AREA 28 POSITION 50° 04' 123° 21'LOCATION OF MOUTH Flows into Squamish River, N. of Turbid CreekLENGTH _____ km WIDTH _____ m DRAINAGE _____ km²DISCHARGE (m³/s) MAX _____ MIN _____

Temperature (°C) _____

COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____

Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 2.4km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	-evenly distributed throughout accessible reach
chum	- " " " "

GENERAL REMARKS

- 1979 First report, formerly included with Squamish River escapements.
 1980 High water in Dec. but little damage.
 1982 Some bed load movement may have taken place during late Oct. when water level was high. Light bird and bear predation.
 1983 Frequent high water over fall and early winter months, but flushes and resumes normal flows quickly. Good gravel and stable stream characteristics.

ESCAPEMENT RECORD FOR (Thirty Seven Mile Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25	25		
80			300	N/O		
81			-	-		
82			175			
83			200	-		
84						
85						

TIMING

ARRIVE			OCT	OCT		
START			NOV	NOV		
PEAK			NOV	DEC		
END			L DEC	DEC		

REMARK Formerly included with Squamish River.

Twenty-Eight Mile Creek

see

Thirty-Seven Mile Creek p.167

NAME OF STREAM (Twenty Eight Mile Creek) RAB NO. _____

LOCAL NAME _____

DISTRICT 2 STATISTICAL AREA 28 POSITION 50° 02' 123° 15'LOCATION OF MOUTH Flows into Squamish River, SW of Cypress PeakLENGTH _____ km WIDTH _____ m DRAINAGE _____ km²DISCHARGE (m³/s) MAX _____ MIN _____

Temperature (°C) _____

COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____

Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable rock falls at 3.2km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

coho	- evenly distributed
chum	- " "

GENERAL REMARKS

- 1979 First report, formerly included with Squamish River escapements.
Light predation.
- 1980 Some silting from flood, little scouring, high water during flood, and the Squamish River deposited silt in this creek when it overflowed.


ESCAPEMENT RECORD FOR (Twenty Eight Mile Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79			25	25		
80			25	N/O		
81			25	N/O		
82						
83						
84						
85						

TIMING

ARRIVE		SEPT-NOV	OCT		
START		SEPT-E DEC	OCT - NOV		
PEAK		OCT-M JAN	NOV - DEC		
END		DEC-L JAN	DEC		

REMARK Formerly included with Squamish River Runs.



NAME OF STREAM (Twin Creek, Archie's Creek) RAB NO. 90-1475
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49°29' 123°29'
 LOCATION OF MOUTH Flows N.W. into Thornbrough Channel, S. of Port Mellon, New
Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Boulders and steep gradient at approx .40km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- in lower river
------	------------------

GENERAL REMARKS

The salmon spawners reported in 1972 were the first observed since the early sixties.

1972 An estimated 50% of the spawn was lost.

1976 Stream subject to the hazards of population and development. Highways Dept. is depositing deleterious substances in creek.

1977 L and K Lumber developing this property and #1 stream to be diverted.

1978 Fish molestation by unknown persons.

1979 Stream was not properly enumerated this year. It is expected that more chum entered the system in Nov. as well as coho. High Dec. flooding.

ESCAPEMENT RECORD FOR (Twin Creek) (Archies Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69		NO RECORDS PRIOR TO 1970				
70				N/O		
71				N/O		
72				200		
73				150		
74				25		
75				N/O		
76				100		
77				400		
78				120		
79				24		
80			N/O	N/O		
81			N/O	N/O		
82			N/O	N/O		
83			N/I	N/I		
84						
85						

TIMING

ARRIVE				M OCT		
START				M OCT-E NOV		
PEAK				L NOV		
END				M DEC		

REMARK

Upper Paradise Channel

see

Cheakamus River p.30

NAME OF STREAM (Upper Paradise Channel) RAB NO. 90-1300-050-012
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 49' 123° 08'
 LOCATION OF MOUTH Flows into Cheakamus River

LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	-throughout channel
coho	-throughout channel

GENERAL REMARKS

Upper Paradise Channel was developed in 1982 in a former flood channel of the Cheakamus River now isolated from the mainstem at its upstream end by a dyke. The groundwater fed channel was designed primarily to enhance chum, but has also proven itself to be an excellent coho producer, yielding approx. 8000 and 6000 smolts in the spring of 1984 and 1985 respectively. The channel provides 2625 m² (460 m long by 5-6 m wide) of spawning and juvenile rearing/overwintering habitat. Design features include size graded (1.6 - 10.2 cm) spawning gravel, rip-rap armouring on the channel banks, and low weirs of wood construction to control channel gradient and maintain a uniform water depth of 25 - 30 cm.


ESCAPEMENT RECORD FOR (Upper Paradise Channel)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70						
71						
72						
73						
74						
75						
76						
77						
78						
79						
80						
81						
82						
83						
84			100	4000		
85						

TIMING

ARRIVE			L NOV	E NOV		
START			E DEC	E DEC		
PEAK			L DEC	M DEC		
END			E FEB	E JAN		

REMARK 1984 Dead pitch count.



NAME OF STREAM WAKEFIELD CREEK RAB NO. 90-1610
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 28' 123° 48'
 LOCATION OF MOUTH Flows SE and S into Straits of Georgai, W. of Sechelt, New
Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Hwy. culvert at approx 91m — chum do not further
 Falls at .40km

Impass rock falls at 2.4km by Anchor Rd. (first reported '84)

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- intertidal spawning -- after removal of culvert chum as far as falls
------	---

GENERAL REMARKS

- 1970 It was recommended that the obstruction and culvert be removed to enable fish to utilize an additional 274m of spawning grounds. Reports from old timers indicate that at one time coho and steelhead migrated as far as 4.8km.
- 1974 All spawning takes place in the intertidal zone.
- 1976 This stream will never support good populations unless the culvert is removed.
- 1978 Plans for local government to make improvements to the culvert. Attempts were made to prevent logs from entering the mouth of creek.
- 1979 90% scouring in late Dec. survival is expected to be poor.
- 1980 Hwy. Department installed a new culvert this year and chum can now make it up to the falls. Sechelt Wildlife Club installed a SEP box with coho eggs from Mixal Creek. Box is located at the falls. High December floods have caused scouring of spawn.

Predation by dogs, birds and residents.


ESCAPEMENT RECORD FOR WAKEFIELD CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			25	200		
48				75		
49				75		
50			25	200		
51			75	400		
52			25	75		
53						
54						
55						
56						
57						
58						
59						
60	NO RECORDS FROM 1953 - 1969					
61						
62						
63						
64						
65						
66						
67						
68						
69						
70				450		
71				750		
72				1500		
73				750		
74				200		
75				200		
76				50		
77				200		
78				200		
79				250		
80				150		
81				100		
82				N/O		
83			200	300		
84			25	600		
85						

TIMING

ARRIVE		M OCT	L-M OCT		
START		E NOV	M OCT-E NOV		
PEAK		M NOV	M NOV		
END		M DEC	M NOV-M DEC		

REMARK



NAME OF STREAM (West Bay Creek) (Riedly's Brook) RAB NO. 90-1500-035
 LOCAL NAME _____
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 26' 123° 24'
 LOCATION OF MOUTH Flows into West Bay, S. side of Gambier Island, Howe Sound,
New Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 40 - 80 km
 Log jams and debris monitored and removed as necessary.

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- even distribution up to falls
------	---------------------------------

GENERAL REMARKS

- 1969 Only .40km of the stream is suitable for spawning, the remainder is too swift.
 1971 This stream is an exceptional producer for its size.
 1977 Some evidence of egg digging.
 1979 Some scouring during mid Dec. Heavy rains destroyed approx 50% of spawn. Driftwood has mouth has been hindering upstream migration at low tide.
 1980 Scouring during December rains. Low water Sept. and Oct. delayed migration. Driftwood was cleared this year.
 1981 15% erosion and silting -- scouring during Oct. rains.

Predation: mergansers, gulls, eagles, and local population.


ESCAPEMENT RECORD FOR (West Bay Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63						
64		NO RECORDS PRIOR TO 1965				
65				200		
66				700		
67				100		
68				450		
69				300		
70				1500		
71				2200		
72				2400		
73				1800		
74				1500		
75				1500		
76				200		
77				750		
78				550		
79				450		
80				900		
81				1500		
82				460		
83				1500		
84				278		
85						

TIMING

ARRIVE				M OCT-L OCT		
START				E-M NOV		
PEAK				M NOV-M DEC		
END				L NOV-L DEC		

REMARK



NAME OF STREAM (Williamson Creek) RAB NO. 90-1510
 LOCAL NAME (YMCA Creek)
 DISTRICT 2 STATISTICAL AREA 28 POSITION 49° 27' 123° 28'
 LOCATION OF MOUTH Flows into Thornbrough Channel, S. of Williamsons Landing, New
Westminster Dist.
 LENGTH _____ km WIDTH _____ m DRAINAGE _____ km²
 DISCHARGE (m³/s) MAX _____ MIN _____
 Temperature (°C) _____
 COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
 Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

Impassable falls at 2.4km

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- evenly distributed in lower reaches.
------	--

GENERAL REMARKS

- 1971 This is an excellent stream, approx 3.65m wide on the average, with a constant water depth of approx 30 - 45cm. The possible capacity of this stream could be 6-8 thousand salmon.
- 1972 Slight erosion 5% — badly scoured during December rains. Estimated loss of spawn is about 50-60%.
- 1979 Flooding in late December may have caused some damage to spawn. This creek was not closely monitored for coho and it is felt there were a few entering this system.
- 1980 This stream appears to be an excellent stream for enhancement. Good gravel throughout.


ESCAPEMENT RECORD FOR (Williamson Creek) (Y.M.C.A. Creek, Jap Creek)

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947						
48						
49						
50						
51						
52						
53						
54						
55						
56						
57						
58						
59						
60						
61						
62						
63		NO RECORDS PRIOR TO 1965				
64						
65				400		
66				700		
67				100		
68				1600		
69				800		
70				1500		
71				3400		
72				4000		
73				2200		
74				75		
75				25		
76				100		
77				100		
78			7	N/O		
79			N/O	20		
80				75		
81				15		
82				25		
83				10		
84			N/O	650		
85						

TIMING

ARRIVE				E-M OCT		
START				M-OCT-E NOV		
PEAK				M-L NOV		
END				E-M DEC		

REMARK



NAME OF STREAM WILSON CREEK RAB NO. 1

LOCAL NAME

DISTRICT 2 STATISTICAL AREA 29 POSITION 49° 26' 123° 43'

LOCATION OF MOUTH Flows S.W. into Straits of Georgia, E. of Wilson Cr. P.O.,
New Westminster Dist.

LENGTH km WIDTH m DRAINAGE km²

DISCHARGE (m³/s) MAX MIN

Temperature ($^{\circ}\text{C}$) _____

COMPOSITION: Bedrock _____ Boulder _____ Coarse _____ Fine _____
Silt & Sand _____ Unclassified _____

Barriers or Points of Difficult Ascent:

impassable falls at 8km

fish ladder passable to coho at 457meters

SPAWNING DISTRIBUTION

Species

Section of Stream Used

chum	- intertidal spawning in estuary
coho	- past fishway to upper reaches

GENERAL REMARKS

- 1952 This long, slow-moving stream with its reliable water supply has good potential. The lower 2.4km is suitable for spawning. The stream bed is 15 - 20 ft. wide and has a gradient of 20' in 1000'. However, this section also contains some heavily silted pools caused by minor debris barriers.
- 1974 The fish ladder at the mouth of the stream enters a log dumping area. The decomposition of organic material and the constant dumping of logs inhibit good production.
- 1976 The lower 91m has been badly abused by logging company. Gibsons Wildlife Club adopted the stream for study and enhancement.
- 1977 Some course changes due to log jams. A stop log at the lowest logging bridge has stopped the lower stream erosion.
- 1978 Fish and Wildlife Club has been doing some stream improvements on Hudson Cr., a tributary of Wilson Cr. SEP may put in some jumping pools along side the fish ladder on the rock face of falls.
- 1979 Extreme erosion and silting from farmlands in upper reaches. Some stream clearance work. Proposed new fishway to allow chum access to upper reaches. Severe scouring from high floods in Dec. Poor survival expected.

ESCAPEMENT RECORD FOR WILSON CREEK

YEAR	SOCKEYE	CHINOOK	COHO	CHUM	PINK	STEELHEAD
1947			75			
48			750	750		25
49			750			25
50			400			
51			400			
52			75			UNK
53						
54		NO RECORDS FROM 1953 - 1969				
55						
56						
57						
58						
59						
60						
61						
62						
63						
64						
65						
66						
67						
68						
69						
70			N/O	N/O		N/O
71			N/O	N/O		N/O
72				60		
73			N/O	N/O		N/O
74			75	75		25
75			75	25		25
76			100	4		
77			50	300		
78			100	325		
79				300		
80			100	250		
81			12	100		
82			N/O	48		
83			75	400		
84			200	500		
85						

TIMING

ARRIVE			L SEPT-E OCT	E-L OCT		
START			M-L OCT	M OCT-E NOV		
PEAK			L OCT-M NOV	M-L NOV		
END			E NOV-E DEC	E DEC-E JAN		

REMARK

Metric Conversions

CONVERSION FACTORS

The following list of convenient equivalents of measure gives the relationship between imperial units and the International System of Units (SI).

1 inch equals 2.54 cm (centimetres)
1 foot equals 0.3048 m (metre)
1 statute mile equals 1.6093 km (kilometres)

1 cm (centimetre) equals 0.393 70 inch
1 m (metre) equals 3.2808 feet
1 km (kilometre) equals 0.621 37 mile

1 acre equals 43 560 square feet
1 acre equals 0.404 69 ha (hectare)
1 square mile equals 640 acres
1 square mile equals 2.5900 km² (square kilometres)
1 square mile equals 259.0 ha (hectares)

1 ha (hectare) equals 10 000 m² (square metres)
1 ha (hectare) equals 2.4710 acres
1 km² (square kilometre) equals 0.386 10 square mile

1 cubic foot equals 6.2288 imperial gallons
1 imperial gallon equals 4.546 09 L (litres)
1 imperial gallon equals 1.2010 U.S. gallons
1 U.S. gallon equals 0.133 68 cubic foot
1 cubic foot equals 0.068 317 m³ (cubic metre)
1 m³ (cubic metre) equals 35.315 cubic feet

1 cubic foot per second for one day equals 1.9835 acre-feet
1 cubic foot per second for one day covers one square mile to a depth of 0.037 19 inch
1 acre-foot equals 1.2335 dam³ (cubic decametres)
1 m³/s (cubic metre per second) for one day equals 86.4 dam³ (cubic decametre)
1 m³/s (cubic metre per second) for one day covers one square kilometre to a depth of 0.0864 m (metre)

1 foot per second equals 0.6818 mile per hour
1 mile per hour equals 1.467 feet per second
1 m/s (metre per second) equals 3.6 km/h (kilometre per hour)
1 km/h (kilometre per hour) equals 0.2778 m/s (metre per second)

1 cubic foot per second equals 0.028 317 m³/s (cubic metre per second)
1 m³/s (cubic metre per second) equals 35.315 cubic feet per second

1 pound equals 0.453 59 kg (kilogram)
1 kg (kilogram) equals 2.2046 pounds

1 short ton (2000 pounds) equals 0.907 18 t (tonne)
1 t (tonne) equals 2204.6 pounds

degrees Celsius = 5/9 (degrees Fahrenheit - 32)
degrees Fahrenheit = 9/5 (degrees Celsius) + 32

FACTEURS DE CONVERSION

Voici une liste des unités de mesure impériales et leurs équivalences dans le Système international d'unités (SI).

1 pouce vaut 2.54 cm (centimètres)
1 pied vaut 0.3048 m (mètre)
1 mille terrestre équivaut à 1.6093 km (kilomètre)

1 cm (centimètre) équivaut à 0.393 70 pouce
1 m (mètre) équivaut à 3.2808 pieds
1 km (kilomètre) équivaut à 0.621 37 mille

1 acre vaut 43 560 pieds carrés
1 acre équivaut à 0.404 69 ha (hectare)
1 mille carré vaut 640 acres
1 mille carré équivaut à 2.5900 km² (kilomètres carrés)
1 mille carré équivaut à 259.0 ha (hectare)

1 ha (hectare) vaut 10 000 m² (mètres carrés)
1 ha (hectare) équivaut à 2.4710 acres
1 km² (kilomètre carré) équivaut à 0.386 10 mille carré

1 pied cube équivaut à 6.2288 gallons impériaux
1 gallon impérial vaut 4.546 09 L (litres)
1 gallon impérial équivaut à 1.2010 gallon américain
1 gallon américain équivaut à 0.133 68 pied cube
1 pied cube vaut 0.068 317 m³ (mètre cube)
1 m³ (mètre cube) vaut 35.315 pieds cubes

1 pied cube par seconde pendant un jour équivaut à 1.9835 acre-pied
1 pied cube par seconde pendant un jour équivaut à un volume d'un mille carré par 0.037 19 pouce
1 acre-pied équivaut à 1.2335 dam³ (décamètres cubes)
1 m³/s (mètre cube par seconde) pour un jour vaut 86.4 dam³ (décamètre cube)
1 m³/s (mètre cube par seconde) pour un jour couvre un kilomètre carré à une profondeur de 0.0864 m (mètre)

1 pied par seconde équivaut à 0.6818 mille par heure
1 mille par heure équivaut à 1.467 pied par seconde
1 m/s (mètre par seconde) vaut 3.6 km/h (kilomètre par heure)
1 km/h (kilomètre par heure) vaut 0.2778 m/s (mètre par seconde)

1 pied cube par seconde équivaut à 0.028 317 m³/s (mètre cube par seconde)
1 m³/s (mètre cube par seconde) équivaut à 35.315 pieds cubes par seconde

1 livre équivaut à 0.453 59 kg (kilogramme)
1 kg (kilogramme) équivaut à 2.2046 livres

1 tonne courte (2000 livres) équivaut à 0.907 18 t (tonne)
1 t (tonne) équivaut à 2204.6 livres

degrés Celsius = 5/9 (degrés Fahrenheit - 32)
degrés Fahrenheit = 9/5 (degrés Celsius) + 32