

Canadian Data Report of
Fisheries and Aquatic Sciences 688

MARCH 1988

NATIVE HARVEST OF WILDLIFE
IN THE KEEWATIN REGION, NORTHWEST TERRITORIES
FOR THE PERIOD OCTOBER 1985 TO MARCH 1986 AND
A SUMMARY FOR THE ENTIRE PERIOD OF THE HARVEST
STUDY FROM OCTOBER 1981 TO MARCH 1986

by

R. L. Gamble¹

Central and Arctic Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

This is the 22nd Data Report
from the Central and Arctic Region, Winnipeg

¹ Present address 20 Amundsen Bay, Winnipeg, Manitoba R3K 0V2

Former address Keewatin Wildlife Federation, Rankin Inlet, N.W.T. X0C 0G0

PREFACE

This report is a supplement to previous technical reports by Gamble (1984, 1987a, 1987b) which cover the period October 1981 to September 1985 and covers the period October 1985 to March 1986. It is presented in fulfillment of Department of Supply and Services Contract DSS 25 S.T.A. 7135-05-0003 let to the Keewatin Wildlife Federation to conduct a wildlife harvest study in the Keewatin Region - Phase III. The work was done on behalf of the Federal Government departments of Environment Canada (Canadian Wildlife Service), Fisheries and Oceans (Central and Arctic Region), and Indian Affairs and Northern Development; the Government of the Northwest Territories Department of Renewable Resources; and the Keewatin Wildlife Federation.

The report is accepted upon recommendation by the steering committee for the study made up of representatives of the agencies noted above (Appendix 3). The harvest study material is published under the auspices of the DFO report series by agreement of the steering committee in order to ensure that the data achieve a wide circulation, be accessible to the interested public, and be published in a standardized format generally recognized as appropriate for the dissemination of such information.

A report of the study in Inuktitut will also be published under the auspices of Indian Affairs and Northern Development.

© Minister of Supply and Services Canada 1988

Cat. no. Fs 97-13/688E ISSN 0706-6465

Correct citation for this publication is:

Gamble, R.L. 1988. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1985 to March 1986 and a summary for the entire period of the harvest study from October 1981 to March 1986. Can. Data Rep. Fish. Aquat. Sci. 688: v + 85 p.

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT/RÉSUMÉ	v
INTRODUCTION	1
MATERIALS AND METHODS	1
General	1
Human resources	1
Materials	1
Data collection and analysis	1
Data processing	2
ACKNOWLEDGMENTS	2
REFERENCES	2

PART I. NATIVE HARVEST FOR THE PERIOD OCTOBER 1985 TO MARCH 1986.

LIST OF TABLES

<u>Table</u>		<u>Page</u>
1	The reported harvest by Baker Lake hunters, for the period October 1985 to March 1986	4
2	The reported harvest by Chesterfield Inlet hunters, for the period October 1985 to March 1986	4
3	The reported harvest by Coral Harbour hunters, for the period October 1985 to March 1986	5
4	The reported harvest by Eskimo Point hunters, for the period October 1985 to March 1986	5
5	The reported harvest by Rankin Inlet hunters, for the period October 1985 to March 1986	6
6	The reported harvest by Repulse Bay hunters, for the period October 1985 to March 1986	6
7	The reported harvest by Whale Cove hunters, for the period October 1985 to March 1986	7
8	The estimated harvest by Baker Lake hunters, for the period October 1985 to March 1986	7
9	The estimated harvest by Chesterfield Inlet hunters, for the period October 1985 to March 1986	8
10	The estimated harvest by Coral Harbour hunters, for the period October 1985 to March 1986	8
11	The estimated harvest by Eskimo Point hunters, for the period October 1985 to March 1986	9
12	The estimated harvest by Rankin Inlet hunters, for the period October 1985 to March 1986	9
13	The estimated harvest by Repulse Bay hunters, for the period October 1985 to March 1986	10

Table

	<u>Page</u>
14 The estimated harvest by Whale Cove hunters, for the period October 1985 to March 1986	10
15 The reported and estimated harvest for Baker Lake hunters	11
16 The reported and estimated harvest for Chesterfield Inlet hunters	11
17 The reported and estimated harvest for Coral Harbour hunters	12
18 The reported and estimated harvest for Eskimo Point hunters	12
19 The reported and estimated harvest for Rankin Inlet hunters	13
20 The reported and estimated harvest for Repulse Bay hunters	13
21 The reported and estimated harvest for Whale Cove hunters	14
22 Monthly theoretical kill factors for seven Keewatin communities	14
23 The harvest by species over the range of age for Baker Lake hunters covering the period October 1985 to March 1986	14
24 The harvest by species over the range of age for Chesterfield Inlet hunters covering the period October 1985 to March 1986	15
25 The harvest by species over the range of age for Coral Harbour hunters covering the period October 1985 to March 1986	15
26 The harvest by species over the range of age for Eskimo Point hunters covering the period October 1985 to March 1986	16
27 The harvest by species over the range of age for Rankin Inlet hunters covering the period October 1985 to March 1986	16
28 The harvest by species over the range of age for Repulse Bay hunters covering the period October 1985 to March 1986	17
29 The harvest by species over the range of age for Whale Cove hunters covering the period October 1985 to March 1986	17
30 Data on the distribution of hunters that were successful in obtaining a harvest, for the period October 1985 to March 1986	18
31 Edible weight values in kilograms for harvested species as calculated from various sources	19

<u>Table</u>	<u>Page</u>	<u>Table</u>	<u>Page</u>
32 Reported and estimated edible weight values for harvested species for the period October 1985 to March 1986	20	46 The estimated harvest of Arctic fox by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	72
33 Reported and estimated edible weight values for four major groups of animals harvested by Keewatin communities, October 1985 to March 1986	22	47 The estimated harvest of wolves by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	73
34 Age distribution of hunters for the seven Keewatin region communities for the period October 1985 to March 1986.	26	48 The estimated harvest of ringed seals by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	74
PART II. SUMMARY OF NATIVE HARVEST FOR THE PERIOD OCTOBER 1981 TO MARCH 1986 (APPENDIX 4 GIVES FOOTNOTES).		49 The estimated harvest of bearded seals by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove, for the period October 1981 to March 1986	75
LIST OF TABLES			
<u>Table</u>	<u>Page</u>	<u>Table</u>	<u>Page</u>
35 The estimated harvest by Baker Lake hunters, for the period November 1981 to March 1986	27	50 The estimated harvest of beluga whales by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove, for the period October 1981 to March 1986	76
36 The estimated harvest by Chesterfield Inlet hunters, for the period January, February and August 1982 through March 1986	33	51 The estimated harvest of Canada geese by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	77
37 The estimated harvest by Coral Harbour hunters, for the period February 1982 and June 1982 through April 1983 and October 1983 to March 1986	37	52 The estimated harvest of Snow geese by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	78
38 The estimated harvest by Eskimo Point hunters, for the period October 1981 to March 1986	44	53 The estimated harvest of charr by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	79
39 The estimated harvest by Rankin Inlet hunters, for the period November 1981 to March 1986	51	54 The estimated harvest of lake trout by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	80
40 The estimated harvest by Repulse Bay hunters, for the period October 1981 to March 1986	57		
41 The estimated harvest by Whale Cove hunters, for the period October 1981 to March 1983 and November 1983 to March 1986	53		
42 The estimated harvest of Kaminuriak caribou by hunters from the seven Keewatin communities, for the period October 1981 to March 1986	68		
43 The estimated harvest of Beverly caribou by Baker Lake and Repulse Bay hunters, for the period October 1981 to March 1986	69		
44 The estimated harvest of Wager Bay caribou by Baker Lake, Coral Harbour, Eskimo Point, Repulse Bay and Whale Cove hunters, for the period October 1981 to March 1986	70		
45 The estimated harvest of polar bears by hunters from the seven Keewatin communities for the period October 1981 to March 1986	71		
LIST OF FIGURES			
<u>Figure</u>	<u>Page</u>		
1 Map of Keewatin District showing the seven communities surveyed during the harvest study and the zonal grid used to locate kills	81		
LIST OF APPENDICES			
<u>Appendix</u>	<u>Page</u>		
1 Monthly percent of hunters interviewed for the seven Keewatin communities for the period October 1981 to March 1986	82		
2 Monthly theoretical kill factors for the seven Keewatin communities for the period October 1981 to March 1986	83		
3 Members of the Steering Committee for the period of the Keewatin Wildlife Federation Harvest Study, 1981 to 1987	84		
4 Footnotes for Tables 35 to 41	85		

ABSTRACT

Gamble, R.L. 1988. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1985 to March 1986 and a summary for the entire period of the harvest study from October 1981 to March 1986. Can. Data Rep. Fish. Aquat. Sci. 688: v + 85 p.

Harvest data have been collected from Inuit hunters of the Keewatin Region from September 1981 until March 1986. The project has been run by an Inuit organization, the Keewatin Wildlife Federation, supported by funding provided through interested federal and territorial government departments. Part I (Tables 1 to 34) of this report covers the period October 1985 to March 1986 and is an update and supplement to the previous reports (Technical Reports 1282, 1543 and 1544) which cover the earlier years of the survey. Results in Part I are presented in the same format as the previous noted. Data were aggregated at a community level and field-workers continued to maintain a high level of performance as measured by participation of hunters in the study and a subjective judgement of the quality of data based on experience. Part II of this report (Tables 35 to 54) is a summary of the entire period of the harvest study. Tables 35 to 41 give the estimated harvest of species by community for the years of survey and the remaining tables give the harvest of selected species aggregated at a regional level.

Key words: resource management; catch statistics; domestic harvest; monitoring; food resources; country foods; terrestrial mammals; marine mammals; birds; fish; computerized harvest study; Inuit organization.

RÉSUMÉ

Gamble, R.L. 1988. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1985 to March 1986 and a summary for the entire period of the harvest study from October 1981 to March 1986. Can. Data Rep. Fish. Aquat. Sci. 688: v + 85 p.

Des données sur les prises/captures ont été recueillies auprès de chasseurs inuit de la région du Keewatin pour la période de septembre 1981 à mars 1986, dans le cadre d'un programme de collecte dont un organisme inuit, la Keewatin Wildlife Federation, assure l'application. Le financement pour le projet vient des ministères fédéral et territorial en cause. La Partie I (tableaux 1 à 34) du rapport porte sur la période d'octobre 1985 à mars 1986 et constitue une mise à jour et un complément aux rapports précédents du même auteur (rapports techniques 1282, 1543 et 1544) sur les premières années de l'étude. Les résultats dans cette Partie sont présentés comme dans les rapports précédents. Les données ont été groupées par collectivité et le travail de collecte a été fait de façon

excellente, comme l'indiquent la participation des chasseurs à l'étude et l'évaluation subjective des données, fondée sur l'expérience. La Partie II du rapport (tableaux 35 à 54) constitue un résumé de toute la période visée par l'étude. Les tableaux 35 à 41 présentent les prises/captures estimatives des espèces par collectivité pour la durée de l'étude tandis que les tableaux 42 à 54 renferment les chiffres des prises/captures de certaines espèces regroupées par région.

Mots-clés: gestion des ressources; statistiques des prises; chasse/pêche de subsistance; contrôle; ressources alimentaires; ressources alimentaires indigènes; mammifères terrestres; mammifères marins; oiseaux; poisson; étude des prises/captures par ordinateur; organisme inuit.

INTRODUCTION

The collection of data for the "Keewatin Region Native Harvest of Wildlife Study" began in September, 1981. Previous results have been published for the following periods: October 1981 to September 1983 (Gamble 1984); October 1983 to September 1984 (Gamble 1987a); and October 1984 to September 1985 (Gamble 1987b). Part I (Tables 1 to 34) of this report supplements and updates these previous reports for the survey period October 1985 to March 1986. Part II of this report is a summary of the entire period of the harvest study from September 1981 to March 1986. In Part II, Tables 35 to 41 give the estimated harvest of species by community for the years of survey, and Tables 42 to 54 give the harvest of selected species aggregated at a regional level. Appendix 4 gives footnotes pertinent to the tables in Part II.

Throughout this report hunter, harvester, trapper and fisherman are used as synonyms. Hunter is defined in the MATERIALS AND METHODS section below.

The main objectives of the study were to:

- 1) determine by survey techniques the hunter kill (i.e. harvest) by Inuit living in District of Keewatin communities and outpost camps;
- 2) develop an approach for the collection of timely, statistically reliable data on wildlife harvesting which could be undertaken by an agency such as the Keewatin Wildlife Federation (KWF) upon completion of the preliminary study;
- 3) determine the number of Inuit directly participating in subsistence harvesting in each community and to compare the proportion of harvest taken by hunters of different ages;
- 4) provide an estimate of the harvest sufficient to determine a measure of its value to each community as food or income, and
- 5) analyze and publish the data collected in a timely report and scientifically acceptable format.

The study area (Fig. 1) remained the same for the entire period of the study as reported in Gamble (1984; 1987a and b) and includes the entire Keewatin district of the Northwest Territories (approximately 386,000 km²). This region contains seven permanent communities. Listed alphabetically (the convention followed throughout this and previous reports) they are Baker Lake, Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove. Current information about these communities including population can be obtained from the NWT Data Book (1986).

MATERIALS AND METHODS

GENERAL

For the entire period of survey fieldworkers tried to include 100% of the region's hunters in their monthly data collection. The study design remained the same as originally described in Gamble (1984).

For the purpose of the study the term hunter includes all Inuit males and females over the age of 16 who hunt (they may or may not have a NWT general hunting licence), Inuit youths under 16 who hunt regularly, and some long-term residents in the area of other ethnic origin who hunt. This latter group makes up less than 1% of the total hunters in the region and also accounts for less than 1% of the animals harvested.

Harvest data were aggregated at the community level. Separate coverage of outpost camps was not necessary because Inuit hunting from such locations visited their home communities frequently during the survey period and it was possible to include their harvest together with that of community based hunters on a consistent basis.

HUMAN RESOURCES AND MATERIALS

Fieldworkers were hired in each of the seven communities to interview hunters and collect data. Duties included explaining the project to hunters; distributing the study materials (calendars and field notebooks) to hunters; keeping an up to date list of hunters; interviewing hunters beginning on the first day of each month to collect harvest statistics for the previous month and recording this information on the appropriate data sheets; making sure the data collected were as accurate as possible; and promptly forwarding a monthly report following an interview period to the Project Biologist located at Rankin Inlet.

Over the period of the survey slight changes were made to data sheets, calendars and field diaries that were distributed to fieldworkers and hunters. The project office organization was also modified. These modifications are described in Gamble 1987a and 1987b.

During the period October 1985 to March 1986 the Project Office organization remained the same as described by Gamble (1987a) and no changes were made to the data sheets, calendars and field diaries that were distributed to fieldworkers and hunters as described in Gamble (1987b).

DATA COLLECTION AND ANALYSIS

The system used to analyze harvest data and to arrive at estimates of the total hunter kill by community required several steps and remained the same as developed during the 1981-1983 preliminary study (Gamble 1984).

Beginning on the first day of each month fieldworkers began interviews so that they could divide the hunter population for each community into the survey categories defined below and list the number of animals killed per species for successful hunters who were interviewed. The monthly interval was defined as an interview period and covered the previous month of hunting. The fieldworker submitted this information to the Project Office where the data were summarized each month against a master list of hunters for individual communities and then entered into the computer. The numbers in some categories were subsequently adjusted the following month (i.e. the second month past the actual hunting episode) if acceptable reports were submitted by fieldworkers on hunters who had been interviewed after a particular interview period had passed. Acceptable reports were determined through a subjective judgement by the Project Biologist based on his experience and a comparison of late reports with the reports submitted on time.

	<u>Definition</u>	<u>Category</u>
1)	The number of hunters who report taking a harvest during an interview period (i.e. successful).	A
2)	The number of hunters who report they were not successful in taking a harvest during an interview period (i.e. unsuccessful).	B
3)	The number of hunters who report they did not hunt during an interview period (i.e. didn't hunt).	C
4)	The number of hunters who were out hunting during the interview period but who were not interviewed (i.e. hunted but not interviewed).	D
5)	The number of hunters who were out of the area of the harvest survey during the interview period for any reason (i.e. out of hunt area).	E
6)	The number of hunters within the harvest study area during the interview period whose activities were unknown (i.e. activities unknown).	F

Subsequently the summarized monthly information contained in categories A through F was used to calculate ratios of participation and hunter success. Participation ratio refers to the percent of hunters in each community who were interviewed as part of the study in relation to the total number of hunters who could have hunted each month. Appendix 1 summarizes the monthly percent of hunters interviewed for the seven Keewatin communities for the period October 1981 to March 1986.

The hunter success ratio was applied to hunters in categories D and F to obtain an estimate of probable hunter success within these groups. The results for all categories were summed to get an estimate of total hunter success and to calculate the theoretical kill factor. This is the value by which the reported kill per species is multiplied to arrive at the estimated harvest. The theoretical kill factors calculated for each month for the seven Keewatin communities for the period October 1981 to March 1986 are given in Appendix 2.

For the purpose of this analysis four main assumptions were made:

- 1) The involvement of hunters in the harvest is the same for those whose activities are unknown as for those that are known.
- 2) The success ratio is the same for hunters who hunted in the unknown categories as for the known categories.
- 3) The probability of a kill of any individual animal is the same for all species when calculating the estimated harvest.
- 4) Reported kills are accurate.

DATA PROCESSING

The study used the basic computer programs described by Gamble (1984; 1987a and b) throughout the period of survey. The processing of harvest data was enhanced after the preliminary survey described by Gamble 1984 by the addition of programmes which allowed the species harvested to be reflected by a geographic zone system (Fig. 1) and a breakdown of the reported kill by species over the range of age groups for the hunters.

The hunter age classes used for the survey were 0-15, 16-30, 31-45, 46-60, 61-75 and 76-99 years. The age group 76-99 was used as a category for hunters with unknown ages because only 8 hunters of known age fell within this group.

ACKNOWLEDGMENTS

I thank the Keewatin Wildlife Federation, which was supportive of this study and provided valuable assistance at a regional and community level.

Thanks are also due to members of the steering committee who provided valuable criticism of my manuscript, in particular R. Peet and B. Hyman of DFO who assisted in the preparation and publication of this and the previous reports (Gamble 1984, 1987a and 1987b) produced for this study. I particularly thank the staff of the Keewatin Wildlife Federation "Native Harvest Study". Their continued efforts in conjunction with the various community fieldworkers allowed this study to be successfully completed.

I especially thank all the hunters who provided data on their harvests and acknowledge their cooperation and understanding in the face of repeated questions that were necessary.

REFERENCES

- BELLROSE, F.C. 1976. Ducks, geese, and swans of North America. Stackpole Books. 540 p.
- BERGER, T. 1977. Northern frontier - northern homeland. The report of the Mackenzie Valley Pipeline Inquiry, 2: 268 p.

- BOND, W.A. 1975. Data on the biology of lake whitefish and lake trout from Kaminuriak Lake, District of Keewatin, N.W.T. Can. Fish. Mar. Serv. Data Rep. Serv. CEN/D-75-4: 28 p.
- CARDER, G.W. 1983. Data from the commercial fishery for Arctic charr, Salvelinus alpinus (Linnaeus), in the Cambridge Bay and Rankin Inlet areas, Northwest Territories, 1981-82. Can. Data Rep. Fish Aquat. Sci. 391: v + 24 p.
- DOME PETROLEUM LTD., ESSO RESOURCES CANADA, LTD., AND GULF CANADA RESOURCES INC. 1982. Beaufort Sea-Mackenzie Delta Environmental Impact Statement. Socio-economic effects 5(5): 42 p.
- EARHART, C.M., and N.K. JOHNSON. 1970. Size dimorphism and food habits of North American owl. Condor 72: 251-264.
- FALK, M.R., and D.V. GILLMAN. 1975. Data on the lake and round whitefish, lake cisco, northern pike, Arctic grayling and longnose sucker from the east arm of Great Slave Lake, N.W.T., 1971-74. Can. Fish. Mar. Serv. Data Rep. Ser. CEN/D-75-2: 95 p.
- GAMBLE, R.L. 1984. A preliminary study of the native harvest of wildlife in the Keewatin Region, Northwest Territories, Canada. Can. Tech. Rep. Fish. Aquat. Sci. 1282: iv + 48 p.
- GAMBLE, R.L. 1987a. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1983 to September 1984. Can. Tech. Rep. Fish. Aquat. Sci. 1543: v + 82 p.
- GAMBLE, R.L. 1987b. Native harvest of wildlife in the Keewatin Region, Northwest Territories for the period October 1984 to September 1985. Can. Tech. Rep. Fish. Aquat. Sci. 1549: v + 59 p.
- KELEHER, J.J. 1964. Round weight conversion factors for Great Slave Lake fish. Fish. Res. Board Can. Manuscr. Rep. Ser. (Biol.) 773: 19 p.
- MACDONALD, G., and R. FUDGE. 1979. Arctic Land Use Research Program 1978: A survey of the fisheries resources of the Kazan Upland (Southeastern District of Mackenzie, Southern District of Keewatin, N.W.T.). Can. Dep. Ind. N. Aff. Environ. Stud. 11: 161 p.
- NATIVE HARVESTING RESEARCH COMMITTEE (NHRC). 1975. Research to establish present levels of harvesting by native peoples of northern Quebec. Phase 1 (1973-1975). Part II. Montreal. 230 p.
- NATIVE HARVESTING RESEARCH COMMITTEE (NHRC). 1976a. Research to establish present levels of harvesting by native peoples of northern Quebec. Part II. A report on the harvesting by the James Bay Cree.
- NATIVE HARVESTING RESEARCH COMMITTEE (NHRC). 1976b. Research to establish present levels of harvesting by native peoples of northern Quebec. Part II. A report on the harvesting by the Inuit of northern Quebec.
- NWT DATA BOOK. 1986. A complete information guide to the Northwest Territories and its communities. M. Devine (ed.) Outcrop Ltd., The Northern Publishers, Yellowknife. 236 p.
- RIEWE, R. 1977. Utilization of wildlife in the Jones Sound region by the Griese Fiord Inuit, p. 623-644. In L.C. Bliss (ed.) True Love Lowlands, Devon Island, Canada: a high Arctic ecosystem. University of Alberta Press, Edmonton.
- SERGEANT, D.E., and P.S. BRODIE. 1969. Body size in white whales, Delphinapterus leucas. J. Fish. Res. Board Can. 26: 2561-2580.
- STEVENS, W.J.D. 1965. Bionomics of the sandhill crane. Ph.D. Thesis. University of Saskatchewan, Saskatoon. 120 p.
- THOMAS, V.G. 1982. Energetic reserves of Hudson Bay willow ptarmigan during the winter and spring. Can. J. Zool. 60: 1618-1623.

Table 1. The reported harvest by Baker Lake hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986			Species	Category ¹	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.			Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
<u>Caribou</u>															
Kaminuriak	M	89	72	10	28	199	199	Kaminuriak	M	8	3	8	8	3	8
	F	74	56	44	48	222	222		F						
	C	8	24	24	5	61	61		U	10	10	10	10	10	10
	U	3	3	3	3	3	3		Subtotal	11	10	10	10	10	10
Subtotal		171	155	78	81	485	485	North of Chesterfield	M	16	15	20	31	23	105
Beverly	M	86	54	8	140	140	140	F	8	5	3	14	4	34	34
	F	232	54	30	286	286	286	U	9	8	4	8	8	21	21
	C	12	18	30	30	30	30	Subtotal	33	20	23	49	35	160	160
	U	11	11	11	11	467	467	Total	11	43	23	49	35	181	181
Subtotal		341	126												
Wager	M	96	14	108	149	178	150	Muskox		1	2		2	2	2
	F	80	10	143	141	157	202	Polar Bear							3
	C	30	9	36	59	81	65	Arctic Fox							30
	U	1	2	35	349	424	418	Red Fox							1
Subtotal		207	287					Wolf							14
Other	F	Subtotal	719	316	365	349	424	Ringed Seal							14
	Total							Bearded Seal							14
Muskox								Walrus							14
Arctic Fox								Eider							14
Wolf								Arctic Charr							14
Arctic Charr								Lake Trout sp.							14
Lake Trout								Arctic Grayling							14
Whitefish sp.								Percent of hunters reporting							14
Percent of hunters reporting		95.4	98.0	94.8	100.0	100.0	100.0	Percent of hunters reporting		100.0	100.0	100.0	100.0	92.0	91.8

¹Categories are as follows: M means male, F means female, C means calf, and U means unknown.

¹See Table 1.

Table 3. The reported harvest by Coral Harbour hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
<u>Caribou</u>							
Southampton	M	1	1	4	6		
	C	1	1	1	1		
	U	3	1	4	8		
Total		7	27	2	1	2	41
Polar Bear		1	13	44	13	2	33
Arctic Fox		1	11	9	5	5	38
Wolf		49	68	47	81	75	73
Arctic Hare		1	2	3	5	5	5
Ringed Seal		1	1	1	5	5	11
Bearded Seal		28	205	282	839	284	222
Harp Seal		216	1009	116	27	36	21
Walrus							
Eider							
Ptarmigan							
Arctic Charr							
Arctic Cod							
Percent of hunters reporting		73.3	85.7	80.0	85.0	84.8	81.0

¹See Table 1.

Table 4. The reported harvest by Eskimo Point hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
<u>Caribou</u>							
Kaminiurjik	M	112	29	26	22	55	75
	F	279	61	53	79	36	219
	C	59	11	8	8	4	727
	U	18	2	2	8	4	35
Total		468	103	87	109	95	323
Moose					1	1	2
Muskox						2	1
Polar Bear						1	1
Arctic Fox						2	9
Red Fox						35	509
Wolf						81	33
Arctic Hare						5	1
Otter						1	8
Arctic Ground Squirrel						1	61
Ringed Seal						44	61
Bearded Seal						1	1
Eider						1	1
Ptarmigan						1	1
Arctic Charr						1	1
Lake Trout						33	39
Whitefish sp.						51	2
Arctic Grayling						20	20
Percent of hunters reporting		87.5	91.9	91.8	94.1	85.8	89.8

¹See Table 1.

Table 5. The reported harvest by Rankin Inlet hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986			Species	Category ¹	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.			Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
Caribou								Polar Bear							
Kamnuritak	M	52	24	21	30	33	30	Wager	M	38	38	31	75	31	249
	F	64	92	28	49	59	32		F	19	19	12	38	56	192
	C	3	5		6	10	6		C	2	3	1	2	3	10
	U			4	3	1	1		U				4	4	4
	Subtotal	119	121	53	88	103	69	Total	59	60	44	86	116	101	466
North of Chesterfield	M							Polar Bear							
	F							Arctic Fox							
	C							Red Fox							
	U							Wolf							
	Subtotal							Wolverine							
	Total	119	121	53	88	103	128	Muskox							
Polar Bear								Polar Bear							
Arctic Fox								Arctic Hare							
Wolf								Ringed Seal							
Arctic Hare								Bearded Seal							
Ringed Seal								Walrus							
Bearded Seal								Narwhal							
Harbour Seal								Ptarmigan							
Harp Seal								Arctic Charr							
Walrus								Lake Trout							
Eider								Percent of hunters reporting							
Ptarmigan								71.4	83.3	97.7	90.3	95.2	100.0		
Searun Arctic Charr		385	586	290	297	104	99								
Landlocked Arctic Charr		30		2		15									
Lake Trout															
Percent of hunters reporting		95.0	98.1	97.5	97.9	96.8	97.2								

¹See Table 1.

Table 6. The reported harvest by Repulse Bay hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986			Species	Category ¹	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.			Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
Caribou								Wager	M	38	38	31	36	75	31
	F								F	19	19	12	48	38	56
	C								C	2	3	1	2	3	10
	U								U				4	4	4
	Subtotal	119	121	53	88	103	128	Total	59	60	44	86	116	101	466
North of Chesterfield	M							Polar Bear							
	F							Arctic Fox							
	C							Red Fox							
	U							Wolf							
	Subtotal							Wolverine							
	Total	119	121	53	88	103	128	Muskox							
Polar Bear								Polar Bear							
Arctic Fox								Arctic Hare							
Wolf								Ringed Seal							
Arctic Hare								Bearded Seal							
Ringed Seal								Walrus							
Bearded Seal								Narwhal							
Harbour Seal								Ptarmigan							
Harp Seal								Arctic Charr							
Walrus								Lake Trout							
Eider								Percent of hunters reporting							
Ptarmigan								71.4	83.3	97.7	90.3	95.2	100.0		
Searun Arctic Charr		385	586	290	297	104	99								
Landlocked Arctic Charr		30		2		15									
Lake Trout															
Percent of hunters reporting		95.0	98.1	97.5	97.9	96.8	97.2								

¹See Table 1.

Table 7. The reported harvest by Whale Cove hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category 1	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
Caribou							
Kaminiurak	M	30	14	10	20	6	87
	F	32	29	33	73	43	251
	C			2	9	4	19
	U				3	1	4
Total		62	43	45	105	53	361
Muskox		10	2	2	1	10	4
Polar Bear		1	1			1	
Arctic Fox					2	3	4
Red Fox						20	3
Arctic Hare		10	1	6	7	4	31
Ringed Seal		10	9	6	40	9	22
Arctic Charr		3			10	9	
Lake Trout							
Percent of hunters reporting		100.0	100.0	100.0	98.3	100.0	100.0

See Table 1.

Table 8. The estimated harvest by Baker Lake hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category 1	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
Caribou							
Kaminiurak	M				89	72	10
	F				74	56	44
	C				8	24	24
	U					3	
Subtotal					171	155	78
Beverly	M				86	54	
	F				232	54	
	C				12	18	
Subtotal					341	126	
Wager	M				96	14	108
	F				80	10	143
	C				30	9	36
Subtotal					207	35	287
Other	F						
Subtotal					719	316	365
Muskox						349	424
Arctic Fox						65	152
Wolf						3	5
Arctic Charr						7	
Lake Trout						520	469
Whitefish sp.						70	65
						35	14
						14	14
						317	261
						7	7
							191
						3	3
						502	2675

See Table 1.

Table 9. The estimated harvest by Chesterfield Inlet hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986			Sum.
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	
<u>Caribou</u>								
Kaminiuriak	M	8						
	F	3						
	U		10					
	Subtotal	11	10					21
North of Chesterfield	M	16	15	20	31	23	105	
	F	8	5	3	14	4	34	
	U	9			4	8	21	
	Subtotal	33	20	23	49	35	160	
	Total	43	20	23	49	35	181	
Muskox								
Polar Bear	1	2						
Arctic Fox		26		1				
Red Fox			7					
Wolf								
Ringed Seal	14	10	9	3	2	14	53	
Bearded Seal	1				2	3	4	
Walrus					1	2	3	
Eider					4		4	
Arctic Charr						1		
Lake Trout							58	
Arctic Grayling							14	

Table 10. The estimated harvest by Coral Harbour hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986			Sum.
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	
<u>Caribou</u>								
Southampton	M							
	C							
	U							
	Total							
Polar Bear					11	35	3	
Arctic Fox					17	69	2	
Wolf					2			
Arctic Hare					14	14	5	
Ringed Seal					79	86	5	
Bearded Seal					2			
Harp Seal					3	4		
Walrus					2			
Eider					18			
Parmigan					45	260	443	
Arctic Charr					350	1281	182	
Arctic Cod								

¹See Table 1.

¹See Table 1.

Table 11. The estimated harvest by Eskimo Point hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
Caribou							
Kaminiutiaq	M	120	35	25	81	90	383
	F	298	74	90	54	264	846
	C	63	13	9	6	42	143
	U	19	2		4	25	
Total		500	124	108	124	141	400
Moose				1	2	2	3
Muskox					3	11	
Polar Bear					40	639	
Arctic Fox		8	255	176	2	7	1
Red Fox					6	2	53
Wolf		4	10				75
Arctic Hare					1	1	
Otter				1		1	
Arctic Ground Squirrel					1	1	
Ringed Seal		16	5	1	12	13	47
Rearded Seal		2				2	
Eider						2	
Ptarmigan		59	119	93	37	76	24
Arctic Charr		75	59	32	2		100
Lake Trout			68	52		4	308
Whitefish sp.						70	242
Arctic Grayling		8				4	120
						12	

¹See Table 1.

Table 12. The estimated harvest by Rankin Inlet hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986		
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
Caribou							
Kaminiutiaq	M	55	25	23	32	36	33
	F	67	97	31	52	65	35
	C	3	5		6	11	7
	U				4	3	1
Subtotal		125	127	58	93	113	592
North of Chesterfield							
Kaminiutiaq	M						
	F						
	C						
Subtotal							
Total		125	127	58	93	113	592
Muskox							
Polar Bear		10				2	2
Arctic Fox		52	119		70	22	10
Wolf					2	4	272
Arctic Hare		2			2	1	7
Ringed Seal		66	25	2	3	2	6
Bearded Seal					9	14	119
Harbour Seal		6				1	9
Harp Seal		2					
Walrus		1					
Eider		8				2	2
Ptarmigan		4					8
Searun Arctic Charr		404	615	316	315	114	4
Landlocked Arctic Charr		32		2	16		32
Lake Trout							18

¹See Table 1.

Table 13. The estimated harvest by Repulse Bay hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986			Sum.
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	
<u>Caribou</u>								
Wager	M	51	44	31	44	83	31	284
	F	25	22	12	59	42	56	216
	C	3	4	1	2	3	10	23
	U						4	4
Total		79	70	44	105	128	101	527
Polar Bear		8		2	4	4	14	
Arctic Fox		2			1	5	8	
Red Fox		4	6		1	1	11	
Wolf		6	8	4	13	5	36	
Wolverine		1					1	
Arctic Hare		62	22	6	3	21	12	126
Ringed Seal		3		1			1	5
Bearded Seal		4					4	
Walrus		13	5		3		5	
Narwhal		27	639		1		16	
Ptarmigan		16					4	21
Arctic Charr								
Lake Trout								

¹See Table 1.

Table 14. The estimated harvest by Whale Cove hunters, expressed as numbers of animals, for the period October 1985 to March 1986.

Species	Category ¹	1985			1986			Sum.
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	
<u>Caribou</u>								
Kaminuriak	M	30	14	10	20	6	7	87
	F	32	29	33	73	43	41	251
	C			2	9	4	4	19
	U				3	3	1	4
Total		62	43	45	105	53	53	361
Muskox		10				2	2	10
Polar Bear								4
Arctic Fox								1
Red Fox								
Arctic Hare							2	4
Ringed Seal						7	3	31
Arctic Charr						40	20	88
Lake Trout						10	9	22

¹See Table 1.

Table 15. The reported and estimated harvest for Baker Lake hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category ¹	Reported Harvest ² Oct. 1985 - March 1986			Estimated Harvest ² Oct. 1985 - March 1986		
		Total	Mean	S.D.	Total	Mean	S.D.
<u>Caribou</u>							
Kaminuriak	M	199	3	2	199	3	2
	F	222	2	1	222	2	1
	C	61	1	1	61	1	1
	U	3	0	3	3	0	3
	Subtotal	485	2	2	485	2	2
Beverly	M	140	3	1	140	3	1
	F	286	4	2	286	4	2
	C	30	2	1	30	2	1
	U	11	0	11	11	0	11
	Subtotal	467	3	2	467	3	2
Wager	M	695	2	1	695	2	1
	F	733	2	1	733	2	1
	C	280	1	1	280	1	1
	U	12	2	2	12	2	2
	Subtotal	1720	2	1	1720	2	1
Other	F	3	3	0	3	3	0
	Subtotal	3	3	0	3	3	0
	Total	2675	2	2	2675	2	2
Muskox	13	1	0	13	1	0	
Arctic Fox	1039	6	4	1039	6	4	
Wolf	42	3	2	42	3	2	
Arctic Charr	11	6	2	11	6	2	
Lake Trout	2615	37	11	2615	37	11	
Whitefish sp.	191	7	3	191	7	3	

¹See Table 1.

²See also Tables 1 and 8.

Table 16. The reported and estimated harvest for Chesterfield Inlet hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category ¹	Reported Harvest ² Oct. 1985 - March 1986			Estimated Harvest ² Oct. 1985 - March 1986		
		Total	Mean	S.D.	Total	Mean	S.D.
<u>Caribou</u>							
Kaminuriak	M	8	3	2	3	1	8
	F	3	1	1	2	1	2
	C	10	5	5	10	5	5
	U	21	3	2	21	3	2
	Subtotal						
North of Chesterfield	M	105	3	1	105	3	1
	F	34	2	2	34	2	2
	C	21	4	2	21	4	2
	U	160	3	1	160	3	1
	Subtotal						
Muskox	M	181	3	1	181	3	1
Polar Bear		2	1	0	2	1	0
Arctic Fox		3	1	0	3	1	0
Red Fox		30	4	1	30	4	1
Wolf		1	1	0	1	1	0
Ringed Seal		14	2	1	14	2	1
Bearded Seal		53	3	2	53	3	2
Walrus		4	1	0	4	1	0
Eider		3	1	0	4	1	0
Arctic Charr		4	1	0	4	1	0
Lake Trout		83	15	4	83	15	4
Arctic Grayling		58	19	7	58	19	7
		14	7	3	14	7	3

¹See Table 1.

²See also Tables 2 and 9.

Table 17. The reported and estimated harvest for Coral Harbour hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category ¹	Reported Harvest ² Oct. 1985 - March 1986			Estimated Harvest ² Oct. 1985 - March 1986		
		Total	Mean	S.D.	Total	Mean	S.D.
<u>Caribou</u>							
Southampton	M	6	1	0	6	1	0
	C	1	1	0	1	1	0
	U	8	1	0	8	1	0
	Total	41	1	0	55	1	0
Polar Bear		105	5	6	136	6	8
Arctic Fox		1	1	0	2	2	0
Wolf		38	2	2	47	3	3
Arctic Hare		393	3	3	481	4	4
Ringed Seal		11	1	0	12	1	0
Bearded Seal		5	2	1	7	2	1
Harp Seal		3	1	0	4	1	0
Walrus		11	6	5	18	9	7
Eider		1860	17	16	2137	19	19
Parmigan		1425	41	37	1903	54	47
Arctic Charr		7	0	8	8	0	0
Arctic Cod							

¹See Table 1.

²See also Tables 3 and 10.

Table 18. The reported and estimated harvest for Eskimo Point hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category ¹	Reported Harvest ² Oct. 1985 - March 1986			Estimated Harvest ² Oct. 1985 - March 1986		
		Total	Mean	S.D.	Total	Mean	S.D.
<u>Caribou</u>							
Kaminuriak	M	319	3	2	383	3	2
	F	727	3	3	844	4	3
	C	125	2	2	143	2	2
	H	23	5	5	25	5	5
	Total	1194	3	3	1395	4	3
Moose		2	1	0	3	1	0
Muskox		1	1	0	2	2	0
Polar Bear		9	1	0	11	1	0
Arctic Fox		509	8	10	638	11	13
Red Fox		8	2	1	11	3	1
Wolf		61	4	4	74	5	5
Arctic Hare		1	1	0	1	1	0
Otter		1	1	0	1	1	0
Arctic Ground Squirrel		1	1	0	1	1	0
Ringed Seal		39	2	1	47	2	1
Bearded Seal		2	1	0	2	2	0
Eider		2	2	0	2	2	0
Parmigan		71	24	20	100	33	30
Arctic Charr		261	11	9	308	13	11
Lake Trout		208	8	10	243	9	11
Whitefish sp.		98	33	16	120	40	20
Arctic Grayling		10	5	2	12	6	2

¹See Table 1.

²See also Tables 4 and 11.

Table 19. The reported and estimated harvest for Rankin Inlet hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category ¹	Reported Harvest ² Oct. 1985 - March 1986			Estimated Harvest ² Oct. 1985 - March 1986		
		Total	Mean	S.D.	Total	Mean	S.D.
<u>Caribou</u>							
Kaminuriak	M	190	3	2	203	3	2
	F	324	3	2	346	3	2
	C	30	2	1	32	2	1
	U	9	2	1	10	2	1
	Subtotal	553	3	2	591	3	2
North of Chesterfield	M	27	2	1	29	3	2
	F	23	2	1	25	3	1
	C	7	2	0	8	3	1
	U	2	2	0	2	0	0
	Subtotal	59	2	1	64	3	1
	Total	612	3	2	655	3	2
Muskox		2	1	0	2	1	0
Polar Bear		10	1	0	10	1	1
Arctic Fox		253	9	12	272	10	13
Wolf		7	2	1	8	2	1
Arctic Hare		6	2	1	6	2	0
Ringed Seal		113	3	3	120	3	3
Bearded Seal		9	2	1	9	2	1
Harbour Seal		2	2	0	2	2	0
Harp Seal		1	1	0	1	1	0
Walrus		2	2	0	2	2	0
Eider		8	4	3	8	4	3
Ptarmigan		4	4	0	4	4	0
Searun Arctic Charr		1761	35	26	1872	37	27
Landlocked Arctic Charr		30	0	0	32	32	0
Lake Trout		17	4	2	18	5	2

¹see Table 1.

²See also Tables 6 and 13.

¹see Table 1.

²See also Tables 5 and 12.

Table 20. The reported and estimated harvest for Repulse Bay hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category ¹	Reported Harvest ² Oct. 1985 - March 1986			Estimated Harvest ² Oct. 1985 - March 1986		
		Total	Mean	S.D.	Total	Mean	S.D.
<u>Caribou</u>							
Wager	M	249	2	1	284	3	2
	F	192	2	1	217	3	2
	C	21	2	1	23	2	1
	U	4	2	1	4	2	1
	Total	466	2	1	528	3	2
Polar Bear		13	1	0	14	1	0
Arctic Fox		8	2	1	8	2	1
Red Fox		10	1	1	11	2	1
Wolf		33	2	1	36	2	1
Wolverine		1	1	0	1	1	0
Arctic Hare		2	2	0	2	2	0
Ringed Seal		105	3	3	126	3	3
Bearded Seal		4	1	0	5	1	0
Walrus		1	1	0	1	1	0
Narwhal		3	2	1	4	2	1
Ptarmigan		4	1	0	5	1	0
Arctic Charr		12	6	4	16	8	5
Lake Trout		70	42	49	729	81	49
		17	3	3	21	4	4

Table 21. The reported and estimated harvest for Whale Cove hunters expressed as numbers of animals. The mean monthly harvest per hunter and standard deviation about the mean are given.

Species	Category 1	Reported Harvest ² Oct. 1985 - March 1986			Estimated Harvest ² Oct. 1985 - March 1986		
		Total	Mean	S.D.	Total	Mean	S.D.
<u>Caribou</u>							
Kaminiutik	M	87	2		87	2	1
	F	251	2		251	2	2
	C	19	1	0	19	1	0
	U	4	1	1	4	1	1
	Total	361	2	1	361	2	1
<u>Muskox</u>							
Polar Bear		1	1	0	1	1	0
Arctic Fox		10	1	0	10	1	0
Red Fox		4	1	1	4	1	1
Arctic Hare		1	1	0	1	1	0
Ringed Seal		4	1	1	4	1	1
Arctic Char		31	1	0	31	1	0
Lake Trout		89	13	12	88	13	12
							6
							22
							3

See Table 1.

22680-2182 Tables 7 and 14

Table 22. Monthly theoretical kill factors for seven Keewatin communities.

	1985			1986		
	Oct.	Nov.	Dec.	Jan.	Feb.	Mar.
Raker Lake	1.00 ¹	1.00	1.00	1.00	1.00	1.00
Chesterfield Inlet	1.00	1.00	1.00	1.00	1.00	1.00
Coral Harbour	1.62	1.27	1.57	1.00	1.10	1.07
Eskimo Point	1.07	1.21	1.24	1.13	1.48	1.21
Rankin Inlet	1.05	1.05	1.09	1.06	1.10	1.08
Repulse Bay	1.33	1.16	1.00	1.23	1.10	1.00
Whale Cove	1.00	1.00	1.00	1.00	1.00	1.00

ה'ב

Table 23. The harvest by species over the range of age for Baker Lake hunters covering the period October 1985 to March 1986.

Species	Category ¹	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5 ²
<u>Caribou</u>						
Kaminuriak	M	6	44	69	75	11
	F		47	93	64	12
	C		24	14	10	8
	U			3		
	Subtotal	11	115	179	149	31
Beverley	M	2	22	56	52	8
	F	7	40	118	94	27
	C		10	7	10	3
	U				11	
	Subtotal	9	72	181	167	38
Wager	M	14	251	262	131	37
	F	10	204	274	180	65
	C	2	123	95	40	20
	U			4	8	
	Subtotal	26	578	635	359	122
Other	F		3			
	Subtotal	46	786	995		
	Total					
Muskox					6	
Arctic Fox					420	122
Wolf						
Arctic Charr						
Lake Trout						
Whitefish sp.						

1

Age classes are as follows:

Muskox Arctic Fox
Wolf Arctic Charr
Lake Trout Whitefish sp.

۱۰۷

Magier

Bevelry

Caribou

Hinter dem Standort der Deviationslinie liegt die Meile 11,5.

Table 24. The harvest by species over the range of age for Chesterfield Inlet hunters covering the period October 1985 to March 1986.

Species	Category ¹	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5 ²
<u>Caribou</u>						
Kamitnurak	M	2	4	2		
	F		1	2		
	U		4	6		
	Subtotal	2	9	10		
North of Chesterfield	M	17	50	30	8	
	F	1	10	23		
	U	2	10	9		
	Subtotal	20	70	62	8	
	Total	22	79	72	8	
Muskox		1		1		
Polar Bear						
Arctic Fox		3	2	1		
Red Fox		12		15		
Wolf			1			
Ringed Seal		1	7	6		
Bearded Seal		6	14	33		
Walrus		1	1	3		
Eider		1				
Ptarmigan		2				
Arctic Charr						
Arctic Cod						
Lake Trout		7	30	46		
Arctic Grayling		10	4	4		

¹see Table 1.

²For age classes see Table 23.

Table 25. The harvest by species over the range of age for Coral Harbour hunters covering the period October 1985 to March 1986.

Species	Category ¹	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5 ²
<u>Caribou</u>						
Southampton	M					
	C	1				
	U	1				
	Total	1				
Polar Bear		1				
Arctic Fox			25	33	44	2
Wolf				1		3
Arctic Hare		20	11	6		
Ringed Seal		75	173	129	1	1
Bearded Seal			5			16
Harp Seal			1	4		
Walrus			2	1		
Eider					10	1
Ptarmigan		2	442	779	580	37
Arctic Charr			459	620	324	22
Arctic Cod				7		

¹see Table 1.

²For age classes see Table 23.

Table 26. The harvest by species over the range of age for Eskimo Point hunters covering the period October 1985 to March 1986.

Species	Category ¹	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5 ²
<u>Caribou</u>						
Kaminuriak	M	96	110	108	5	
	F	127	373	223	4	
	C	32	62	27	4	
	U	5	18			
Total		260	563	358	13	
Moose		1	1	1		
Muskox		3	3	2	1	
Polar Bear		93	220	196		
Arctic Fox		1	4	3		
Red Fox		19	40	2		
Wolf		1				
Arctic Hare		1		1		
Otter				1		
Arctic Ground Squirrel		1				
Ringed Seal		10	22	7		
Bearded Seal		1	1			
Eider		2				
Ptarmigan		59	135	67		
Arctic Charr		85	54	69		
Lake Trout						
Whitefish sp.		3	7			
Arctic Grayling						

¹see Table 1.

²For age classes see Table 23.

Table 27. The harvest by species over the range of age for Rankin Inlet hunters covering the period October 1985 to March 1986.

Species	Category ¹	Number of Animals Harvested Per Age Class of Hunter				
		1	2	3	4	5 ²
<u>Caribou</u>						
Kaminuriak	M					
	F					
	C					
	U					
	Total					
North of Chesterfield	M					
	F					
	C					
	U					
	Total					
Muskox						
Polar Bear						
Arctic Fox						
Red Fox						
Wolf						
Arctic Hare						
Otter						
Arctic Ground Squirrel						
Ringed Seal						
Bearded Seal						
Eider						
Ptarmigan						
Arctic Charr						
Lake Trout						
Whitefish sp.						
Arctic Grayling						

¹see Table 1.

²For age classes see Table 23.

Table 28. The harvest by species over the range of age for Repulse Bay hunters covering the period October 1985 to March 1986.

Species	Number of Animals Harvested Per Age Class of Hunter				
	Category 1	2	3	4	5 ²
<u>Caribou</u>					
Wager	M	73	78	68	30
	F	72	88	19	13
	C	12	1	8	
	U	1	3		
Total		158	170	95	43
Polar Bear		2	5	4	2
Arctic Fox		1	4	1	2
Red Fox		4	3	3	
Wolf		12	18	2	1
Holverine				1	
Arctic Hare					
Ringed Seal		13	45	29	2
Bearded Seal					
Walrus		1	2	1	2
Narwhal		2	2		
Ptarmigan					
Arctic Charr		63	487	64	12
Lake Trout		5	9	1	20

Table 29. The harvest by species over the range of age for Whale Cove hunters covering the period October 1985 to March 1986.

Species	Number of Animals Harvested Per Age Class of Hunter					
	Category ¹	1	2	3	4	
<u>Caribou</u>						
Kamtnuritak	M					23
	F					80
	C					4
	U					1
Total						111
Muskox						1
Polar Bear						4
Arctic Fox						2
Red Fox						1
Wolf						1
Holverine						3
Arctic Hare						13
Ringed Seal						8
Bearded Seal						5
Walrus						12
Narwhal						55
Ptarmigan						16
Arctic Charr						1
Lake Trout						21

¹See Table 1.

²For age classes see Table 23.

¹See Table 1.

²For age classes see Table 23.

Table 30. Data on the distribution of hunters that were successful in obtaining a harvest expressed as a percentage over the range of age of hunters for the period October 1985 to March 1986.

Community	Range of Ages	Distribution of Successful Hunters by Month (%)						Total by Harvest Year
		Oct.	Nov.	Dec.	Jan.	Feb.	Mar.	
Baker Lake	0-15	3.9	1.0	2.1	1.8	2.2	2.8	1.8
	16-30	33.7	37.5	34.2	37.9	38.7	40.1	43.4
	31-45	32.6	30.7	31.6	30.2	33.9	31.1	27.6
	46-60	21.5	22.4	22.1	21.3	19.4	17.9	18.4
	61-75	8.3	8.3	10.0	8.9	5.9	8.0	8.8
	Number of successful hunters	181	192	190	169	186	212	272
Chesterfield Inlet	0-15	0.0	5.9	0.0	0.0	0.0	0.0	2.9
	16-30	37.5	41.2	14.3	0.0	27.8	31.3	42.9
	31-45	37.5	23.5	42.9	42.9	33.3	37.5	28.6
	46-60	25.0	23.5	28.6	42.9	38.9	31.3	22.9
	61-75	0.0	5.9	14.3	14.3	0.0	0.0	2.9
	Number of successful hunters	8	17	7	7	18	16	35
Coral Harbour	0-15	0.0	2.1	0.0	0.0	0.0	0.0	1.1
	16-30	48.1	29.2	16.0	31.6	21.2	26.9	40.2
	31-45	29.6	43.8	40.0	38.6	48.5	50.0	34.8
	46-60	18.5	18.8	32.0	21.1	27.3	19.2	15.2
	61-75	3.7	6.3	12.0	8.8	3.0	3.8	8.7
	Number of successful hunters	27	48	25	57	33	26	92
Eskimo Point	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	21.4	31.1	23.7	18.4	29.3	39.4	30.6
	31-45	47.6	28.9	39.5	47.4	39.0	43.9	40.8
	46-60	28.6	35.6	36.8	34.2	29.3	19.5	25.9
	61-75	2.4	4.4	0.0	0.0	2.4	1.2	2.7
	Number of successful hunters	84	45	38	38	41	82	147
Rankin Inlet	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	28.2	30.6	9.5	14.3	26.7	19.4	28.6
	31-45	35.9	30.6	33.3	25.0	40.0	41.7	35.7
	46-60	30.8	36.1	42.9	53.6	30.0	33.3	28.6
	61-75	5.1	2.8	14.3	7.1	3.3	5.6	7.1
	Number of successful hunters	39	36	21	28	30	36	84
Repulse Bay	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	25.0	35.3	37.0	30.0	36.7	48.7	44.1
	31-45	29.2	29.4	33.3	46.7	36.7	33.3	30.9
	46-60	25.0	20.6	18.5	16.7	23.3	12.8	14.7
	61-75	20.8	14.7	11.1	6.7	3.3	5.1	10.3
	Number of successful hunters	24	34	27	30	30	39	68
Whale Cove	0-15	0.0	0.0	0.0	0.0	0.0	0.0	0.0
	16-30	43.3	46.4	34.8	51.9	40.9	50.0	47.8
	31-45	20.0	28.6	26.1	25.9	27.3	20.0	21.7
	46-60	26.7	21.4	26.1	18.5	27.3	30.0	19.6
	61-75	10.0	3.6	13.0	3.7	4.5	0.0	10.9
	Number of successful hunters	30	28	23	27	22	20	46
Regional Total	0-15	1.8	1.0	1.2	.8	1.1	1.4	.9
	16-30	31.8	35.8	29.9	32.5	34.4	37.6	39.1
	31-45	34.9	31.5	33.2	34.3	36.1	35.5	32.0
	46-60	24.4	24.5	26.0	25.0	23.9	20.2	20.6
	61-75	7.1	7.3	9.7	7.3	4.4	5.3	7.4
	Number of successful hunters	393	400	331	356	360	431	744

Table 31. Edible weight values in kilograms for harvested species as calculated from various sources.

Species	Estimated Individual Weight (kg)	Reference ¹
Caribou	48.0	Berger 1977
Moose	199.0	Berger 1977
Muskox	110.0	Riewe 1977
Polar bear	158.8	Native Harvesting Research Committee 1975, 1976a or b Dome et al. 1982 "
Black bear	45.4	"
Grizzly bear	45.4	"
Arctic hare	2.3	Native Harvesting Research Committee 1975, 1976a or b
Ringed seal	14.3	"
Bearded seal	98.4	"
Harbour seal	27.7	"
Harp seal	43.1	"
Walrus	185.1	"
Beluga ²	(M) 555.0 (F) 407.9	Sergeant and Brodie 1969
Narwhal	(M) 595.2 (F) 397.0	Hay (personal communication, DFO, St. Johns's, NF); Sergeant and Brodie 1969 Bellrose 1976
Canada geese (Hutchinsii)	2.4	"
Snow geese (Lesser)	1.6	"
Ross's geese	1.0	"
Eider (Hudson Bay)	1.5	"
Old squaw	0.5	"
mallard	0.7	"
Ptarmigan	0.4	Thomas 1982
Sandhill crane	4.1	Stevens 1965
Snowy owl	1.8	Earhart and Johnson 1970 Bellrose 1976
Swan	6.8	Carder 1983
Arctic charr	2.5	Rond 1975; Keleher 1964
Lake trout	2.4	MacDonald and Fudge 1979; Keleher 1964
Whitefish sp.	2.8	Falk and Gillman 1975; Keleher 1964
Northern pike	2.1	"
Arctic grayling	0.9	"

¹These references are listed in detail in the reference section of the report.

²"M" means male, "F" means female.

Table 32. Reported and estimated edible weight values (kg) for harvested species for the period October, 1985 to March, 1986. The reported and estimated edible weight values for Baker Lake, Chesterfield Inlet and Whale Cove are the same as all six theoretical kill values equalled one.

Community and Species	1985-86 Reported Harvest (kg)	1985-86 Estimated Harvest (kg)
	Total	Total
<u>Baker Lake</u>		
Caribou	128400	128400
Muskox	1430	1430
Arctic Charr	28	28
Lake Trout	6276	6276
Whitefish sp.	535	535
Total	136669	136669
<u>Chesterfield Inlet</u>		
Caribou	8688	8688
Muskox	220	220
Polar Bear	476	476
Ringed Seal	758	758
Bearded Seal	394	394
Walrus	555	555
Eider	6	6
Arctic Charr	208	208
Lake Trout	139	139
Arctic Grayling	13	13
Total	11457	11457
<u>Coral Harbour</u>		
Caribou	384	402
Polar Bear	6511	8577
Arctic Hare	87	109
Ringed Seal	5620	6980
Bearded Seal	1082	1193
Harp Seal	216	304
Walrus	555	696
Eider	17	27
Ptarmigan	744	855
Arctic Charr	3563	4756
Total	18779	23799
<u>Eskimo Point</u>		
Caribou	57312	67141
Moose	398	519
Muskox	110	163
Polar Bear	1429	1815
Arctic Hare	2	3
Ringed Seal	558	676
Bearded Seal	197	211
Eider	3	4
Ptarmigan	28	40
Arctic Charr	653	769
Lake Trout	499	584
Whitefish sp.	274	336
Arctic Grayling	9	11
Total	61472	72272

Table 32 Cont'd.

Community and Species	1985-86 Reported Harvest (kg)	1985-86 Estimated Harvest (kg)
	Total	Total
<u>Rankin Inlet</u>		
Caribou	29376	31420
Muskox	220	238
Polar Bear	1588	1667
Arctic Hare	14	15
Ringed Seal	1616	1710
Bearded Seal	886	933
Harbour Seal	55	58
Harp Seal	43	45
Walrus	370	400
Eider	12	13
Ptarmigan	2	2
Arctic Charr	4403	4680
Lake Trout	41	43
Total	38626	41224
<u>Repulse Bay</u>		
Caribou	22368	25270
Polar Bear	1747	2020
Arctic Hare	5	5
Ringed Seal	1502	1801
Bearded Seal	394	459
Walrus	555	739
Narwhal	1984	2302
Ptarmigan	5	6
Arctic Charr	1585	1822
Lake Trout	41	50
Total	30186	34474
<u>Whale Cove</u>		
Caribou	17328	17328
Muskox	110	110
Polar Bear	1588	1588
Arctic Hare	9	9
Ringed Seal	443	443
Arctic Charr	163	163
Lake Trout	53	53
Total	19694	19694

*Please note that rounding has caused small discrepancies in column totals.

Table 33. Reported and estimated edible weight values (kg) for four major groups of animals harvested by Keewatin communities, October, 1985 to March, 1986.

Baker Lake (reported edible wt.)						Baker Lake (estimated edible wt.)						
Period	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Terrestrial	Marine	Fowl	Fish
		Terrestrial	Marine	Fowl		Fish						
1985												
Oct	36633 ¹	34512 (94.2)			2121 (5.8)	36633 ¹	34512 (94.2)		2121 (5.8)			
Nov	16598	15168 (91.4)			1430 (8.6)	16598	15168 (91.4)		1430 (8.6)			
Dec	18761	17520 (93.4)			1241 (6.6)	18761	17520 (93.4)		1241 (6.6)			
1986												
Jan	17418	16752 (96.2)			666 (3.8)	17418	16752 (96.2)		666 (3.8)			
Feb	21113	20352 (96.4)			761 (3.6)	21113	20352 (96.4)		761 (3.6)			
Mar	26146	25526 (97.6)			620 (2.4)	26146	25526 (97.6)		620 (2.4)			
Total	136669	129830 (95.0)			6839 (5.0)	136669	129830 (95.0)		6839 (5.0)			

¹The theoretical kill factor (Table 22) is the value by which the reported kill per species is multiplied to arrive at the estimated harvest. In cases where this value is one then 100% of the hunters have been interviewed and the reported and estimated harvests are equal.

Table 33 Cont'd.

Chesterfield Inlet (reported edible wt.)						Chesterfield Inlet (estimated edible wt.)						
Period	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Terrestrial	Marine	Fowl	Fish
		Terrestrial	Marine	Fowl		Fish						
1985												
Oct	985	687 (69.7)		299 (30.3)		985	687 (69.7)		299 (30.3)			
Nov	2852	2382 (83.5)		143 (5.0)		2852	2382 (83.5)		143 (5.0)			
Dec	960	960 (100.0)				960	960 (100.0)					
1986												
Jan	1233	1104 (89.6)		129 (10.4)		1233	1104 (89.6)		129 (10.4)			
Feb	2849	2572 (90.3)		271 (9.5)		2849	2572 (90.3)		271 (9.5)			
Mar	2579	1680 (65.2)		666 (33.6)		2579	1680 (65.2)		866 (33.6)			
Total	11458	9385 (81.9)		1708 (14.9)		6 (.1)	360 (3.1)		11458			
										1708 (14.9)		
										6 (.1)		
											360 (3.1)	

Table 33 Cont'd.

Coral Harbour (reported edible wt.)						Coral Harbour (estimated edible wt.)						
Period	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Terrestrial	Marine	Fowl	Fish
		Terrestrial	Marine	Fowl		Fish						
1985												
Oct	2750	1112 (40.4)	1070 (38.9)	28 (1.0)	540 (19.6)	4455	1801 (40.4)	1734 (38.9)	45 (1.0)	875 (19.6)		
Nov	8020	4313 (53.8)	1102 (13.7)	82 (1.0)	2523 (31.5)	10184	5477 (53.8)	1399 (13.7)	104 (1.0)	3204 (31.5)		
Dec	1413	338 (23.9)	672 (47.6)	113 (8.0)	29n (20.5)	12218	531 (23.9)	1055 (47.6)	177 (8.0)	455 (20.5)		
1986												
Jan	2527	473 (18.7)	1650 (65.3)	336 (13.3)	68 (2.7)	2527	473 (18.7)	1650 (65.3)	336 (13.3)	68 (2.7)		
Feb	1994	226 (11.3)	1565 (78.5)	114 (5.7)	90 (4.5)	2193	248 (11.3)	1721 (78.5)	125 (5.7)	99 (4.5)		
Mar	2077	521 (25.1)	1414 (68.1)	89 (4.3)	53 (2.5)	2222	558 (25.1)	1513 (68.1)	95 (4.3)	56 (2.5)		
Total	18781	6982 (37.2)	7473 (39.8)	762 (4.0)	3564 (19.0)	23799	9088 (38.2)	9073 (38.1)	882 (3.7)	4757 (20.0)		

Table 33 Cont'd.

Eskimo Point (reported edible wt.)						Eskimo Point (estimated edible wt.)						
Period	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Terrestrial	Marine	Fowl	Fish
		Terrestrial	Marine	Fowl		Fish						
1985												
Oct	23187	22464 (96.9)	411 (1.8)	312 (1.3)	24811	24037 (96.9)	440 (1.R)					
Nov	6575	6056 (92.1)	57 (1.2)	519 (7.9)	7956	7327 (92.1)						
Dec	4601	4176 (90.3)	57 (1.2)	368 (8.0)	5705	5178 (90.8)	71 (1.2)					
1986												
Jan	5532	5431 (98.2)	14 (.3)	87 (1.6)	6252	6137 (98.2)	16 (.3)					
Feb	5331	5187 (97.3)	114 (2.1)	20 (.4)	7890	7676 (97.3)	169 (2.1)					
Mar	16245	15938 (98.1)	157 (1.0)	11 (.1)	19656	19285 (98.1)	190 (1.0)					
Total	61471	59252 (96.4)	753 (1.2)	31 (.1)	1435 (2.3)	72270	69640 (96.4)	886 (1.2)				
									43 (.1)	1701 (2.4)		

Table 33 Cont'd.

Rankin Inlet (estimated edible wt.)

Period	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)		
		Terrestrial	Marine	Fowl		Terrestrial	Marine	Fowl
1985								
Oct	8481	5717 (67.4)	1787 (21.1)	14 (.2)	963 (11.4)	8903	6002 (67.4)	1876 (21.1)
Nov	9204	7396 (80.4)	343 (3.7)	29 (.9)	1465 (15.9)	9664	7766 (80.4)	360 (3.7)
Dec	3303	2544 (77.0)			730 (22.1)	3601	2773 (77.0)	31 (.9)
1986								
Jan	5051	4229 (83.7)	43 (.8)		779 (15.4)	5353	4482 (83.7)	46 (.8)
Feb	5323	4949 (93.0)	114 (2.1)		260 (4.9)	5956	5444 (93.0)	126 (2.1)
Mar	7267	6364 (87.6)	655 (9.0)		248 (3.4)	7847	6873 (87.6)	707 (9.0)
Total	38629	31199 (80.8)	2971 (7.7)	14 (.0)	4445 (11.5)	41223	33340 (80.9)	3146 (7.6)
								14 (.0)
								4723 (11.5)

Table 33 Cont'd.

Repulse Bay (estimated edible wt.)

Period	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)		
		Terrestrial	Marine	Fowl		Terrestrial	Marine	Fowl
1985								
Oct	5771	3767 (65.3)	1894 (32.8)	5 (.1)	105 (1.8)	4339	2832 (65.3)	1424 (32.8)
Nov	8671	4446 (51.3)	2617 (30.2)		1598 (18.4)	7467	3833 (51.3)	2256 (30.2)
Dec	2298	2112 (91.9)	184 (8.0)		2 (1)	2298	2112 (91.9)	184 (8.0)
1986								
Jan	5504	5468 (99.3)	35 (.6)	1 (.0)		4474	4446 (99.3)	27 (.6)
Feb	6948	6649 (95.7)	299 (4.3)			6316	6044 (95.7)	272 (4.3)
Mar	5290	4853 (91.7)	270 (5.1)			5290	4853 (91.7)	270 (5.1)
Total	34472	27295 (79.2)	5299 (15.4)	6 (.0)	1872 (5.4)	30184	24120 (79.9)	4433 (14.7)
								5 (.0)
								1626 (5.4)

Table 33 Cont'd.

Period	Whale Cove (reported edible wt.)			Whale Cove (estimated edible wt.)		
	Total Edible Weight (kg)	Weight (kg) per Category (bracketed figures are % of total)			Weight (kg) per Category (bracketed figures are % of total)	
		Terrestrial	Marine	Fowl	Fish	
1985						
Oct	3151	2976 (94.4)	143 (4.5)	32 (.1)	3151	2976 (94.4)
Nov	3691	3654 (99.0)	14 (.4)	23 (.6)	3691	3654 (99.0)
Dec	2263	2162 (95.5)	86 (3.8)	15 (.7)	2263	2162 (95.5)
1986						
Jan	5264	5040 (95.7)	100 (1.9)	124 (2.4)	5264	5040 (95.7)
Feb	2614	2549 (97.5)	43 (1.6)	22 (.8)	2614	2549 (97.5)
Mar	2711	2654 (97.9)	57 (2.1)		2711	2654 (97.9)
Total	19694	19035 (96.7)	443 (2.3)	216 (1.1)	19694	19035 (96.7)
						443 (2.3)
						216 (1.1)

Table 34. Age distribution of hunters for the seven Keewatin region communities for the period October 1985 to March 1986.

Community	Percentage of Hunters Per Age Category						Total Known Hunters
	0-15	16-30	31-45	46-60	61-75	76+ ¹	
Baker Lake	1.6	41.3	27.6	20.2	6.8	2.5	366
Chesterfield Inlet	1.2	51.2	25.6	17.4	3.5	1.2	86
Coral Harbour	2.4	47.6	25.6	15.2	8.5	0.6	164
Eskimo Point	0.3	42.4	33.3	17.8	6.1	0.0	309
Rankin Inlet	0.3	40.6	34.0	17.7	6.6	0.7	288
Repulse Bay	0.8	47.5	29.5	13.1	7.4	1.6	122
Whale Cove	0.0	42.0	28.4	15.9	11.4	2.3	88
Total hunters for the Keewatin District	1.0	43.3	30.0	17.6	7.0	1.2	1423

¹This category includes hunters of unknown ages. There are only 8 hunters of known age in this group.

Table 35. The estimated harvest by Baker Lake hunters, expressed as number of animals, for the period November, 1981 to March, 1986. Data were not available for October 1981 and August and September 1982. Results for November and December, 1981 show the actual reported harvest as data were insufficient to provide good a estimate. Therefore the actual harvest was considered to be the best estimate for these months.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	\bar{X}	NUM
CARIBOU KAMINURIAK ⁶ (H,A)	5	C	1982	1	2	0	0	0	0	0	0	3	6	0	0	6	0.6	10
	1985	0	0	31	24	9	0	0	0	0	0	24	24	120	10	10	10	
	1986	0	0	5	4	4	4	4	3	4	3	5	5	5	5	1.7	3	
	N	5	5	36	24	9	0	0	0	0	6	8	24	24	24	24	24	
	SUM	0	0	0	7	6	2	0	0	2	3	5	5	5	5	5	5	
	MEAN	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
F	1981	267	147	112	14	0	2	0	23	41	143	195	122	236	118	0	2	
	1982	353	0	42	102	33	0	3	28	37	21	66	27	63	129	6	10	
	1983	201	0	0	28	0	0	0	14	0	61	74	0	28	76	63	8	
	1984	0	0	87	42	34	0	14	0	0	0	56	44	44	119	9.9	12	
	1985	0	0	48	0	0	0	0	0	0	0	0	0	0	412	34.3	12	
	1986	0	0	5	5	4	4	4	3	4	3	3	5	5	48	16.0	3	
N	SUM	554	267	324	284	81	3	67	78	428	140	394	394	257	257	257	257	
	MEAN	111	53	65	71	20	1	17	26	107	47	79	79	51	51	51	51	
	5	161	169	129	264	10	451	0	0	0	0	0	0	0	0	0	0	
	1982	255	0	29	232	128	0	218	175	209	26	44	44	44	44	44	44	
	1983	85	0	0	65	0	45	0	275	26	10	0	0	0	0	0	0	
	1984	0	0	104	46	44	0	64	0	104	89	72	72	72	72	72	72	
M	1985	0	0	28	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1986	0	0	5	5	4	4	4	0	0	0	0	0	0	0	0	0	
	N	5	161	330	472	436	55	1008	201	670	115	311	311	311	311	311	311	
	SUM	340	32	66	118	109	14	252	67	168	38	62	62	62	62	62	62	
	MEAN	68	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	5	34	14	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
U	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1985	0	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	
	N	5	5	4	4	4	4	4	4	4	4	4	4	4	4	4	4	
	SUM	0	18	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	MEAN	0	7	4	0	0	0	0	0	0	0	0	0	0	0	0	0	

Table 35. (cont'd)

CARIBOU WAGER (H,A)		C		F		M		U	
1984	0	0	0	0	0	0	0	0	0
1985	0	1	0	19	12	4	2	35	30
1986	59	81	65	5	4	4	3	3	9
N	5	5	5	19	12	4	2	6	3
SUM	59	82	65	5	3	1	1	9	13
MEAN	12	16	13						
1983	0	0	0	0	100	23	57	151	20
1984	2	0	0	0	88	17	9	242	191
1985	0	9	0	33	42	57	63	227	325
1986	141	157	202	4	4	4	3	4	80
N	5	5	5	33	130	174	95	526	667
SUM	143	166	202	8	33	44	24	175	167
MEAN	29	33	40						
1982	0	0	0	0	0	0	0	6	0
1983	0	0	0	0	301	169	188	329	111
1984	0	0	0	4	388	116	111	267	310
1985	0	5	0	45	36	70	170	325	488
1986	149	178	150	5	4	4	3	3	96
N	5	5	5	12	49	424	487	450	780
SUM	149	183	150	30	12	106	113	260	1133
MEAN	30	37	30						
1983	0	0	0	0	0	0	0	0	0
1984	0	0	0	0	32	21	14	2	1
1985	0	0	0	0	0	3	2	0	2
1986	0	8	1	-	-	-	-	-	0
N	5	5	5	1	4	4	4	4	4
SUM	0	8	1	0	32	24	16	5	4
MEAN	0	2	0						

For footnotes see Appendix 4.

Table 35. (cont'd)

CARIBOU BEVERLY (H.A.)		C		F		M		U	
YR	NUM	0	0	0	0	0	0	0	0
1983	0	5	0	0	0	0	0	0	0
1984	0	0	0	0	0	0	0	0	0
1985	N	73	65	13	6	0	0	0	0
	SUM	73	70	13	6	0	0	0	0
	MEAN	15	14	3	2	0	0	0	0
1981	1982	11	27	36	22	0	0	0	0
	1983	0	270	98	84	38	0	21	7
	1984	100	247	292	234	8	0	3	0
	1985	N	191	148	26	10	0	0	0
	SUM	302	692	452	350	46	0	24	3
	MEAN	60	138	90	88	12	0	6	2
1981	1982	0	0	5	8	34	9	0	10
	1983	0	0	87	153	174	281	0	163
	1984	151	337	457	424	50	0	73	40
	1985	N	179	164	25	13	0	0	0
	SUM	330	593	643	645	4	0	4	4
	MEAN	66	119	129	161	85	0	246	40
1981	1983	0	0	0	0	0	0	0	0
	1984	0	0	9	0	0	0	2	0
	1985	N	17	5	3	1	0	0	1
	SUM	17	14	3	1	1	0	2	0
	MEAN	3	3	1	1	1	0	1	0

For footnotes see Appendix 4.

Table 35. (cont'd)

For footnotes see Appendix 4.

Table 35. (cont'd)

WOLF (H,S)	1982	4	16	2	0	0	0	0	28	2.8	10
	1983	0	0	3	0	2	0	11	20	1.7	12
	1984	0	12	25	1	0	0	9	58	4.8	12
	1985	7	16	18	1	0	0	0	21	63	5.3
	1986	3	5	13	1	0	0	0	21	7.0	3
N	SUM	5	5	4	4	4	3	4	5	5	5
	MEAN	14	33	75	4	2	0	0	41	1	1
RINGED SEAL (H,S)	1983	0	0	0	0	1	0	0	0	0	0
	1984	0	0	0	0	2	3	0	0	1	0.1
CANADA GEESE (L,S)	1984	0	0	0	0	154	142	0	0	296	24.7
	1985	0	0	0	0	479	144	0	0	623	51.9
	N	5	5	5	4	4	4	3	5	5	12
	SUM	0	0	0	0	633	286	0	0	0	0
	MEAN	0	0	0	0	158	72	0	0	0	0
SNOW GEESE (L,S)	1984	0	0	0	0	149	201	0	0	350	29.2
	1985	0	0	0	0	30	0	0	0	30	2.5
	N	5	5	5	4	4	4	3	5	5	12
	SUM	0	0	0	0	149	231	0	0	0	0
	MEAN	0	0	0	0	37	58	0	0	0	0
PTARMIGAN (L,A)	1981	0	0	0	0	0	0	0	4	0	4
	1984	0	0	0	0	0	0	0	0	752	62.7
SWAN (L,S)	1985	0	0	0	0	2	4	0	0	0	0
CANADA GOOSE EGGS	1985	0	0	0	0	151	265	0	0	0	0
GOOSE EGGS	1984	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	773	1695	0	0	0	0
	N	5	5	5	4	4	4	3	5	5	12
	SUM	0	0	0	0	773	4417	0	0	0	0
	MEAN	0	0	0	0	193	1104	0	0	0	0

For footnotes see Appendix 4.

Table 35. (cont'd)

ARCTIC CHARR (H,S)	1981	0	0	0	138	0	0	65	0	128	0	128	64.0	2
	1984	0	0	0	100	57	0	0	4	42	2	247	20.6	12
	1985	0	0	0	4	4	0	0	3	0	7	168	14.0	12
N	5	5	5	0	100	195	0	0	4	5	5	.	.	.
SUM	0	0	0	0	25	49	0	0	16	1	34	2	.	.
MEAN	0	0	0	0
LAKE TROUT (H,A)	1981	0	0	380	400	0	0	0	0	0	0	8744	2154	10898
	1982	0	0	0	796	0	1886	188	0	1732	545	178	780	10
	1983	0	0	0	241	32	175	72	87	182	196	144	5325	443.8
	1984	76	157	268	0	264	596	333	59	25	798	520	192	151.8
	1985	181	43	0	0	469	3288	274.0
	1986	261	317	250	4	4	4	3	3	3	3	828	276.0	12
N	5	5	5	5	1092	771	2291	334	207	2726	9953	5	.	.
SUM	518	517	898	641	160	273	193	573	111	52	909	1991	2993	.
MEAN	104	103	180	64.1	160	273	193	573	111	52	909	1991	599	.
WHITEFISH (H,A)	1983	0	0	224	53	0	0	0	0	0	0	72	349	29.1
	1984	50	151	144	136	19	0	17	27	22	114	37	65	65.2
	1985	129	29	0	0	0	21	54	25	29	70	65	35	437
	1986	14	0	7	0	21	36.4
N	5	5	5	5	4	4	4	3	3	3	3	5	5	12
SUM	193	180	151	360	72	21	71	52	31	184	102	172	172	3
MEAN	39	36	30	90	18	5	18	17	8	61	20	34	34	.
ARCTIC GRAYLING (L,A)	1984	0	0	0	0	0	0	0	25	0	3	0	28	2.3
	1985	0	0	0	0	0	56	30	0	0	0	0	86	12
NORTHERN PIKE (L,S)	1984	0	0	0	0	0	0	0	25	0	0	0	25	2.1
FRESHWATER FISH	1982	0	142	0	0	0	0	0	0	0	0	0	142	14.2

For footnotes see Appendix 4.

Table 36. The estimated harvest by Chesterfield Inlet hunters, expressed as number of animals for the period January, February and August, 1982 through March, 1986. The actual harvest for August, 1982 is used as it is the best estimate of the harvest for this particular month.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	\bar{X}	NUM	
CARIBOU KAMINURIAK ⁵ (H.A.) 6	F	1982	1	3							2	0	0	15	0	28	4.0	7	
	1983	2	11	0	0	0	0	0	0	0	0	0	0	0	6	26	2.2	12	
	1984	17	0	1	0	6	0	0	1	0	0	0	0	0	10	35	2.9	12	
	1985	0	2	0	0	0	0	0	0	0	0	0	0	0	5	0.4	12		
	SUM	30	5	4	3	3	3	3	4	4	4	4	4	15	16				
	MEAN	6	22	1	0	0	0	0	1	0	0	1	4	16					
CARIBOU N. CHEST. ⁴	M	1981	2	6	0	0	0	0	0	0	0	0	0	0	0	27	3.9	7	
	1982	1	1	0	0	0	0	0	0	0	0	0	0	0	11	28	2.3	12	
	1983	6	0	0	0	0	0	0	0	0	0	0	0	0	10	63	5.3	12	
	1984	14	0	0	0	0	0	0	0	0	0	0	0	0	0	52	4.3	12	
	1985	0	7	30	3	2	0	0	0	0	0	0	0	0	0				
	SUM	22	5	4	3	3	3	3	4	4	4	4	4	13	21				
CARIBOU N. CHEST. ⁴	MEAN	4	13	3	3	3	2	1	1	1	1	1	1	5	13				
	U	1983	0	7	0	0	0	0	0	0	0	0	0	0	0	7	0.6	12	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	8	0.7	12	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	10	0.8	12	
	SUM	74	5	5	4	3	3	3	4	4	4	4	4	19	35				
	MEAN	15	18	8	8	8	8	8	8	8	8	8	8	5	9	26			
CARIBOU N. CHEST. ⁴	(H.A.)	F	1982	34	2	19	50	2	2	1	2	9	3	9	1	21	70	10.0	7
	1983	37	50	9	4	0	0	0	0	0	0	5	8	16	0	191	15.9	12	
	1984	0	23	9	3	0	0	0	0	0	0	0	0	10	0	70	5.8	12	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	5	18	1.5	12	
	1986	3	14	4	3	3	3	3	3	3	3	4	4	4	4	21	7.0	3	
	SUM	72	5	5	4	3	3	3	4	4	4	4	4	19	35				
CARIBOU N. CHEST. ⁴	MEAN	15	18	8	8	8	8	8	8	8	8	8	8	5	9	26			
	M	1982	18	4	1	107	26	9	56	4	10	12	3	10	4	27	124	17.7	7
	1983	31	21	31	38	23	5	13	2	8	10	35	9	4	0	310	25.8	12	
	1984	0	21	31	9	5	3	3	3	3	4	15	0	0	0	190	15.8	12	
	1985	3	7	23	9	5	3	3	3	3	4	4	4	4	0	97	8.1	12	
	1986	20	31	23	9	5	3	3	3	3	4	4	4	4	0	74	24.7	3	
CARIBOU N. CHEST. ⁴	SUM	5	4	3	3	3	3	3	4	4	4	4	4	31	27				
	MEAN	14	17	24	50	21	5	15	2	8	11	24	8	7	11				
	1983	0	2	0	2	0	0	0	0	0	0	0	0	0	0	0	4	0.3	12
	1984	0	6	0	0	0	0	0	0	0	0	0	0	0	0	0	15	1.3	12
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	18	1.5	12
	1986	0	4	8	4	3	3	3	3	3	3	3	3	3	0	0	12	4.0	3
CARIBOU N. CHEST. ⁴	N	5	5	4	3	3	3	3	3	3	3	3	3	3	0	0	0	0	0
	SUM	0	12	8	2	2	2	2	2	2	2	2	2	2	0	0	0	0	0
CARIBOU N. CHEST. ⁴	MEAN	0	2	2	2	2	2	2	2	2	2	2	2	2	0	0	0	0	0

Table 36. (cont'd)

Table 36. (cont'd)

For footnotes see Appendix 4.

Table 36. (cont'd)

ARCTIC CHARR (H,A)	1982	76	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	0	0	0	0	0	0	0
	1986	0	0	13	3	3	3	4	4	4	4	3
N	5	5	4	0	0	1	63	58	469	50	70	13
SUM	76	0	13	0	0	0	19	117	117	0	18	0
MEAN	15	0	3	0	0	0	21	19	117	13	0	0
LAKE TROUT (H,A)	1982	0	19	0	131	41	72	8	202	0	26	0
	1983	0	0	30	0	59	26	0	0	0	43	0
	1984	0	0	0	0	2	31	22	0	0	111	0
	1985	0	0	0	0	4	3	3	0	0	0	0
N	5	5	4	3	30	134	131	120	8	202	0	197
SUM	0	19	0	8	45	44	40	3	51	0	137	113
MEAN	0	4	0	0	0	0	0	0	0	34	127	94
ARCTIC GRAYLING (L,A)	1985	0	0	0	0	0	0	0	0	0	14	0
SCULPINS (L,A)	1984	0	0	0	0	0	1	0	0	0	0	1

For footnotes see Appendix 4.

Table 37. The estimated harvest by Coral Harbour hunters, expressed as number of animals, for the period February, 1982 and June 1982 through April, 1983, and October, 1983 to March, 1986. The actual harvest was used as the best estimate of the harvest for October, November, December, 1982, and October, November, December, 1983, and January, 1984.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	\bar{X}	NUM
CARIBOU KAMINURIAK	F	1985	0 ²	0	0	6	0	0	0	0	0	0	0	0	0	6	0.5	12
	⁶ (H,S)	M	1985	0	0	3	0	0	0	0	0	0	0	0	0	3	0.3	12
	U	1984 ⁴	0	0	4	0	0	0	0	0	0	0	0	0	4	0.3	12	
CARIBOU WAGER	F	1984 ¹	0	0	78	0	0	3	0	0	0	0	0	0	0	81	6.8	12
	1985	N	4	0	50	63	0	0	0	0	0	0	0	0	0	113	9.4	12
	SUM	0	0	0	128	63	3	3	4	4	4	4	4	4	4	.	.	.
	MEAN	0	0	0	43	32	0	0	1	0	0	0	0	0	0	.	.	.
	M	1984	0	0	60	0	0	9	0	0	0	0	0	0	0	69	5.8	12
	1985	N	4	0	4	27	0	0	0	0	0	0	0	0	0	31	2.6	12
	SUM	0	0	0	64	27	3	3	4	4	4	4	4	4	4	.	.	.
	MEAN	0	0	0	21	14	0	0	2	0	0	0	0	0	0	.	.	.
	U	1984	0	0	55	0	0	0	0	0	0	0	0	0	0	55	4.6	12
	1985	N	4	0	30	36	0	0	0	0	0	0	0	0	0	66	5.6	12
	SUM	0	0	0	85	36	3	2	3	4	4	4	4	4	4	.	.	.
	MEAN	0	0	0	28	18	0	0	0	0	0	0	0	0	0	.	.	.

For footnotes see Appendix 4.

Table 37. (cont'd)

Table 37. (cont'd)

CARIBOU OTHER		M	1985	1	0	0	0	0	0	0	0	1	0.1	12					
POLAR BEAR (H,A)		1982	0	0	0	0	0	0	0	0	0	6	14	1.8	8				
1983		1984	4	0	0	0	0	0	0	0	0	36	40.0	9					
1985		1	0	3	4	0	0	0	0	0	0	45	3.8	12					
1986		2	1	2	3	3	3	3	3	3	3	61	5.1	12					
N	SUM	4	5	4	3	2	3	4	4	4	4	5	1.7	3					
MEAN		2	1	5	4	4	2	0	0	1	16	59	9						
ARCTIC FOX (H,A)		1982	50	224	17	0	0	0	0	0	0	166	167	103	486	60.8	B		
1983		117	160	55	89	123	0	0	0	0	0	21	0	51	739	82.1	9		
1984		85	29	115	76	74	12	0	0	0	0	0	0	123	584	48.7	12		
1985		29	13	2	35	35	0	0	0	0	0	0	0	69	392	32.7	12		
1986		13	4	5	4	3	2	3	3	3	3	4	4	4	50	16.7	3		
N	SUM	4	382	382	424	214	12	0	0	0	0	23	21	166	419	346			
MEAN		6	76	106	71	6	0	0	0	0	0	42	5	105	87				
WOLF (H,S)		1984	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12		
1985		0	0	0	4	4	0	0	0	0	0	0	0	0	10	0.8	12		
ARCTIC HARE (L,S)		1982	0	0	0	0	0	0	0	0	0	0	0	0	9	9	18	2.3	8
1983		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	9	1.0	9
1984		0	2	0	0	1	0	0	0	0	0	0	0	0	3	3	3	0.3	12
1985		2	5	9	5	7	1	0	0	0	0	0	0	0	14	14	39	3.3	12
1986		5	4	5	4	3	2	3	3	4	4	4	4	4	4	19	6.3	3	
N	SUM	4	15	7	9	15	7	1	0	0	0	0	0	0	23	26			
MEAN		2	2	2	4	2	1	0	0	0	0	0	0	0	6	7			

For footnotes see Appendix 4.

Table 37. (cont'd)

RINGED SEAL (H,A)	1982	10	22	108	52	42	30	52	679	84.9	8	
	1983	67	40	192	49	59	87	13	591	65.7	9	
	1984	134	101	55	41	57	142	20	756	63.0	12	
	1985	89	93	19	22	30	113	18	739	61.6	12	
	1986	81	83	78	3	2	3	3	242	80.7	3	
N		4	5	4	3	87	533	3				
SUM		371	349	192	85	264	338	179	226			
MEAN		93	70	48	28	44	178	88	160			
BEARDED SEAL (H,S)	1982	16	6	2	3	3	10	1	0	35	4.4	8
	1983	0	0	17	2	0	5	20	33	1	69	7.7
	1984	4	7	5	2	0	0	12	4	0	3	66
	1985	6	5	6	2	0	0	14	6	0	49	5.5
	1986	5	6	0	3	2	3	3	2	0	11	12
N		4	5	4	3	10	0	7	4	4	4	3.7
SUM		15	34	26	10	0	7	21	41	7	4	3
MEAN		4	7	7	3	0	2	7	10	15	1	1
HARBOUR SEAL (H,S)	1983	0	0	0	0	0	0	3	0	0	0	0
	1985	0	0	0	0	0	0	4	2	0	0	0
								3	0	0	3	0
								0	0	6	0.3	9
HARP SEAL (H,S)	1982	0	0	0	0	0	0	10	58	37	0	105
	1983	0	0	0	0	0	0	1	46	18	0	0
	1984	4	3	0	0	0	2	1	13	3	5	64
	1985	0	0	0	0	0	3	2	3	0	4	28
N		4	5	4	3	0	2	3	120	4	4	0
SUM		4	3	0	0	0	1	4	30	58	4	14
MEAN		1	1	0	0	0	0	1	15	2	0	1
UNKNOWN SEAL	1982	32	0	0	0	0	0	0	0	0	0	32
	1984	0	1	0	0	0	0	0	0	0	0	1
WALRUS (H,A)	1982	2	9	6	6	9	10	19	12	5	1	59
	1983	3	0	0	0	0	0	21	12	5	0	59
	1984	4	0	1	0	0	0	3	11	14	2	41
	1985	1	0	1	2	0	0	6	9	8	0	41
	1986	0	0	2	3	0	0	3	3	4	4	29
N		4	5	4	3	2	3	4	4	4	4	2
SUM		8	11	7	8	6	9	19	60	46	1	0
MEAN		2	2	2	2	3	3	6	15	12	1	0

For footnotes see Appendix 4.

Table 37. (cont'd)

BELUGA (H,A)	1982	0	0	0	0	0	0	0	1	125	15.6	8
	1983	0	0	0	0	0	0	0	0	128	14.2	9
	1984	2	1	0	0	3	33	50	2	0	0	12
	1985	0	0	0	0	0	15	60	24	0	0	12
N	4	5	4	3	2	3	3	4	13	0	0	88
SUM	2	1	0	0	0	3	65	262	117	4	4	7.3
MEAN	1	0	0	0	0	1	22	66	29	1	1	12
CANADA GEESE (L,S)	1982	0	0	0	0	0	41	407	199	0	0	9
	1983	0	0	0	0	0	13	101	9	10	0	19
	1984	0	0	0	0	0	30	25	0	23	0	0
	1985	0	0	0	0	0	43	167	3	0	0	0
N	4	5	4	3	2	3	43	407	3	4	4	4
SUM	2	1	0	0	0	0	22	56	136	10	4	4
MEAN	0	0	0	0	0	0	22	56	58	3	0	0
SNOW GEESE (H,A)	1982	0	0	0	0	0	0	4345	0	0	0	22
	1983	0	0	0	0	0	5	322	5063	21	0	0
	1984	3	2	0	0	0	0	118	3798	12	0	0
	1985	0	0	0	0	0	4	5	24	301	0	0
N	4	5	4	3	2	3	4	4	3	4	4	4
SUM	3	2	1	0	0	0	2	440	13206	534	0	0
MEAN	1	0	0	0	0	0	2	220	4402	158	0	0
ROSS'S GEESE (L,S)	1982	-	0	-	-	-	-	248	7	11	1	0
BRANT GEESE (L,S)	1984	0	0	0	0	0	4	1	0	0	0	0
UNKNOWN GEESE	1984	0	0	0	0	0	3	76	0	0	0	0
	1985	0	0	0	0	0	0	309	0	1	0	0
OLD SQUAW (L,S)	1982	-	0	-	-	-	-	0	0	1	0	1
GUILLEMOT (L,S)	1982	-	0	-	-	-	0	0	0	3	0	0

For footnotes see Appendix 4.

Table 37. (cont'd)

EIDER	(L,A)	1982	0	0	0	0	0	0	327	40.9	8
		1983	0	0	0	0	0	0	66	7.3	9
		1984	0	0	0	0	0	0	49	4.1	12
		1985	0	0	0	0	0	0	29	2.4	12
N	4	5	4	3	2	3	3	4	4	4	4
SUM	0	0	0	0	14	223	0	59	131	41	3
MEAN	0	0	0	0	7	74	0	15	33	10	1
PTARMIGAN	(H,A)	1982	113	0	0	171	0	17	165	137	151
		1983	82	28	96	283	183	98	90	134	99
		1984	158	152	33	254	206	0	0	100	127
		1985	33	29	175	312	10	0	17	45	66
		1986	839	312	238	537	2	3	4	260	443
N	4	5	4	3	3	389	181	0	4	4	4
SUM	1112	634	542	136	179	195	60	115	372	319	576
MEAN	278	127	127	127	179	195	60	29	93	80	845
SNOWY OWL	(L,S)	1982	-	2	-	-	0	0	0	0	211
UNKNOWN FOWL		1982	-	0	-	-	5	0	0	0	0
		1983	0	0	0	0	0	0	0	0	0
		1984	0	2	0	0	0	0	0	0	0
SWAN	(L,S)	1982	0	0	0	0	2	0	0	0	0
		1984	0	0	0	0	0	0	0	0	0
CANADA GOOSE EGGS		1984	0	0	0	0	71	0	0	0	0
SNOW GOOSE EGGS		1984	0	0	0	0	10290	0	0	0	0
BRANT EGGS		1983	0	0	0	0	0	3	0	0	0
GOOSE EGGS		1984	0	0	0	0	30	0	0	0	0
		1985	0	0	0	0	3954	0	0	0	3954

Table 37. (cont'd)

ARCTIC CHARR (H,A)	1982	318	0	0	1504	882	1546	26	12	53	0	4341	542.6	8		
	1983	0	0	0	1066	1066	1066	140	616	366	57	2245	249.4	9		
	1984	300	12	2	210	108	370	278	690	29	1110	140	484	3733	311.1	12
	1985	0	202	331	0	138	1093	867	817	11	350	1281	182	5272	439.3	12
	1986	27	40	23	3	2	3	3	4	4	4	4	90	30.0	3	
N		4	5	4	3	2	3	3	4	4	4	4	.	.	.	
SUM		327	572	356	210	246	2967	2027	4119	206	2088	1840	723	.	.	.
MEAN		82	114	89	70	123	989	676	1030	52	522	460	181	.	.	.
LAKE TROUT (L,S)	1982	0	0	0	0	0	0	0	0	0	0	0	9	9	1.1	8
	1983	0	0	410	0	0	0	0	0	0	0	0	0	410	45.6	9
	1985	0	0	0	0	0	0	7	0	0	0	0	0	0	0.6	12
FRESHWATER FISH	1984	0	0	0	0	0	0	0	19	0	0	0	0	19	1.6	12
ARCTIC COD (L,A)	1982	0	0	0	0	0	18	0	0	0	0	0	0	18	2.3	8
	1984	0	0	0	0	0	150	16	4	0	0	0	0	170	14.2	12
	1986	0	8	0	0	0	0	0	0	0	0	0	0	8	2.7	3
SCULPINS (L,A)	1985	0	0	0	0	0	5	0	0	0	0	0	0	5	0.4	12

For footnotes see Appendix 4.

Table 38. The estimated harvest by Eskimo Point hunters, expressed as number of animals, for the period October, 1981 to March, 1986.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	\bar{X}	NUM		
CARIBOU KAMINURIAK ⁶	C ⁵	1981	1	3	0	0	0	0	0	3	27	91	0 ²	62	11	73	24.3	3		
		1982	0	0	0	0	1	2	2	1	20	16	24	3	164	13.7	12			
		1983	16	4	4	0	0	0	0	15	47	38	16	8	0	85	7.1	12		
		1984	0	0	2	2	0	0	0	0	32	24	36	6	145	12.1	12			
		1985	18	0	7	4	6	6	1	15	47	38	13	10	211	17.6	12			
		1986	9	6	42	1	1	1	1	1	32	24	63	13	57	19.0	3			
N		5	5	5	4	4	4	4	4	4	4	4	4	5	5	5	5			
SUM		43	10	53	7	9	4	4	36	85	182	133	143	30	30	30	30			
MEAN		9	2	11	2	2	1	1	9	21	46	27	29	6	6	6	6			
F		1981	1	1	1	1	1	1	1	1	35	121	226	78	350	62	647	215.7	3	
		1982	58	67	136	184	28	0	0	0	22	72	155	50	57	58	1048	87.3	12	
		1983	105	94	235	160	101	2	2	2	17	22	148	65	60	115	1306	109.8	12	
		1984	157	292	333	255	14	4	4	4	8	11	62	298	74	55	1422	118.5	12	
		1985	98	136	102	156	27	8	7	7	11	62	298	74	66	1045	87.1	12		
		1986	90	54	264	4	4	4	4	4	4	4	4	5	5	408	136.0	3		
N		5	5	5	5	4	4	4	4	4	4	4	4	5	5	5	5	5		
SUM		508	643	1070	755	170	14	81	226	591	871	591	591	356	356	356	356	356		
MEAN		102	129	214	189	43	4	20	57	148	174	174	174	71	71	71	71	71		
H		1981	1	1	1	1	1	1	1	1	10	265	302	337	196	121	80	397	132.3	3
		1982	102	131	90	84	30	25	98	136	128	128	128	31	19	1574	131.2	12		
		1983	23	10	46	34	36	25	67	151	67	234	23	44	33	737	61.4	12		
		1984	18	46	119	61	6	7	56	97	101	187	120	35	36	51	819	68.3	12	
		1985	44	43	27	54	54	56	56	56	56	56	56	32	32	552	71.0	12		
		1986	25	81	90	1	1	1	1	1	1	1	1	1	1	196	65.3	3		
N		5	5	5	5	4	4	4	4	4	4	4	4	5	5	5	5	5		
SUM		212	311	372	233	130	98	611	606	886	634	634	634	267	215	215	215	215		
MEAN		42	62	74	58	33	25	153	152	222	127	127	127	53	43	43	43	43		
U		1981	1	1	1	1	1	1	1	1	1	1	1	24	6	3	33	111.0	3	
		1982	25	18	32	28	12	0	1	1	23	19	10	22	9	23	336	28.0	12	
		1983	45	27	25	41	5	1	1	9	5	9	31	32	10	5	243	20.3	12	
		1984	18	17	13	18	12	9	14	17	14	8	0	2	2	43	179	14.9	12	
		1985	6	32	7	14	34	17	14	14	14	8	0	19	2	0	153	12.8	12	
		1986	0	0	4	4	4	4	4	4	4	4	4	5	5	4	4	4	3	
N		5	5	5	5	4	4	4	4	4	4	4	4	5	5	5	5	5		
SUM		94	94	81	101	63	27	53	116	117	99	99	99	29	29	29	29	29		
MEAN		19	19	16	25	16	7	13	29	29	29	29	29	6	6	6	6	6		
CARIBOU WAGER (H.S)	M	1983 ⁴	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

For footnotes see Appendix 4.

Table 38. (cont'd)

MOOSE (H,S)	1982	0	0	1	0	1	0.1	12
	1984	0	0	2	0	5	0.4	12
	1986	1	2	0	0	3	1.0	3
MUSKOX (H,S)	1985	0	1	0	0	1	0.1	12
	1986	0	2	0	0	2	0.7	3
POLAR BEAR (H,A)	1981	-	0	4	-	0	10	3
	1982	0	0	1	0	0	17	3
	1983	0	0	2	0	12	1.4	12
	1984	0	0	1	0	0	21	1.8
	1985	0	0	0	0	13	1.3	12
	1986	0	0	0	0	8	0.7	12
N	5	5	5	5	5	5	1.0	3
SUM	0	0	0	0	0	66	-	-
MEAN	0	0	0	0	0	11	-	-
ARCTIC FOX (H,A)	1981	-	0	0	-	0	137	75.7
	1982	40	58	48	172	90	227	3
	1983	210	62	80	27	469	2298	12
	1984	117	142	163	55	40	191.5	12
	1985	28	51	262	186	0	528	44.0
	1986	40	120	40	40	227	263	12
N	5	5	5	5	4	8	968	80.7
SUM	435	433	593	440	7	255	176	80.5
MEAN	87	87	119	110	2	0	200	66.7
RED FOX (H,S)	1981	-	0	0	0	0	0	3
	1982	0	0	2	6	0	13	3
	1983	5	0	17	9	2	35	12
	1984	1	0	1	0	0	0	1.3
	1985	0	2	7	1	7	16	12
	1986	2	7	1	1	0	39	3.3
N	5	5	5	4	4	0	0	12
SUM	8	25	15	14	0	0	0	3.3
MEAN	2	25	3	4	0	0	0	3
WOLF (H,S)	1982	0	0	0	0	0	0	12
	1983	0	0	2	22	0	10	2.1
	1984	0	0	0	12	0	25	12
	1985	0	2	12	30	0	31	2.6
	1986	6	2	3	32	2	50	4.2
N	6	6	53	5	5	0	10	12
SUM	6	4	70	96	5	0	51	20.3
MEAN	1	1	12	19	2	0	61	3
MARTIN (H,S)	1982	0	0	0	0	0	0	12

For footnotes see Appendix 4.

Table 38. (cont'd)

For footnotes see Appendix 4.

Table 38. (cont'd)

HARBOUR SEAL (H,S)	1981	0	2	0.7	3
	1982	0	1	0.1	12
	1983	0	3	0.3	12
	1984	1	2	0.2	12
HARP SEAL (H,S)	1982	0	0	0.4	12
	1983	0	0	0.5	12
	1984	0	2	0.2	12
	1985	0	2	0.2	12
UNKNOWN SEAL	1983	0	0	0	1
	1984	0	0	0	1
	1985	0	0	0	1
WALRUS (H,S)	1985	0	0	0	1
BELUGA (H,A)	1982	0	0	0	12
	1983	0	0	0	12
	1984	0	0	0	12
	1985	0	0	0	12
N	SUM	5	0	0	12
	MEAN	0	0	0	12
CANADA GEESE (H,A)	1982	0	0	4	12
	1983	0	0	4	12
	1984	0	0	4	12
	1985	0	0	4	12
N	SUM	5	0	0	12
	MEAN	0	0	0	12
SNOW GEESE (H,A)	1982	0	0	9	12
	1983	0	0	3	12
	1984	0	0	1	12
	1985	0	0	0	12
N	SUM	5	0	0	12
	MEAN	0	0	0	12
UNKNOWN GEESE	1984	0	0	0	0
	1985	0	0	0	0

For footnotes see Appendix 4.

Table 38. (cont'd)

OLD SQUAW (L,S)	1983	0	0	0	0	8	0	8	0.7	12
EIDER (L,S)	1981	0	0	0	0	0	0	0	0	3
	1983	0	0	0	0	0	0	0	19	12
	1984	0	0	0	0	0	0	0	4	12
	1985	0	0	0	0	0	0	0	0	0.3
N	5	5	5	4	4	4	5	5	0	0
SUM	0	0	0	0	0	9	10	0	1	0
MEAN	0	0	0	0	0	2	0	0	0	0
MALLARD (L,S)	1982	0	0	0	0	0	0	0	0	2
	1984	0	0	0	0	0	0	0	1	0.1
PTARMIGAN (L,A)	1981	14	18	15	48	26	0	12	32	3
	1982	4	7	0	2	39	16	6	11	191
	1983	5	0	20	111	7	0	0	16	12
	1984	2	0	0	40	72	8	0	1	190
	1985	0	76	24	1	1	0	0	45	15.8
	1986	5	5	5	4	4	4	4	0	0
N	5	101	59	201	144	24	18	24	129	323
SUM	25	12	50	36	36	6	5	46	82	26.9
MEAN	5	20	12	12	12	6	5	6	16	12
SNOWY OWL (L,A)	1981	0	0	0	0	0	0	0	0	0
SWAN (L,S)	1984	0	0	0	1	0	0	0	0	1
CANADA GOOSE EGGS	1984	0	0	0	0	384	0	0	0	0
SNOW GOOSE EGGS	1984	0	0	0	5	0	0	0	0	0
OLD SQUAW EGGS	1983	0	0	0	0	7	0	0	0	0
									7	0.6
										12

For footnotes see Appendix 4.

Table 38. (cont'd.)

GOOSE EGGS	1983	0	0	0	1112	0	0	0	1112	92.7	12	
1984	0	0	0	0	61	0	0	0	61	5.1	12	
1985	0	0	0	0	141	204	0	0	0	345	28.8	12
N	5	5	5	4	4	4	4	5	5	5	.	.
SUM	0	0	0	0	141	1377	0	0	0	0	.	.
MEAN	0	0	0	0	35	344	0	0	0	0	.	.
DUCK EGGS	1983	0	0	0	0	8	0	0	0	0	0.7	12
OTHER WATERFOWL EGGS	1983	0	0	0	0	7	1	0	0	0	0	0.7
ARCTIC CHARR (H,A)	1981	4	6	0	0	72	1080	1042	169	47	24	35.0
1982	3	0	0	0	61	364	562	694	258	57	12	2479
1983	0	0	0	2	163	172	1249	608	152	38	2095	206.6
1984	0	0	0	1	41	289	659	1779	357	109	68	174.6
1985	1	0	0	0	0	0	0	0	0	119	93	2601
1986	37	0	0	0	4	4	4	4	4	0	0	283.5
N	5	5	5	1	6	266	897	3550	4123	936	5	12.3
SUM	45	6	1	0	2	67	224	888	1031	234	347	3
MEAN	9	1	0	0	0	0	0	0	0	61	47	.
LAKE TROUT (H,A)	1981	12	79	182	237	38	9	65	75	879	128	686
1982	48	11	213	115	281	72	0	24	130	0	0	562.0
1983	0	0	4	183	62	138	8	6	111	159	231	68.4
1984	0	0	0	0	0	0	0	0	0	22	32	112.7
1985	4	0	0	0	0	0	0	0	0	75	59	568
1986	2	4	70	4	4	4	4	4	4	0	0	49.0
N	5	5	5	1	27	366	480	580	248	95	5	14.2
SUM	145	27	5	73	120	145	62	4	24	316	1072	12
MEAN	29	5	0	0	0	0	0	0	0	79	214	3
WHITEFISH (H,S)	1981	13	0	20	248	424	57	0	0	23	0	27
1982	0	0	0	0	0	0	0	0	0	0	0	356
1983	0	0	0	0	0	0	0	0	0	101	0	783
1984	0	0	0	0	0	0	0	0	0	0	2	65.4
1985	0	0	0	0	0	0	0	0	0	0	68	109
N	5	5	5	5	20	4	4	4	4	4	4	9.1
SUM	13	0	0	4	248	424	57	0	0	68	5	120
MEAN	3	0	0	4	62	106	14	0	0	17	0	10.0
										39	418	.
										84	.	.

For footnotes see Appendix 4.

Table 38. (cont'd)

ARCTIC GRAYLING (L,A)	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
N	5	5	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4
SUM	0	4	0	67	20	1	0	0	29	114	27	750	128	53	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11
MEAN	0	1	0	17	5	0	0	0	29	114	27	750	128	53	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11	11
NORTHERN PIKE (L,S)	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
FRESHWATER FISH	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ARCTIC COD (L,A)	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
SCULPIN (L,A)	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
SALTWATER FISH	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

For footnotes see Appendix 4.

Table 39. The estimated harvest by Rankin Inlet hunters, expressed as number of animals, for the period November, 1981 to March, 1986. The actual harvest was used as the best estimate of the harvest for August, 1982 and June, 1983.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	\bar{X}	NUM	
CARIBOU KAMINURIAK	C	5	1982	0	0	0	0	0	0	0	53	7	2	10	74	6.2	12		
(H,A)		6	1983	0	0	1	0	0	0	0	0	0	0	1	4	0.3	12		
			1984	0	4	0	0	0	0	0	2	0	0	0	6	0.5	12		
			1985	0	0	0	1	0	0	0	4	1	3	5	0	14	1.2	12	
			1986	6	11	7	4	4	4	4	4	4	4	4	24	8.0	3		
N	SUM	5	5	5	5	4	4	4	4	4	54	10	9	11	5	.	.		
	MEAN	1	6	15	8	1	0	0	0	0	2	14	3	2	2	.	.		
F	1981	3	1982	116	29	80	56	10	0	3	19	93	24	37	223	355	177.5	2	
			1983	65	57	85	42	80	2	0	27	20	49	41	69	506	42.2	12	
			1984	33	65	61	71	25	0	1	5	7	11	89	99	537	44.8	12	
			1985	132	65	37	114	38	25	0	15	13	67	97	31	634	38.9	12	
			1986	52	65	35	4	4	4	4	4	4	4	4	152	52.8	12		
N	SUM	5	398	281	298	283	153	27	4	66	133	151	38	38	461	50.7	3		
	MEAN	80	56	60	71	38	7	1	17	33	33	38	79	92	.	.	.		
M	1981	1	1982	28	114	107	163	127	7	74	163	320	91	43	94	137	68.5	2	
			1983	50	39	47	48	275	15	12	126	130	51	60	29	1260	105.0	12	
			1984	79	138	90	113	132	8	64	39	37	20	44	61	914	76.2	12	
			1985	100	47	78	103	106	14	57	181	124	55	25	68	832	69.3	12	
			1986	32	36	33	4	4	4	4	4	4	4	4	23	913	76.1	12	
N	SUM	5	289	374	355	427	640	44	4	207	509	611	217	209	101	33.7	3		
	MEAN	58	75	71	107	160	11	52	127	153	54	42	55	55	.	.	.		
U	1982	0	21	3	0	0	0	0	0	0	0	0	0	36	11	72	6.0	12	
			1983	0	0	2	0	0	0	9	2	25	6	0	26	70	5.8	12	
			1984	0	2	22	14	19	13	1	1	0	0	14	0	86	7.2	12	
			1985	9	0	0	0	0	12	4	4	0	0	0	0	4	33	2.8	12
			1986	3	1	1	1	1	1	1	1	1	1	1	1	4	5	1.7	3
N	SUM	5	12	24	28	14	9	5	4	4	4	4	4	4	42	25	31	6	
	MEAN	2	5	6	4	5	6	4	5	6	4	5	6	4	2	11	.	.	

For footnotes see Appendix 4.

Table 39. (cont'd)

CARIBOU N. CHEST.	C	1986 ⁴	0	0	8	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
(H,A)	F	1984	0	21	0	25	0	15	0	60	0	29	0	0	0	0	40	3.3	12	
	M	1986	0	0	15	0	0	0	3	0	0	0	0	0	0	0	25	8.3	3	
	M	1984	0	0	15	0	0	0	0	0	0	0	0	0	0	0	45	3.8	12	
	M	1985	0	0	60	0	0	0	0	0	0	0	0	0	0	0	67	5.6	12	
	M	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	29	9.7	3	
U	U	1984	0	0	0	2	-	-	-	-	-	-	-	-	-	-	0.6	12	3	
U	U	1986	0	0	0	0	-	-	-	-	-	-	-	-	-	-	0.7	3	3	
MUSKOX (H,S)		1985	3	0	0	0	-	-	-	-	-	-	-	-	-	-	0.3	12	3	
POLAR BEAR (H,S)		1981	-	0	1	1	0	1	1	0	0	0	0	0	0	0	5	2.5	2	
		1982	0	0	1	0	1	0	1	0	0	0	0	0	0	0	9	0.8	12	
		1983	0	0	1	0	1	0	1	0	0	0	0	0	0	0	20	1.7	12	
		1984	0	0	2	0	0	0	0	0	0	0	0	0	0	0	4	0.3	12	
		1985	0	0	5	5	5	4	4	4	0	0	0	0	0	0	10	1.8	12	
	N	SUM	0	0	4	3	3	1	1	5	0	0	0	0	0	0	5	-	-	
	MEAN		0	0	1	1	1	1	1	5	0	0	0	0	0	0	1	-	-	
ARCTIC FOX (H,A)		1981	-	6	3	12	0	1	0	0	0	0	0	0	0	0	7	3.5	2	
		1982	1	0	1	0	1	0	1	0	0	0	0	0	0	0	23	61.0	12	
		1983	26	21	56	20	10	0	0	0	0	0	0	0	0	0	33	13.4	12	
		1984	17	23	25	21	10	0	0	0	0	0	0	0	0	0	33	13.4	12	
		1985	0	0	5	5	5	4	4	4	0	0	0	0	0	0	52	11.2	12	
	N	SUM	70	77	22	9	4	4	4	5	0	0	0	0	0	0	119	14.8	12	
	MEAN		114	15	5	3	1	1	1	5	0	0	0	0	0	0	101	33.7	3	
WOLF (H,S)		1982	4	0	2	7	14	9	6	3	1	1	1	1	1	1	5	-	-	
		1983	2	0	2	3	3	4	3	4	4	4	4	4	4	4	14	1.2	12	
		1984	0	0	2	2	1	1	1	1	1	1	1	1	1	1	9	2.7	12	
		1985	1	1	2	2	1	1	1	1	1	1	1	1	1	1	9	0.8	12	
	N	SUM	2	2	5	5	5	5	5	5	5	5	5	5	5	5	0	1.4	12	
	MEAN		114	15	101	33	20	8	18	30	30	30	30	30	30	30	0	7	2.3	3
WOLVERINE (H,S)		1983	0	0	0	1	0	0	0	0	0	0	0	0	0	0	9	0.8	12	
		1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	1	0.1	12	

For footnotes see Appendix 4.

Table 39. (cont'd)

For footnotes see Appendix 4.

Table 39. (cont'd)

WALRUS	(H,S)	1982	0	0	2	0.2	12
		1983	0	0	47	3.9	12
		1984	0	0	1	0.1	12
		1985	0	0	5	0.4	12
		1986	0	0	2	0.7	3
BELUGA	(H,A)	1982	0	0	34	2.8	12
		1983	0	0	29	2.4	12
		1984	0	0	69	5.8	12
		1985	0	0	47	3.9	12
N		5	0	0	5	·	·
SUM		0	5	4	4	·	·
MEAN		0	0	2	0	·	·
CANADA GEESE	(L,S)	1982	0	0	0	0	12
		1983	0	0	20	1.7	12
		1984	0	0	401	33	12
		1985	0	0	315	26.3	12
N		5	5	4	4	·	·
SUM		0	0	10	0	·	·
MEAN		0	0	2	0	·	·
SNOW GEESE	(L,S)	1982	0	0	41	0	12
		1983	0	0	18	0	12
		1984	0	0	41	0	12
		1985	0	0	0	0	12
N		5	5	4	4	·	·
SUM		0	0	2	100	0	0
MEAN		0	0	1	0	0	0
BRANT GEESE	(L,S)	1984	0	0	0	0	12
UNKNOWN GEESE		1982	0	0	0	0	12
		1985	0	0	3	0	12
SANDHILL CRANE	(L,S)	1982	0	0	0	0	12
		1984	0	0	0	0	12

For footnotes see Appendix 4.

Table 39. (cont'd)

Table 39. (cont'd)

LANDLOCKED ARCTIC CHAR (L,A)		1985	0	0	0	0	0	0	0	32	0	0	32	2.7	12		
LAKE TROUT (H,A)		1981	15	78	59	0	0	0	0	0	0	0	33	16.5	2		
		1982	0	20	0	289	0	0	0	0	0	45	0	16.4	12		
		1983	0	20	0	120	232	0	10	21	0	54	0	30.3	12		
		1984	0	20	0	14	0	173	127	0	15	0	103	0	42.2	12	
		1985	1	16	0	0	0	0	0	0	0	0	2	348	29.0	12	
		1986	16	0	5	4	4	4	4	4	4	5	16	5.3	3		
N		5	5	92	179	694	127	10	21	15	0	202	35	0	0		
SUM		17	71	18	45	174	32	3	5	4	0	40	7	0	0		
MEAN		3	14														
WHITEFISH (H,S)		1983	0	0	0	0	0	0	0	0	0	0	7	0	0.6	12	
		1984	0	0	0	1	0	0	0	0	0	0	9	0	0.8	12	
ARCTIC GRAYLING (L,A)		1982	0	0	0	0	0	0	0	0	0	0	0	10	0.8	12	
		1984	0	0	0	0	0	0	0	0	0	49	0	49	4.1	12	
FRESHWATER FISH		1982	0	0	0	30	117	0	0	0	0	0	0	147	12.3	12	
		1983	0	0	0	104	0	0	0	0	0	0	0	104	8.7	12	
ARCTIC COD (L,A)		1985	0	0	0	0	0	0	12	0	0	0	0	0	12	1.0	12
SCULPINS (L,A)		1985	0	0	0	0	0	0	0	0	0	0	0	13	1.1	12	
SALTWATER FISH		1983	0	0	0	52	0	0	0	0	0	0	0	52	4.3	12	

For footnotes see Appendix 4.

Table 40. The estimated harvest by Repulse Bay hunters, expressed as number of animals for the period October, 1981 to March, 1986. Data were not available for December, 1982. The best estimate of the harvest for December 1981, March 1982, and April 1983 was the actual harvest.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	\bar{x}	NUM
CARIBOU KAMINURIAK		F	1984	4	2	0	0	0	6	0	0	0	0	0	0	6	0.5	12
(H,S)	M	1982	3	0	0	0	0	0	0	0	0	0	0	0	3	0.3	11	
	1983	0	0	0	0	0	0	0	2	0	0	0	0	0	2	0.2	12	
	1984	0	0	0	0	0	0	0	34	0	0	0	0	0	34	2.8	12	
CARIBOU WAGER (H,A)	U	1984	0	0	0	0	1	0	0	0	0	0	0	0	1	0.1	12	
C	1982	10	0	0	0	0	0	0	9	24	9	0	0	0	52	4.7	11	
	1983	0	0	0	0	0	0	0	0	7	6	0	0	0	16	1.3	12	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	5	0.4	12	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	59	4.9	12	
	1986	2	3	10	4	4	4	4	4	4	4	4	4	4	15	5.0	3	
N	SUM	5	5	10	0	0	0	0	9	56	45	3	7	2	·	·	·	
	MEAN	2	3	2	0	0	0	0	2	14	11	1	1	1	·	·	·	
F	1981	3	·	·	·	·	·	·	·	·	·	·	·	·	22	137	45.7	3
	1982	46	74	17	44	10	0	0	3	90	25	24	53	386	35.1	11		
	1983	41	32	36	28	9	8	3	29	15	24	11	17	253	21.1	12		
	1984	20	47	60	20	6	9	7	79	36	25	72	38	419	34.9	12		
	1985	48	79	72	55	13	2	4	24	61	26	22	12	417	34.8	12		
	1986	59	42	56	·	·	·	·	·	·	·	·	·	157	52.3	3		
N	SUM	5	5	5	4	4	4	4	4	4	4	4	4	4	·	·	·	
	MEAN	214	274	241	147	38	19	17	222	137	164	56	34	207	89	·	·	
F	1981	3	·	·	·	·	·	·	·	·	·	·	·	·	22	·	·	
	1982	46	74	17	44	10	0	0	3	90	25	24	53	386	35.1	11		
	1983	41	32	36	28	9	8	3	29	15	24	11	17	253	21.1	12		
	1984	20	47	60	20	6	9	7	79	36	25	72	38	419	34.9	12		
	1985	48	79	72	55	13	2	4	24	61	26	22	12	417	34.8	12		
	1986	59	42	56	·	·	·	·	·	·	·	·	·	157	52.3	3		
N	SUM	5	5	5	4	4	4	4	4	4	4	4	4	4	·	·	·	
	MEAN	214	274	241	147	38	19	17	222	137	164	56	34	207	89	·	·	
M	1981	·	·	·	·	·	·	·	·	·	·	·	·	·	19	167	55.7	3
	1982	34	30	25	42	42	22	22	192	88	16	46	46	685	62.3	11		
	1983	70	27	43	33	41	62	63	68	81	52	9	9	17	566	47.2	12	
	1984	15	65	39	34	30	91	59	194	94	69	49	29	768	64.0	12		
	1985	18	46	48	48	75	44	125	121	141	51	44	31	792	66.0	12		
	1986	44	83	31	·	·	·	·	·	·	·	·	·	158	52.7	3		
N	SUM	5	5	5	4	4	4	4	4	4	4	4	4	4	·	·	·	
	MEAN	181	251	186	164	188	338	269	575	404	285	199	96	24	·	·	·	
F	1981	36	50	37	41	47	85	67	144	101	57	40	40	·	·	·	·	
	1982	0	5	14	3	0	8	22	6	0	0	0	0	0	58	5.3	11	
	1983	0	0	6	0	0	0	0	5	1	16	4	1	0	39	3.3	12	
	1984	5	12	37	46	0	10	5	1	15	4	0	0	0	140	11.7	12	
	1985	1	0	4	0	0	0	15	7	15	4	0	0	0	46	3.8	12	
	1986	0	0	4	·	·	·	·	·	·	·	·	·	0	4	1.3	3	
N	SUM	5	5	4	4	4	4	4	4	4	4	4	4	4	4	5	1	
	MEAN	17	65	49	0	33	34	22	20	23	15	3	3	3	4	5	3	
U	1982	0	5	14	3	0	8	22	6	0	0	0	0	0	58	5.3	11	
	1983	0	0	6	0	0	0	0	5	1	16	4	1	0	39	3.3	12	
	1984	5	12	37	46	0	10	5	1	15	4	0	0	0	140	11.7	12	
	1985	1	0	4	0	0	0	15	7	15	4	0	0	0	46	3.8	12	
	1986	0	0	4	·	·	·	·	·	·	·	·	·	0	4	1.3	3	
N	SUM	5	5	4	4	4	4	4	4	4	4	4	4	4	4	5	1	
	MEAN	1	3	13	12	0	8	9	12	12	12	12	12	12	4	5	1	

Table 40. (cont'd)

CARIBOU SOUTHAMPTON (H,S)	U	1985	0	0	0	0	0	0	1	0	0	0	1	0.1	12	
CARIBOU BEVERLY (H,S)	F	1981	5	2	0	7	2.3	3		
CARIBOU N. CHEST. (H,S)	M	1981	3	0	2	5	1.7	3		
CARIBOU OTHER (H,S)	M	1984	0	0	0	0	23	0	0	0	0	23	1.9	12		
MUSKOX (H,S)	U	1984	0	0	0	0	7	0	0	0	0	7	0.6	12		
POLAR BEAR (H,S)	1981	0	0	1	0	0	0	0	0	0	0	15	1	16	5.3	3
	1982	0	0	1	0	0	0	0	0	0	0	15	1	15	1.4	11
	1983	0	2	2	0	0	0	0	0	0	0	8	2	13	1.1	12
	1984	0	3	0	0	0	0	0	0	0	0	4	1	9	0.8	12
	1985	2	4	0	4	0	0	0	0	0	0	8	0	11	0.9	12
	1986	5	5	4	4	4	4	4	4	4	4	5	4	6	2.0	3
N	SUM	2	10	3	1	0	0	0	0	0	0	50	4	1
MEAN	0	2	1	0	0	0	0	0	0	0	0	10	1
BLACK BEAR (L,S)	1983	0	1	0	0	0	0	0	0	0	0	0	0	1	0.1	12
GRIZZLY BEAR (H,S)	1982	0	0	5	0	0	0	5	0	0	0	0	0	5	0.5	11
N	SUM	5	5	4	4	4	4	4	4	4	4	5	5	4
MEAN	0	0	0	0	0	0	0	0	0	0	0	0	0	0

For footnotes see Appendix 4.

Table 40. (cont'd.)

ARCTIC FOX	(H,A)	1981		1982		1983		1984		1985		1986		N		SUM		MEAN	
		0	3	0	3	0	3	0	3	0	3	0	3	0	3	0	3	0	3
RED FOX	(H,S)	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1984	0	2	0	3	0	3	0	3	0	3	0	3	0	3	0	3	0
		1985	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0	1	0
		1986	0	5	0	5	0	5	0	5	0	5	0	5	0	5	0	5	0
		N	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
		SUM	115	97	93	70	18	19	19	19	19	19	19	19	19	19	19	19	19
		MEAN	23	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19	19
WOLF	(H,S)	1982	5	18	0	4	18	2	18	2	10	4	13	3	13	3	13	3	13
		1983	6	0	0	0	0	0	0	0	9	0	4	0	4	0	4	0	4
		1984	2	4	0	3	0	3	0	3	0	3	0	3	0	3	0	3	0
		1985	2	10	0	9	0	9	0	9	0	9	0	9	0	9	0	9	0
		1986	4	13	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
		N	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
		SUM	19	45	9	8	18	5	42	8	18	5	24	6	24	6	24	6	24
		MEAN	4	9	1	2	3	1	5	2	3	1	4	1	4	1	4	1	4
WOLVERINE	(H,S)	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
ARCTIC HARE	(L,S)	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
		N	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
		SUM	0	9	2	1	5	1	4	1	5	1	4	1	4	1	4	1	4
		MEAN	0	1.8	0.4	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0

For footnotes see Appendix 4.

Table 40. (cont'd)

Table 40. (cont'd)

CANADA GEESE (L,S)	1983	0	0	0	0	3	0	0	0	0	2
	1984	0	0	0	0	7	1	0	0	0	7
	1985	0	0	0	0	12	1	0	0	0	23
SNOW GEESE (L,S)	1982	0	0	0	0	19	8	0	0	0	27
	1984	0	0	0	0	4	0	0	0	4	2.5
ROSS'S GEESE (L,S)	1982	0	0	0	0	3	5	0	0	0	8
	1983	0	0	0	0	0	0	10	0	0	10
UNKNOWN GEESE	1985	0	0	0	0	4	0	0	0	0	0
SANDHILL CRANE (L,S)	1984	0	0	0	0	1	0	0	0	0	0
OLD SQUAW (L,S)	1984	0	0	0	0	6	0	0	0	0	6
GUILLEMOT (L,S)	1982	0	0	0	0	0	6	3	0	0	9
EIDER (L,S)	1982	0	0	0	0	0	0	9	0	0	9
	1983	0	0	0	0	0	0	15	7	0	22
	1984	0	0	0	0	0	0	0	0	0	5
	1985	0	0	0	0	0	0	0	0	0	4

For footnotes see Appendix 4.

Table 40. (cont'd)

PTARMIGAN (L.S.)	1982	0	0	0	96	29	0	0	117	10	0	0	252	22.9	11		
	1983	0	0	0	5	0	4	0	6	61	0	4	0	86	0.3	12	
	1984	0	6	0	0	5	1	0	26	13	0	0	0	67	7.2	12	
	1985	2	0	0	0	20	5	0	0	0	0	0	0	3	5.6	12	
	1986	3	0	0	4	4	4	4	4	4	5	5	4	3	1.0	3	
N	5	5	5	5	5	119	38	1	6	204	23	4	0	0	.	.	
SUM	5	6	0	1	30	10	0	2	51	5	1	0	0	.	.	.	
MEAN	1	1	0	1	30	10	0	2	51	5	1	0	0	.	.	.	
UNKNOWN FOWL	1983	0	0	0	0	0	0	7	0	0	0	0	0	7	0.6	12	
ARCTIC CHARR (H.A.)	1981	0	23	0	157	313	47	390	29	542	418	1919	283	705	235.0	3	
	1982	0	14	0	0	63	76	109	2	147	564	52	1027	174.5	11		
	1983	0	0	0	0	14	655	265	9	631	426	350	2843	85.6	12		
	1984	0	0	0	30	195	501	1017	749	0	27	639	0	3418	236.9	12	
	1985	260	0	0	63	0	0	0	0	0	0	0	0	284.8	12		
	1986	0	0	5	5	4	4	4	4	4	5	4	0	63	21.0	3	
N	5	5	0	100	30	366	1532	1405	1741	40	1480	2336	685	0	.	.	
SUM	260	0	0	20	8	92	383	351	435	10	296	467	171	0	.	.	
MEAN	52	0	0	20	8	92	383	351	435	10	296	467	171	0	.	.	
LAKE TROUT (H.A.)	1981	0	1	15	199	345	0	0	11	0	5	0	0	750	250.0	3	
	1982	75	0	8	46	8	2	0	0	0	2	0	0	651	59.2	11	
	1983	0	0	0	0	60	0	0	0	0	0	0	0	0	66	5.5	12
	1984	0	0	0	0	68	115	8	0	0	0	0	0	225	285	12	
	1985	0	0	0	4	4	4	4	4	4	5	4	1	0	208	17.3	12
	1986	0	0	5	5	4	4	4	4	4	5	4	0	4	1.3	3	
N	5	5	0	13	61	335	462	8	0	11	763	235	1	0	.	.	
SUM	75	0	0	13	61	335	462	8	0	3	153	47	0	0	.	.	
MEAN	15	0	0	3	15	84	116	2	0	1	763	235	1	0	.	.	
ARCTIC GRAYLING (L.A.)	1982	0	0	0	0	0	0	0	0	14	0	0	0	0	14	1.3	11
FRESHWATER FISH	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	216	0	0

For footnotes see Appendix 4.

Table 41. The estimated harvest by Whale Cove hunters, expressed as number of animals for the period October 1981 to March 1983 and November 1983 to March 1986. The best estimate of the harvest for October, 1981, 1982, 1983, and November 1983 and December 1983 was the actual reported harvest.

SPECIES	CATEGORY	SEX	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC	SUM	\bar{X}	NUM	
CARIBOU KAMINURIAK	5	C	1982	2	0	0	1	0	0	0	13	36	0	2	0	52	4.3	12	
(H,A)	6	C	1984	0	0	0	0	0	0	0	0	0	0	4	1	5	0.4	12	
		C	1985	7	1	0	0	0	1	1	1	1	0	0	2	14	1.2	12	
		C	1986	9	4	4	3	3	3	3	3	4	5	5	17	5.7	3		
N			SUM	5	5	5	3	3	3	3	3	4	5	5	17	5.7	3		
			MEAN	16	5	4	1	0	1	14	37	0	6	1	1	1	1	1	
F			1981	3	1	1	0	0	0	0	0	0	0	0	0	0	0	0	
			1982	39	88	58	57	27	0	4	5	34	30	71	91	192	64.0	3	
			1983	18	32	65	11	11	10	0	0	13	6	36	54	395	32.9	12	
			1984	10	32	77	38	11	10	0	0	0	6	33	15	166	33.2	5	
			1985	97	44	91	46	20	3	5	13	0	32	29	33	58	288	24.0	12
			1986	73	43	41	41	1	1	1	1	1	1	1	1	33	413	34.4	12
N			SUM	5	5	5	3	3	3	3	3	3	4	5	5	157	52.3	3	
			MEAN	237	239	332	141	58	13	9	18	47	68	198	251	50	50	50	
M			1981	47	48	66	47	19	4	3	6	16	17	40	40	40	12	40	
			1982	29	42	44	52	65	0	19	97	101	12	12	12	40	13.3	3	
			1983	47	11	15	16	22	18	12	17	21	22	28	23	34	523	43.6	12
			1984	52	46	52	52	7	52	28	46	45	42	30	17	10	111	22.2	5
			1985	11	11	20	7	7	52	52	52	45	42	30	14	6	301	25.1	12
			1986	20	6	7	3	3	3	3	3	3	3	3	3	10	316	26.3	12
N			SUM	5	5	5	3	3	3	3	3	3	3	3	3	3	33	11.0	3
			MEAN	159	116	138	75	139	46	77	159	164	81	55	20	20	20	20	20
U			1982	0	0	4	0	0	0	0	0	0	0	0	0	0	6	35	2.9
			1983	5	0	20	0	0	0	0	0	0	0	0	0	2	34	6.8	5
			1984	0	0	0	0	0	0	0	0	0	0	0	0	1	1	0.1	12
			1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	12
			1986	3	0	1	0	0	0	0	0	0	0	0	0	0	0	0	1.0
N			SUM	5	5	5	3	3	3	3	3	3	3	3	3	3	4	1.3	3
			MEAN	8	0	25	9	1	0	0	2	25	0	0	0	0	7	9	2

For footnotes see Appendix 4.

Table 41. (cont'd)

CARIBOU WAGER (H,S)	M	1981 ⁴	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012
MUSKOX (H,S)																																	
POLAR BEAR (H,A)																																	
	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
N		5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	
SUM		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
MEAN		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
BLACK BEAR (L,S)																																	
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
ARCTIC FOX (H,A)																																	
	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1983	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32	32		
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1986	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	
N		5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	
SUM		34	10	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30	30			
MEAN		7	2	6	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
RED FOX (H,S)																																	
	1983	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
WOLF (H,S)																																	
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
ARCTIC HARE (L,S)																																	
	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1982	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1983	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1985	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
N		5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	
SUM		7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	
MEAN		1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	

For footnotes see Appendix 4.

Table 41. (cont'd.)

RINGED SEAL (H,A)	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1983	10	11	8	5	5	21	29	13	6	14	29	40	40	28	200	167	12	113	0	32	64	5	142	118	12	3	113	3		
	1984	6	6	5	4	5	21	29	13	6	14	29	40	40	28	200	167	12	12	0	32	64	5	142	118	12	3	113	3		
	1985	9	11	16	9	9	34	17	36	16	1	10	1	1	1	6	166	138	12	12	12	19	0	5	0	0	0	0	0	0	0
	1986	7	3	4	3	3	3	3	3	3	3	3	3	3	3	3	54	54	5	5	5	5	5	5	5	5	5	5	5	5	5
N		5	5	5	5	5	49	22	84	66	52	53	23	47	50	45	45	45	45	45	45	45	45	45	45	45	45	45	45	45	
SUM		32	35	49	10	7	28	22	17	18	8	12	10	9	10	9	9	9	9	9	9	9	9	9	9	9	9	9	9	9	
MEAN		6	6	7	10	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	7	
BEARDED SEAL (H,S)	1982	0	1	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1984	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HARBOUR SEAL (H,S)	1982	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
HARP SEAL (H,S)	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
WALRUS (H,S)	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
BELUGA (H,A)	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
N		5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
SUM		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MEAN		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
NAR		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
NARWHAL (H,S)	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
CANADA GEESE (L,S)	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
N		5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5	5
SUM		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
MEAN		0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

For footnotes see Appendix 4.

Table 41. (cont'd)

SNOW GEESE (L,S)	1982	0	109	40	0	0	0	0	149	12.4
	1984	0	226	273	37	0	29	0	540	45.0
	1985	0	55	0	0	3	3	5	84	7.0
N	5	5	3	3	3	3	4	5	·	12
SUM	0	0	390	313	37	0	33	0	·	·
MEAN	0	0	130	104	12	0	11	0	·	·
ROSS'S GEESE (L,S)	1982	0	0	0	0	0	0	0	0	0.2
UNKNOWN GEESE	1985	0	0	0	338	197	1	0	62	0
EIDER (L,S)	1981	·	·	·	·	·	·	1	0	0.3
	1982	0	0	0	0	0	0	0	0	3
	1984	0	0	0	0	0	0	0	8	0.7
	1985	0	3	0	0	1	0	0	17	1.4
PTARMIGAN (L,A)	1981	·	·	·	·	·	·	1	0	0.3
	1982	0	0	0	0	0	0	0	0	3
	1983	0	9	0	0	0	0	0	21	1.8
	1984	0	0	5	7	0	0	0	9	5
	1985	0	10	0	0	0	0	0	12	1.1
CANADA GOOSE EGGS	1985	0	0	0	0	2	0	0	0	0.2
GOOSE EGGS	1984	0	0	0	24	599	0	0	0	12
	1985	0	0	0	124	0	0	0	723	60.3
ARCTIC CHARR (H,A)	1981	·	·	·	·	·	·	42	11	3
	1982	111	42	79	58	86	630	6583	69	219
	1983	2	25	0	7	3	261	467	63	673.6
	1984	23	0	0	56	76	117	416	117	12
	1985	4	5	0	0	0	3	3	30	24.0
	1986	40	20	3	5	3	3	3	16	58.5
N	5	5	5	3	3	3	3	3	6	12
SUM	180	92	82	65	145	496	1008	7466	5	21.0
MEAN	36	18	16	22	48	165	336	2489	54	3

For footnotes see Appendix 4.

Table 41. (cont'd)

LAKE TROUT (H.A.)	1981	12	14	71	52	15	23	120	3	4	33	156	193	64.3	3
1982	59	18	14	92	93	80	18	0	0	0	26	58	468	39.0	12
1983	11	0	0	0	0	6	52	44	7	1	0	9	0	85	17.0
1984	17	14	92	93	92	80	18	0	0	3	0	0	0	323	26.9
1985	0	0	0	0	0	6	52	44	7	1	0	0	0	113	9.4
1986	10	9	0	5	5	3	3	3	3	3	0	0	0	19	6.3
N	5	5	124	170	184	77	30	121	3	4	5	5	5	19	3
SUM	97	91	25	57	61	26	10	40	1	6	14	43	43	11	12
MEAN	19	18													
WHITEFISH (H.S.)	1982	0	0	0	0	0	0	0	0	18	15	43	76	6.3	12
ARCTIC GRAYLING (L.A.)	1981	0	0	0	5	0	0	0	0	0	0	0	0	2	0.7
1985	0	0	0	0	0	0	0	0	0	0	0	0	0	5	0.4
NORTHERN PIKE (L.S.)	1982	0	0	0	0	2	0	0	0	0	0	0	0	2	0.2
FRESHWATER FISH	1982	0	0	3	8	0	0	0	0	0	0	0	0	11	0.9
SALTWATER FISH	1982	0	0	0	0	6	0	0	0	0	0	0	0	6	0.5

For footnotes see Appendix 4.

Table 42. The estimated harvest of Kaminuriak caribou by hunters from the seven Keewatin communities expressed as numbers of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	X	NUM	
BAKER LAKE	1981	608	462	330	241	278	10	453	.	496	.	295	266	561	280.5	2	
	1982	286	0	71	334	161	0	241	216	412	92	239	110	3227	322.7	10	
	1983	0	0	93	0	51	311	63	33	0	2	41	79	1933	193.3	10	
	1984	0	0	226	112	87	0	78	0	165	171	155	0	563	92.2	6	
	1985	0	0	81	78	1072	134.0	8	
CORAL HARBOUR	1984	0	0	4	0	0	0	0	0	0	0	0	0	81.0	1		
	1985	0	0	0	9	0	0	0	0	0	0	0	0	81.0	1		
CHESTERFIELD INLET	1981	4	17	0	0	0	0	0	0	5	2	0	0	0	0.0	0	
	1982	17	16	0	0	40	0	0	0	0	0	1	17	61	8.7	7	
	1983	31	0	1	0	2	0	0	0	0	0	0	26	106	21.2	5	
	1984	0	9	30	3	2	0	0	0	0	0	11	10	0	67	7	
	1985	0	9	30	3	2	0	0	0	0	0	0	0	0	0.0	0	
ESKIMO POINT	1981	185	216	258	296	70	10	314	530	730	289	121	103	3122	260.2	12	
	1982	189	135	310	236	146	30	144	238	313	365	112	153	2371	197.6	12	
	1983	193	355	465	336	33	20	173	113	460	128	134	155	2565	213.8	12	
	1984	166	211	143	228	123	83	150	152	273	500	124	108	2261	188.4	12	
	1985	124	141	400	665	221.7	3	
REPULSE BAY	1982	3	0	0	0	0	0	0	0	0	0	0	0	0	0.0	1	
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	1	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	1	
RANKIN INLET	1981	144	164	190	219	137	7	77	184	466	158	87	79	1912	159.3	12	
	1982	115	96	135	90	355	17	21	155	175	106	103	157	1525	127.1	12	
	1983	112	209	173	198	176	21	66	47	44	31	147	167	1391	115.9	12	
	1984	241	112	115	218	144	51	61	204	138	125	127	58	1594	132.8	12	
	1985	93	113	76	282	94.0	3	
WHALE COVE	1981	130	106	110	92	.	0	23	140	171	42	87	103	232	77.3	3	
	1982	70	43	100	54	33	.	.	12	17	54	71	27	311	62.2	5	
	1983	62	78	129	54	33	28	12	17	34	28	54	66	595	49.6	12	
	1984	115	56	111	62	73	32	54	59	43	62	43	45	755	62.9	12	
	1985	105	53	53	211	70.3	3	
KEEWATIN	1981	1012	989	884	866	577	27	867	860	1864	464	497	1096	842	2435	811.7	3
	1982	677	290	616	660	662	52	406	614	902	563	528	386	9324	777.0	12	
	1983	398	642	772	681	323	120	562	248	571	187	328	433	6203	516.9	12	
	1984	522	388	625	632	429	166	345	415	619	869	459	289	5758	437.9	12	
	1985	322	307	610	1239	413.0	3	

For footnotes see Appendix 4.

Table 43. The estimated harvest of Beverly caribou by Baker Lake and Repulse Bay hunters expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM	
BAKER LAKE	1981	1												2		3	
	1982	11	32	44	56	9	0	10		19			102	0	102.0	1	
	1983	0	362	251	258	319	0	184	47	92	151	467	0	222	27.8	8	
	1984	251	593	749	658	60	0	77	0	437	569	0	340	230.6	230.6	10	
	1985	460	382	67	30	0	0	0	0	363	341	126	0	1769	3734	414.9	9
REPULSE BAY	1981															252.7	7
	1982																
KEEWATIN	1981																
	1982	11	32	44	56	9	0	10		19			104	2	114	38.0	3
	1983	0	362	251	258	319	0	184	47	92	151	467	0	222	27.8	8	
	1984	251	593	749	658	60	0	77	0	437	569	0	340	230.6	230.6	10	
	1985	460	382	67	30	0	0	0	0	363	341	126	0	1769	3734	414.9	9
																252.7	7

For footnotes see Appendix 4.

Table 44. The estimated harvest of Wager Bay caribou by Baker Lake, Coral Harbour, Eskimo Point, Repulse Bay and Whale Cove hunters, expressed as numbers of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM	
BAKER LAKE	1982	0	0	0	0	0	0	0	248	480	31	0	0	6	6.0	3	
	1983	0	0	0	0	401	192	134	505	631	279	8	68	1428	204.0	7	
	1984	2	0	0	4	508	154	511	505	560	207	35	2763	276.3	10		
	1985	0	15	0	97	90	134	237	560	848	35	287	2510	251.0	10		
	1986	349	424	418	1191	397.0	3			
CORAL HARBOUR	1984	0	0	0	0	193	0	0	0	12	0	0	0	0	205	102.5	2
	1985	0	0	0	0	84	126	0	0	0	0	0	0	0	210	105.0	2
ESKIMO POINT	1983	0	0	0	0	0	0	0	0	1	0	0	0	0	1	1.0	1
REPULSE BAY	1981	90	109	56	96	52	149	56	312	122	40	99	41	304	101.3	3	
	1982	59	85	61	50	70	66	66	104	102	95	37	1181	107.4	11		
	1983	111	124	136	100	36	110	71	278	146	98	122	34	874	72.8	12	
	1984	40	125	103	88	61	136	161	236	79	70	44	71	1332	111.0	12	
	1985	67	105	128	101	44	1314	109.5	12	
	1986												334	111.3	3		
WHALE COVE	1981	4	0	0	4	4.0	1	
											167	100	41	308	102.7	3	
KEEWATIN	1981	40	99	0	1187	107.9	11	
	1982	90	109	56	96	52	149	56	312	128	583	126	45	102	2303	191.9	12
	1983	111	59	85	61	50	471	258	352	651	729	401	106	4300	358.3	12	
	1984	42	124	136	297	544	264	205	801	1084	286	105	331	4034	336.2	12	
	1985	67	140	124	284	304	195	373	741	.	.	.	1625	508.3	3		
	1986	454	552	519		

For footnotes see Appendix 4.

Table 45. The estimated harvest of polar bears by hunters from the seven Keeewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM		
BAKER LAKE	1985	0	2	1	0	0	0	0	0	0	0	0	0	1	1.0	1		
CORAL HARBOUR	1982	1	0	0	0	0	0	2	0	0	0	1	6	5	14	3.5	4	
	1983	0	0	0	0	0	0	0	0	0	2	20	9	1	36	7.2	5	
	1984	4	0	0	0	0	0	0	0	0	0	32	9	0	45	15.0	3	
	1985	1	0	3	4	0	0	0	0	0	1	11	35	3	61	8.7	7	
	1986	2	1	2	0	0	0	0	0	0	0	0	0	5	5	1.7	3	
CHESTERFIELD INLET	1982	3	0	1	7	0	0	0	0	0	0	0	0	1	4	2.0	2	
	1983	2	0	0	4	0	0	0	0	0	0	0	0	0	14	3.5	4	
	1984	1	0	1	2	0	0	0	0	0	0	0	0	0	5	2.5	2	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	8	1.6	5	
ESKIMO POINT	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	10	10.0	1	
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	17	8.5	2	
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	21	5.3	4	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	15	5.0	3	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	6	6.0	1	
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	3	3.0	1	
REPULSE BAY	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	15	1.0	2	
	1982	0	0	0	1	0	0	0	0	0	0	0	0	0	15	15.0	1	
	1983	0	0	1	2	0	0	0	0	0	0	0	0	0	2	13	2.6	5
	1984	0	0	2	2	0	0	0	0	0	0	0	0	0	4	1.0	4	
	1985	0	0	3	4	0	0	0	0	0	0	0	0	0	9	2.3	4	
	1986	2	0	0	0	0	0	0	0	0	0	0	0	0	0	5.5	2	
RANKIN INLET	1981	0	0	0	1	1	0	0	0	0	0	0	0	0	3	2	2.5	2
	1982	0	0	0	1	1	0	0	0	0	0	0	0	0	3	0	3	0
	1983	0	0	0	1	1	0	0	0	0	0	0	0	0	3	0	3	0
	1984	0	0	1	0	0	0	0	0	0	0	0	0	0	4	1.0	4	1
	1985	0	0	2	0	0	0	0	0	0	0	0	0	0	0	6	3.0	2
	1986	2	4	0	0	0	0	0	0	0	0	0	0	0	0	6	3.0	2
WHALE COVE	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1982	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
KEEWIN	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
	1982	3	0	1	8	0	0	2	0	0	0	0	0	0	2	41	8	65
	1983	2	4	9	12	0	0	4	2	0	0	0	0	0	27	41	6	109
	1984	5	3	10	0	2	1	0	0	0	0	0	0	0	32	34	1	88
	1985	1	6	4	13	3	0	0	0	0	0	0	0	0	12	73	3	121
	1986	4	5	5	5	5	5	5	5	5	5	5	5	5	5	14	4.7	3

For footnotes see Appendix 4.

Table 46. The estimated harvest of Arctic fox by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	X	NUM						
BAKER LAKE	1981	1	50	71	39	0	0	0	0	0	0	0	0	13	13.0	1						
	1982	0	50	0	0	0	0	0	0	0	0	0	417	149	726	145.2	5					
	1983	30	7	180	5	0	0	0	0	0	0	0	17	275	329	82.3	4					
	1984	124	156	216	113	0	0	0	0	0	0	0	526	1090	2081	346.8	6					
	1985	758	585	75	0	0	0	0	0	0	0	0	497	250	2419	403.2	6					
	1986	65	152	0	0	0	0	0	0	0	0	0	0	292	97	97.3	3					
CORAL HARBOUR	1982	117	50	224	17	0	0	0	0	0	0	0	166	167	103	486	121.5	4				
	1983	85	55	89	123	0	0	0	0	0	0	0	21	0	126	739	92.4	8				
	1984	29	115	76	74	12	0	0	0	0	0	0	0	0	123	584	97.3	6				
	1985	13	2	35	0	0	0	0	0	0	0	0	0	0	17	69	392	56.0	7			
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	50	16.7	3			
CHESTERFIELD INLET	1982	10	15	34	69	0	0	0	0	0	0	0	0	0	0	223	92	340	85.0	4		
	1983	85	73	3	2	0	0	0	0	0	0	0	0	0	0	24	8	26	287	57.4	5	
	1984	2	5	6	0	0	0	0	0	0	0	0	0	0	0	0	0	0	59	8.4	7	
	1985	0	0	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	37	12.3	3	
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	4	4.0	1	
ESKIMO POINT	1981	0	58	48	172	0	0	0	0	0	0	0	0	0	0	137	90	227	113.5	2		
	1982	40	62	80	27	6	0	0	0	0	0	0	0	0	0	24	1487	469	2298	328.3	7	
	1983	210	142	163	55	1	0	0	0	0	0	0	0	0	0	0	40	103	528	75.4	7	
	1984	117	51	262	186	0	0	0	0	0	0	0	0	0	0	0	227	263	968	138.3	7	
	1985	28	40	120	40	0	0	0	0	0	0	0	0	0	0	0	255	176	966	138.0	7	
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	200	66.7	3	
REPULSE BAY	1981	13	19	8	34	0	0	0	0	0	0	0	0	0	0	0	3	0	0	3.0	1	
	1982	37	22	22	16	0	0	0	0	0	0	0	0	0	0	0	7	81	272	45.3	6	
	1983	35	20	34	16	0	0	0	0	0	0	0	0	0	0	0	78	77	520	74.3	7	
	1984	30	35	24	4	0	0	0	0	0	0	0	0	0	0	0	337	0	95	19.0	5	
	1985	0	1	5	0	0	0	0	0	0	0	0	0	0	0	0	2	0	6	3.0	2	
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
RANKIN INLET	1981	1	6	15	21	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1982	26	21	56	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1983	17	23	20	10	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1984	0	5	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1985	70	22	9	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
WHALE COVE	1981	0	1	0	4	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1982	32	8	30	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1983	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1984	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	
KEEWATIN	1981	64	199	142	270	0	0	0	0	0	0	0	0	0	0	0	153	97	250	125.0	2	
	1982	537	353	446	131	6	0	0	0	0	0	0	0	0	0	0	2956	1020	4842	605.3	0	
	1983	380	399	488	211	1	0	0	0	0	0	0	0	0	0	0	21	0	313	592	2422	269.1
	1984	845	797	585	377	12	0	0	0	0	0	0	0	0	0	0	25	0	1358	1608	4470	558.8
	1985	190	297	168	0	0	0	0	0	0	0	0	0	0	0	0	0	8	849	616	4089	511.1
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

For footnotes see Appendix 4.

Table 47. The estimated harvest of wolves by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	X	NUM				
BAKER LAKE	1982	4	0	16	2	0	0	0	0	0	0	2	28	7.0	4	3				
	1983	0	0	3	0	2	0	0	0	0	1	3	11	20	4.0	5				
	1984	0	12	25	1	0	0	0	0	0	0	0	9	58	11.6	5				
	1985	7	16	18	1	0	0	0	0	0	0	0	21	63	12.6	5				
	1986	3	5	13	-	-	-	-	-	-	-	-	21	7.0	3					
CORAL HARBOUR	1984	0	0	0	0	1	0	0	0	0	0	0	0	1	1.0	1				
	1985	0	0	0	0	4	4	0	0	0	0	0	0	10	3.3	3				
CHESTERFIELD INLET	1982	0	0	3	0	0	0	0	0	0	0	0	4	4	4.0	1				
	1983	0	2	0	0	0	0	0	0	0	0	0	2	9	17	3.4	5			
	1984	6	5	0	0	0	0	0	0	0	0	0	4	15	5.0	3				
	1985	0	4	0	0	0	0	0	0	0	0	0	0	5	4.3	3				
	1986	0	3	2	-	-	-	-	-	-	-	-	5	2.5	2					
ESKIMO POINT	0	0	0	0	0	0	0	0	0	0	0	0	10	10	10.0	1				
	1982	0	0	2	22	0	0	0	0	0	0	0	1	0	25	8.3	3			
	1983	0	0	0	12	8	0	0	0	0	0	0	5	31	6.2	5				
	1984	0	2	12	30	2	0	0	0	0	0	0	1	50	8.3	6				
	1985	0	0	3	32	2	0	0	0	0	0	0	4	10	10.2	5				
	1986	6	2	53	-	-	-	-	-	-	-	-	10	51	20.3	3				
REPULSE BAY	1982	5	18	4	2	13	0	0	0	0	0	0	0	0	42	8.4	5			
	1983	6	0	6	4	3	0	0	0	0	0	0	4	10	2	35	5.0	7		
	1984	2	4	18	2	4	0	0	0	0	0	0	1	8	0	39	5.6	7		
	1985	2	10	9	10	4	0	0	0	0	0	0	6	8	6	52	6.5	8		
	1986	4	13	5	-	-	-	-	-	-	-	-	-	-	22	7.3	3			
RANKIN INLET	1982	4	0	1	9	0	0	0	0	0	0	0	0	0	0	14	4.7	3		
	1983	2	2	7	14	6	0	0	0	0	0	0	0	0	32	5.3	6			
	1984	0	2	3	3	1	0	0	0	0	0	0	0	0	9	2.3	4			
	1985	1	2	6	4	3	1	0	0	0	0	0	0	0	3	4	2.0	2		
	1986	2	4	1	-	-	-	-	-	-	-	-	0	0	6	2.0	3			
WHALE COVE	1982	0	0	2	3	0	0	0	0	0	0	0	0	0	0	14	7.0	2		
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	10	10.0	1		
	1984	0	3	0	2	0	0	0	0	0	0	0	0	0	7	4	11.8	6.9	7	
	1985	0	2	12	0	0	0	0	0	0	0	0	0	0	21	30	139	17.4	8	
	1986	0	2	12	0	0	0	0	0	0	0	0	0	0	2	14	178	22.3	8	
KEEWATIN	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	116	220	24.4	9	
	1982	13	18	25	38	13	0	0	0	0	0	0	0	0	7	4	118	16.9	7	
	1983	8	4	16	33	20	0	0	0	0	0	0	0	0	21	30	139	17.4	8	
	1984	8	28	58	38	8	0	0	0	0	0	0	0	0	2	12	46	220	24.4	9
	1985	10	34	48	51	13	4	0	0	0	0	0	0	0	2	12	46	116	38.7	3
	1986	15	27	74	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

For footnotes see Appendix 4.

Table 48. The estimated harvest of ringed seals by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM	
BAKER LAKE	1983	0	2	0	0	1	0	0	0	0	0	0	0	1	1.0	1	
	1984	0	0	0	0	2	3	0	0	1	0	0	0	6	2.0	3	
CORAL HARBOUR	1982	10	22	22	22	278	107	108	52	42	30	52	679	84.9	6		
	1983	62	40	57	57	142	74	192	49	59	87	13	591	65.7	9		
CHESTERFIELD INLET	1984	101	55	41	57	113	83	20	45	46	20	21	756	63.0	12		
	1985	93	19	22	30	113	83	18	33	79	86	74	739	61.6	12		
ESKIMO POINT	1986	81	83	78	242	80.7	3		
	1987	7	6	7	19	8	16	20	6	11	6	5	2	104	8.7	12	
REPULSE BAY	1981	4	3	10	5	37	20	149	70	70	4	6	379	31.6	12		
	1982	3	4	0	2	2	82	31	27	12	110	36	0	309	30.9	10	
RANKIN INLET	1983	29	20	39	36	40	44	13	132	133	0	0	0	503	50.3	10	
	1984	17	0	3	1	35	42	10	3	7	5	0	0	122	13.6	9	
WHALE COVE	1985	1	12	13	26	8.7	3		
	1986	9	6	14	29	9.7	3		
KEEWATIN	1981	0	0	2	29	6	172	71	114	45	20	10	4	371	123.7	3	
	1982	0	0	8	14	10	12	134	30	69	87	14	0	462	51.3	9	
WILDLANDS	1983	0	0	10	9	16	19	122	126	85	57	123	17	0	423	42.3	10
	1984	8	10	9	18	180	90	145	64	95	62	22	6	592	53.8	11	
FOR FOOTNOTES SEE APPENDIX 4.	1985	14	0	4	18	6	800	72.7	11	
	1986	3	21	12	36	12.0	3		

Table 49. The estimated harvest of bearded seals by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	X	NUM	
CORAL HARBOUR	1982	1	16	3	6	5	2	3	3	10	1	2	0	35	5.8	6	
	1983	0	0	17	2	0	12	4	10	33	2	4	1	69	9.9	7	
	1984	4	7	2	0	0	6	14	6	2	0	0	3	66	6.6	10	
	1985	6	5	2	0	0	0	6	2	0	0	0	0	49	6.1	8	
	1986	5	6	0	-	-	-	-	-	-	-	-	-	11	5.5	2	
CHESTERFIELD INLET	1982	0	0	0	0	0	0	0	0	0	0	0	0	2	2.0	1	
	1983	0	0	0	0	0	0	0	0	3	0	0	0	3	3.0	1	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	1	1.0	1	
	1985	1	0	0	1	1	0	0	0	0	1	0	0	5	1.0	5	
	1986	0	3	0	-	-	-	-	-	-	-	-	-	3	3.0	1	
ESKIMO POINT	1981	-	-	2	0	1	0	0	6	0	3	0	0	21	21.0	1	
	1982	0	0	0	0	3	6	11	1	0	1	2	0	12	3.0	4	
	1983	0	0	0	2	14	6	2	3	9	1	0	0	34	5.7	6	
	1984	0	0	0	0	0	5	4	0	3	1	2	0	49	6.1	8	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	15	3.0	5	
REPULSE BAY	1981	-	-	-	0	0	0	0	0	12	2	2	0	5	7	3.5	2
	1982	0	0	0	0	0	0	0	0	16	6	2	0	0	14	7.0	2
	1983	0	0	0	0	0	0	0	0	10	4	3	0	0	18	4.5	4
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	24	6.0	4
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	18	4.5	4
	1986	0	0	0	0	0	0	0	0	0	0	0	0	0	30	6.0	5
RANKIN INLET	1982	0	0	0	3	0	2	1	3	3	1	6	2	0	21	3.0	7
	1983	0	0	0	4	0	0	1	2	3	0	2	1	0	13	2.2	6
	1984	0	0	0	2	4	3	2	3	3	1	0	1	0	26	3.7	7
	1985	0	0	0	0	0	0	0	0	8	6	6	0	0	30	6.0	5
	1986	0	0	0	1	-	-	-	-	-	-	-	-	1	1.0	1	
WHALE COVE	1982	0	1	0	4	0	0	2	0	0	2	0	0	0	9	2.3	4
	1983	0	0	0	1	0	0	7	0	2	0	17	0	0	27	6.8	4
	1984	0	1	0	1	0	0	0	2	0	0	3	0	0	7	1.8	4
	1985	0	1	0	1	0	0	0	0	0	0	0	0	0	28	14.0	2
KEEWATIN	1981	-	-	17	2	7	1	4	7	26	15	12	2	0	93	9.3	10
	1982	0	0	3	10	3	7	15	30	41	20	7	1	137	13.7	10	
	1983	0	0	21	9	16	23	25	22	42	0	3	1	193	17.5	11	
	1984	4	7	6	9	3	6	8	16	33	19	16	0	1	124	11.3	11
	1985	7	6	5	6	5	-	-	-	-	-	-	-	16	5.3	3	
	1986	5	6	-	-	-	-	-	-	-	-	-	-	-	-	-	

For footnotes see Appendix 4.

Table 50. The estimated harvest of beluga whales by hunters from Chesterfield Inlet, Coral Harbour, Eskimo Point, Rankin Inlet, Repulse Bay and Whale Cove expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM	
CORAL HARBOUR	1982	1	2	0	0	0	0	17	75	32	0	0	1	125	31.3	4	
	1983	0	0	0	0	0	0	3	77	48	2	1	0	128	32.0	4	
	1984	2	1	0	0	0	0	15	50	24	8	0	0	121	17.3	7	
	1985	0	0	0	0	0	0	0	60	13	0	0	0	88	29.3	3	
CHESTERFIELD INLET	1981	0	0	0	0	0	0	0	7	0	0	0	0	0	0.0	0	
	1982	0	0	0	0	0	0	0	4	4	0	0	0	0	0.0	0	
	1983	0	0	0	0	0	0	0	16	16	0	0	0	0	7	0.0	
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	12	4.0	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	32	16.0	
ESKIMO POINT	1982	0	0	0	0	0	0	9	74	2	0	0	0	0	85	28.3	3
	1983	0	0	0	0	0	0	8	48	2	0	0	0	0	58	19.3	3
	1984	0	0	0	0	0	0	35	15	0	0	0	0	0	50	25.0	2
	1985	0	0	0	0	0	0	10	84	0	0	0	0	0	94	47.0	2
REPULSE BAY	1981	0	0	0	0	0	0	0	15	20	3	0	0	0	3	3.0	1
	1982	0	0	0	0	0	0	8	16	17	0	0	0	0	35	17.5	2
	1983	0	0	0	0	0	0	2	12	11	0	0	0	0	41	13.7	3
	1984	0	0	0	0	0	0	1	3	0	0	0	0	0	25	8.3	3
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	4	2.0	2
RANKIN INLET	1982	0	0	0	0	0	0	6	17	11	0	0	0	0	34	11.3	3
	1983	0	0	0	0	0	0	0	29	0	0	0	0	0	29	29.0	1
	1984	0	0	0	0	0	0	2	13	49	5	0	0	0	69	17.3	4
	1985	0	0	0	0	0	0	4	37	6	0	0	0	0	47	15.7	3
WHALE COVE	1982	0	0	0	0	0	0	0	1	2	0	0	0	0	3	1.5	2
	1983	0	0	0	0	0	0	0	19	5	0	0	0	0	24	12.0	2
	1984	0	0	0	0	0	0	0	17	2	0	0	0	0	19	9.5	2
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
KEEWATIN	1981	0	0	0	0	0	0	0	32	187	70	0	0	0	3	3.0	1
	1982	0	0	0	0	0	0	0	16	177	67	2	1	0	290	72.5	4
	1983	0	0	0	0	0	0	0	149	49	8	0	0	0	263	52.6	5
	1984	2	1	0	0	0	0	5	87	149	0	0	0	0	301	43.0	7
	1985	0	0	0	0	0	0	0	30	217	37	0	0	0	284	94.7	3

For footnotes see Appendix 4.

Table 51. The estimated harvest of Canada Geese by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM	
BAKER LAKE	1984	0	0	2	0	154	142	0	0	0	0	0	0	296	148.0	3	
	1985	0	0	0	0	479	144	0	0	0	0	0	0	623	311.5	2	
CORAL HARBOUR	1982	1	0	0	0	41	407	199	0	0	0	0	9	656	164.0	4	
	1983	0	0	0	0	13	101	9	10	0	0	0	0	19	9.5	2	
	1984	0	0	0	0	30	25	0	23	0	0	0	0	137	45.7	3	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	55	27.5	2	
CHESTERFIELD INLET	1984	0	0	0	0	0	8	0	0	0	0	0	0	0	0	0	
	1985	0	0	0	0	0	30	0	0	0	0	0	0	30	30.0	1	
ESKIMO POINT	1982	0	0	0	0	49	4	0	4	2	0	0	0	0	59	14.8	4
	1983	0	0	0	0	86	452	0	0	7	0	0	0	0	545	181.7	3
	1984	0	0	0	0	449	191	0	0	9	0	0	0	0	649	216.3	3
	1985	0	0	0	0	113	43	0	0	0	0	0	0	0	156	78.0	2
REPULSE BAY	1983	0	0	0	0	0	0	0	0	2	0	0	0	0	0	0	
	1984	0	0	0	0	4	3	0	0	0	0	0	0	0	0	0	
	1985	0	0	0	0	7	12	1	0	3	0	0	0	0	0	0	
RANKIN INLET	1982	0	0	0	0	76	1112	0	0	0	0	0	0	0	1188	594.0	2
	1983	0	0	0	0	6	14	0	0	0	0	0	0	0	20	10.0	2
	1984	0	0	0	0	16	376	0	9	0	0	0	0	0	401	133.7	3
	1985	0	0	0	0	65	247	2	1	0	0	0	0	0	315	78.8	4
WHALE COVE	1982	0	0	0	0	71	29	0	0	0	0	0	0	0	100	50.0	2
	1984	0	0	0	0	11	13	0	0	0	0	0	0	0	24	12.0	2
	1985	0	0	0	0	68	12	19	0	5	0	0	0	0	104	26.0	4
KEEWATIN	1982	0	0	0	0	196	1186	407	203	2	0	0	0	9	2003	333.8	6
	1983	0	0	0	0	92	466	0	9	19	0	0	0	0	586	146.5	4
	1984	0	0	0	0	647	834	0	32	9	0	0	0	0	1522	380.5	4
	1985	0	0	0	0	762	513	22	1	8	0	0	0	0	1306	261.2	5

For footnotes see Appendix 4.

Table 52. The estimated harvest of snow geese by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM
BAKER LAKE	1984	0	0 ²	0	0	149	201	0	0	0	0	0	0	350	175.0	2
	1985	0	0	0	0	0	30	0	0	0	0	0	0	30	30.0	1
CORAL HARBOUR	1982	0	0	0	0	4345	0	0	21	0	0	0	22	4388	1462.7	3
	1983	0	0	0	5	322	5063	0	203	12	0	0	0	215	107.5	2
CHESTERFIELD INLET	1984	3	21	0	0	118	3798	24	301	113	0	0	0	5557	793.9	7
	1985	0	0	0	0	0	0	12	0	0	0	0	0	4253	850.6	5
ESKIMO POINT	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0
	1982	0	0	0	0	0	0	0	0	0	0	0	0	39	13.0	3
REPULSE BAY	1982	0	0	0	0	0	0	0	0	0	0	0	0	9	9.0	1
	1984	0	0	0	0	0	0	0	0	0	0	0	0	705	352.5	2
RANKIN INLET	1982	0	0	0	0	0	0	0	0	0	0	0	0	52	17.3	3
	1983	0	0	0	0	0	0	0	0	0	0	0	0	98	49.0	2
WHALE COVE	1984	0	0	0	0	0	0	0	0	0	0	0	0	301	100.3	3
	1985	0	0	0	0	0	0	0	0	0	0	0	0	519	259.5	2
KEEWATIN	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0.0	0
	1982	0	0	0	0	0	0	0	0	0	0	0	0	22	5369	671.1
	1983	0	0	0	0	0	0	0	0	0	0	0	0	406	135.3	3
	1984	3	21	0	5	1018	5602	0	37	30	159	0	0	0	6875	859.4
	1985	0	0	0	0	0	0	0	0	0	0	0	0	5600	1120.0	5

For footnotes see Appendix 4.

Table 53. The estimated harvest of caribou by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM				
BAKER LAKE	1981	0	0	0	0	0	0	138	0	0	65	0	128	0	2	128	3			
	1984	0	0	0	0	0	100	57	0	0	0	4	42	2	247	61.8	4			
	1985	0	0	0	0	0	0	0	0	0	0	0	7	168	42.0	4				
CORAL HARBOUR	1982	318	0	0	0	0	1504	882	1546	26	12	53	0	4341	620.1	7				
	1983	0	0	0	0	0	210	108	370	278	616	366	57	2245	449.0	5				
	1984	300	12	2	0	0	138	1093	867	690	29	1110	140	484	3733	311.1	12			
	1985	0	202	331	0	0	0	0	0	0	11	350	1281	182	5272	527.2	10			
	1986	27	40	23	0	0	0	0	0	0	0	0	0	0	90	30.0	3			
CHESTERFIELD INLET	1982	76	0	0	0	0	0	55	43	0	0	0	0	0	76	76.0	1			
	1983	0	0	0	0	0	0	0	1	0	15	414	50	0	0	153	51.0	3		
	1984	0	0	0	0	0	0	0	0	0	0	0	0	0	480	120.0	4			
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	78	39.0	2			
	1986	0	0	0	0	13	0	0	0	0	0	0	0	0	13	13.0	1			
ESKIMO POINT	1981	4	6	0	0	0	1	72	1080	1042	169	47	24	34	105	35.0	3			
	1982	3	0	0	0	0	61	364	562	694	258	36	57	12	2479	247.9	10			
	1983	0	0	0	0	2	163	172	1249	608	152	78	38	30	2095	232.8	9			
	1984	0	0	0	0	4	41	289	659	1779	357	59	109	68	2601	289.0	9			
	1985	1	0	0	1	0	0	0	0	0	0	0	0	93	3402	309.3	11			
	1986	37	0	0	0	0	0	0	0	0	0	0	0	0	37	37.0	1			
REPULSE BAY	1981	0	0	0	0	0	0	157	313	47	390	29	133	289	283	705	235.0	3		
	1982	0	0	0	0	0	0	0	63	76	109	2	147	418	1919	239.9	8			
	1983	0	0	0	0	0	0	0	14	655	265	493	9	631	564	52	1027	128.4	8	
	1984	0	0	0	0	0	0	0	30	195	501	1017	749	0	27	350	350	355.4	8	
	1985	260	0	0	0	0	0	0	0	0	0	0	0	0	0	3418	427.3	8		
	1986	0	0	0	0	63	0	0	0	0	0	0	0	0	0	63	63.0	1		
RANKIN INLET	1981	0	0	0	0	0	0	134	82	913	7366	1115	791	0	719	719	719.0	1		
	1982	0	0	0	0	0	0	0	0	420	788	2117	65	102	443	727	359	12226	1222.6	10
	1983	214	0	0	0	0	0	57	74	861	724	1804	45	70	763	492	4641	580.1	8	
	1984	333	86	93	0	0	0	114	139	1290	1448	2873	0	404	615	205	5115	426.3	12	
	1985	124	44	96	0	0	0	107	0	0	0	0	0	0	0	316	7463	678.5	11	
	1986	315	114	0	0	0	0	0	0	0	0	0	0	0	0	536	178.7	3		
WHALE COVE	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	166	219	73.0	3	
	1982	111	42	79	0	0	0	0	0	0	0	0	0	0	0	69	13	8083	673.6	12
	1983	2	25	0	0	0	0	0	0	0	0	0	0	0	0	63	30	120	30.0	4
	1984	23	0	0	0	0	0	0	0	0	0	0	0	0	0	117	16	1197	119.7	10
	1985	4	5	0	0	0	0	0	0	0	0	0	0	0	0	6	702	70.2	10	
	1986	40	20	3	0	0	0	0	0	0	0	0	0	0	0	63	63	21.0	3	
KEEWINATIN	1981	0	0	0	0	0	0	0	0	0	0	0	0	0	0	222	452	1202	1876	3
	1982	191	719	102	0	0	0	0	0	0	0	0	0	0	0	1417	1324	3812	29124	12
	1983	219	25	14	0	0	0	0	0	0	0	0	0	0	0	465	950	1474	661	10281
	1984	656	98	95	276	363	2284	2792	4476	370	2084	1597	2733	854	2733	604	1125	16216	1351.3	12
	1985	369	251	148	0	0	0	0	0	0	0	0	0	0	0	0	0	20503	1708.6	12
	1986	419	174	209	0	0	0	0	0	0	0	0	0	0	0	0	0	802	267.3	3

For footnotes see Appendix 4.

Table 54. The estimated harvest of lake trout by hunters from the seven Keewatin communities expressed as number of animals harvested for the period October, 1981 to March, 1986.

AREA	YEAR	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEPT	OCT	NOV	DEC	SUM	\bar{X}	NUM		
BAKER LAKE	1981	1	0	380	400	0	0	0	0	0	0	0	0	8744	2154	10898	5449.0	
	1982	0	0	0	796	0	1886	188	0	0	1732	545	178	5325	390.0	2		
	1983	0	0	241	32	175	72	87	0	182	196	144	192	469	887.5	6		
	1984	76	157	265	0	596	333	59	25	798	520	469	3288	328.8	10			
	1985	181	43	0	264	·	·	·	·	·	·	828	828	276.0	3			
	1986	261	317	250	·	·	·	·	·	·	·	·	·	·	·	·		
CORAL HARBOUR	1982	0	0	410	0	0	0	0	0	0	0	0	0	0	410	410.0		
	1983	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0		
	1985	0	0	30	131	41	72	8	0	0	0	0	0	0	0	7		
CHESTERFIELD INLET	1982	0	19	·	·	·	·	202	0	0	26	0	0	273	68.3	4		
	1983	0	0	0	1	59	26	0	0	0	0	43	0	325	54.2	6		
	1984	0	0	0	2	31	22	0	0	0	0	111	0	0	197	49.3		
	1985	0	0	0	0	0	0	0	0	0	58	0	0	113	28.3	4		
ESKIMO POINT	1981	91	12	79	182	237	38	9	65	75	78	28	5	1686	562.0	3		
	1982	48	11	213	115	281	72	0	24	130	68	159	231	0	821	74.6	11	
	1983	0	0	4	183	62	138	8	6	111	22	22	32	588	58.8	10		
	1984	0	0	0	0	0	0	0	0	0	75	59	32	170	42.5	4		
	1985	4	0	0	0	0	0	0	0	0	0	0	0	76	25.3	3		
	1986	2	4	70	·	·	·	·	·	·	·	·	·	·	·	·		
REPULSE BAY	1981	·	1	15	199	345	0	0	11	745	5	0	0	750	375.0	2		
	1982	75	0	8	46	8	2	0	0	0	0	5	0	651	93.0	7		
	1983	0	0	0	0	60	0	0	0	0	2	0	0	66	13.2	5		
	1984	0	0	0	0	0	0	0	0	0	0	0	0	285	142.5	2		
	1985	0	0	0	0	68	115	8	0	0	16	0	0	1	208	41.6	5	
	1986	0	0	4	·	·	·	·	·	·	·	·	·	4	4.0	1		
RANKIN INLET	1981	15	76	59	0	0	0	0	0	0	0	0	0	0	33	33.0	1	
	1982	0	20	0	289	0	0	0	0	0	0	54	0	197	49.3	4		
	1983	0	20	0	120	232	0	10	21	0	0	103	0	363	121.0	3		
	1984	0	16	14	0	173	127	0	0	15	0	0	0	506	84.3	6		
	1985	1	16	0	0	·	·	·	·	·	·	·	2	348	49.7	7		
	1986	16	0	0	0	0	0	0	0	0	0	0	0	16	16.0	1		
WHALE COVE	1981	59	12	14	71	52	15	23	120	3	4	33	4	156	193	64.3	3	
	1982	11	56	18	93	80	18	0	0	15	26	58	58	468	468	39.0	12	
	1983	17	14	92	0	52	44	7	1	0	0	0	0	0	85	85	3	
	1984	0	0	0	6	52	44	7	1	0	3	0	0	0	323	46.1	7	
	1985	0	0	0	0	0	0	0	0	0	0	0	0	0	0	113	18.8	6
	1986	10	9	0	·	·	·	·	·	·	·	·	·	19	19	9.5	2	
KEEWATIN	1981	225	58	552	727	488	398	32	387	89	69	107	67	3199	266.6	12		
	1982	59	87	679	292	1415	146	1894	212	130	1802	801	409	660.5	660.5	12		
	1983	191	364	638	525	357	90	114	293	329	503	224	3721	3721	310.1	12		
	1984	93	14	588	911	348	60	40	892	637	504	4247	353.9	353.9	3	3		
	1985	186	59	14	8	324	·	·	·	·	·	·	943	314.3	3	3		
	1986	289	330	·	·	·	·	·	·	·	·	·	·	·	·	·		

For footnotes see Appendix 4.




Fig. 1. Map of Keewatin District showing the seven communities surveyed during the harvest study and the zonal grid used to locate kills.

Appendix 1. Monthly percent of hunters interviewed for the seven Keewatin communities for the period October 1981 to March 1986.

	Year	Oct.	Nov.	Dec.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.
Baker Lake	1981-82	* ¹	27.3	18.6	45.5	56.4	49.5	40.9	57.3	39.5	41.8	*	90.9
	1982-83	*	94.1	94.5	99.6	85.5	91.8	95.7	93.2	94.5	92.3	95.9	92.7
	1983-84	98.7 ²	93.2	97.8	96.1	95.2	97.4	94.4	97.0	95.9	96.6	95.6	95.8
	1984-85	97.6	97.7	98.0	96.8	94.9	96.5	100.0	98.6	99.0	95.0	95.0	96.7
	1985-86	95.4	98.0	94.8	100.0	100.0	100.0						
Chesterfield Inlet	1981-82	*	*	*	68.0	48.0	*	*	*	*	*	24.0	88.0
	1982-83	78.0	86.0	90.0	62.0	56.0	28.0	58.0	88.0	100.0	90.0	100.0	100.0
	1983-84	100.0	88.3	98.3	93.3	92.7	100.0	100.0	78.1	96.9	87.0	98.4	100.0
	1984-85	100.0	100.0	100.0	100.0	100.0	98.5	98.3	95.8	95.7	97.2	93.2	97.1
Coral Harbour	1981-82	*	*	*	49.5	*	*	*	*	*	56.2	59.0	36.2
	1982-83	18.1	27.6	26.7	31.4	32.4	35.2	18.1	*	*	*	24.0	88.0
	1983-84	27.6	22.9	16.2	32.4	95.2	85.7	82.9	69.5	96.3	73.3	67.6	60.0
	1984-85	99.1	61.9	62.9	68.6	79.0	93.3	94.3	78.1	59.0	62.9	75.2	86.7
	1985-86	73.3	85.7	80.0	85.0	84.8	81.0						
Eskimo Point	1981-82	57.5	62.8	61.1	79.6	78.3	66.4	50.0	81.4	79.2	80.5	80.1	88.2
	1982-83	97.3	94.7	95.6	97.8	97.0	100.0	97.7	96.7	95.5	96.2	98.4	97.7
	1983-84	98.8	98.4	97.0	98.3	94.6	93.2	93.5	99.1	98.2	99.2	98.4	84.1
	1984-85	86.3	94.5	79.2	93.8	89.3	90.1	95.2	88.3	89.4	82.3	95.2	94.3
	1985-86	87.5	91.9	91.8	94.1	85.8	89.8						
Rankin Inlet	1981-82	*	92.3	57.5	92.7	66.8	71.0	33.7	31.6	45.6	63.7	22.8	96.9
	1982-83	61.7	64.8	100.0	84.5	89.1	84.5	95.9	33.7	19.2	34.2	55.4	61.1
	1983-84	51.3	81.5	92.2	90.4	89.5	91.0	97.5	74.4	90.3	82.4	100.0	85.8
	1984-85	84.7	82.1	86.0	86.7	85.4	91.4	94.8	87.6	87.3	88.6	93.5	96.2
	1985-86	95.0	98.1	97.5	97.9	96.8	97.2						
Repulse Bay	1981-82	61.1	58.9	26.7	38.9	60.0	26.7	50.0	31.1	37.8	32.2	33.3	44.4
	1982-83	51.1	56.7	*	70.0	73.3	71.1	*	53.3	53.3	73.3	57.8	53.3
	1983-84	57.8	66.7	41.1	58.9	44.4	54.4	51.1	73.3	71.1	51.1	76.7	58.9
	1984-85	67.9	72.6	70.2	79.8	66.7	73.8	73.8	67.9	61.9	72.6	73.6	75.0
	1985-86	71.4	83.3	97.7	90.3	95.2	100.0						
Whale Cove	1981-82	20.0	86.0	82.0	82.0	74.0	94.0	76.0	52.0	100.0	100.0	72.0	92.0
	1982-83	28.0	32.0	54.0	62.0	80.0	40.0	*	*	*	*	*	*
	1983-84	*	30.0	14.0	52.0	98.0	98.0	100.0	77.6	70.7	69.9	71.0	88.7
	1984-85	*	68.0	85.9	87.5	93.5	93.3	87.3	83.9	82.5	98.3	98.3	100.0
	1985-86	100.0	100.0	98.3	100.0	100.0	100.0						

¹Data not provided.

²These figures are the percentage of hunters in each community who were interviewed each month in relation to the total number of hunters who could have hunted that month.

Appendix 2. Monthly theoretical kill factors for seven Keewatin communities for the period October 1981 to March 1986.

	Year	Oct.	Nov.	Dec.	Jan.	Feb.	March	April	May	June	July	Aug.	Sept.
Baker Lake	1981-82	* ³	(3.67) ¹	(5.36) ⁴	2.19	1.77	2.02	2.44	1.75	2.52	2.38	*	1.11
	1982-83	*	1.06	1.06	1.00	1.19	1.08	1.01	1.03	1.03	1.04	1.00	1.06
	1983-84	1.00	1.07	1.00	1.00	1.00	1.00	1.00	1.08	1.00	1.00	1.00	1.00
	1984-85	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.01	1.00	1.00	1.00	1.00
	1985-86	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Chesterfield Inlet	1981-82	*	*	*	1.47	2.08	*	*	*	*	*	(4.17)	1.13
	1982-83	1.28	1.16	1.11	1.61	1.78	(3.57)	1.72	1.14	1.00	1.18	1.00	1.00
	1983-84	1.00	1.00	1.04	1.14	1.27	1.00	1.00	1.35	1.07	1.39	1.03	1.00
	1984-85	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.13	1.11	1.07	1.21	1.08
	1985-86	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Coral Harbour	1981-82	*	*	*	2.01	*	*	*	*	1.78	1.69	2.76	1.36
	1982-83	(5.53)	(3.62)	(3.75)	3.17	3.08	2.83	5.53	*	*	*	*	*
	1983-84	(3.62)	(4.38)	(6.18)	(3.09)	1.05	1.17	1.21	1.44	1.01	1.35	1.44	1.61
	1984-85	1.01	1.54	1.40	1.22	1.59	1.04	1.02	1.28	1.65	1.52	1.29	1.09
	1985-86	1.62	1.27	1.57	1.00	1.10	1.07	1.07	1.07	1.07	1.07	1.07	1.07
Eskimo Point	1981-82	1.74	1.59	1.64	1.25	1.28	1.51	2.00	1.23	1.18	1.24	1.23	1.14
	1982-83	1.03	1.06	1.02	1.07	1.11	1.11	1.05	1.04	1.08	1.06	1.03	1.05
	1983-84	1.38	1.01	1.00	1.00	1.05	1.01	1.01	1.01	1.02	1.01	1.02	1.12
	1984-85	1.09	1.01	1.22	1.04	1.14	1.31	1.17	1.14	1.11	1.40	1.07	1.01
	1985-86	1.07	1.21	1.24	1.13	1.48	1.21	1.21	1.21	1.21	1.21	1.21	1.21
Rankin Inlet	1981-82	*	1.08	1.74	1.05	1.50	1.41	2.97	3.16	2.19	1.56	(4.39)	1.03
	1982-83	1.62	1.54	1.00	1.18	1.06	1.17	1.04	2.95	(5.22)	2.92	1.80	1.63
	1983-84	1.95	1.15	1.22	1.16	1.18	1.02	1.06	1.42	1.00	1.38	1.00	1.06
	1984-85	1.18	1.36	1.36	1.33	1.63	1.37	1.35	1.39	1.30	1.96	1.25	1.16
	1985-86	1.05	1.05	1.09	1.06	1.10	1.08	1.08	1.08	1.08	1.08	1.08	1.08
Repulse Bay	1981-82	1.64	1.70	(3.75)	2.57	1.66	(3.75)	1.70	3.21	2.65	3.10	3.00	2.25
	1982-83	1.95	1.70	*	1.43	1.36	1.40	*	1.52	1.67	1.36	1.73	1.87
	1983-84	1.78	1.42	2.36	1.66	2.04	1.76	1.76	1.36	1.25	1.80	1.27	1.55
	1984-85	1.38 ²	1.34	1.40	1.20	1.38	1.27	1.31	1.39	1.51	1.35	1.33	1.31
	1985-86	1.33	1.16	1.07	1.23	1.10	1.00	1.00	1.00	1.00	1.00	1.00	1.00
Whale Cove	1981-82	{5.00}	1.16	1.21	1.35	1.06	1.31	1.92	1.00	11.00	1.39	1.09	*
	1982-83	{3.57}	(3.13)	1.85	1.61	1.25	2.50	*	*	*	*	*	*
	1983-84	*	(3.33)	(7.14)	7.92	1.01	1.02	1.00	1.13	1.47	1.48	1.45	1.07
	1984-85	*	1.32	1.04	1.00	1.00	1.03	1.00	1.06	1.05	1.00	1.00	1.00
	1985-86	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

¹Bracketed figures were not used because they were based on insufficient data.

²The figure for known hunters in Repulse Bay was lowered to 84 from 90 beginning October, 1985. This figure is important when determining the number of hunters not contacted on a given month with in a community.

³*Data not provided.

()Data provided on successful hunters only. Estimated harvest = Reported harvest.

Appendix 3. Members of the Steering Committee for the period of the Keewatin Wildlife Federation Harvest Study, 1981 to 1987.

Chairperson

Mr. F. McFarland and Ms. D. Stewart	Northern Affairs Program, Department of Indian Affairs and Northern Development.	1981-85 1985-87
----------------------------------------	----------------------------------------------------------------------------------	--------------------

Members

Mr. R. Bailey	Canadian Wildlife Service, Department of the Environment.	1981-82
Mr. R. Cole		1982-87

Mr. K. Lloyd	Department of Renewable Resources,	1981-82
Mr. R. Graf	Government of the Northwest Territories.	1982-87

Mr. R. Peet	Department of Fisheries and Oceans.	1981-87
-------------	-------------------------------------	---------

Mr. L. Pilikapsi	President, Keewatin Wildlife Federation.	1981-82
Mr. P. Kritterdiluk	President, Keewatin Wildlife Federation.	1982-83
Mr. D. Milortok	President, Keewatin Wildlife Federation.	1983-85
Mr. A. Anyootealuk	President, Keewatin Wildlife Federation.	1985-86
Mr. J. Kaunak	President, Keewatin Wildlife Federation.	1986-87

Mr. L. Gamble	Regional Resource Manager, Keewatin Harvest Study.	1981-87
---------------	----------------------------------------------------	---------

Mr. L. Guluk	Project Manager, Keewatin Harvest Study.	1981-83
--------------	------------------------------------------	---------

Ms. V. Curley	Assistant Regional Resource Manager, Keewatin Harvest Study.	1983-85
---------------	--------------------------------------------------------------	---------

Mr. J. Kusugak	Assistant Regional Resource Manager, Keewatin Harvest Study.	1985-86
----------------	--------------------------------------------------------------	---------

Appendix 4. Footnotes for Tables 35 to 54.

A) Footnotes for Tables 35 to 41.

- 1 In the case where there was zero harvest for all months this row is not represented in the table but is reflected in the N row.
- 2 Zero means there was no harvest for that particular month.
- 3 A dot means the data were not available for that month. When this occurred that month is not counted in the row N for that year or in the column N for that month.
- 4 In cases where there was minimal harvest column N, sums and means have not been included to conserve space.
- 5 Categories are as follows: M = males, F = females, C = calf, and U = unknown.
- 6 Bracketted letters indicate the following: H = high profile species, L = low profile species, A = abundant, S = scarce. These categories are defined as follows:
 1. A high profile species is one which is of economic and cultural importance in a particular community. This may vary between communities because the availability for harvest influences the importance of specific species.
 2. A low profile species is one which has little economic or cultural significance no matter what its abundance.
 3. Abundant refers primarily to availability for harvest. This may vary with distance or with time. For example some species, such as caribou may be available to a community year-round but their harvest may involve considerable travel at certain periods of the season. Other species such as beluga whale and geese are only abundant at specific locations and periods in each year.
 4. Scarce indicates the availability of a species for harvest is limited or that the effort involved in harvesting it is too great. In many instances the harvest of such species is incidental to more preferred species which are actively sought.

B) Footnotes for Tables 42 to 54.

- 1 A dot means the data were not available for that month.
- 2 Zero means there was no harvest for that particular month.
- 3 \bar{X} was calculated using only those months showing a harvest.

