

Scientific Excellence • Resource Protection & Conservation • Benefits for Canadians
Excellence scientifique • Protection et conservation des ressources • Bénéfices aux Canadiens

Marine Ichthyoplankton Data From the Canadian Beaufort Sea Shelf, July and September, 1984

D.B. Chiperzak, G.E. Hopky, M.J. Lawrence
and G. Lacho

Central and Arctic Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

February 1990

**Canadian Data Report of
Fisheries and Aquatic Sciences
No. 779**



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Data Report of
Fisheries and Aquatic Sciences 779

February 1990

MARINE ICHTHYOPLANKTON DATA FROM THE CANADIAN BEAUFORT SEA SHELF,
JULY AND SEPTEMBER, 1984

by

D.B. Chiperzak, G.E. Hopky, M.J. Lawrence and G. Lacho

Central and Arctic Region
Department of Fisheries and Oceans
Winnipeg, Manitoba R3T 2N6

This is the 43rd Data Report
from the Central and Arctic Region, Winnipeg

PREFACE

This study was funded by the Northern Oil and Gas Action Program (NOGAP), through the Department of Fisheries and Oceans, Central and Arctic Region. It is one of a series of projects being executed under NOGAP B.2, to provide background data for assessing the implications of hydrocarbon development and production on critical estuarine and marine habitats of the Canadian Arctic Coastal Shelf. This document constitutes NOGAP Report B2.41.

(c) Minister of Supply and Services Canada 1990

Cat. no. FS 97-13/779E

ISSN 0706-6465

Correct citation for this publication is:

Chiperzak, D.B., G.E. Hopky, M.J. Lawrence, and G. Lacho. 1990. Marine ichthyoplankton data from the Canadian Beaufort Sea Shelf, July and September, 1984. Can. Data Rep. Fish. Aquat. Sci. 779: v + 45 p.

TABLE OF CONTENTS

	<u>Page</u>
PREFACE	ii
ABSTRACT/RESUME	iv
INTRODUCTION	1
METHODS	1
RESULTS	2
ACKNOWLEDGMENTS	2

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1 Location and detailed map of Beaufort Sea study area	5
2 Stations sampled in July and September, 1984	6

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1 Summary of plankton stations sampled in the Beaufort Sea, 1984.	7
2 Plankton summary of Wisconsin and neuston tows conducted in the Canadian Beaufort Sea, July and September, 1984	8
3 Stomach fullness codes used in Table 15	9
4 Scientific and common names of fishes captured in the Canadian Beaufort Sea Shelf, 1984	9
5 Number, percent by number, and frequency of occurrence for larval fish captured by Wisconsin tows in the Beaufort Sea Shelf during July and September, 1984.	10
6 Larval fish catch summary for Wisconsin tows on the Beaufort Sea Shelf, 1984	11
7 Larval fish number 100 m^{-3} for Wisconsin tows on the Beaufort Sea Shelf, 1984	12
8 Number, percent by number, and frequency of occurrence for larval fish captured by neuston tows on the Beaufort Sea Shelf, 1984.	13
9 Larval fish catch summary for neuston tows on the Beaufort Sea Shelf, 1984.	14

<u>Table</u>	<u>Page</u>
10 Larval fish number 100 m^{-3} for neuston tows on the Beaufort Sea Shelf, 1984	15
11 Standard and total lengths, and wet weight of larval fish captured in the Beaufort Sea, 1984	16
12 Mean length and weight of larval <i>B. saida</i> for July and September, 1984.	30
13 Mean length and weight of larval <i>G. tricuspidis</i> for July and September, 1984	30
14 Mean length and weight of larval <i>M. quadricornis</i> for July, 1984.	30
15 Standard length and stomach fullness for larval fish captured in the Beaufort Sea, 1984	31
16 Food item totals for <i>B. saida</i> captured in the Canadian Beaufort Sea Shelf, 1984	34
17 Food item totals for <i>M. quadricornis</i> captured in the Canadian Beaufort Sea Shelf, 1984	40
18 Food item totals for <i>G. tricuspidis</i> captured in the Canadian Beaufort Sea Shelf, 1984	41
19 Food item totals for <i>Liparis</i> sp. captured in the Canadian Beaufort Sea Shelf, 1984	42
20 List of comment codes used in Tables 16 to 19	42
21 Frequency of occurrence and percent by number of food items found in <i>B. saida</i> stomachs	43
22 Frequency of occurrence and percent by number of food items found in <i>M. quadricornis</i> stomachs, July 1984	44
23 Frequency of occurrence and percent by number of food items found in <i>G. tricuspidis</i> stomachs, July and September 1984	44
24 Frequency of occurrence and percent by number of food items found in <i>Liparis</i> sp. stomachs, July 1984	45
25 Parasites found in larval fish stomachs	45

ABSTRACT

Chiperzak, D.B., G.E. Hopky, M.J. Lawrence, and G. Lacho. 1990. Marine ichthyoplankton data from the Canadian Beaufort Sea Shelf, July and September, 1984. Can. Data Rep. Fish. Aquat. Sci. 779: v + 45 p.

Ichthyoplankton were collected from 760 μm Wisconsin net and 500 μm neuston net plankton tows in July and September of 1984 along the Canadian Beaufort Sea Shelf. A total of 1408 larval fish representing six families and nine species groups were caught in the 34 plankton tows. Arctic cod (Boreogadus saida) was the most abundant larval fish species caught both in the Wisconsin and neuston tows.

Standard length, total length, and wet-weight measurements were made on 1123 larval fish of which 1097 were Arctic cod. The July mean standard length for Arctic cod was 13.4 mm with a standard deviation of 2.7 mm, and a mean weight of 0.015 g with a standard deviation of 0.010 g. In September, Arctic cod mean standard length was 26.3 mm, standard deviation of 3.8 mm, and the mean weight was 0.130 g with a standard deviation of 0.058 g.

A total of 126 fish were analyzed for stomach contents of which four were found to be empty. Copepods were the most frequently occurring food item in all four species groups analysed with Calanoida being the most frequently occurring and the most abundant food item for both B. saida and the fourhorn sculpin, Myoxcephalus quadricornis.

Nine stomachs, all from B. saida, contained the parasite (Trematoda: Hemiuridae).

Key words: Arctic; coastal waters; ichthyoplankton; larval fishes; Boreogadus saida; Arctic cod; distribution; diet.

RÉSUMÉ

Chiperzak, D.B., G.E. Hopky, M.J. Lawrence, and G. Lacho. 1990. Marine ichthyoplankton data from the Canadian Beaufort Sea Shelf, July and September, 1984. Can. Data Rep. Fish. Aquat. Sci. 779: v + 45 p.

En juillet et septembre 1984, on a récolté l'ichtyoplancton à l'aide de filets Wisconsin (760 µm) et de filets à neuston (500 µm) le long de la partie canadienne de la plate-forme de la mer de Beaufort. Trente-quatre traits ont permis de capturer un total de 1408 larves de poisson représentant six familles et neuf groupes d'espèces. Parmi les larves de poisson capturées, la saïda franc (Boreogadus saïda) a été l'espèce la plus abondante et ce, indépendamment du filet utilisé (à neuston ou Wisconsin).

Des mesures de la longueur standard, de la longueur totale et du poids frais ont été effectuées pour 1123 larves de poisson, dont 1097 de saïda franc. En juillet, la longueur standard moyenne des larves de saïda franc était de 13,4 mm, avec un écart-type de 2,7 mm, et le poids moyen était de 0,015 g, avec un écart-type de 0,010 g. En septembre, la longueur standard moyenne des larves de saïda franc était de 26,3 mm, avec un écart-type de 3,8 mm, et le poids moyen était de 0,130 g, avec un écart-type de 0,058 g.

Le contenu stomacal a été analysé chez 126 poissons; quatre de ces poissons avaient l'estomac vide. Les copépodes constituaient l'article alimentaire le plus fréquent dans les quatre groupes d'espèces analysés. Calanoida était l'article alimentaire le plus fréquent et le plus abondant chez B. saïda et chez Myoxcephalus quadricornis (chabosseau à quatre cornes).

Neuf estomacs, appartenant tous à des poissons de l'espèce B. saïda, étaient parasités (Trematoda: Hemiuridae).

Mots-clés: Arctique; eaux côtières; ichtyoplancton; larves de poisson; Boreogadus saïda; saïda franc; distribution; régime alimentaire.

INTRODUCTION

Ichthyoplankton data presented in this report were collected as part of the Beaufort Shelf Fish Habitat Research Subproject (B.2.1). This Subproject was one component of the Critical Arctic Estuarine and Marine Habitat Project (B.2) of the Northern Oil and Gas Action Program (NOGA-P). The objectives of the subproject were:

1. To conduct research towards identifying, in spatial and temporal terms, areas of significance to marine and estuarine fish species of the Beaufort Sea Shelf (from Herschel Island to Cape Bathurst) (Fig. 1).
2. To characterize these areas of important marine and estuarine habitat in terms of community composition, and in terms of chemical, physical and biological parameters.
3. To describe the feeding habits of selected species of pelagic and demersal fish species of the Beaufort Sea Shelf in relation to habitat and season (open water vs ice covered).

Ichthyoplankton samples were collected for four consecutive summers in the Beaufort Sea beginning in 1984. The data in this report are from 1984. Temperature, salinity and density data for 1984 are reported by Hopky et al. (1986). Stomach data for Boreogadus saida (Arctic cod) captured from otter trawl tows done in 1984 are reported by Lacho (1986).

Past studies along the Canadian Beaufort Sea Shelf have had their emphasis on juvenile and adult fish (Galbraith and Hunter 1975), and have been mostly nearshore (Kendel et al. 1975; Percy 1975; Lawrence et al. 1984; Bond and Erickson 1989). The study of larval fish in the Canadian Beaufort Sea has been to date largely ignored, except for Tuktoyaktuk Harbour which has been reported on by Ratynski (1983); yet the larval fish stage is an important time in any fish population because the mortality rate at this stage is so high that even small increases in this rate can severely affect a year class strength (Bagenal and Braum 1978).

Sampling was done on board the chartered M.V. Sequel, a 13 m, wooden hull vessel. Due to the size of the vessel and the nature of its hull, site selection for sampling stations was often at the mercy of ice and weather conditions present at the time of sailing.

METHODS

Coordinates for sampling stations (Table 1) were obtained using a combination of radar (Furuno Model CR240) fixes to shore, bathymetric charts and a Magnavox Model MX4102 satellite navigator with a nominal accuracy of ± 0.1 km.

Plankton samples were collected between July 14 to 25 and September 7 to 9 of 1984 (Table 2). Stations sampled are shown in Fig. 2.

Midwater plankton collections were made using a Wisconsin design plankton net with an opening diameter of 0.5 m and a 763 μm Nitex mesh net. A General Oceanic flowmeter, model #2030, with a standard rotor was affixed to the centre of the opening. One horizontal tow of 20-40 minutes duration was made at each station. Tow depth was calculated by taking the cosine of the wire angle multiplied by the length of wire spooled out. Ship speed was held at approximately 3 knots for the duration of the tow.

One surface tow using a neuston sampler was made at each station except at stations 84A11 and 84A12 where no surface tows were conducted. The neuston sampler has a square opening of 45 cm x 45 cm with a net constructed of 500 μm Nitex mesh. A more detailed description of the neuston sampler used is given by Mason and Phillips (1986). A General Oceanic flowmeter, model #2030, with a standard rotor was affixed to the opening of the neuston sampler offset 17 cm vertically from the center of the mouth opening. For volumetric calculations an opening of 45 cm x 35 cm was used, as the top 10 cm of the sampler rode out of the water during the tow periods. The sampler was towed 30 m off the starboard side of the vessel in front of the vessel wake using a polyethylene rope. Ship speed was approximately 4-5 knots for the duration of the tow.

Densities were calculated as the number of fish per 100 m^3 of water filtered through either the Wisconsin or neuston nets.

All plankton samples were preserved in a solution of seawater and 5% non-buffered formalin. Larval fish were extracted later in the laboratory, placed in 70% ethanol and identified to the lowest possible taxon.

All larval fish excepting those extracted for stomach analysis or those in too poor a condition for further examination were measured for standard length, total length and wet

weight. Length measurements were made using a Nikon Type 102 dissecting microscope and a small ruler marked in 1 mm graduations. Measurements were made to the nearest \pm 0.5 mm. Wet weights were obtained by first blotting the larval fish with a paper towel and then weighing them on a Mettler PC440 digital scale with an accuracy of \pm 0.001 g. No adjustments for shrinkage were made in any of the measurements.

Larval fish for stomach analysis were arbitrarily selected to provide a representative cross-section of samples by station and season. These fish were first measured for standard length, rounded to the nearest \pm 0.5 mm, using the micrometer eyepiece of a Wild MSA dissecting microscope. The stomach was then removed or in cases where the stomach had not yet fully formed, the entire intestine was taken. A code (Table 3) was assigned to each stomach or intestine to indicate the degree of fullness and the content's state of digestion. Stomach contents were then identified and enumerated. Parasites found in larval fish stomachs were identified and enumerated whenever possible.

RESULTS

A total of 1408 larval fish consisting of nine species groups representing six families (Table 4) were captured from plankton tows in July and September of 1984. Eighteen Wisconsin tows were made (Table 2), capturing 1186 larval fish of which eight species groups in five families were represented. Only 16 neuston tows were made (Table 2), capturing 222 larval fish of which three species groups in three families were represented.

Boreogadus saida was the most abundant species found in Wisconsin tow catches making up 93.0% of the catch in July and 95.8% in September (Table 5). B. saida was also the most frequently occurring species found in Wisconsin tow catches being caught in 66.7% of the catches in July and 100% of the catches in September. All Gadidae captured at station 84B09 (Table 6) were too badly damaged or decomposed to positively identify past family level but are probably B. saida and thus would increase B. saida's percent by number and frequency of occurrence for the month of July. The staghorn sculpin, Gymnoanthus tricuspidis, was the next most abundant species captured in Wisconsin tows making up 2% of the catch in July and 4.2% in September (Table 5). It occurred in 50% of the catches both in July and September. Although six other species were caught intermittently through July, only B. saida and G. tricuspidis were caught in Wisconsin tows during September (Table 6 - "B" stations after 84A12).

Densities (number \cdot 100 m $^{-3}$) of fish by species and station for Wisconsin tow catches are given in Table (7).

B. saida was also the most abundant species captured using neuston tows, making up 99.1% of the catch in July and 50% of the catch in September (Table 8). Note that only two fish were caught by the six neuston tows done in September (Table 9 - "B" stations after 84A12). The second fish was an unidentified Cottidae captured at station 84B11. In July, B. saida occurred in 70% of the neuston catches. Only two other species were caught in July by the neuston sampler, one Osmerus mordax and one Myoxcephalus quadricornis both of which were captured at station 84A01 (Table 9).

Densities (number \cdot 100 m $^{-3}$) of fish by species and station for all neuston tows are given in Table 10.

Standard length, total length and wet weight measurements were taken on 1123 larval fish (Table 11). Only one specimen each of O. mordax, Anisarchus medius, Triglops pingeli and Aspidophoroides olrikii were measured (Table 11). Virtually all specimens sampled were B. saida, G. tricuspidis or M. quadricornis.

B. saida captured in July had a mean standard length of 13.4 mm with a range of 5.5 to 30.0 mm and a standard deviation of 2.7 while in September the mean standard length was 26.3 mm with a range from 19.5 to 34.0 mm and a standard deviation of 3.8 mm (Table 12). Mean wet weight of B. saida in July was 0.015 g with a standard deviation of 0.010 g while in September the mean wet weight was 0.130 g with a standard deviation of 0.058 g.

G. tricuspidis in July had a mean standard length of 13.6 mm with a range of 11.5 to 15.5 mm and a standard deviation of 1.6 (Table 13). In September G. tricuspidis had a mean standard length of 20.5 mm with a range from 20.0 mm to 21.0 mm and a standard deviation 0.5 mm. The mean wet weight in July was 0.020 g with a standard deviation of 0.009 g and in September the mean wet weight was 0.116 g with a standard deviation of 0.011 g.

Ten M. quadricornis, all in July from station 84A11, were measured (Table 14). The mean standard length was 13.3 mm with a range of 7.0 to 15.5 mm and a standard deviation of 2.3 mm. The mean wet weight was 0.012 g with a range from <0.001 to 0.021 g and a standard deviation of 0.006 g.

A total of 126 fish representing four species groups were examined for their stomach content (Table 15). Of these 99 were B. saida

of which three had empty stomachs. As well, 13 G. tricuspidis, six M. quadricornis and four Liparis sp. were examined and all exhibited varying degrees of fullness except one M. quadricornis whose stomach was empty.

A detailed list of food items with totals and comments for the B. saida specimens examined is given in Table 16, for M. quadricornis in Table 17, for G. tricuspidis in Table 18 and for Liparis sp. in Table 19. Comment codes are defined in Table 20.

Copepods were the most dominant food group found in the stomachs of all four fish species groups examined (Tables 21 - 24). Calanoids were the most frequently observed food item as well as being the most numerous food item found in both B. saida and M. quadricornis stomachs (Tables 21 and 22).

In B. saida (Table 21), unidentified Calanoida occurred in 83.6% of the stomachs analysed from July and occurred in all stomachs analysed from September. Unidentified Calanoida made up 61.6% by number from July and 63.8% by number from September of food items counted. Pseudocalanus minutus was the next most frequently occurring food item at 26.9% from July as well as the next most numerous food item counted at 31.9% by number, while in September Acartia clausi occurred slightly more often than P. minutus (59.4% as compared to 56.3%) but P. minutus was the next most abundant species found after Calanoida at 26.2% as compared to A. clausi at 2.2%.

Only copepods were found in the six M. quadricornis stomachs examined (Table 22). Calanoida occurred the most frequently in 50% of the stomachs and was the most abundant food item making up 84.8% by number.

P. minutus was the most frequently occurring food item found in G. tricuspidis stomachs, occurring 84.6% of the time, but the diatom Coscinodiscus sp. contributed the greatest percent by number at 49.0% (Table 23). In the four Liparis sp. stomachs the copepod P. minutus occurred the most frequently, being found in 75% of the stomachs analysed and made up the greatest percent by number at 87% (Table 24).

Nine fish, all B. saida, contained the parasite (Trematoda: Hemiuridae) in their stomachs (Table 25).

ACKNOWLEDGMENTS

The authors would like to thank Mr. L. deMarch and Mr. T. Maddeford for field assistance as well as Dr. R. Korczynski who assisted during the September survey. Thanks to Mr. J. Ostrick

and staff at the Inuvik Research Laboratory for logistic support, to Polar Continental Shelf Project for their field operations support and Mr. Ray Ratynski of North South Consultants Ltd. for larval fish identification. Nell Stallard of LGL Ltd. did the stomach content analysis and Margaret Keast provided laboratory assistance; we thank both. The skipper of the M.V. Sequel was L. Pearson. Mr. R. Gervais provided computer assistance and advice. Final manuscript preparation was completed by Barb Hyman and Karla DeCaigny.

REFERENCES

- BAGENAL, T.B., and E. BRAUM. 1978. Eggs and early life history, p. 165-201. In T.B. Bagenal (ed.) Methods for assessment of fish production in fresh waters. Blackwell Scientific Publications, Oxford. 365 p.
- BOND, W.A., and R.N. ERICKSON. 1989. Summer studies of the nearshore fish community at Phillips Bay, Beaufort Sea coast, Yukon. Can. Tech. Rep. Fish. Aquat. Sci. 1676: vi + 102 p.
- GALBRAITH, D.F., and J.G. HUNTER. 1975. Fishes of offshore waters and Tuktoyaktuk vicinity. Beaufort Sea Project, Tech. Rep. 7: 47 p.
- HOPKY, G.E., D.B. CHIPERZAK, and M.J. LAWRENCE. 1986. Seasonal salinity, temperature and density data for the Canadian Beaufort Sea Shelf, 1984-1985. Can. Data Rep. Fish. Aquat. Sci. 593: iv + 249 p.
- KENDEL, R.E., R.A.C. JOHNSTON, U. LOBSIGER, and M.D. KOZAK. 1975. Fishes of the Yukon coast. Beaufort Sea Project, Tech. Rep. 6: 114 p.
- LACHO, G. 1986. Analysis of Arctic cod stomach contents from the Beaufort Shelf, July and August, 1984. Can. Data Rep. Fish. Aquat. Sci. 614: iv + 10 p.
- LAWRENCE, M.J., G. LACHO, and S. DAVIES. 1984. A survey of the coastal fishes of the southeastern Beaufort Sea. Can. Tech. Rep. Fish. Aquat. Sci. 1220: x + 178 p.
- MASON, J.C., and A.C. PHILLIPS. 1986. An improved otter surface sampler. Fish. Bull. 84: 480-484.
- PERCY, R. 1975. Fishes of the outer Mackenzie Delta. Beaufort Sea Project Tech. Rep. 8: 114 p.

RATYNSKI, R.A. 1983. Mid-summer ichthyoplankton populations of Tuktoyaktuk Harbour. N.W.T. Can. Tech. Rep. Fish. Aquat. Sci. 1218: iv + 21 p.




Fig. 1. Location and detailed map of Beaufort Sea study area.




Fig. 2. Stations sampled in July and September, 1984.

Table 1. Summary of plankton stations sampled in the Beaufort Sea, 1984.

Station	Latitude		Longitude		Date Sampled	Time* GMT	Total Depth (m)
	Deg	Min	Deg	Min			
84A01	69	49.5	132	37.0	Jul 14	0100	9
84A02	70	2.0	132	20.0	Jul 15	1625	20
84A03	70	12.4	132	21.6	Jul 15	2045	30
84A04	70	17.3	131	53.5	Jul 15	2300	33
84A05	70	6.4	131	0.5	Jul 16	0600	12
84A06	69	34.1	136	22.5	Jul 23	2000	10
84A07	69	33.2	137	0.1	Jul 23	2335	17
84A08	69	33.6	137	6.4	Jul 23	0220	34
84B09	69	31.7	138	44.6	Jul 24	2000	14
84A10	69	36.5	138	49.0	Jul 24	2330	36
84A11	69	40.0	135	14.0	Jul 25	1200	5
84A12	69	43.5	133	55.0	Jul 25	1650	4
84B01	69	51.8	132	54.0	Sep 7	2300	12
84B02	70	4.0	132	26.0	Sep 7	0335	24
84B03	70	13.0	132	22.0	Sep 8	1900	34
84B04	70	20.0	131	50.0	Sep 8	2325	36
84B05	70	6.0	131	0.0	Sep 9	1640	12
84B11	70	13.0	131	16.0	Sep 9	2100	25

* GMT - Greenwich Mean Time.

Table 2. Plankton summary of Wisconsin and neuston tows conducted in the Canadian Beaufort Sea, July and September, 1984.

Plankton Sample Number	Station	Date	Gear Type	Mesh Size (um)	Tow Type	Maximum Tow Depth (m)	Volume Filtered (m³)
1	84A01	Jul 14	NEUST	500	HOR	0.0	693
2	84A01	Jul 14	WISCN	760	HOR	6.2	531
3	84A02	Jul 15	NEUST	500	HOR	0.0	488
4	84A02	JUL 15	WISCN	760	HOR	13.7	240
5	84A03	Jul 15	WISCN	760	HOR	10.2	427
6	84A03	Jul 15	NEUST	500	HOR	0.0	544
7	84A04	JUL 15	NEUST	500	HOR	0.0	409
8	84A04	Jul 15	WISCN	760	HOR	11.3	370
9	84A05	Jul 16	WISCN	760	HOR	5.1	428
10	84A05	Jul 16	NEUST	500	HOR	0.0	538
11	84A06	Jul 23	NEUST	500	HOR	0.0	515
12	84A06	Jul 23	WISCN	760	HOR	5.1	361
13	84A07	Jul 23	NEUST	500	HOR	0.0	498
14	84A07	Jul 23	WISCN	760	HOR	13.7	448
15	84A08	Jul 23	NEUST	500	HOR	0.0	594
16	84A08	Jul 23	WISCN	760	HOR	11.1	477
17	84B09	Jul 24	NEUST	500	HOR	0.0	321
18	84B09	Jul 24	WISCN	760	HOR	10.2	248
19	84A10	Jul 24	NEUST	500	HOR	0.0	497
20	84A10	Jul 24	WISCN	760	HOR	8.2	383
21	84A11	Jul 25	WISCN	760	HOR	N/A*	N/A
22	84A12	Jul 25	WISCN	760	HOR	4.0	555
29	84B01	Sep 07	NEUST	500	HOR	0.0	532
30	84B01	Sep 07	WISCN	760	HOR	4.7	395
31	84B02	Sep 07	NEUST	500	HOR	0.0	637
32	84B02	Sep 07	WISCN	760	HOR	11.0	483
33	84B03	Sep 08	NEUST	500	HOR	0.0	567
34	84B03	Sep 08	WISCN	760	HOR	11.0	497
35	84B04	Sep 08	NEUST	500	HOR	0.0	496
36	84B04	Sep 08	WISCN	760	HOR	11.7	392
37	84B05	Sep 09	NEUST	500	HOR	0.0	579
38	84B05	Sep 09	WISCN	760	HOR	7.5	442
39	84B11	Sep 09	NEUST	500	HOR	0.0	609
40	84B11	Sep 09	WISCN	760	HOR	12.4	534

* N/A = Not available

Table 3. Stomach fullness codes used in Table 15.

Code	Description
0	Empty
1	1/3 full, digested
2	1/3 full, good condition
3	2/3 full, digested
4	2/3 full, good condition
5	Full, digested
6	Full, good condition

Table 4. Scientific and common names of fishes captured in the Canadian Beaufort Sea Shelf, 1984.

	Scientific Name	Common Name
Family Osmeridae		
	<u>Osmerus mordax</u> (Mitchill)	Rainbow smelt
Family Gadidae		
	<u>Boreogadus saida</u> (Lepechin)	Arctic cod
	<u>Eliginus gracilis</u> (Tilesius)	Saffron cod
Family Cottidae		
	<u>Myoxcephalus quadricornis</u> (Linnaeus)	Fourhorn sculpin
	<u>Gymnophanthis tricuspidis</u> (Reinhardt)	Staghorn sculpin
	<u>Triglops pingeli</u> Reinhardt	Ribbed sculpin
Family Cyclopteridae		
	<u>Liparis</u> sp.	Snailfish
Family Stichaeidae		
	<u>Anisarchus medius</u> (Reinhardt)	Stout eelblenny
Family Agonidae		
	<u>Aspidophoroides olriki</u> Lutken	Arctic alligatorfish

Table 5. Number, percent by number, and frequency of occurrence for larval fish captured by Wisconsin tows in the Beaufort Sea Shelf during July and September, 1984.

Species	JULY			SEPT		
	Number	%	Freq Occ	Number	%	Freq Occ
Gadidae (unid)	24	2.2	8.3	0	0	0
<u>B. saida</u>	1037	93.0	66.7	68	95.8	100
<u>E. gracilis</u>	1	0.1	8.3	0	0	0
Cottidae (unid)	2	0.2	16.7	0	0	0
<u>G. tricuspidis</u>	22	2.0	50.0	3	4.2	50
<u>M. quadricornis</u>	16	1.4	41.7	0	0	0
<u>T. pingeli</u>	1	0.1	8.3	0	0	0
<u>A. olriki</u>	1	0.1	8.3	0	0	0
Liparis sp.	8	0.7	33.3	0	0	0
<u>A. medius</u>	1	0.1	8.3	0	0	0
Unid. fish	2	0.2	16.7	0	0	0

Table 6. Larval fish catch summary for Wisconsin tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS																				
	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	T	
	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	0	
	A	A	A	A	A	A	A	B	A	A	B	B	B	B	B	B	B	B	B	T	
	0	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	1	A	
	1	2	3	4	5	6	7	8	9	0	1	2	1	2	3	4	5	1	L		
O. mordax																				0	
Gadidae (unid)																				24	
B. saida	412	16	154	10	314	64	43				24				1	26	20	19	1	1	1105
E. gracilis	1																			1	
Cottidae (unid)												1	1							2	
G. tricuspis	14	1	3		2		1		1						1	1	1	1	1	25	
M. quadricornis	2	1					1		2	10										16	
T. pingelli							1													1	
A. olriki			1																	1	
Liparis sp.		3					1	1	3											8	
A. medius		1																		1	
Unid. fish	1	1																		2	
TOTAL	4	432	17	158	10	316	65	47	24	31	10	1	1	1	27	20	20	1	2	1186	

Table 7. Larval fish number·100 m⁻³ for Wisconsin tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS																	
	8 4 A 0 1	8 4 A 0 2	8 4 A 0 3	8 4 A 0 4	8 4 A 0 5	8 4 A 0 6	8 4 A 0 7	8 4 A 0 8	8 4 A 0 9	8 4 A 1 0	8 4 A 1 1	8 4 A 1 2	8 4 B 0 1	8 4 B 0 2	8 4 B 0 3	8 4 B 0 4	8 4 B 0 5	8 4 B 0 6
<i>O. mordax</i>																		
Gadidae (unid)																		
<i>B. sarda</i>	171.7	3.7	41.6	2.3	87.0	14.3	9.0			6.3			0.3	5.4	4.0	4.8	0.2	0.2
<i>E. gracilis</i>	0.2																	
Cottidae (unid)										0.3			0.2					
<i>G. tricuspidis</i>	5.8	0.2	0.8		0.6			0.2		0.2			0.2		0.3		0.2	
<i>M. quadricornis</i>	0.4	0.4						0.2		0.5	*							
<i>T. pingelli</i>							0.2											
<i>A. olrikii</i>			0.2															
Liparis sp.		1.3					0.2	0.2		0.8								
<i>A. medius</i>		0.4																
Unid. fish	0.2	0.4	0.7															
TOTAL	0.8	180.0	4.6	42.6	2.3	87.6	14.5	9.8	9.7	8.1		0.2	0.3	5.6	4.0	5.1	0.2	0.4

* - Not available

Table 8. Number, percent by number and frequency of occurrence of larval fish captured by neuston tows on the Beaufort Sea Shelf during July and September, 1984.

Species	JULY			SEPT		
	Number	%	Freq Occ	Number	%	Freq Occ
O. <u>mordax</u>	1	0.5	10.0	0	0	0
B. <u>saida</u>	218	99.1	70.0	1	50	16.7
Cottidae (unid.)	0	0.0	0.0	1	50	16.7
M. <u>quadricornis</u>	1	0.5	10.0	0	0	0

Table 9. Larval fish catch summary for neuston tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS																		T
	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	8	
	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	4	0
	A	A	A	A	A	A	A	B	A	B	B	B	B	B	B	B	B	T	
	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	1	A	
	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	1	L		
<i>O. mordax</i>			1															1	
Gadidae (unid)																		0	
<i>B. saida</i>			5	166	1	2	.26	10		8		1						219	
<i>E. gracilis</i>																		0	
Cottidae (unid)																1	1		
<i>G. tricuspis</i>																		0	
<i>M. quadricornis</i>	1																	1	
<i>T. pingelli</i>																		0	
<i>A. olriki</i>																		0	
Liparis sp.																		0	
<i>A. medius</i>																		0	
Unid. fish																		0	
TOTAL	2	0	5	166	1	2	26	10	0	8	0	1	0	0	0	1	222		

Table 10. Larval fish number·100 m⁻³ for neuston tows on the Beaufort Sea Shelf, 1984.

SPECIES	STATIONS															
	1	2	3	4	5	6	7	8	9	0	1	2	3	4	5	6
<i>O. mordax</i>	0.1															
Gadidae (unid)																
<i>B. sarda</i>	0.9	40.6	0.2	0.4	5.2	1.7				1.6		0.2				
<i>E. gracilis</i>																
Cottidae (unid)																
<i>G. tricuspidis</i>																
<i>M. quadricornis</i>	0.1															
<i>T. pingelli</i>																
<i>A. olrikii</i>																
Liparis sp.																
<i>A. medius</i>																
Unid. fish																
TOTAL	0.2	0.0	0.9	40.6	0.2	0.4	5.2	1.7	0.0	1.6	0.0	0.2	0.0	0.0	0.0	0.2

Table 11. Standard and total lengths, and wet weight of larval fish captured in the Beaufort Sea, 1984.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
1	40	<i>O. mordax</i>	9.0	9.0	0.002	4	537	<i>B. saida</i>	14.0	15.0	0.017
4	270	<i>B. saida</i>	15.5	17.0	0.028	4	538	<i>B. saida</i>	10.0	10.5	0.006
4	271	<i>B. saida</i>	13.0	14.0	0.015	4	539	<i>B. saida</i>	12.0	13.0	0.013
4	273	<i>B. saida</i>	12.0	13.0	0.009	4	540	<i>B. saida</i>	12.0	13.0	0.013
4	274	<i>B. saida</i>	11.0	12.0	0.007	4	541	<i>B. saida</i>	11.5	12.5	0.010
4	275	<i>B. saida</i>	15.0	16.0	0.023	4	542	<i>B. saida</i>	13.0	14.0	0.014
4	277	<i>B. saida</i>	15.0	16.0	0.018	4	543	<i>B. saida</i>	13.0	14.0	0.012
4	278	<i>B. saida</i>	10.0	11.0	0.007	4	544	<i>B. saida</i>	11.5	12.5	0.009
4	281	<i>B. saida</i>	13.0	14.0	0.015	4	545	<i>B. saida</i>	11.0	12.0	0.006
4	282	<i>B. saida</i>	12.5	13.5	0.013	4	546	<i>B. saida</i>	11.0	12.0	0.007
4	283	<i>B. saida</i>	14.0	15.0	0.019	4	547	<i>B. saida</i>	12.0	13.0	0.011
4	284	<i>B. saida</i>	13.5	14.5	0.021	4	548	<i>B. saida</i>	12.5	13.5	0.011
4	285	<i>B. saida</i>	13.0	14.0	0.017	4	549	<i>B. saida</i>	11.0	12.0	0.011
4	288	<i>B. saida</i>	9.5	10.5	0.007	4	550	<i>B. saida</i>	13.0	14.0	0.012
4	289	<i>B. saida</i>	12.0	13.0	0.006	4	551	<i>B. saida</i>	16.0	17.0	0.029
4	291	<i>B. saida</i>	10.0	11.0	0.006	4	552	<i>B. saida</i>	13.0	14.0	0.014
4	292	<i>B. saida</i>	17.5	20.0	0.040	4	553	<i>B. saida</i>	12.0	13.0	0.008
4	293	<i>B. saida</i>	13.5	14.5	0.019	4	554	<i>B. saida</i>	12.5	13.5	0.013
4	294	<i>B. saida</i>	10.0	11.0	0.006	4	555	<i>B. saida</i>	11.0	12.0	0.007
4	295	<i>B. saida</i>	17.0	18.0	0.030	4	556	<i>B. saida</i>	12.0	13.0	0.009
4	513	<i>B. saida</i>	16.0	17.0	0.025	4	557	<i>B. saida</i>	13.0	14.0	0.014
4	514	<i>B. saida</i>	16.0	17.0	0.031	4	558	<i>B. saida</i>	14.0	15.0	0.018
4	515	<i>B. saida</i>	12.0	13.0	0.009	4	559	<i>B. saida</i>	12.0	13.0	0.010
4	516	<i>B. saida</i>	14.0	15.0	0.024	4	560	<i>B. saida</i>	11.0	12.0	0.013
4	517	<i>B. saida</i>	13.5	14.5	0.016	4	561	<i>B. saida</i>	12.5	13.5	0.015
4	518	<i>B. saida</i>	19.5	20.5	0.024	4	562	<i>B. saida</i>	11.0	12.0	0.006
4	519	<i>B. saida</i>	5.5	6.5	0.007	4	563	<i>B. saida</i>	12.0	13.0	0.010
4	520	<i>B. saida</i>	15.5	16.5	0.025	4	564	<i>B. saida</i>	15.5	16.5	0.028
4	521	<i>B. saida</i>	12.0	13.0	0.012	4	565	<i>B. saida</i>	11.5	12.5	0.011
4	522	<i>B. saida</i>	11.0	12.0	0.010	4	566	<i>B. saida</i>	12.0	13.0	0.010
4	523	<i>B. saida</i>	12.0	13.0	0.013	4	568	<i>B. saida</i>	12.0	13.0	0.011
4	524	<i>B. saida</i>	17.0	18.5	0.036	4	569	<i>B. saida</i>	12.0	13.0	0.009
4	525	<i>B. saida</i>	14.5	15.5	0.019	4	570	<i>B. saida</i>	12.0	13.0	0.012
4	526	<i>B. saida</i>	15.0	16.0	0.026	4	571	<i>B. saida</i>	10.5	11.5	0.008
4	527	<i>B. saida</i>	12.5	13.5	0.014	4	572	<i>B. saida</i>	14.0	15.0	0.018
4	528	<i>B. saida</i>	14.5	15.5	0.022	4	573	<i>B. saida</i>	11.0	12.0	0.009
4	529	<i>B. saida</i>	12.0	13.0	0.011	4	574	<i>B. saida</i>	12.0	13.0	0.009
4	530	<i>B. saida</i>	13.5	14.5	0.011	4	575	<i>B. saida</i>	14.0	15.0	0.018
4	531	<i>B. saida</i>	12.0	13.0	0.010	4	576	<i>B. saida</i>	9.5	10.0	0.007
4	532	<i>B. saida</i>	14.0	15.0	0.013	4	577	<i>B. saida</i>	11.0	12.0	0.005
4	533	<i>B. saida</i>	12.0	13.0	0.011	4	578	<i>B. saida</i>	12.0	13.0	0.012
4	534	<i>B. saida</i>	14.0	15.0	0.020	4	795	<i>B. saida</i>	16.0	17.0	0.021
4	535	<i>B. saida</i>	13.0	14.0	0.018	4	796	<i>B. saida</i>	16.0	17.0	0.022
4	536	<i>B. saida</i>	12.0	13.0	0.009	4	797	<i>B. saida</i>	14.5	15.5	0.020

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
4	798	B. saida	10.5	11.5	0.009	4	1014	B. saida	14.0	15.0	0.018
4	799	B. saida	14.0	15.0	0.018	4	1015	B. saida	13.0	14.0	0.013
4	800	B. saida	15.0	16.0	0.028	4	1016	B. saida	14.5	15.5	0.022
4	801	B. saida	6.5	7.5	0.009	4	1017	B. saida	14.0	15.0	0.020
4	802	B. saida	10.0	11.0	0.007	4	1018	B. saida	13.5	14.5	0.014
4	803	B. saida	15.5	16.5	0.021	4	1019	B. saida	12.5	13.5	0.014
4	804	B. saida	13.0	14.0	0.014	4	1020	B. saida	12.0	13.0	0.013
4	805	B. saida	12.5	13.5	0.013	4	1021	B. saida	13.5	14.5	0.016
4	806	B. saida	17.0	18.0	0.018	4	1022	B. saida	14.0	15.0	0.019
4	807	B. saida	17.0	18.0	0.012	4	1023	B. saida	12.0	13.0	0.013
4	808	B. saida	14.0	15.0	0.019	4	1024	B. saida	11.0	11.5	0.007
4	809	B. saida	11.5	12.5	0.011	4	1025	B. saida	15.0	16.0	0.022
4	810	B. saida	13.0	14.0	0.016	4	1026	B. saida	13.0	14.0	0.015
4	812	B. saida	13.5	14.5	0.017	4	1027	B. saida	14.5	15.5	0.024
4	813	B. saida	11.0	12.0	0.010	4	1028	B. saida	15.0	16.0	0.022
4	814	B. saida	12.0	13.0	0.010	4	1029	B. saida	11.5	12.5	0.008
4	815	B. saida	14.0	15.0	0.013	4	1030	B. saida	12.0	13.0	0.014
4	816	B. saida	13.5	14.5	0.015	4	1031	B. saida	15.0	16.0	0.020
4	818	B. saida	12.0	13.0	0.013	4	1032	B. saida	14.5	15.5	0.021
4	819	B. saida	14.5	15.5	0.021	4	1033	B. saida	14.0	15.0	0.015
4	820	B. saida	14.5	15.5	0.021	4	1034	B. saida	13.0	14.0	0.017
4	821	B. saida	12.0	13.0	0.014	4	1035	B. saida	15.0	16.0	0.022
4	822	B. saida	10.0	11.0	0.010	4	1036	B. saida	12.5	13.5	0.014
4	823	B. saida	13.0	14.0	0.019	4	1037	B. saida	11.0	12.0	0.010
4	824	B. saida	12.5	13.5	0.012	4	1038	B. saida	10.5	11.0	0.006
4	825	B. saida	13.0	14.0	0.016	4	1039	B. saida	14.0	15.0	0.016
4	826	B. saida	16.0	17.0	0.030	4	1040	B. saida	13.0	14.0	0.013
4	827	B. saida	13.5	14.5	0.018	4	1041	B. saida	12.0	13.0	0.013
4	828	B. saida	11.0	12.0	0.009	4	1042	B. saida	14.0	15.0	0.015
4	829	B. saida	11.0	12.0	0.009	4	1043	B. saida	12.0	13.0	0.011
4	830	B. saida	12.0	13.0	0.012	4	1044	B. saida	13.5	14.5	0.015
4	831	B. saida	14.0	15.0	0.017	4	1045	B. saida	15.0	16.0	0.020
4	832	B. saida	13.0	14.0	0.016	4	1046	B. saida	14.0	15.0	0.018
4	1002	B. saida	14.0	15.0	0.015	4	1047	B. saida	12.5	13.5	0.011
4	1003	B. saida	7.5	8.5	0.011	4	1048	B. saida	12.5	13.5	0.010
4	1004	B. saida	16.0	17.0	0.025	4	1049	B. saida	10.5	11.5	0.006
4	1005	B. saida	15.0	16.0	0.017	4	1050	B. saida	13.0	14.0	0.013
4	1006	B. saida	14.0	15.0	0.016	4	1051	B. saida	12.5	13.5	0.012
4	1007	B. saida	13.5	14.5	0.017	4	1052	B. saida	12.0	13.0	0.008
4	1008	B. saida	14.0	15.0	0.017	4	1053	B. saida	14.5	15.5	0.017
4	1009	B. saida	12.5	13.5	0.014	4	1054	B. saida	14.0	15.0	0.020
4	1010	B. saida	11.5	12.0	0.010	4	1055	B. saida	11.0	12.0	0.008
4	1011	B. saida	13.0	14.0	0.015	4	1056	B. saida	13.5	14.5	0.012
4	1013	B. saida	17.0	18.0	0.028	4	1057	B. saida	15.0	16.0	0.019

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
4	1058	B. saida	14.5	15.5	0.017	4	1111	B. saida	15.0	16.0	0.019
4	1059	B. saida	13.0	14.0	0.017	4	1112	B. saida	15.0	16.0	0.021
4	1060	B. saida	12.0	13.0	0.009	4	1113	B. saida	13.5	14.5	0.013
4	1061	B. saida	14.0	15.0	0.020	4	1114	B. saida	11.0	12.0	0.008
4	1062	B. saida	11.0	12.0	0.008	4	1115	B. saida	13.0	14.0	0.014
4	1063	B. saida	13.5	14.5	0.014	4	1116	B. saida	14.5	15.5	0.020
4	1064	B. saida	12.5	13.5	0.013	4	1117	B. saida	13.0	14.0	0.013
4	1065	B. saida	11.0	12.0	0.007	4	1118	B. saida	12.0	13.0	0.013
4	1066	B. saida	14.5	15.5	0.017	4	1119	B. saida	13.5	14.5	0.015
4	1067	B. saida	12.0	13.0	0.013	4	1121	B. saida	12.0	13.0	0.015
4	1068	B. saida	12.5	13.5	0.012	4	1122	B. saida	14.5	15.5	0.014
4	1069	B. saida	12.0	13.0	0.010	4	1123	B. saida	12.0	13.0	0.010
4	1070	B. saida	13.0	14.0	0.013	4	1124	B. saida	13.5	14.5	0.014
4	1071	B. saida	13.0	14.0	0.011	4	1125	B. saida	14.0	15.0	0.020
4	1072	B. saida	16.0	17.0	0.021	4	1126	B. saida	15.0	16.0	0.025
4	1073	B. saida	15.0	16.0	0.028	4	1127	B. saida	15.5	17.0	0.024
4	1074	B. saida	13.0	14.0	0.012	4	1128	B. saida	15.5	16.5	0.019
4	1075	B. saida	12.5	13.5	0.015	4	1129	B. saida	12.0	13.0	0.011
4	1076	B. saida	11.5	12.5	0.012	4	1130	B. saida	13.5	14.5	0.016
4	1077	B. saida	15.0	16.0	0.021	4	1131	B. saida	14.0	15.0	0.020
4	1079	B. saida	12.5	13.5	0.011	4	1132	B. saida	16.0	17.0	0.022
4	1080	B. saida	10.0	11.0	0.009	4	1133	B. saida	11.0	12.0	0.008
4	1081	B. saida	14.0	15.0	0.016	4	1134	B. saida	14.0	15.0	0.014
4	1082	B. saida	15.0	16.0	0.023	4	1135	B. saida	12.5	13.5	0.010
4	1083	B. saida	10.0	11.0	0.008	4	1136	B. saida	13.0	14.0	0.014
4	1084	B. saida	11.0	12.0	0.013	4	1137	B. saida	12.0	13.0	0.013
4	1085	B. saida	13.0	14.0	0.015	4	1138	B. saida	12.0	13.0	0.010
4	1094	B. saida	15.0	16.0	0.023	4	1139	B. saida	12.0	13.0	0.011
4	1095	B. saida	13.0	14.0	0.013	4	1140	B. saida	11.0	12.0	0.007
4	1096	B. saida	14.0	15.0	0.022	4	1141	B. saida	13.5	14.5	0.014
4	1097	B. saida	16.5	17.5	0.026	4	1142	B. saida	11.5	12.5	0.012
4	1098	B. saida	16.5	17.5	0.026	4	1143	B. saida	12.0	13.0	0.013
4	1099	B. saida	13.5	14.5	0.015	4	1144	B. saida	12.5	13.5	0.012
4	1100	B. saida	15.0	16.0	0.023	4	1145	B. saida	14.0	15.0	0.017
4	1101	B. saida	12.0	13.0	0.011	4	1146	B. saida	13.0	14.0	0.013
4	1102	B. saida	15.0	16.0	0.018	4	1147	B. saida	13.5	14.5	0.020
4	1103	B. saida	12.0	13.0	0.014	4	1148	B. saida	13.5	14.5	0.014
4	1104	B. saida	11.5	12.5	0.009	4	1149	B. saida	10.5	11.5	0.010
4	1105	B. saida	15.0	16.0	0.019	4	1150	B. saida	12.0	13.0	0.014
4	1106	B. saida	13.0	14.0	0.016	4	1151	B. saida	15.0	16.0	0.023
4	1107	B. saida	13.0	14.0	0.015	4	1152	B. saida	12.0	13.0	0.009
4	1108	B. saida	14.5	15.5	0.021	4	1153	B. saida	10.0	11.0	0.008
4	1109	B. saida	13.0	14.0	0.019	4	1154	B. saida	12.0	13.0	0.014
4	1110	B. saida	11.5	12.5	0.009	4	1155	B. saida	14.0	15.0	0.017

Table 11. Continued.

	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)		PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
4	1156	B. saida		12.5	13.5	0.014	4	1202	B. saida		12.0	13.0	0.010
4	1157	B. saida		13.5	14.5	0.012	4	1203	B. saida		14.5	15.5	0.019
4	1158	B. saida		9.0	9.5	0.003	4	1204	B. saida		12.0	13.0	0.009
4	1159	B. saida		15.5	16.5	0.021	4	1205	B. saida		11.0	12.0	0.008
4	1160	B. saida		13.5	14.5	0.018	4	1207	B. saida		13.0	14.0	0.015
4	1161	B. saida		14.0	15.0	0.018	4	1208	B. saida		10.0	10.5	0.005
4	1162	B. saida		12.0	13.0	0.010	4	1210	B. saida		13.0	14.0	0.012
4	1163	B. saida		12.0	13.0	0.010	4	1211	B. saida		10.5	11.5	0.007
4	1164	B. saida		13.0	14.0	0.016	4	1212	B. saida		12.5	13.5	0.011
4	1165	B. saida		13.5	14.5	0.018	4	1213	B. saida		10.5	11.5	0.008
4	1166	B. saida		12.0	13.0	0.014	4	1214	B. saida		12.0	13.0	0.014
4	1167	B. saida		12.5	13.5	0.013	4	1215	B. saida		10.0	11.0	0.006
4	1169	B. saida		9.0	9.5	0.004	4	1216	B. saida		12.0	13.0	0.011
4	1170	B. saida		12.0	13.0	0.012	4	1217	B. saida		11.0	12.0	0.008
4	1171	B. saida		10.0	11.0	0.007	4	1218	B. saida		11.0	12.0	0.007
4	1172	B. saida		16.5	17.5	0.023	4	1219	B. saida		16.0	17.0	0.023
4	1173	B. saida		16.0	17.0	0.028	4	1220	B. saida		13.0	14.0	0.016
4	1174	B. saida		13.5	14.5	0.013	4	1221	B. saida		13.0	14.0	0.014
4	1175	B. saida		10.0	10.5	0.008	4	1222	B. saida		16.0	17.0	0.019
4	1176	B. saida		14.0	15.0	0.016	4	1223	B. saida		13.0	14.0	0.015
4	1177	B. saida		10.0	11.0	0.009	4	1224	B. saida		12.0	13.0	0.011
4	1178	B. saida		18.0	19.0	0.034	4	1225	B. saida		13.0	14.0	0.017
4	1179	B. saida		10.0	11.0	0.008	4	1226	B. saida		13.0	14.0	0.017
4	1180	B. saida		11.0	12.0	0.008	4	1227	B. saida		14.0	15.0	0.021
4	1181	B. saida		12.5	13.5	0.011	4	1228	B. saida		15.0	16.0	0.023
4	1182	B. saida		11.5	12.5	0.009	4	1229	B. saida		15.0	16.0	0.017
4	1183	B. saida		14.0	15.0	0.021	4	1230	B. saida		14.5	15.5	0.020
4	1184	B. saida		10.5	11.5	0.007	4	1231	B. saida		14.0	15.0	0.020
4	1186	B. saida		11.0	12.0	0.006	4	1232	B. saida		12.0	13.0	0.012
4	1187	B. saida		12.0	13.0	0.014	4	1233	B. saida		14.0	15.0	0.020
4	1188	B. saida		13.0	14.0	0.016	4	1234	B. saida		17.5	18.5	0.038
4	1189	B. saida		15.0	16.0	0.023	4	1235	B. saida		13.0	14.0	0.019
4	1190	B. saida		12.5	13.5	0.014	4	1236	B. saida		11.0	12.0	0.010
4	1191	B. saida		11.5	12.5	0.012	4	1237	B. saida		14.5	15.5	0.026
4	1192	B. saida		14.5	15.5	0.024	4	1238	B. saida		12.0	13.0	0.012
4	1193	B. saida		12.5	13.5	0.014	4	1239	B. saida		11.0	11.5	0.008
4	1194	B. saida		13.5	14.5	0.016	4	1240	B. saida		15.0	16.0	0.023
4	1195	B. saida		11.5	12.5	0.007	4	1241	B. saida		16.0	17.0	0.035
4	1196	B. saida		15.5	16.5	0.019	4	1242	B. saida		13.0	14.0	0.018
4	1197	B. saida		12.0	13.0	0.009	4	1243	B. saida		13.5	14.5	0.015
4	1198	B. saida		15.5	16.5	0.022	4	1244	B. saida		12.0	13.0	0.014
4	1199	B. saida		11.5	12.5	0.007	4	1245	B. saida		12.5	13.5	0.013
4	1200	B. saida		13.5	14.5	0.014	4	1246	B. saida		13.0	14.0	0.014
4	1201	B. saida		10.5	11.5	0.008	4	1247	B. saida		12.0	13.0	0.010

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
4	1248	B. saida	11.5	12.5	0.011	4	21	G. tricuspidis	15.0	17.5	0.026
4	1249	B. saida	11.5	12.0	0.010	4	24	G. tricuspidis	15.0	17.0	0.033
4	1250	B. saida	14.0	15.0	0.021	4	1325	G. tricuspidis	11.5	13.0	0.011
4	1251	B. saida	11.5	12.5	0.013	4	1326	G. tricuspidis	13.0	15.0	0.015
4	1252	B. saida	13.5	14.5	0.014	4	1327	G. tricuspidis	15.0	17.5	0.030
4	1253	B. saida	10.5	11.0	0.006	4	1339	A. medius	19.5	21.0	0.013
4	1254	B. saida	12.0	13.0	0.014	5	244	B. saida	12.0	13.0	0.009
4	1255	B. saida	10.0	10.5	0.007	5	245	B. saida	9.5	10.0	0.002
4	1257	B. saida	9.5	10.0	0.003	5	246	B. saida	10.0	11.0	0.006
4	1258	B. saida	10.0	10.5	0.005	5	247	B. saida	12.5	13.5	0.014
4	1284	B. saida	14.0	15.0	0.019	5	248	B. saida	9.0	10.0	0.003
4	1285	B. saida	12.5	13.5	0.012	5	249	B. saida	13.0	14.0	0.014
4	1286	B. saida	15.5	17.0	0.031	5	250	B. saida	10.5	11.0	0.003
4	1287	B. saida	12.0	13.0	0.013	5	251	B. saida	9.5	10.5	0.003
4	1288	B. saida	13.0	14.0	0.015	5	252	B. saida	9.5	10.5	0.003
4	1289	B. saida	15.5	16.5	0.021	5	253	B. saida	10.0	11.0	0.005
4	1290	B. saida	14.5	15.5	0.019	5	254	B. saida	10.0	11.0	0.005
4	1291	B. saida	17.5	18.5	0.027	5	256	B. saida	12.0	13.0	0.010
4	1292	B. saida	15.0	16.0	0.023	5	257	B. saida	8.0	8.5	0.000
4	1294	B. saida	15.5	16.5	0.023	5	258	B. saida	7.5	8.0	0.000
4	1295	B. saida	18.0	19.0	0.042	5	259	B. saida	7.5	8.0	0.000
4	1296	B. saida	15.0	16.0	0.023	5	1328	G. tricuspidis	11.5	13.0	0.011
4	1297	B. saida	12.0	13.0	0.014	7	833	B. saida	12.0	13.0	0.009
4	1298	B. saida	16.0	17.0	0.030	7	834	B. saida	9.5	9.5	0.003
4	1299	B. saida	11.0	12.0	0.011	7	835	B. saida	11.0	11.0	0.006
4	1300	B. saida	14.0	15.0	0.023	7	836	B. saida	10.0	10.0	0.005
4	1301	B. saida	17.5	19.0	0.037	7	837	B. saida	10.0	10.0	0.006
4	1302	B. saida	14.0	15.0	0.020	7	838	B. saida	9.0	9.0	0.003
4	1303	B. saida	11.0	12.0	0.011	7	839	B. saida	11.5	12.0	0.007
4	1304	B. saida	15.0	16.0	0.026	7	840	B. saida	10.5	11.0	0.004
4	1305	B. saida	15.0	16.0	0.021	7	841	B. saida	10.5	11.0	0.007
4	1306	B. saida	14.0	15.0	0.020	7	842	B. saida	9.0	9.0	0.002
4	1307	B. saida	15.0	16.0	0.030	7	843	B. saida	11.0	12.0	0.011
4	1308	B. saida	14.5	15.5	0.025	7	844	B. saida	11.0	11.0	0.006
4	1309	B. saida	14.0	15.0	0.021	7	845	B. saida	10.0	10.0	0.005
4	1310	B. saida	15.5	16.5	0.027	7	846	B. saida	11.0	12.0	0.007
4	1311	B. saida	12.5	13.5	0.011	7	847	B. saida	11.0	11.0	0.006
4	1312	B. saida	14.5	15.5	0.025	7	848	B. saida	7.0	7.0	0.002
4	1313	B. saida	14.5	15.5	0.022	7	849	B. saida	8.5	8.5	0.004
4	1314	B. saida	14.0	15.0	0.022	7	850	B. saida	9.0	9.5	0.006
4	1315	B. saida	15.0	16.0	0.021	7	851	B. saida	9.0	9.0	0.005
4	1316	B. saida	14.0	15.0	0.022	7	852	B. saida	9.0	9.5	0.009
4	1317	B. saida	12.5	13.5	0.016	7	853	B. saida	14.0	15.0	0.016
4	20	G. tricuspidis	15.0	17.5	0.027	7	854	B. saida	9.5	10.0	0.005

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
7	855	B. saida	9.5	9.5	0.002	7	901	B. saida	8.0	8.0	0.003
7	856	B. saida	10.5	11.0	0.008	7	902	B. saida	9.5	10.0	0.006
7	857	B. saida	10.0	10.0	0.006	7	903	B. saida	10.5	11.0	0.003
7	858	B. saida	10.5	11.0	0.006	7	904	B. saida	9.5	10.0	0.007
7	859	B. saida	10.0	10.5	0.004	7	905	B. saida	8.5	8.5	0.002
7	861	B. saida	11.0	11.5	0.011	7	906	B. saida	9.0	9.0	0.004
7	862	B. saida	10.5	11.0	0.007	7	907	B. saida	9.0	9.0	0.003
7	863	B. saida	10.0	10.5	0.006	7	908	B. saida	12.0	13.0	0.010
7	864	B. saida	10.0	10.5	0.006	7	909	B. saida	10.0	10.0	0.003
7	865	B. saida	9.5	9.5	0.005	7	910	B. saida	12.0	12.5	0.009
7	866	B. saida	10.5	11.0	0.007	7	911	B. saida	9.0	9.0	0.003
7	867	B. saida	10.0	10.0	0.003	7	912	B. saida	9.0	9.5	0.004
7	868	B. saida	9.0	9.0	0.002	7	913	B. saida	9.5	10.0	0.005
7	869	B. saida	9.5	10.0	0.007	7	914	B. saida	9.5	10.0	0.005
7	870	B. saida	8.5	8.5	0.002	7	915	B. saida	9.0	9.0	0.004
7	871	B. saida	9.0	9.0	0.004	7	916	B. saida	9.0	9.0	0.004
7	872	B. saida	11.0	11.0	0.005	7	917	B. saida	9.0	9.5	0.003
7	873	B. saida	9.0	9.0	0.007	7	918	B. saida	10.0	10.5	0.003
7	874	B. saida	9.5	10.0	0.004	7	919	B. saida	9.5	10.0	0.004
7	875	B. saida	9.5	10.0	0.004	7	920	B. saida	9.5	9.5	0.005
7	876	B. saida	10.5	11.0	0.005	7	921	B. saida	9.0	9.5	0.004
7	877	B. saida	9.0	9.0	0.003	7	922	B. saida	8.0	8.0	0.002
7	878	B. saida	9.0	9.0	0.002	7	923	B. saida	9.5	10.0	0.004
7	879	B. saida	10.5	11.0	0.009	7	924	B. saida	8.5	8.5	0.001
7	880	B. saida	9.0	9.0	0.003	7	926	B. saida	11.0	11.5	0.008
7	881	B. saida	10.0	10.0	0.004	7	927	B. saida	10.0	10.5	0.007
7	882	B. saida	11.0	11.5	0.008	7	928	B. saida	9.0	9.5	0.005
7	883	B. saida	10.0	10.5	0.007	7	930	B. saida	11.5	12.5	0.011
7	884	B. saida	10.0	10.5	0.004	7	931	B. saida	9.0	9.5	0.004
7	885	B. saida	9.0	9.0	0.003	7	933	B. saida	11.0	11.5	0.005
7	886	B. saida	10.0	10.5	0.005	7	934	B. saida	7.5	7.5	0.000
7	887	B. saida	8.5	9.0	0.004	7	935	B. saida	11.0	12.0	0.010
7	888	B. saida	9.5	10.0	0.006	7	936	B. saida	9.5	10.0	0.005
7	889	B. saida	10.5	11.0	0.008	7	937	B. saida	11.0	12.0	0.009
7	890	B. saida	9.0	9.0	0.004	7	938	B. saida	10.0	10.5	0.006
7	891	B. saida	9.5	10.0	0.004	7	939	B. saida	8.0	8.0	0.002
7	893	B. saida	9.5	10.0	0.003	7	940	B. saida	10.5	11.0	0.009
7	894	B. saida	13.0	14.0	0.013	7	941	B. saida	10.5	11.0	0.008
7	895	B. saida	12.0	13.0	0.010	7	942	B. saida	10.0	10.0	0.002
7	896	B. saida	10.0	10.5	0.006	7	943	B. saida	9.5	10.0	0.005
7	897	B. saida	8.0	8.0	0.002	7	944	B. saida	10.0	10.5	0.006
7	898	B. saida	8.0	8.0	0.002	7	945	B. saida	11.0	12.0	0.010
7	899	B. saida	9.5	10.0	0.008	7	946	B. saida	9.5	10.0	0.004
7	900	B. saida	8.0	8.0	0.002	7	947	B. saida	12.0	13.0	0.011

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
7	948	B. saida	9.0	9.5	0.003	7	994	B. saida	9.5	10.0	0.003
7	949	B. saida	10.0	10.5	0.004	7	995	B. saida	9.5	10.0	0.004
7	950	B. saida	9.5	10.0	0.005	7	996	B. saida	8.0	8.0	0.000
7	951	B. saida	9.0	9.0	0.004	7	997	B. saida	9.0	9.0	0.002
7	952	B. saida	10.5	11.0	0.007	8	47	B. saida	13.5	14.0	0.011
7	953	B. saida	9.0	9.5	0.005	8	48	B. saida	14.0	15.0	0.011
7	954	B. saida	8.0	8.0	0.001	8	49	B. saida	12.5	13.5	0.008
7	955	B. saida	9.0	9.5	0.004	8	50	B. saida	15.0	16.0	0.016
7	956	B. saida	9.0	9.0	0.005	8	52	B. saida	14.0	15.0	0.017
7	957	B. saida	8.0	8.5	0.002	8	53	B. saida	11.0	12.0	0.011
7	958	B. saida	12.0	13.0	0.011	8	54	B. saida	15.0	16.0	0.016
7	959	B. saida	9.0	9.0	0.003	8	55	B. saida	14.0	15.0	0.016
7	960	B. saida	9.0	9.5	0.003	8	56	B. saida	11.5	12.5	0.010
7	961	B. saida	10.0	10.0	0.002	8	57	B. saida	10.5	10.5	0.006
7	962	B. saida	10.5	11.0	0.005	8	58	B. saida	14.0	15.0	0.013
7	963	B. saida	9.5	10.0	0.007	8	59	B. saida	12.5	13.5	0.009
7	964	B. saida	9.5	10.0	0.006	8	60	B. saida	12.0	13.0	0.009
7	965	B. saida	10.0	10.5	0.004	8	61	B. saida	14.0	15.0	0.012
7	966	B. saida	10.5	11.0	0.005	8	62	B. saida	14.0	15.0	0.015
7	967	B. saida	9.0	9.0	0.004	8	63	B. saida	12.0	12.5	0.007
7	968	B. saida	9.5	10.0	0.005	8	64	B. saida	12.5	13.5	0.010
7	969	B. saida	10.0	10.0	0.003	8	65	B. saida	14.0	15.0	0.012
7	970	B. saida	10.0	10.0	0.004	8	66	B. saida	12.0	13.0	0.007
7	971	B. saida	9.0	9.0	0.002	8	67	B. saida	11.5	12.0	0.008
7	972	B. saida	10.0	10.5	0.005	8	68	B. saida	12.0	13.0	0.010
7	973	B. saida	10.0	10.5	0.005	8	69	B. saida	11.5	12.5	0.008
7	974	B. saida	9.5	10.0	0.004	8	70	B. saida	13.0	14.0	0.009
7	975	B. saida	9.0	9.0	0.002	8	71	B. saida	12.0	13.0	0.008
7	976	B. saida	11.5	12.5	0.010	8	72	B. saida	11.5	12.5	0.009
7	977	B. saida	8.0	8.0	0.002	8	73	B. saida	14.0	15.0	0.016
7	978	B. saida	8.0	8.0	0.001	8	74	B. saida	11.0	11.0	0.018
7	979	B. saida	12.0	13.0	0.011	8	75	B. saida	13.0	14.0	0.013
7	980	B. saida	8.0	8.0	0.002	8	76	B. saida	13.0	14.0	0.012
7	981	B. saida	9.0	9.0	0.003	8	77	B. saida	15.0	16.0	0.021
7	982	B. saida	10.0	11.0	0.006	8	78	B. saida	12.0	13.0	0.009
7	983	B. saida	12.0	13.0	0.009	8	79	B. saida	14.0	15.0	0.017
7	984	B. saida	11.0	12.0	0.008	8	80	B. saida	14.0	15.0	0.016
7	985	B. saida	10.5	11.0	0.007	8	81	B. saida	12.0	13.0	0.012
7	987	B. saida	11.5	12.0	0.008	8	82	B. saida	12.0	13.0	0.007
7	988	B. saida	10.0	10.5	0.006	8	83	B. saida	13.5	14.0	0.008
7	989	B. saida	10.0	10.5	0.005	8	84	B. saida	12.0	13.0	0.009
7	991	B. saida	9.5	9.5	0.006	8	85	B. saida	13.0	14.0	0.010
7	992	B. saida	9.5	9.5	0.003	8	86	B. saida	13.0	14.0	0.011
7	993	B. saida	9.5	10.0	0.004	8	87	B. saida	13.0	14.0	0.013

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
8	88	B. saida	13.5	14.5	0.010	8	140	B. saida	12.0	13.0	0.008
8	89	B. saida	12.0	13.0	0.010	8	141	B. saida	11.5	12.5	0.008
8	90	B. saida	14.0	15.0	0.015	8	143	B. saida	10.0	10.5	0.008
8	91	B. saida	13.0	14.0	0.012	8	144	B. saida	10.5	11.0	0.007
8	92	B. saida	12.5	13.5	0.009	8	145	B. saida	9.5	10.0	0.004
8	93	B. saida	10.0	10.5	0.006	8	146	B. saida	11.5	12.5	0.008
8	94	B. saida	15.5	16.5	0.018	8	147	B. saida	11.0	12.0	0.007
8	95	B. saida	13.5	14.5	0.011	8	148	B. saida	13.5	14.5	0.014
8	97	B. saida	12.5	13.5	0.010	8	149	B. saida	12.5	14.0	0.013
8	99	B. saida	11.5	12.5	0.009	8	150	B. saida	11.0	12.0	0.006
8	100	B. saida	12.0	13.0	0.010	8	151	B. saida	10.5	11.5	0.009
8	101	B. saida	11.0	12.0	0.008	8	152	B. saida	12.0	13.0	0.008
8	102	B. saida	13.0	14.0	0.011	8	153	B. saida	12.5	14.0	0.013
8	103	B. saida	12.5	13.5	0.010	8	154	B. saida	13.0	14.0	0.015
8	104	B. saida	13.0	14.0	0.014	8	155	B. saida	10.5	11.5	0.009
8	105	B. saida	13.0	14.0	0.013	8	156	B. saida	14.0	15.0	0.017
8	106	B. saida	11.5	12.5	0.011	8	157	B. saida	11.0	12.0	0.007
8	107	B. saida	12.5	13.5	0.008	8	158	B. saida	14.0	15.0	0.013
8	109	B. saida	9.0	10.0	0.004	8	159	B. saida	13.5	14.5	0.013
8	111	B. saida	12.0	13.0	0.011	8	160	B. saida	10.0	11.0	0.005
8	112	B. saida	13.0	14.0	0.012	8	161	B. saida	12.5	13.5	0.011
8	113	B. saida	12.5	13.5	0.010	8	162	B. saida	14.0	15.0	0.017
8	114	B. saida	16.0	17.5	0.024	8	163	B. saida	9.0	10.0	0.004
8	116	B. saida	12.0	13.0	0.010	8	164	B. saida	12.0	13.0	0.011
8	117	B. saida	14.0	15.0	0.013	8	165	B. saida	12.0	13.0	0.008
8	120	B. saida	12.0	13.0	0.008	8	166	B. saida	13.0	14.0	0.012
8	121	B. saida	13.0	14.0	0.015	8	167	B. saida	10.5	11.5	0.007
8	122	B. saida	13.5	14.5	0.013	8	169	B. saida	11.0	12.0	0.005
8	123	B. saida	12.0	13.0	0.009	8	170	B. saida	10.0	10.5	0.004
8	124	B. saida	15.5	17.0	0.020	8	171	B. saida	12.0	13.0	0.007
8	126	B. saida	12.0	13.0	0.009	8	172	B. saida	11.0	12.0	0.006
8	127	B. saida	12.0	13.0	0.011	8	173	B. saida	11.0	12.0	0.007
8	128	B. saida	14.0	15.0	0.013	8	174	B. saida	13.0	14.0	0.010
8	129	B. saida	13.5	15.0	0.015	8	175	B. saida	11.5	12.5	0.010
8	130	B. saida	11.0	12.0	0.006	8	176	B. saida	11.5	12.5	0.007
8	131	B. saida	11.0	12.0	0.007	8	178	B. saida	10.5	11.5	0.007
8	132	B. saida	13.0	14.0	0.013	8	179	B. saida	11.0	12.0	0.006
8	133	B. saida	13.0	14.0	0.012	8	182	B. saida	11.0	12.0	0.008
8	134	B. saida	12.0	13.0	0.007	8	183	B. saida	15.0	16.0	0.017
8	135	B. saida	13.0	14.0	0.011	8	184	B. saida	12.0	13.0	0.009
8	136	B. saida	14.0	15.0	0.013	8	186	B. saida	11.0	12.0	0.008
8	137	B. saida	14.0	15.0	0.012	8	188	B. saida	13.5	14.5	0.013
8	138	B. saida	12.0	13.0	0.009	8	189	B. saida	8.0	9.0	0.003
8	139	B. saida	10.5	11.0	0.004	8	190	B. saida	12.0	13.0	0.009

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
8	191	B. saida	12.0	13.0	0.008	12	335	B. saida	17.0	18.0	0.022
8	192	B. saida	10.5	11.5	0.005	12	337	B. saida	14.0	15.0	0.010
8	193	B. saida	12.0	13.0	0.008	12	338	B. saida	17.0	18.5	0.024
8	194	B. saida	10.5	11.5	0.005	12	339	B. saida	17.5	19.0	0.024
8	197	B. saida	6.0	7.0	0.002	12	341	B. saida	15.0	16.0	0.014
8	198	B. saida	8.0	8.5	0.003	12	342	B. saida	17.0	18.0	0.020
8	1329	G. tricuspidis	11.5	12.5	0.006	12	343	B. saida	13.0	14.0	0.013
8	1330	G. tricuspidis	15.5	17.0	0.024	12	344	B. saida	19.0	21.0	0.036
8	1331	G. tricuspidis	13.0	14.5	0.015	12	345	B. saida	17.5	19.0	0.024
8	1337	A. olrikii	8.5	9.0	0.001	12	346	B. saida	17.0	18.5	0.020
10	605	B. saida	30.0	32.5	0.188	12	347	B. saida	16.5	17.5	0.018
11	1000	B. saida	17.0	18.5	0.021	12	348	B. saida	15.0	16.0	0.015
11	1001	B. saida	15.5	16.5	0.017	12	349	B. saida	14.5	15.5	0.014
12	296	B. saida	19.0	20.0	0.035	12	350	B. saida	15.0	16.0	0.014
12	297	B. saida	16.5	17.5	0.023	12	351	B. saida	15.0	16.0	0.014
12	298	B. saida	19.0	21.5	0.030	12	352	B. saida	12.0	13.0	0.009
12	299	B. saida	17.0	18.0	0.030	12	354	B. saida	17.5	18.5	0.022
12	301	B. saida	16.5	17.5	0.023	12	355	B. saida	17.5	19.0	0.024
12	302	B. saida	17.0	18.0	0.025	12	356	B. saida	15.5	16.5	0.017
12	303	B. saida	15.0	16.0	0.017	12	357	B. saida	14.0	15.5	0.014
12	304	B. saida	16.5	17.5	0.020	12	358	B. saida	18.0	19.5	0.027
12	305	B. saida	16.0	17.0	0.019	12	359	B. saida	14.0	15.0	0.011
12	306	B. saida	16.0	17.0	0.017	12	360	B. saida	16.5	17.5	0.020
12	307	B. saida	16.0	17.0	0.019	12	361	B. saida	18.0	19.0	0.026
12	308	B. saida	17.0	18.5	0.020	12	362	B. saida	15.0	16.0	0.017
12	310	B. saida	13.0	14.0	0.015	12	363	B. saida	16.5	17.5	0.020
12	311	B. saida	16.5	17.5	0.023	12	365	B. saida	18.0	19.0	0.028
12	312	B. saida	13.0	14.0	0.010	12	366	B. saida	16.0	17.0	0.021
12	313	B. saida	18.0	19.0	0.032	12	368	B. saida	13.0	14.0	0.008
12	314	B. saida	19.0	20.0	0.039	12	369	B. saida	14.0	15.0	0.014
12	316	B. saida	15.0	16.0	0.015	12	370	B. saida	18.0	19.5	0.026
12	318	B. saida	17.0	18.0	0.026	12	371	B. saida	15.0	16.0	0.019
12	319	B. saida	17.0	18.0	0.028	12	373	B. saida	16.0	17.0	0.019
12	321	B. saida	12.5	13.5	0.008	12	374	B. saida	16.5	17.5	0.018
12	322	B. saida	17.5	19.0	0.026	12	375	B. saida	14.0	15.0	0.012
12	323	B. saida	11.5	12.5	0.020	12	376	B. saida	12.0	13.0	0.007
12	324	B. saida	14.0	15.0	0.014	12	377	B. saida	15.0	16.0	0.013
12	325	B. saida	17.5	19.0	0.030	12	378	B. saida	16.0	17.0	0.024
12	326	B. saida	17.0	18.0	0.028	12	379	B. saida	16.0	17.0	0.020
12	328	B. saida	13.5	14.5	0.010	12	380	B. saida	14.0	15.0	0.010
12	329	B. saida	16.0	17.0	0.019	12	381	B. saida	17.0	18.0	0.028
12	330	B. saida	16.0	17.0	0.018	12	382	B. saida	17.0	18.0	0.021
12	331	B. saida	16.0	17.0	0.023	12	383	B. saida	15.0	16.0	0.015
12	332	B. saida	18.0	19.5	0.034	12	384	B. saida	16.0	17.0	0.022

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
12	385	B. saida	13.0	14.0	0.011	12	616	B. saida	10.0	11.0	0.011
12	387	B. saida	17.0	18.0	0.021	12	617	B. saida	15.0	16.0	0.017
12	388	B. saida	18.0	19.5	0.035	12	618	B. saida	14.0	15.0	0.013
12	389	B. saida	17.0	18.5	0.023	12	620	B. saida	18.5	20.0	0.035
12	390	B. saida	15.0	16.0	0.014	12	621	B. saida	17.5	18.5	0.026
12	393	B. saida	15.5	16.5	0.016	12	622	B. saida	15.0	16.0	0.016
12	394	B. saida	16.0	17.0	0.016	12	623	B. saida	16.5	18.0	0.031
12	396	B. saida	18.0	19.0	0.026	12	624	B. saida	15.0	16.0	0.015
12	397	B. saida	19.0	20.5	0.027	12	625	B. saida	14.0	15.0	0.009
12	398	B. saida	15.0	16.0	0.016	12	628	B. saida	16.0	17.0	0.016
12	399	B. saida	19.0	20.5	0.033	12	629	B. saida	17.5	19.0	0.029
12	401	B. saida	16.5	17.5	0.020	12	630	B. saida	14.5	15.5	0.016
12	402	B. saida	15.0	16.0	0.010	12	632	B. saida	13.5	14.5	0.009
12	403	B. saida	16.5	17.5	0.019	12	633	B. saida	14.0	15.0	0.012
12	404	B. saida	17.0	18.0	0.018	12	634	B. saida	17.0	18.0	0.022
12	405	B. saida	14.5	15.5	0.017	12	635	B. saida	13.0	14.0	0.010
12	406	B. saida	14.0	15.0	0.009	12	636	B. saida	16.5	17.5	0.020
12	407	B. saida	16.5	17.5	0.025	12	637	B. saida	18.0	19.0	0.026
12	408	B. saida	17.5	19.0	0.029	12	638	B. saida	15.5	16.5	0.017
12	409	B. saida	17.0	18.5	0.022	12	639	B. saida	15.0	16.0	0.016
12	410	B. saida	18.0	19.5	0.026	12	640	B. saida	13.5	14.5	0.015
12	411	B. saida	14.0	15.0	0.012	12	641	B. saida	14.5	15.5	0.017
12	412	B. saida	13.0	14.0	0.010	12	642	B. saida	19.0	20.5	0.031
12	413	B. saida	15.0	16.0	0.015	12	643	B. saida	16.5	17.5	0.021
12	414	B. saida	13.0	14.0	0.009	12	644	B. saida	18.0	19.0	0.029
12	415	B. saida	17.0	18.0	0.022	12	645	B. saida	17.0	18.5	0.026
12	416	B. saida	14.5	15.5	0.015	12	646	B. saida	16.0	17.0	0.022
12	417	B. saida	17.0	18.5	0.021	12	647	B. saida	17.0	18.0	0.024
12	418	B. saida	18.5	20.0	0.031	12	648	B. saida	16.0	17.0	0.017
12	419	B. saida	12.0	13.0	0.008	12	649	B. saida	15.0	16.0	0.020
12	422	B. saida	15.0	16.0	0.016	12	651	B. saida	16.0	17.0	0.020
12	423	B. saida	16.0	17.0	0.025	12	652	B. saida	18.5	20.0	0.033
12	426	B. saida	15.0	16.0	0.015	12	653	B. saida	18.0	19.0	0.032
12	427	B. saida	15.0	16.0	0.014	12	654	B. saida	13.5	14.5	0.010
12	606	B. saida	14.0	15.0	0.016	12	655	B. saida	14.0	15.0	0.012
12	607	B. saida	15.0	16.0	0.015	12	656	B. saida	16.5	17.5	0.022
12	608	B. saida	15.0	16.0	0.015	12	657	B. saida	16.0	17.0	0.025
12	609	B. saida	18.0	20.0	0.038	12	658	B. saida	13.0	14.0	0.012
12	610	B. saida	13.0	14.0	0.015	12	659	B. saida	19.0	20.0	0.038
12	611	B. saida	17.0	18.0	0.017	12	660	B. saida	16.0	17.0	0.024
12	612	B. saida	18.0	19.0	0.026	12	661	B. saida	15.0	16.0	0.011
12	613	B. saida	18.0	20.0	0.037	12	662	B. saida	19.0	20.5	0.040
12	614	B. saida	16.5	17.5	0.026	12	663	B. saida	16.0	17.0	0.021
12	615	B. saida	9.5	10.5	0.017	12	664	B. saida	16.5	17.5	0.020

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
12	665	B. saida	16.0	17.0	0.021	12	713	B. saida	14.0	15.0	0.013
12	667	B. saida	14.5	15.5	0.015	12	714	B. saida	15.0	16.0	0.016
12	668	B. saida	16.0	17.0	0.014	12	715	B. saida	13.0	14.0	0.013
12	669	B. saida	15.0	16.0	0.012	12	716	B. saida	18.5	20.0	0.034
12	670	B. saida	14.0	15.0	0.016	12	717	B. saida	15.0	16.0	0.019
12	671	B. saida	18.0	19.5	0.031	12	718	B. saida	16.0	17.5	0.023
12	672	B. saida	13.5	14.5	0.013	12	719	B. saida	15.0	16.0	0.013
12	673	B. saida	15.0	16.0	0.015	12	720	B. saida	15.5	16.5	0.016
12	674	B. saida	13.0	14.0	0.014	12	721	B. saida	18.0	19.0	0.032
12	675	B. saida	16.5	18.0	0.024	12	722	B. saida	18.0	19.0	0.027
12	676	B. saida	14.0	15.0	0.009	12	723	B. saida	20.5	22.0	0.045
12	677	B. saida	16.0	17.0	0.021	12	724	B. saida	18.0	19.5	0.032
12	678	B. saida	14.0	15.0	0.018	12	725	B. saida	18.5	19.5	0.030
12	680	B. saida	16.0	17.0	0.021	12	726	B. saida	18.5	20.0	0.034
12	681	B. saida	15.5	16.5	0.018	12	727	B. saida	15.0	16.0	0.016
12	682	B. saida	15.5	16.5	0.016	12	729	B. saida	17.0	18.0	0.021
12	683	B. saida	18.5	20.0	0.039	12	730	B. saida	15.0	16.0	0.015
12	684	B. saida	18.5	20.0	0.035	12	731	B. saida	14.0	15.0	0.018
12	685	B. saida	20.0	22.0	0.041	12	732	B. saida	14.0	15.0	0.016
12	686	B. saida	14.0	15.0	0.015	12	733	B. saida	16.0	17.0	0.024
12	687	B. saida	17.0	18.0	0.032	12	734	B. saida	16.0	17.0	0.021
12	688	B. saida	16.0	17.0	0.020	12	736	B. saida	16.5	17.5	0.019
12	689	B. saida	15.0	16.0	0.018	12	737	B. saida	15.5	17.0	0.019
12	690	B. saida	15.5	16.5	0.016	12	738	B. saida	14.5	15.5	0.015
12	691	B. saida	12.5	13.5	0.008	12	739	B. saida	14.5	15.5	0.020
12	692	B. saida	17.0	18.0	0.027	12	740	B. saida	17.5	19.0	0.027
12	693	B. saida	16.5	18.0	0.020	12	741	B. saida	14.0	15.0	0.014
12	694	B. saida	14.5	15.5	0.014	12	743	B. saida	16.0	17.0	0.018
12	695	B. saida	14.5	15.5	0.015	12	744	B. saida	16.5	17.5	0.020
12	696	B. saida	14.0	15.0	0.014	12	745	B. saida	17.0	18.0	0.021
12	697	B. saida	18.5	19.5	0.031	12	746	B. saida	16.0	17.0	0.023
12	698	B. saida	13.5	14.5	0.012	12	747	B. saida	15.5	16.5	0.016
12	699	B. saida	12.0	13.0	0.007	12	748	B. saida	18.0	19.5	0.031
12	700	B. saida	15.0	16.0	0.014	12	749	B. saida	15.5	16.5	0.017
12	701	B. saida	13.5	14.5	0.013	12	750	B. saida	19.0	20.0	0.032
12	702	B. saida	17.0	18.5	0.028	12	752	B. saida	14.0	15.0	0.016
12	703	B. saida	15.5	16.5	0.016	12	753	B. saida	19.5	21.0	0.042
12	704	B. saida	16.0	17.0	0.018	12	754	B. saida	17.0	18.0	0.022
12	705	B. saida	18.0	20.0	0.030	12	755	B. saida	17.0	18.0	0.032
12	707	B. saida	17.5	18.5	0.020	12	756	B. saida	13.5	14.5	0.007
12	708	B. saida	17.0	18.0	0.023	12	757	B. saida	17.0	18.0	0.022
12	709	B. saida	14.0	15.0	0.009	12	758	B. saida	14.0	15.0	0.015
12	710	B. saida	18.0	19.5	0.031	12	759	B. saida	15.5	16.5	0.018
12	711	B. saida	19.0	20.5	0.030	12	761	B. saida	18.5	20.0	0.029

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
12	762	B. saida	15.5	16.5	0.019	14	462	B. saida	15.0	16.0	0.017
12	763	B. saida	15.5	16.5	0.016	14	463	B. saida	16.0	17.0	0.025
12	764	B. saida	18.5	20.0	0.030	14	464	B. saida	14.0	15.0	0.017
12	765	B. saida	16.0	17.0	0.016	14	465	B. saida	17.0	18.0	0.029
12	766	B. saida	16.0	17.0	0.018	14	466	B. saida	14.5	15.5	0.013
12	767	B. saida	15.0	16.0	0.017	14	467	B. saida	17.5	19.0	0.030
12	768	B. saida	17.5	19.0	0.028	14	468	B. saida	15.5	16.5	0.024
12	769	B. saida	19.0	20.5	0.036	14	469	B. saida	17.0	18.0	0.034
12	770	B. saida	18.0	19.5	0.036	14	470	B. saida	16.5	17.5	0.021
12	772	B. saida	16.0	17.0	0.018	14	471	B. saida	12.0	13.0	0.010
12	773	B. saida	14.0	15.0	0.017	14	472	B. saida	16.0	17.0	0.017
12	774	B. saida	17.0	18.0	0.024	14	473	B. saida	18.0	19.5	0.031
12	775	B. saida	16.0	17.0	0.020	14	474	B. saida	14.5	15.5	0.020
12	776	B. saida	13.5	14.5	0.011	14	475	B. saida	15.5	16.5	0.023
12	778	B. saida	16.0	17.0	0.018	14	477	B. saida	16.0	17.0	0.025
12	779	B. saida	17.0	18.0	0.020	14	478	B. saida	16.0	17.0	0.024
12	780	B. saida	14.5	15.5	0.013	14	479	B. saida	15.0	16.0	0.019
12	781	B. saida	16.0	17.0	0.019	14	480	B. saida	19.0	21.0	0.037
12	783	B. saida	16.0	17.0	0.025	14	481	B. saida	18.5	20.0	0.040
12	784	B. saida	16.0	17.0	0.021	14	483	B. saida	16.0	17.0	0.021
12	785	B. saida	14.0	15.0	0.015	14	484	B. saida	12.5	13.5	0.012
12	786	B. saida	17.0	18.0	0.026	14	485	B. saida	17.0	18.0	0.029
13	1259	B. saida	14.5	15.5	0.013	14	486	B. saida	15.0	16.0	0.024
13	1260	B. saida	14.5	15.5	0.018	14	487	B. saida	17.5	19.0	0.031
13	1261	B. saida	15.0	16.0	0.014	14	488	B. saida	14.0	15.0	0.015
13	1262	B. saida	17.5	18.5	0.022	14	489	B. saida	15.5	16.5	0.016
13	1263	B. saida	15.0	16.0	0.014	14	490	B. saida	16.5	17.5	0.024
13	1266	B. saida	16.0	17.0	0.020	14	491	B. saida	15.0	16.0	0.022
13	1268	B. saida	16.0	17.0	0.016	14	492	B. saida	14.0	15.0	0.015
13	1270	B. saida	16.0	17.0	0.016	14	493	B. saida	14.5	15.5	0.020
13	1281	B. saida	14.5	15.5	0.014	14	494	B. saida	17.0	18.5	0.028
14	449	B. saida	16.0	17.0	0.013	14	495	B. saida	12.0	13.0	0.008
14	450	B. saida	16.0	17.0	0.024	14	496	B. saida	17.0	19.5	0.031
14	451	B. saida	14.0	15.0	0.018	14	497	B. saida	13.5	14.5	0.016
14	452	B. saida	15.0	16.0	0.020	14	498	B. saida	16.0	17.0	0.022
14	453	B. saida	15.5	16.5	0.022	14	499	B. saida	13.0	14.0	0.012
14	454	B. saida	16.5	17.5	0.021	14	500	B. saida	18.0	19.0	0.038
14	455	B. saida	17.0	18.0	0.026	14	501	B. saida	14.0	15.0	0.014
14	456	B. saida	14.0	15.0	0.012	14	502	B. saida	16.0	17.0	0.028
14	457	B. saida	15.0	16.0	0.014	14	503	B. saida	12.0	13.0	0.008
14	458	B. saida	16.5	17.5	0.033	14	504	B. saida	19.0	20.5	0.039
14	459	B. saida	15.0	16.0	0.021	14	505	B. saida	11.0	12.0	0.007
14	460	B. saida	16.0	17.0	0.022	14	506	B. saida	16.0	17.0	0.022
14	461	B. saida	15.0	16.0	0.016	14	507	B. saida	15.0	16.0	0.021

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
12	762	B. saida	15.5	16.5	0.019	14	462	B. saida	15.0	16.0	0.017
12	763	B. saida	15.5	16.5	0.016	14	463	B. saida	16.0	17.0	0.025
12	764	B. saida	18.5	20.0	0.030	14	464	B. saida	14.0	15.0	0.017
12	765	B. saida	16.0	17.0	0.016	14	465	B. saida	17.0	18.0	0.029
12	766	B. saida	16.0	17.0	0.018	14	466	B. saida	14.5	15.5	0.013
12	767	B. saida	15.0	16.0	0.017	14	467	B. saida	17.5	19.0	0.030
12	768	B. saida	17.5	19.0	0.028	14	468	B. saida	15.5	16.5	0.024
12	769	B. saida	19.0	20.5	0.036	14	469	B. saida	17.0	18.0	0.034
12	770	B. saida	18.0	19.5	0.036	14	470	B. saida	16.5	17.5	0.021
12	772	B. saida	16.0	17.0	0.018	14	471	B. saida	12.0	13.0	0.010
12	773	B. saida	14.0	15.0	0.017	14	472	B. saida	16.0	17.0	0.017
12	774	B. saida	17.0	18.0	0.024	14	473	B. saida	18.0	19.5	0.031
12	775	B. saida	16.0	17.0	0.020	14	474	B. saida	14.5	15.5	0.020
12	776	B. saida	13.5	14.5	0.011	14	475	B. saida	15.5	16.5	0.023
12	778	B. saida	16.0	17.0	0.018	14	477	B. saida	16.0	17.0	0.025
12	779	B. saida	17.0	18.0	0.020	14	478	B. saida	16.0	17.0	0.024
12	780	B. saida	14.5	15.5	0.013	14	479	B. saida	15.0	16.0	0.019
12	781	B. saida	16.0	17.0	0.019	14	480	B. saida	19.0	21.0	0.037
12	783	B. saida	16.0	17.0	0.025	14	481	B. saida	18.5	20.0	0.040
12	784	B. saida	16.0	17.0	0.021	14	483	B. saida	16.0	17.0	0.021
12	785	B. saida	14.0	15.0	0.015	14	484	B. saida	12.5	13.5	0.012
12	786	B. saida	17.0	18.0	0.026	14	485	B. saida	17.0	18.0	0.029
13	1259	B. saida	14.5	15.5	0.013	14	486	B. saida	15.0	16.0	0.024
13	1260	B. saida	14.5	15.5	0.018	14	487	B. saida	17.5	19.0	0.031
13	1261	B. saida	15.0	16.0	0.014	14	488	B. saida	14.0	15.0	0.015
13	1262	B. saida	17.5	18.5	0.022	14	489	B. saida	15.5	16.5	0.016
13	1263	B. saida	15.0	16.0	0.014	14	490	B. saida	16.5	17.5	0.024
13	1266	B. saida	16.0	17.0	0.020	14	491	B. saida	15.0	16.0	0.022
13	1268	B. saida	16.0	17.0	0.016	14	492	B. saida	14.0	15.0	0.015
13	1270	B. saida	16.0	17.0	0.016	14	493	B. saida	14.5	15.5	0.020
13	1281	B. saida	14.5	15.5	0.014	14	494	B. saida	17.0	18.0	0.028
14	449	B. saida	16.0	17.0	0.013	14	495	B. saida	12.0	13.0	0.008
14	450	B. saida	16.0	17.0	0.024	14	496	B. saida	17.0	19.5	0.031
14	451	B. saida	14.0	15.0	0.018	14	497	B. saida	13.5	14.5	0.016
14	452	B. saida	15.0	16.0	0.020	14	498	B. saida	16.0	17.0	0.022
14	453	B. saida	15.5	16.5	0.022	14	499	B. saida	13.0	14.0	0.012
14	454	B. saida	16.5	17.5	0.021	14	500	B. saida	18.0	19.0	0.038
14	455	B. saida	17.0	18.0	0.026	14	501	B. saida	14.0	15.0	0.014
14	456	B. saida	14.0	15.0	0.012	14	502	B. saida	16.0	17.0	0.028
14	457	B. saida	15.0	16.0	0.014	14	503	B. saida	12.0	13.0	0.008
14	458	B. saida	16.5	17.5	0.033	14	504	B. saida	19.0	20.5	0.039
14	459	B. saida	15.0	16.0	0.021	14	505	B. saida	11.0	12.0	0.007
14	460	B. saida	16.0	17.0	0.022	14	506	B. saida	16.0	17.0	0.022
14	461	B. saida	15.0	16.0	0.016	14	507	B. saida	15.0	16.0	0.021

Table 11. Continued.

PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)	PLANKTON SAMPLE NUMBER	FISH SAMPLE NUMBER	SPECIES	STANDARD LENGTH (mm)	TOTAL LENGTH (mm)	WET WEIGHT (g)
16	45	<i>B. saida</i>	9.5	9.5	0.000	32	601	<i>B. saida</i>	30.0	32.5	0.170
16	46	<i>B. saida</i>	12.5	13.5	0.007	32	602	<i>B. saida</i>	34.0	37.0	0.277
16	1338	<i>T. pingelli</i>	20.5	23.0	0.062	32	603	<i>B. saida</i>	22.5	24.5	0.071
20	220	<i>B. saida</i>	13.0	13.0	0.006	32	604	<i>B. saida</i>	25.0	27.0	0.094
20	221	<i>B. saida</i>	17.0	18.5	0.031	32	1318	<i>G. tricuspis</i>	20.0	25.0	0.105
20	222	<i>B. saida</i>	17.0	18.5	0.026	34	430	<i>B. saida</i>	29.0	32.0	0.129
20	223	<i>B. saida</i>	18.0	19.0	0.021	34	431	<i>B. saida</i>	31.0	34.0	0.221
20	225	<i>B. saida</i>	15.0	16.0	0.012	34	433	<i>B. saida</i>	25.0	27.0	0.086
20	228	<i>B. saida</i>	17.0	18.0	0.021	34	438	<i>B. saida</i>	22.0	24.0	0.066
20	229	<i>B. saida</i>	16.5	17.5	0.017	34	439	<i>B. saida</i>	26.5	29.5	0.112
20	230	<i>B. saida</i>	13.0	13.0	0.008	34	440	<i>B. saida</i>	27.0	29.5	0.123
20	233	<i>B. saida</i>	13.0	13.0	0.004	34	441	<i>B. saida</i>	31.0	34.0	0.194
20	1086	<i>B. saida</i>	18.0	19.0	0.023	34	442	<i>B. saida</i>	22.5	25.0	0.080
20	1087	<i>B. saida</i>	13.5	14.5	0.012	34	447	<i>B. saida</i>	21.5	23.5	0.059
20	1088	<i>B. saida</i>	14.0	15.0	0.015	36	199	<i>B. saida</i>	28.0	31.0	0.155
20	1090	<i>B. saida</i>	13.5	14.5	0.012	36	205	<i>B. saida</i>	22.0	24.0	0.093
20	1091	<i>B. saida</i>	16.0	17.0	0.021	36	206	<i>B. saida</i>	29.0	32.0	0.172
20	1092	<i>B. saida</i>	15.5	16.5	0.021	36	208	<i>B. saida</i>	26.0	28.0	0.120
20	1093	<i>B. saida</i>	15.0	16.0	0.015	36	210	<i>B. saida</i>	20.0	22.0	0.062
21	2	<i>M. quadricornis</i>	15.0	17.5	0.019	36	211	<i>B. saida</i>	26.0	28.0	0.128
21	3	<i>M. quadricornis</i>	14.5	16.0	0.016	36	212	<i>B. saida</i>	26.0	28.0	0.121
21	4	<i>M. quadricornis</i>	15.5	17.5	0.021	36	214	<i>B. saida</i>	19.5	21.5	0.050
21	5	<i>M. quadricornis</i>	14.0	16.0	0.013	36	217	<i>B. saida</i>	20.0	22.0	0.061
21	6	<i>M. quadricornis</i>	14.0	16.0	0.011	40	1336	<i>G. tricuspis</i>	21.0	24.5	0.127
21	7	<i>M. quadricornis</i>	12.5	14.0	0.009						
21	8	<i>M. quadricornis</i>	14.0	16.0	0.014						
21	9	<i>M. quadricornis</i>	13.5	15.0	0.011						
21	10	<i>M. quadricornis</i>	13.0	14.0	0.005						
21	11	<i>M. quadricornis</i>	7.0	7.0	0.000						
29	219	<i>B. saida</i>	31.0	34.0	0.187						
31	235	<i>B. saida</i>	23.0	25.0	0.071						
32	580	<i>B. saida</i>	27.0	29.0	0.122						
32	582	<i>B. saida</i>	27.0	29.0	0.154						
32	583	<i>B. saida</i>	28.0	30.0	0.147						
32	584	<i>B. saida</i>	22.0	24.0	0.068						
32	585	<i>B. saida</i>	33.0	37.0	0.270						
32	586	<i>B. saida</i>	24.0	26.0	0.089						
32	590	<i>B. saida</i>	27.0	30.0	0.137						
32	593	<i>B. saida</i>	30.5	33.5	0.218						
32	594	<i>B. saida</i>	24.0	26.0	0.104						
32	595	<i>B. saida</i>	25.0	27.0	0.121						
32	598	<i>B. saida</i>	30.0	33.0	0.188						
32	600	<i>B. saida</i>	30.5	33.0	0.168						

Table 12. Mean length and weight of larval B. saida for July and September, 1984.

	JULY				SEPT			
	Mean	Range	Stnd Dev	N	Mean	Range	Stnd Dev	N
Stnd length	13.4	5.5-30.0	2.7	1061	26.3	19.5-34.0	3.8	36
Total length	14.3	6.5-32.5	3.0	1061	28.7	21.5-37.0	4.2	36
Wet weight	0.015	0.007-0.188	0.010	1061	0.130	0.050-0.277	0.058	36

Table 13. Mean length and weight of larval G. tricuspis for July and September, 1984.

	JULY				SEPT			
	Mean	Range	Stnd Dev	N	Mean	Range	Stnd Dev	N
Stnd length	13.6	11.5-15.5	1.6	10	20.5	20.0-21.0	0.5	2
Total length	15.5	12.5-17.5	2.0	10	24.8	24.5-25.0	0.25	2
Wet weight	0.020	0.006-0.033	0.009	10	0.116	0.105-0.127	0.011	2

Table 14. Mean length and weight of larval M. quadricornis for July, 1984.

	JULY			
	Mean	Range	Stnd Dev	N
Stnd length	13.3	7.0-15.5	2.26	10
Total length	14.9	7.0-17.5	2.87	10
Wet weight	0.119	<0.001-0.021	0.006	10

Table 15. Standard length and stomach fullness for larval fish captured in the Beaufort Sea, 1984.

Plankton Sample Number	Fish Sam No	Species	Stnd Length (mm)	Stomach Taken	Stomach Fullness
1	1681	<u>M. quadricornis</u>	11.0	W	1
2	1682	<u>M. quadricornis</u>	10.0	W	0
2	1683	<u>M. quadricornis</u>	11.0	W	3
4	1501	<u>B. saida</u>	10.5	W	3
4	1502	<u>B. saida</u>	11.5	W	5
4	1503	<u>B. saida</u>	14.5	W	5
4	1504	<u>B. saida</u>	14.0	W	5
4	1505	<u>B. saida</u>	11.5	W	3
4	1506	<u>B. saida</u>	11.0	W	3
4	1507	<u>B. saida</u>	14.0	W	5
4	1508	<u>B. saida</u>	11.5	W	1
4	1509	<u>B. saida</u>	15.5	S	6
4	1669	<u>G. tricuspidis</u>	11.5	W	5
4	1670	<u>G. tricuspidis</u>	11.5	W	3
4	1671	<u>G. tricuspidis</u>	11.0	W	3
4	1672	<u>G. tricuspidis</u>	13.5	W	3
4	1673	<u>G. tricuspidis</u>	14.0	W	3
4	1674	<u>G. tricuspidis</u>	14.5	W	5
4	1684	<u>G. tricuspidis</u>	13.0	W	3
4	1689	<u>Liparis sp.</u>	20.5	S	6
4	1690	<u>Liparis sp.</u>	13.0	W	5
4	1691	<u>Liparis sp.</u>	9.0	W	3
5	1675	<u>G. tricuspidis</u>	11.0	W	3
6	1528	<u>B. saida</u>	8.5	W	1
6	1529	<u>B. saida</u>	7.5	W	3
6	1530	<u>B. saida</u>	9.0	W	3
6	1531	<u>B. saida</u>	8.5	W	3
6	1532	<u>B. saida</u>	8.5	W	3
8	1533	<u>B. saida</u>	11.5	W	5
8	1534	<u>B. saida</u>	12.5	W	5
8	1535	<u>B. saida</u>	13.0	W	5
8	1536	<u>B. saida</u>	14.0	W	5
8	1537	<u>B. saida</u>	14.5	S	6
8	1538	<u>B. saida</u>	14.5	S	4
8	1539	<u>B. saida</u>	15.0	S	6
8	1540	<u>B. saida</u>	14.5	S	4
8	1541	<u>B. saida</u>	15.0	S	4
8	1542	<u>B. saida</u>	16.0	S	4
9	1543	<u>B. saida</u>	*	W	5
9	1544	<u>B. saida</u>	10.5	W	3
9	1545	<u>B. saida</u>	9.0	W	0
9	1546	<u>B. saida</u>	10.0	W	1
9	1547	<u>B. saida</u>	9.5	W	5

Table 15. Continued.

Plankton Sample Number	Fish Sam No	Species	Stnd Length (mm)	Stomach Taken	Stomach Fullness
9	1548	B. <u>saida</u>	9.5	W	3
9	1549	B. <u>saida</u>	10.5	W	3
9	1550	B. <u>saida</u>	11.5	W	3
9	1551	B. <u>saida</u>	10.5	W	5
9	1552	B. <u>saida</u>	12.5	W	3
12	1553	B. <u>saida</u>	15.0	W	3
12	1554	B. <u>saida</u>	17.0	W	0
12	1555	B. <u>saida</u>	16.0	W	3
12	1556	B. <u>saida</u>	16.0	W	3
12	1557	B. <u>saida</u>	16.5	W	1
12	1558	B. <u>saida</u>	18.5	S	4
12	1559	B. <u>saida</u>	17.0	S	4
12	1560	B. <u>saida</u>	17.5	S	0
12	1561	B. <u>saida</u>	17.0	S	2
12	1562	B. <u>saida</u>	18.5	S	4
12	1676	G. <u>tricuspidis</u>	12.0	W	1
12	1677	G. <u>tricuspidis</u>	11.0	W	1
13	1563	B. <u>saida</u>	15.5	W	4
13	1565	B. <u>saida</u>	15.5	W	1
13	1567	B. <u>saida</u>	9.0	W	1
13	1569	B. <u>saida</u>	12.5	W	1
13	1570	B. <u>saida</u>	12.5	W	3
13	1572	B. <u>saida</u>	14.0	W	1
13	1574	B. <u>saida</u>	15.5	W	3
13	1577	B. <u>saida</u>	14.5	W	3
13	1581	B. <u>saida</u>	17.0	W	5
13	1583	B. <u>saida</u>	15.5	W	1
14	1692	Liparis sp.	10.5	W	3
15	1589	B. <u>saida</u>	10.0	W	5
15	1590	B. <u>saida</u>	11.0	W	5
15	1591	B. <u>saida</u>	11.0	W	5
15	1592	B. <u>saida</u>	11.5	W	5
15	1593	B. <u>saida</u>	11.5	W	5
15	1594	B. <u>saida</u>	13.0	W	5
15	1595	B. <u>saida</u>	13.5	W	3
15	1596	B. <u>saida</u>	15.0	W	5
16	1678	G. <u>tricuspidis</u>	14.0	W	1
16	1688	M. <u>quadricornis</u>	11.5	W	1
19	1597	B. <u>saida</u>	12.5	W	3
19	1598	B. <u>saida</u>	13.0	W	3
19	1599	B. <u>saida</u>	14.0	W	3
19	1600	B. <u>saida</u>	14.5	W	5
19	1601	B. <u>saida</u>	14.5	W	3

Table 15. Continued.

Plankton Sample Number	Fish Sam No	Species	Stnd Length (mm)	Stomach Taken	Stomach Fullness
19	1602	B. <u>saida</u>	16.0	W	6
19	1603	B. <u>saida</u>	16.5	S	2
19	1604	B. <u>saida</u>	17.0	S	3
20	1679	G. <u>tricuspidis</u>	14.0	W	3
20	1686	M. <u>quadricornis</u>	11.5	W	1
20	1687	M. <u>quadricornis</u>	12.0	W	1
21	1685	M. <u>quadricornis</u>	15.5	W	1
32	1605	B. <u>saida</u>	11.0	W	3
32	1608	B. <u>saida</u>	24.5	S	6
32	1610	B. <u>saida</u>	25.5	S	4
32	1613	B. <u>saida</u>	25.5	S	6
32	1615	B. <u>saida</u>	23.5	S	6
32	1617	B. <u>saida</u>	26.0	S	5
32	1619	B. <u>saida</u>	28.5	S	3
32	1621	B. <u>saida</u>	*	S	4
32	1623	B. <u>saida</u>	30.0	S	6
32	1627	B. <u>saida</u>	34.0	S	2
34	1628	B. <u>saida</u>	18.5	S	1
34	1630	B. <u>saida</u>	21.0	S	4
34	1632	B. <u>saida</u>	25.0	S	5
34	1634	B. <u>saida</u>	20.5	S	4
34	1636	B. <u>saida</u>	24.5	S	4
34	1638	B. <u>saida</u>	25.5	S	6
34	1639	B. <u>saida</u>	26.0	S	2
34	1641	B. <u>saida</u>	38.5	S	6
34	1645	B. <u>saida</u>	28.5	S	2
34	1647	B. <u>saida</u>	32.0	S	4
36	1648	B. <u>saida</u>	21.0	S	5
36	1650	B. <u>saida</u>	21.0	S	4
36	1652	B. <u>saida</u>	22.0	S	6
36	1654	B. <u>saida</u>	21.0	S	6
36	1656	B. <u>saida</u>	26.5	S	4
36	1658	B. <u>saida</u>	25.0	S	4
36	1660	B. <u>saida</u>	26.0	S	4
36	1662	B. <u>saida</u>	27.5	S	4
36	1664	B. <u>saida</u>	28.5	S	4
36	1666	B. <u>saida</u>	30.0	S	4
36	1680	G. <u>tricuspidis</u>	*	S	3
38	1667	B. <u>saida</u>	20.0	S	1
40	1668	B. <u>saida</u>	24.0	S	2

* - No measurement taken

W - Intestine (stomach not yet developed)

S - Stomach

Table 16. Food items totals for *B. solda* captured in the Canadian Beaufort Sea Shelf, 1984.

Plankton Fish		Som. Sem.												Eggs (Unit d.)												
No.	No.	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL		
4	1501																Ce 1					Le 2	Ju 10			
4	1502																Ps 3					Le 1				
4	1503																C3 1					Le 1				
4	1504																Cf 2									
																	Ce 28									
																	Ce 80									
																	C3 1									
																	Cf 6									
																	C3 1									
4	1505																Cf 1	Cf 1								
4	1506																Cf 2									
4	1507																C7 1									
																	C8 7									
																	Ce 17									
																	Ce 10									
4	1508																Ce 10		C3 1							
4	1509																C8 2		C5 2							
																	C4 1		Cf 3							
																	Ce 35									
6	1528																					N2 2				
6	1529																					N2 3				
6	1530																					N2 2				
6	1551																					N2 4				
6	1532																					Ce 1				
8	1533																C7 1									
8	1534																					C2 1				
8	1535																					Cf 2				
8	1536																					Ce 50				
8	1537																					Ce 32	N3 1	N2 1		
8	1538																					Cf 2		N2 18		
8	1539																					Ce 23		N1 1		
8	1540																					Cf 3		N2 33		
																						C2 2		N1 4		
																						C1 2		N1 14		
																						Ce 25		N2 35		
																						Cf 3		N1 3		
																						Ce 3		N2 90		
																						C1 4		N3 4		
																						Ce 4		N3 1		
																						Cf 1				

Table 16. Continued.

Plankton Fish																			
Som. Som. No. No.	CC TL																		
<i>Acanthocephala</i>																			
<i>Ascidia</i> sp.																			
<i>Cocconotus</i> sp.																			
<i>Pardinium</i> sp.																			
<i>Ratifer</i> (Unid.)																			
<i>Ceratium</i> sp.																			
<i>Polychaete</i> (Unid.)																			
<i>Arcteria</i> clausi																			
<i>Calanus</i> hyperboreus																			
<i>Calanus</i> glacialis																			
<i>Eurytemora</i> sp.																			
<i>Doridogymna</i> rotifera																			
<i>Cetanus</i> sp.																			
<i>Oithona</i> stali																			
<i>Microsetella</i> rosea																			
<i>Limnocalanus macrurus</i>																			
<i>Oncasae</i> borealis																			
<i>Cyclopoidae</i>																			
<i>Calanoidae</i>																			
<i>Apherus</i> glacialis																			
<i>Ciliata</i> <i>lilacina</i>																			
<i>Limacina</i> <i>helicina</i>																			
<i>Bivalvia</i> (Unid.)																			
<i>Oikopleura</i> <i>ventosa</i>																			
Eggs (Unid.)																			
<i>Fritillaria</i> <i>borealis</i>																			

Table 16. Continued.

		Eggs (Unid.)												Fertilization												
		Bivalvia (Unid.)						Okoplaera ventrofemoralis						Fritillaria borealis						Leucina helicina						
		Clione limacina						Aphorura glacialis						Cyclopoida						Calanoida						
Plankton Fish		CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL	CC TL		
Sem. Sem.	No. No.																									
13	1567																									
13	1569																									
13	1570																									
15	1572																									
13	1574																									
13	1577	Ps 1																							Ce 2	
13	1581																									
13	1583																									
15	1589																									
15	1590																									
15	1591																									
15	1592																									
15	1593																									
15	1594																									
15	1595	Ps 1																								
15	1596																									
19	1597																									
19	1598																									
19	1599															Cs 1										
19	1600																Cf 1									
19	1601						C1 1										Cs 1									
19	1602																	Cf 5								
19	1603																	Ce 157								
19	1604																	Cf 1								
32	1605																		Ce 34						La 2	

Table 16. Continued.

Table 16. Continued.

Table 16. Continues.

CC = comment code

TL • total

Table 17. Food item totals for M. quadricornis captured in the Canadian Beaufort Sea Shelf, 1984.

Planton Sam. No.	Fish Sam. No.	<u>Limnocalanus</u> <u>macrurus</u>		<u>Calanoida</u>		<u>Cyclopoida</u>	
		CC	TL	CC	TL	CC	TL
1	1681					N2	3
2	1683			N1	6		
				N3	19		
16	1688	Ps	1				
20	1686			Ps	2		
20	1687			Ps	1		
21	1685	Ps	1				

CC = comment code

TL = total

Table 18. Food item totals for *G. tricuspidis* captured in the Canadian Beaufort Sea Shelf, 1984.

CC = comment code

Tl = total

Table 19. Food item totals for Liparis sp. captured in the Canadian Beaufort Sea Shelf, 1984.

Plankton Sam. No.	Fish Sam. No.	Polychaete (Unid.)		Oncaea borealis		Pseudocalanus minutus		Apherusa glacialis		Gammarus sp.	
		CC	TL	CC	TL	CC	TL	CC	TL	CC	TL
4	1689					C2	1			8	Ju 1
4	1690					Cf	14				
						Ce	125				
4	1691	Ju	1	Cm	1	C5	1				
14	1692	Ju	10								

CC = comment code

TL = total

Table 20. List of comment codes used in Tables 16 to 19.

Code	Comment
Ps	poor condition
P1	Polychaete larvae
La	larva - no specific taxon
Na	nauplii - no specific taxon
Ju	juvenile - no specific taxon
C1	copepodite I
C2	copepodite II
C3	copepodite III
C4	copepodite IV
C5	copepodite V
Cf	adult female (copepodite VI)
Cm	adult male (copepodite VI)
N1	nauplius < 200um
N2	nauplius 200 - 400um
N3	nauplius > 400um
C7	copepodite < 400um (Cyclopoids and Harpacticoids only)
C8	copepodite 400 - 800um (Cyclopoids and Harpacticoids only)
Ce	copepod eggs

Table 21. Frequency of occurrence and percent by number of food items found in B. saida stomachs.

		<u>July</u>			<u>Sept</u>		
	Food Item	Freq Occur	% Num		Freq Occur	% Num	
Diatoms:	<u>Coscinodiscus</u> sp.	1.5	0.1		12.5	0.2	
	<u>Peridinium</u> sp.				9.4	0.1	
	<u>Ceratium</u> sp.	1.5	0.1				
Rotifers:	Unid.	3.0	0.1		34.4	0.6	
Polychaetes:	Unid.				9.4	0.2	
Copepoda:	<u>Acartia clausi</u>				59.4	2.2	
	<u>Calanus glacialis</u>	3.0	0.1		46.9	1.0	
	<u>Calanus hyperboreus</u>				3.1	<0.1	
	<u>Calanus</u> sp.				3.1	<0.1	
	<u>Eurytemora</u> sp.				3.1	<0.1	
	<u>Derjuginia tolli</u>	1.5	0.2				
	<u>Limnocalanus macrurus</u>	17.9	1.1		3.1	<0.1	
	<u>Microcalanus pygmaeus</u>				3.1	<0.1	
	<u>Microstella rosea</u>				9.4	0.1	
	<u>Oithona similis</u>	17.9	2.8		34.4	0.7	
	<u>Oncaea borealis</u>	4.5	0.2		43.8	0.5	
	<u>Pseudocalanus minutus</u>	26.9	31.9		56.3	26.2	
	<u>Calanoida</u> (Unid.)	83.6	61.6		100.0	63.8	
	<u>Cyclopoida</u> (Unid.)	6.0	0.3		21.9	0.3	
Amphipoda:	<u>Apherusa glacialis</u>				3.1	0.1	
Gastropoda:	<u>Clione limacina</u>				3.1	<0.1	
	<u>Limacina helicina</u>				3.1	<0.1	
Bivalvia:	Unid.	9.0	0.5		15.6	0.2	
Larvacea:	<u>Oikopleura vanhoeffeni</u>	6.0	1.0		40.6	1.6	
	<u>Fritillaria borealis</u>				18.8	0.9	
Eggs:	Unid.	3.0	0.2		50.0	1.3	

Table 22. Frequency of occurrence and percent by number of food items found in M. quadricornis stomachs, July, 1984.

	Food Item	Freq Occur	% Num
Copepoda:	<u>Limnocalanus macrurus</u>	33.3	6.1
	Calanoida (Unid.)	50.0	84.8
	Cyclopoida (Unid.)	16.7	9.1

Table 23. Frequency of occurrence and percent by number of food items found in G. tricuspis stomachs, July and September, 1984.

	Food Item	Freq Occur	% Num
Diatoms:	<u>Coscinodiscus</u> sp.	61.5	49.0
	<u>Peridinium</u> sp.	7.7	0.3
	Unid.	7.7	1.7
Polychaetes:	Unid.	15.4	0.6
Copepoda:	<u>Cyclopina</u> sp.	15.4	10.1
	<u>Limnocalanus macrurus</u>	15.4	0.8
	<u>Microsetella rosea</u>	7.7	0.3
	<u>Oithona similis</u>	7.7	0.3
	<u>Oncaeа borealis</u>	7.7	0.3
	<u>Pseudocalanus minutus</u>	84.6	20.8
	Calanoida	15.4	0.8
	Harpacticoida	7.7	0.3
Cirripedia:	Unid.	38.5	1.4
Amphipoda:	<u>Apherusa glacialis</u>	7.7	0.6
Bivalvia:	Unid.	23.1	3.4
Stelleroide:	Unid.	7.7	1.1
Larvacea:	<u>Oikopleura vanhoeffeni</u>	7.7	2.8
Eggs:	Unid.	30.8	5.6

Table 24. Frequency of occurrence and percent by number of food items found in Liparis sp. stomachs, July, 1984.

	Food Item	Freq Occur	% Num
Polychaetes:	Unid.	50.0	6.8
Copepoda:	<u>Oncaeae borealis</u>	25.0	0.6
	<u>Pseudocalanus minutus</u>	75.0	87.0
Amphipoda:	<u>Gammarus</u> sp.	25.0	0.6
	<u>Apherusa glacialis</u>	25.0	4.9

Table 25. Parasites found in larval fish stomachs.

Fish Sam No	Species	Station	Parasite	Count
1507	<u>B. saida</u>	84A02	Trematoda: Hemiuridae	1
1542	<u>B. saida</u>	84A04	Trematoda: Hemiuridae	2
1608	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	2
1619	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	1
1621	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	2
1623	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	1
1627	<u>B. saida</u>	84B02	Trematoda: Hemiuridae	1
1647	<u>B. saida</u>	84B03	Trematoda: Hemiuridae	1
1660	<u>B. saida</u>	84B04	Trematoda: Hemiuridae	1

