



Scientific Excellence • Resource Protection & Conservation • Benefits for Canadians  
Excellence scientifique • Protection et conservation des ressources • Bénéfices aux Canadiens

# **The Domestic Beluga (*Delphinapterus leucas*) Fishery in the Mackenzie River Estuary, Northwest Territories, 1981-1986**

J.T. Strong

Central and Arctic Region  
Department of Fisheries and Oceans  
Winnipeg, Manitoba R3T 2N6

October 1990

## **Canadian Data Report of Fisheries and Aquatic Sciences No. 800**



Fisheries  
and Oceans

Pêches  
et Océans

Canada

## **Canadian Data Report of Fisheries and Aquatic Sciences**

Data reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of data reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was introduced with the publication of report number 161.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

## **Rapport statistique canadien des sciences halieutiques et aquatiques**


Les rapports statistiques servent à classer et à archiver les compilations de données pour lesquelles il y a peu ou point d'analyse. Ces compilations auront d'ordinaire été préparées à l'appui d'autres publications ou rapports. Les sujets des rapports statistiques reflètent la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports statistiques ne sont pas destinés à une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans autorisation écrite préalable de l'établissement auteur. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 25 de cette série ont été publiés à titre de relevés statistiques, Services des pêches et de la mer. Les numéros 26 à 160 ont été publiés à titre de rapports statistiques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 161.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Data Report of  
Fisheries and Aquatic Sciences 800



October 1990

THE DOMESTIC BELUGA (Delphinapterus leucas) FISHERY  
IN THE MACKENZIE RIVER ESTUARY, NORTHWEST TERRITORIES,  
1981-1986

by

J.T. Strong

Central and Arctic Region  
Department of Fisheries and Oceans  
Winnipeg, Manitoba R3T 2N6

This is the 47th Data Report  
from the Central and Arctic Region, Winnipeg

© Minister of Supply and Services Canada 1990

Cat. No. Fs 97-13-800E

ISSN 0706-6465

Correct citation for this publication is:

Strong, J.T. 1990. The domestic beluga (Delphinapterus leucas) fishery in the Mackenzie River estuary, Northwest Territories, 1981-1986. Can. Data Rep. Fish. Aquat. Sci. 800: iv + 52 p.

TABLE OF CONTENTS

	<u>Page</u>
ABSTRACT/RÉSUMÉ . . . . .	iv
INTRODUCTION . . . . .	1
MATERIALS AND METHODS . . . . .	1
RESULTS . . . . .	1
Current hunting techniques . . . . .	1
Harvest . . . . .	1
Biology . . . . .	2
ACKNOWLEDGMENTS . . . . .	3
REFERENCES . . . . .	3

LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1 Map of Mackenzie River Estuary, Northwest Territories . . . . .	4
2 Temporal distribution of beluga whale harvest in the Mackenzie River Estuary, 1981-1986 . . . . .	5
3 Comparison of mean length, range and standard deviation for male and female beluga whale harvested in the Mackenzie River Estuary 1981-1986 . . . . .	6
4 Length frequencies of beluga whale harvested in the Mackenzie River Estuary 1981-1986 . . . . .	7

LIST OF TABLES

<u>Table</u>	<u>Page</u>
1 Cartridge use and shots fired during beluga hunts in the Mackenzie River estuary . . . . .	8
2 Numbers of beluga whales reported as struck/landed in the Mackenzie River Estuary area, by location 1981-1986 . . . . .	9
3 Reported loss rates from the Mackenzie River estuary beluga fishery 1981-1986 . . . . .	10
4 Annual breakdown, by sex, of Mackenzie River Estuary beluga whale harvest 1981-1986 . . . . .	11
5 Range, mean and standard deviation of various measurements from beluga taken in the Mackenzie River Estuary 1981-1986 . . . . .	12

Table

	<u>Page</u>
6 Total lengths of female beluga known to be currently reproductive, Mackenzie River Estuary, 1981-1986 . . . . .	14
7 Total lengths (cm) and date of sampling for fetuses and neonates collected in the Mackenzie River Estuary Beluga Fishery, 1981-1986 . . . . .	15
8 Mean, range and standard deviations from total lengths of fetuses and neonates collected in the Mackenzie River Estuary Beluga Fishery, 1981-1986 . . . . .	16
9 Range and mean of the number of teeth recorded from beluga harvested in the Canadian Beaufort Sea . . . . .	17

LIST OF APPENDICES

<u>Appendix</u>	<u>Page</u>
1 Monitor data sheet and DFO data sheet . . . . .	18
2 Temporal distribution of the beluga whale harvest in the Mackenzie River Estuary, annually, by area and cumulative, 1981-1986 . . . . .	21
3 Range, mean, and standard deviation of total lengths of beluga from the Mackenzie Estuary, annually, 1981-1986 . . . . .	34
4 Length frequencies of beluga harvested in the Mackenzie River Estuary, annually, by area and cumulative, 1981-1986 . . . . .	39
5 Number of teeth recorded from beluga harvested in the Canadian Beaufort Sea . . . . .	52

## ABSTRACT

Strong, J.T. 1990. The domestic beluga (Delphinapterus leucas) fishery in the Mackenzie River estuary, Northwest Territories, 1981-1986. Can. Data Rep. Fish. Aquat. Sci. 800: iv + 52 p.

Biological data and harvest information collected from the 1981-1986 domestic beluga in the Mackenzie River estuary are presented. The landed beluga reported by the contract monitors for each of six years were: 137, 107, 85, 134, 122 and 149. The average annual loss rate was 17 percent, with a range of 10-26 percent. Average annual mean lengths were: males 426 cm and females 369 cm. Sex ratios for each of the six years were 109, 267, 453, 694, 281 and 112 males per 100 females. The harvest is biased towards larger animals, and the greatest harvest occurs in the Kugmallit Bay area.

Key words: Western Arctic; white whales; harvest; morphometry; population dynamics.

## RÉSUMÉ

Strong, J.T. 1990. The domestic beluga (Delphinapterus leucas) fishery in the Mackenzie River estuary, Northwest Territories, 1981-1986. Can. Data Rep. Fish. Aquat. Sci. 800: iv + 52 p.

On présente les données biologiques et l'information sur les prises relatives à la pêche intérieure au béluga dans l'estuaire du fleuve Mackenzie pour la période de 1981 à 1986. Le nombre de bélugas débarqués signalé par les surveillants à contrat pour chacune des six années a été le suivant : 137, 107, 85, 134, 122 et 149. Le taux des pertes annuelles moyennes a été de 17 p. cent, l'étendue étant de 10-26 p. cent. La moyenne des longueurs moyennes annuelles ont été de 426 cm pour les mâles et de 369 cm pour les femelles. Les sex-ratios pour les six années sont respectivement de 109, 267, 453, 694, 281 et 112 mâles pour 100 femelles. La récolte est biaisées en faveur des animaux de plus grande taille et la récolte la plus importante a eu lieu dans la région de la baie Kugmallit.

Mots-clés: Arctique occidental; béluga; récolte; morphométrie; dynamique des populations.

## INTRODUCTION

The belugas (*Delphinapterus leucas*) summering in the Canadian Beaufort Sea (Fig. 1) are considered to be part of a population which is distributed through the Bering, Beaufort, Chukchi and Eastern Siberian seas, and Amundsen Gulf (Burns and Seaman 1986). The first recorded European contact with beluga in the Mackenzie Delta was July 3, 1789 when Alexander Mackenzie saw white whales on the west side of "Whale Island" (Daniells 1971), probably in what is locally called Niakunak Bay. The association between Delta residents and beluga predates European contact with North America; archaeological investigations (McGhee 1974) have revealed a well-developed beluga hunting culture in existence as early as the 14th century.

Fraker (1976) identified three "concentration areas" all located near major outflows of the Mackenzie River in Niakunak Bay, East Mackenzie Bay and Kugmallit Bay, where beluga gather annually from late June to August. The Mackenzie River estuary domestic beluga fishery, carried out by local Inuvialuit, is centered on these concentration areas and major hunting camps are located adjacent to them.

There are apparently no records of the western Arctic domestic beluga harvest prior to 1954, although recollected estimates such as those of Nuligak (1966) indicate that historical harvests may have been, at least occasionally, as high as 300 animals. Available catch statistics are reviewed by Smith and Taylor (1977) and Strong (1989).

This study was initiated to gather data on the harvest rate and to collect morphological data from landed specimens to be used in comparisons with other stocks at some later time.

## MATERIALS AND METHODS

### HARVEST DATA COLLECTION

Harvest information was collected annually by both observation and hunter interview. Two hunters from each community (Aklavik, Inuvik and Tuktoyaktuk) were hired on contract to monitor the beluga harvest of the camps in which they were seasonal residents. The Aklavik monitors recorded information from camps on West Mackenzie and Shallow Bays; the Inuvik monitors recorded information from Kendall Island, East Whitefish Station and camps in their vicinity; the Tuktoyaktuk monitors recorded information from hunts originating in the settlement and from Hendrickson Island (Fig. 1). Our data collection focussed on Kugmallit Bay, where most of the annual harvest occurs.

Data was recorded on either of two forms; one for use by contractors, the other by DFO staff (Appendix 1). Landed whales were assigned a sequential catalogue number, which identified location, species and year of harvest.

Various standard measurements (Norris 1961) were collected from each specimen. Measurements were most often recorded in imperial units then converted to metric and rounded. Total length was recorded as the straight line distance between nose and tail notch, parallel to the body surface. Half-girth was measured along the axillary and umbilical circumferences from dorsal ridge to mid-belly, and this value was doubled to obtain a full girth value. Other measurements including the number of teeth present were taken opportunistically (see Appendix 1 - DFO data sheet), but the completeness of the record depended on the progress of butchering, the time of observer arrival, and hunter cooperation.

Estimates of hunting losses from 1981-1986 are based on information provided by the contract observers. To derive our estimate, we assumed that all whales struck and lost died of their wounds (since we could not observe all hunting activity), and used the formula:

$$\frac{\# \text{ Struck} - \# \text{ Landed}}{\text{Total} \# \text{ Struck}}$$

## RESULTS

### CURRENT HUNTING TECHNIQUES

Current hunting techniques differ little from those described by Slaney (1974), Fraker (1976, 1977) and Hunt (1979), although speedboats have largely replaced freighter canoes and some hunters favour fluorescent plastic buoys over 10 gallon gas drum floats. The 30/30 is still the most common rifle caliber used for beluga hunting (Table 1), as noted by Fraker and Fraker (1979).

Residents of Tuktoyaktuk generally hunt from the settlement, and take their catch home for processing. The catch may be towed whole or the muktuk (skin) and meat removed and transported. It is common for hunters from Tuktoyaktuk to hunt on weekends or when "off-shift" from their employment. Residents of Aklavik and Inuvik travel greater distances to the hunting area, generally hunt from and process their catch in a seasonal hunting camp and must transport their catch a considerable distance back to their homes. A few individuals from these settlements travel to the coast for short-term or weekend hunts and many of them utilize the established family camps. In 1981, four families from Holman participated in the whale hunt at Bird Camp in West Mackenzie Bay. Cooperation between hunting parties is still a feature of the hunt, but it is common for parties to hunt singly.

### HARVEST

The landed catch of belugas from the four main hunting areas during the period 1981-1986 is presented in Table 2. The majority of beluga are harvested in July although successful hunts

as early as June 24 and as late as August 23 were reported during this study. Figure 2 shows the temporal distribution of the pooled reported harvest 1981-1986, annual data by hunting area and combined is provided in Appendix 2.

Discussions with long term beluga hunters suggest the number of individuals involved in the harvest has declined over time and that fewer families depend on the beluga for a winter food supply, even though the number of beluga harvested has remained fairly constant during recent times.

Harvest values reflect actual landings by residents in seasonal camps, and by most transient hunters. The harvest for Tuktoyaktuk may be under-reported due to the difficulty of keeping in contact with all active hunters. Six years of data (Table 2) show the number of beluga taken in Kugmallit Bay has consistently exceeded the number taken from Niakunak Bay. The greatest number of beluga landed, approximately 45% of total, are taken by hunters from Tuktoyaktuk likely because the settlement is adjacent to a major hunting area.

Total annual loss rates between 1981 and 1986 ranged from 10 to 26% (Table 3), and average annual loss for the 5 year period is 17%. We believe the whale monitors accurately report the information they receive, but we are also aware that some hunters may not report landed or lost animals.

Most cartridges used today contain soft point bullets. Experienced hunters are of the opinion that fewer whales were lost when hard point ammunition was common and attribute a portion of the animals struck but lost to the inefficiency of soft-point ammunition compared to hard-point ammunition.

The sex composition of the harvest varied over the period of this study from 48% female in 1981 to 13% female in 1984 and back to 47% female in 1986 (Table 4). Hunters from Tuktoyaktuk took over 50% of all female beluga landed.

It is possible that some hunters report females as males and thus bias the harvest data. The male/female ratio reported by the contract whale observers from direct observation is correct, based on biological samples collected for other programs. Misrepresentation may have occurred with whales reported through hunter interviews and not seen by the observer or DFO staff. The sex ratio of the observed harvest is similar to that of the reported harvest, suggesting that most hunters report the sex of their catch accurately.

## BIOLOGY

Standard total lengths (Norris 1961) were collected from 400 male and 186 female beluga between 1981 and 1986. The total lengths reported in this study were collected by various individuals. Although minor inconsistencies

among individuals are likely, we are confident that the lengths reported are suitable for distinguishing between length classes of beluga. This study includes all animals larger than 200 cm in calculations of mean total length. Table 5 provides the range, mean and standard deviation of those total lengths; annual data by area and combined is provided in Appendix 3. Figure 3 graphically compares the mean length, range and standard deviation of the total lengths of female and male belugas collected during the study. Figure 4 illustrates the annual combined total length data for 1981 to 1986, annual data is in Appendix 4. Table 5 provides the range, mean and standard deviation of axillary girths, fluke widths and flipper length and widths collected between 1983 and 1986.

During the period 1981-1986, 82 of 125 female beluga examined were in reproductive condition; of these, 43 were with a neonate and lactating or carrying a term foetus and lactating, five were carrying a non-term foetus but not lactating. No specific information was given for the remaining 26 reproductive females. Total lengths from reproductive female beluga all of which were longer than 320 cm, are provided in Table 6.

Lengths were collected from three neonates, six term foetuses and 11 non-term foetuses during this study. All lengths were measured as standard total lengths, and with the exception of a 22.5 cm non-term foetus collected August 2, were collected in July. Table 7 provides the total length and date of collection for each sample.

The mean length of non-term foetuses was 19.23 cm, of term foetuses 159.20 cm, and of neonates 198.91 cm. Ranges and standard deviation are provided in Table 8. The mean length of the combined sample of neonates and term foetuses is 172.44 cm.

Three non-term foetuses collected on July 13 were 12.7, 17.8 and 38.1 cm, and there is great variation in length over the sampling period for both categories of foetuses and for neonates. The data support an extended calving period for beluga lasting well into August.

The dental formula of 23 male and four female adult beluga was examined in the field (Table 9, Appendix 5). The formulae are similar, but not identical to those of other researchers. Tomilin (1957) gives 9-9/8-8 or 10-10/9-9 as the most common dental formula for adult beluga in the Soviet Union and stated that only one of 69 animals examined had 11 pair of teeth in the upper jaw. The data show a variable number of teeth in each jaw, and in each quadrant. This has been attributed to tooth loss or incomplete eruption (Tomilin 1957; Kleinenberg et al. 1964). In total we found 14 different formulae among 27 whales with the number of teeth ranging from 28-40. Dental formulae 11-11/9-9, 10-10/10-10, and 10-10/9-9 occurred in three animals each; 8-8/8-8 and 7-7/7-7 were each found in two animals. All other combinations were found in one animal only.



## ACKNOWLEDGMENTS

We would like to thank the contractors who collected much of the data for this report. They are: Elijah Allen, Wayne Allen, Joseph Avik, Alex Aviugana, Jonah Carpenter, Willie Carpenter, Henry Chicksi, Billy Day, William Day, Alex Elanik, Frank Elanik, Walter Elias, Colin Harry, Walter Malegana, Elsie Nuttal, John Nuttal, Jessie Panaktalok, Sam Pingo, the late Big Jim Rogers, and Larry Sittichinli. We also thank the Hunters and Trappers Associations of Aklavik, Inuvik and Tuktoyaktuk who have cooperated with and contributed to the program since its inception.

Steve Tinker and Dave Barber coordinated the field program in 1982 and 1983 respectively. Tommy Chicksi, our guide for three years and Forrest Day for one year assisted in sample collection and made many valuable contributions. Technical assistance was provided by Larry Dueck, Barb Glassey, Elaine Murkin, Wouter Pleuniss, Kim Wazura, and in particular Patt Weaver who made a major contribution.

Richard Barnes, William Day, Bill Ferguson, Leonard Harry and Wally Kowal of DFO, Inuvik and Ed Henderson of The Government of the Northwest Territories, Department of Renewable Resources, Tuktoyaktuk provided valuable advice and logistic support.

## REFERENCES

- DANIELLS, R. 1971. Alexander Mackenzie and the Northwest. Oxford University Press, Toronto. 219 p.
- FRAKER, M.A. 1976. The 1975 white whale study. In Summer environmental program, Mackenzie River estuary. Vol. 2 White whale studies. Unpublished report for Imperial Oil Ltd. by F.F. Slaney and Co. Ltd. 62 p. + Appendix and maps.
- FRAKER, M.A. 1977. The 1977 whale monitoring program, Mackenzie Estuary, N.W.T. Unpublished report for Imperial Oil Ltd. by F.F. Slaney and Co. Ltd., Vancouver. 47 p. + Appendix.
- FRAKER, M.A., and P.N. FRAKER. 1979. The 1979 white whale monitoring program, Mackenzie estuary. Unpublished report for Esso Resources Canada by L.G.L. Ltd. 51 p.
- HUNT, W.J. 1979. Domestic whaling in the Mackenzie Estuary, Northwest Territories. Can. Fish. Mar. Serv. Tech. Rep. 769: 14 p.
- KLEINENBERG, S.E., A.V. YABLOKOV, B.M. BELKOVICH, and M.N. TARASEVICH. 1964. Beluga (Delphinapterus leucas) Investigation of the Species. Academy of Sciences of the U.S.S.R. A.N. Severtson Institute of Animal Morphology. vi + 376 p. (IPST (Israel Program for Scientific Translations 1969)).
- McGHEE, R. 1974. Beluga hunters. An archaeological reconstruction of the history and culture of the Mackenzie Delta Kittygaryumiut. Newfoundland social and economic studies No. 13. Institute of Social and Economic Research. Memorial University of Newfoundland. 124 p.
- NORRIS, K.S. 1961. Committee on Marine Mammals Standardized Methods for measuring and recording data on the smaller cetaceans. J. Mammal. 42: 471-476.
- NULIGAK, 1971. I Nuligak. The autobiography of a Canadian Eskimo. edited and translated by Maurice Metayer. Pocket book edition. Simon and Schuster of Canada. 191 p.
- SLANEY, F.F., and COMPANY LTD. 1974. White whale study, Herschel Island - Cape Dalhousie, coastal region of the Beaufort Sea. Unpublished report for Imperial Oil Ltd. 29 p. + maps and appendices.
- SMITH, T.G., and D. TAYLOR. 1977. Notes on marine mammal, fox and polar bear harvests in the Northwest Territories, 1940-1972. Can. Fish. Mar. Serv. Tech. Rep. 694: 37 p.
- STRONG, J.T. 1989. Reported harvests of narwhal, walrus and beluga in the Northwest Territories, 1948-1987. Can. Data Rep. Fish. Aquat. Sci. 794: iv + 14 p.
- TOMILIN, A.G. 1957. Volume IX. Cetacea. In V.G. Heptner (ed). Mammals of the U.S.S.R. and adjacent countries. 717 p. (IPST (Israel Program for Scientific Translations 1967)).




Fig. 1 Map of Mackenzie River estuary, Northwest Territories (showing hunting camps).




Fig. 2. Temporal distribution of the beluga whale harvest in the Mackenzie River estuary, 1981-1986.




Fig. 3. comparison of mean length, range and standard deviation for male and female beluga whale harvested in the Mackenzie River estuary, 1981-1986.




Fig. 4. Length frequencies of beluga whale harvested in the Mackenzie River estuary, 1981-1986.

Table 1. Cartridge use and shots fired during successful beluga hunts in the Mackenzie River Estuary. Data reported by hunters. Calibers ranked in order of use.

Caliber	Number of Hunts	Number of Shots/Landed Whale	
		Average	Range
30/30 Winchester	418	6	1-50
.270 Winchester	69	7	1-20
.243 Winchester	68	6	1-15
.303 British	44	6	1-40
30/06 Spring field	44	6	1-20
.308 Winchester	35	5	1-15
.300 Savage	17	6	1-14
6 mm Remington	10	9	3-10
.30 Remington	3	11	8-13
.250 Savage	2	4	3-4
444 Marlin	2	6	2-10
12 GA.	2	1	-
7 mm Remington Magnum	1	1	-
.357 <sup>1</sup>	1	5	-
.300 <sup>1</sup>	1	5	-
25/06 Winchester	1	1	-
25/35 Winchester	1	NR <sup>2</sup>	-

<sup>1</sup> no other information provided on data sheet and more than one cartridge possibility exists. .357 may be an error.

<sup>2</sup> NR = not recorded

Table 2. Numbers of beluga reported as struck/landed, by hunting area 1981-1986, from observer records.

Year		West Whitefish	Kendall Island	East Whitefish	Tuktoyaktuk	Total
1981	Struck	41	25	35	76	177
	Landed	40 <sup>1</sup>	23	27	62	152
1982	Struck	28	28	27	44	127
	Landed	20	25	23	39	107
1983	Struck	16	28	15	43	102
	Landed	13	25	13	35	86
1984	Struck	21	31	37	67	156
	Landed	20	30	32	59	141
1985	Struck	15	35	28	67	149 <sup>2</sup>
	Landed	12	25	21	59	121 <sup>2</sup>
1986	Struck	22	25	35	117	199 <sup>3</sup>
	Landed	22	14	25	89	150 <sup>3</sup>
Totals	Struck	143	172	177	414	910
	Landed	127	142	141	343	757

<sup>1</sup>15 of the whales reported landed were utilized by Holman families.

<sup>2</sup>1985 Totals include 3 whales reported from Paulatuk and 1 whale reported from Coppermine which are not included in the hunting area totals.

<sup>3</sup>1986 Totals include 40 whales (20 landed) taken from Kugmallit Bay but not reported to monitors, and 6 whales taken in a net in the West Whitefish area which are not included in the struck total.

Table 3. Reported loss rates, (struck-landed/struck) from the Mackenzie River estuary beluga fishery 1981-1986.

Year	WWF	KI	EFW	TUK	Yearly Totals
1981	0.02	0.08	0.23	0.18	0.14
1982	0.23	0.11	0.15	0.11	0.16
1983	0.19	0.11	0.13	0.19	0.16
1984	0.05	0.03	0.14	0.12	0.10
1985	0.20	0.29	0.25	0.12	0.19
1986 <sup>1</sup>	0.14	0.44	0.29	0.24	0.26
1981-1986 mean	0.13	0.17	0.20	0.17	0.17

<sup>1</sup> 1986 loss rates do not include 6 beluga taken in a net.



Table 4. Annual breakdown, by sex of Mackenzie Delta beluga whale harvest 1981-1986 and 1974-1981.

Year	Male	Female	% of Harvest Reported
1981	60 (52%)	55 (48%)	76
1982	72 (73%)	27 (27%)	93
1983	68 (82%)	15 (18%)	97
1984	111 (87%)	16 (13%)	90
1985	76 (74%)	27 (26%)	85
1986	64 (53%)	57 (47%)	81

Table 5. Range, mean and standard deviation of various measurements from beluga harvested in the Mackenzie River estuary 1981-1986. From observers record sheets.

	sex	n.	size range cm.	mean	s.d.
1981 <u>Total Length</u>	M	51	208 - 549	421.16	51.33
	F	53	152 - 429	361.32	43.53
1982 <u>Total Length</u>	M	69	274 - 505	430.75	48.62
	F	27	190 - 450	357.56	54.71
1983 <u>Total Length</u>	M	60	154 - 473	423.35	43.40
	F	15	353 - 452	378.53	28.43
<u>Axillary Girth</u>	M	38	88 - 398	260.13	44.36
	F	9	200 - 254	225.44	18.85
<u>Fluke Width</u>	M	20	36 - 117	97.35	17.77
	F	10	62 - 102	78.80	13.28
<u>Flipper Length</u>	M	28	23 - 53	40.68	7.74
	F	9	30 - 46	36.67	5.12
<u>Flipper Width</u>	M	27	12 - 48	33.63	7.89
	F	9	25 - 41	29.67	4.92
1984 <u>Total Length</u>	M	85	310 - 495	429.73	29.81
	F	15	310 - 442	371.00	31.30
<u>Axillary Girth</u>	M	51	168 - 356	258.16	31.45
	F	12	183 - 264	219.33	21.92
<u>Fluke Width</u>	M	29	79 - 117	100.21	9.91
	F	6	53 - 104	77.17	16.24
<u>Flipper Length</u>	M	34	36 - 52	45.24	4.04
	F	8	38 - 46	41.00	2.83
<u>Flipper Width</u>	M	33	20 - 37	31.67	4.06
	F	8	24 - 33	27.88	2.53

cont'd

Table 5. Continued

	Sex	n.	size range cm.	mean	s.d.
1985	<u>Total Length</u>				
	M	76	338 - 513	432.00	30.95
	F	26	295 - 434	366.69	28.25
	<u>Axillary Girth</u>				
	M	71	190 - 321	251.86	29.51
	F	22	178 - 244	203.68	17.74
	<u>Fluke Width</u>				
	M	70	71 - 121	100.91	10.75
	F	23	71 - 104	83.22	7.61
	<u>Flipper Length</u>				
	M	68	32 - 58	44.03	4.83
	F	21	36 - 46	39.71	2.99
<u>Flipper Width</u>					
M	68	27 - 40	32.91	3.04	
F	21	24 - 33	27.76	2.83	
1986	<u>Total Length</u>				
	M	59	343 - 493	421.64	28.01
	F	50	320 - 457	378.48	28.14
	<u>Axillary Girth</u>				
	M	59	190 - 328	247.37	30.58
	F	44	137 - 259	210.48	26.17
	<u>Fluke Width</u>				
	M	56	84 - 127	100.94	9.13
	F	45	51 - 99	81.16	9.11
	<u>Flipper Length</u>				
	M	58	30 - 53	43.48	4.55
	F	46	30 - 48	38.63	3.93
<u>Flipper Width</u>					
M	58	23 - 51	32.98	4.71	
F	46	18 - 36	27.85	3.20	

Table 6. Total lengths of female beluga known to be currently reproductive, Mackenzie River Delta 1981-1986.

Year	n	Range (cm)	Mean	S.D.
1981	27	320-419	374	21.90
1982	4	320-381	356	27.21
1983	8	356-383	368	9.91
1984	10	320-385	365	19.20
1985	12	333-434	363	18
1986	13	320-457	369	33
Total	49	320-457	364	21.44

Table 7. Total lengths (cm) and date of collection for fetuses and neonates collected in the Mackenzie River Estuary Beluga Fishery 1981-1986.

Date	July														August	
	2	6	10	12	13	16	17	18	20	21	22	23	24	25	27	2
Non-Term	17.78		21.25	12.7	17.78	38.1	15.0	17.78		10.8				20.32	17.5	22.5
Term				165.10	172.72			154.5		120.0	190.5	152.4				
Neonate	197.95	208.28	190.50													

Table 8. Mean, range and standard deviation from total lengths of fetuses and neonates collected in the Mackenzie River Estuary Beluga Fishery 1981-1986.

Category	n	Range (cm)	Mean	S.D.
1 Non-term Foetus	11	10.8-38.1	19.23	7.16
2 Term Foetus	6	120.0-190.50	159.20	23.67
3 Neonates	3	190.50-208.28	198.91	8.93
2 & 3	9	120.0-208.28	172.44	27.65

Table 9. Range and average of the number of teeth recorded from beluga harvested in the Canadian Beaufort sea.

---

<u>Upper Jaw</u>		<u>Lower Jaw</u>	
number	23	number	23
range	13-20	range	12-20
average	17.6	average	16.5
<u>Upper Left</u>		<u>Lower Left</u>	
number	21	number	21
range	7-11	range	6-10
average	17.6	average	16.5
<u>Upper Right</u>		<u>Lower Right</u>	
number	21	number	21
range	6-11	range	6-10
average	17.6	average	16.5
<u>OVERALL AVERAGES</u>			
	<u>Upper</u>		<u>Lower</u>
left	9		8
right	9		9

---

APPENDIX 1  
Data Sheets





BELUGA WHALE STUDY - DAILY RECORD  
 - FISH AND MARINE MAMMAL MANAGEMENT DIVISION

NO: \_\_\_\_\_

AREA: \_\_\_\_\_ DATE: \_\_\_\_\_

FIELD WORKER/MONITOR: \_\_\_\_\_

A. THE HUNT:

WEATHER: \_\_\_\_\_ SUNNY \_\_\_\_\_ CLOUDY \_\_\_\_\_ WINDY \_\_\_\_\_ RAIN  
 WATER: \_\_\_\_\_ RIPPLES (1-6 INCHES) \_\_\_\_\_ ROUGH (1-2 FEET)  
 \_\_\_\_\_ SMALL WAVES (1/2-1 FOOT) \_\_\_\_\_ STORM (OVER 2 FEET)

HUNTER NAME(S): \_\_\_\_\_  
 \_\_\_\_\_ COMMUNITY \_\_\_\_\_

TIME OUT OF CAMP \_\_\_\_\_ TIME RETURNED TO CAMP \_\_\_\_\_

DID SEE WHALES? \_\_\_\_\_ YES \_\_\_\_\_ NO IF YES, HOW MANY? \_\_\_\_\_

HOW MANY WHALES STRUCK (SHOT)? \_\_\_\_\_ HOW MANY WHALES LANDED? \_\_\_\_\_


HOW MANY WHALES WOUNDED, LOST OR SUNK? \_\_\_\_\_

CALIBRE OF RIFLE? \_\_\_\_\_ TOTAL NUMBER OF SHOTS FIRED? \_\_\_\_\_

GENERAL COMMENTS: \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_

B. SAMPLE INFORMATION:

"STRAIGHT" LINE FROM TIP OF SNOUT TO NOTCH  
 IN TAIL



COLOR:

\_\_\_\_\_ BROWN \_\_\_\_\_ GREY OR BLUISH  
 \_\_\_\_\_ WHITE WITH GREY ON FLUKE AND FLIPPERS  
 \_\_\_\_\_ WHITE \_\_\_\_\_ YELLOW

STOMACH: \_\_\_\_\_ FULL \_\_\_\_\_ 1/2 FULL \_\_\_\_\_ EMPTY

IF FOOD PRESENT: \_\_\_\_\_ MOSTLY FISH  
 \_\_\_\_\_ MOSTLY SHRIMP OTHER: \_\_\_\_\_  
 \_\_\_\_\_

LENGTH: \_\_\_\_\_ FEET \_\_\_\_\_ INCHES

SEX: \_\_\_\_\_ MALE \_\_\_\_\_ FEMALE

IF FEMALE: WAS SHE WITH A NEWBORN CALF? \_\_\_\_\_ YES \_\_\_\_\_ NO

WAS SHE GIVING MILK? \_\_\_\_\_ YES \_\_\_\_\_ NO

WAS SHE PREGNANT? \_\_\_\_\_ YES \_\_\_\_\_ NO

IF PREGNANT, MEASURE LENGTH OF FETUS (UNBORN CALF). \_\_\_\_\_ FEET \_\_\_\_\_ INCHES

SAMPLES TAKEN: \_\_\_\_\_ JAW (1/2 OF LOWER JAW WITH TEETH)

\_\_\_\_\_ TESTIS

\_\_\_\_\_ UTERUS AND OVARIES

\_\_\_\_\_ FETUS

\_\_\_\_\_ STOMACH CONTENTS

\_\_\_\_\_ OTHER: \_\_\_\_\_

NOTE: PLEASE INSURE THAT THE SAMPLE NUMBER ON THE SPECIMEN TAG IS THE SAME NUMBER AS SHOWN ON THE TOP RIGHT OF THIS PAGE.



SAMPLE NO. \_\_\_\_\_ SEX \_\_\_\_\_ DATE \_\_\_\_\_ SAMPLERS \_\_\_\_\_  
sp. mo. yr. no. yr. mo. day  
 SPECIES \_\_\_\_\_ Location \_\_\_\_\_ lat. \_\_\_\_\_ long. \_\_\_\_\_  
deg. min. sec. deg. min.  
 Hunter(s) \_\_\_\_\_ Community \_\_\_\_\_ TIME AFTER DEATH \_\_\_\_\_  
hrs.  
 Weather: wind \_\_\_\_\_ sky \_\_\_\_\_ water/ice \_\_\_\_\_

PIGMENTATION:

CAUSE OF DEATH  
bullet entry and path

SCARS

TOTAL LENGTH 1 \_\_\_\_\_ cm  
 FLUKE WIDTH 2 \_\_\_\_\_ cm  
 GIRTH AT ARMPIT 3 \_\_\_\_\_ cm  
 GIRTH AT NAVEL 4 \_\_\_\_\_ cm  
 FLIPPER LENGTH  
 anterior origin to tip 5 \_\_\_\_\_ cm  
 axilla to tip 6 \_\_\_\_\_ cm  
 maximum width 7 \_\_\_\_\_ cm  
 TUSK  
 exposed length 8 \_\_\_\_\_ cm  
 total length \_\_\_\_\_ cm  
 basal circ. \_\_\_\_\_ cm  
 condition \_\_\_\_\_

MILK amt: none/little/lots  
 (tick boxes)  
 lt. green  thick   
 dk. green  thin   
 yellow  oily   
 cream  sticky   
 white  watery   
 clear  other   
 opaque

STOMACH CONTENTS: description

AMOUNT  
SAMPLES

BLUBBER THICKNESS

at armpit MUKTUK BLUBBER  
 back \_\_\_\_\_ cm  
 side \_\_\_\_\_ cm  
 belly \_\_\_\_\_ cm  
 at navel  
 back \_\_\_\_\_ cm  
 side \_\_\_\_\_ cm  
 belly \_\_\_\_\_ cm

FOETUS

if FULL TERM — sampled? yes/no  
 sample no. \_\_\_\_\_  
 if NON TERM  
 which horn? left/right  
 sex ♂ ♀  
 length \_\_\_\_\_ cm  
 weight \_\_\_\_\_ kg  
 placental wt. \_\_\_\_\_ kg

snout or Tail  
 to anus 11 \_\_\_\_\_ cm  
 to mid-point of genital slit 12 \_\_\_\_\_ cm  
 to navel 13 \_\_\_\_\_ cm  
 to anterior origin of flipper 14 \_\_\_\_\_ cm  
 to ear 15 \_\_\_\_\_ cm  
 to eye 16 \_\_\_\_\_ cm  
 to blowhole 17 \_\_\_\_\_ cm  
 to angle of mouth 18 \_\_\_\_\_ cm  
 depth of tail notch 19 \_\_\_\_\_ cm  
 girth at anus 20 \_\_\_\_\_ cm  
 girth at eye 21 \_\_\_\_\_ cm  
 eye to ear (surface) \_\_\_\_\_ cm  
 eye to blowhole (surface) \_\_\_\_\_ cm  
 eye to angle of mouth (surface) \_\_\_\_\_ cm  
 blowhole width \_\_\_\_\_ cm

TESTIS left/right  
 weight \_\_\_\_\_ kg  
 length \_\_\_\_\_ cm  
 width \_\_\_\_\_ cm  
 height \_\_\_\_\_ cm  
 sperm? no/little/lots

MAMMARY GLAND left right  
 thickness \_\_\_\_\_ cm  
 UTERINE CORNUA  
 diameter \_\_\_\_\_ cm  
 CORPORA LUTEA  
 present? \_\_\_\_\_

COMMENTS-


APPENDIX 2





Temporal distribution of the beluga whale harvest in the  
Mackenzie River Estuary: Annually, by area and cumulative,  
1981-1986.

## TEMPORAL DISTRIBUTION 1981

FEMALES

MALES










# TEMPORAL DISTRIBUTION 1982

FEMALES



MALES










## TEMPORAL DISTRIBUTION 1983

FEMALES

MALES












## TEMPORAL DISTRIBUTION 1984

FEMALES



MALES










## TEMPORAL DISTRIBUTION 1985

females








males





## TEMPORAL DISTRIBUTION 1986





APPENDIX 3

Range, mean and standard deviation of total lengths of  
beluga from the Mackenzie River Estuary, annually, 1979-1986



Table A3.1. Range, mean and standard deviation of total lengths from the 1979 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean	S.D. (cm)
Kendall Island	male	2	360.68-421.64	391.16	43.11
	female	8	345.44-383.54	363.22	11.76
		10			
East Whitefish	male	10	365.76-462.28	423.67	26.61
	female	4	347.98-426.72	377.83	35.52
		14			
Delta Totals	male	12	360.68-462.28	418.25	30.14
	female	12	345.44-426.72	368.09	22.00
		24			

Table A3.2. Range, mean and standard deviation of total lengths from the 1980 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean (cm)	S.D. (cm)
West Whitefish	male	9	426.72-548.64	472.72	38.49
	female	5	365.76-457.20	384.05	40.89
	total	14			
Kendall Island	male	14	365.76-500.38	433.61	41.37
	female	7	381.00-434.34	402.05	17.58
	total	21			
East Whitefish	male	2	338.62-472.44	450.53	59.27
	female	9	355.60-447.04	384.95	35.95
	total	11			
Tuktoyaktuk	male	10	411.48-513.08	470.41	32.48
	female	13	320.04-457.20	374.75	41.21
	total	23			
Delta Totals	male	35	365.76-548.64	454.01	
	female	34	335.28-457.20	385.23	
	total	69			

Table A3.3. Range, mean and standard deviation of total lengths from the 1981 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean (cm)	S.D. (cm)
West Whitefish	male	13	208.28-548.64	427.94	78.61
	female	7	281.94-378.46	344.90	40.73
Kendall Island	male	6	391.16-464.82	426.30	29.68
	female	15	335.28-406.40	379.81	18.81
East Whitefish	male	8	365.76-487.68	432.75	37.08
	female	16	152.40-419.10	347.19	61.01
Tuktoyaktuk	male	24	248.92-480.06	412.43	41.49
	female	15	312.42-429.26	365.68	34.94
Delta Totals	male	51	208.28-487.68	421.20	51.27
	female	53	152.40-429.26	361.35	43.48

Table A3.4. Range, mean and standard deviation of total lengths from the 1982 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean (cm)	S.D. (cm)
West Whitefish	male	13	386.08-480.06	436.32	26.46
	female	3	365.76-449.58	398.78	44.65
Kendall Island	male	23	335.28-469.90	424.44	33.23
	female	2	335.28-365.76	350.52	21.55
East Whitefish	male	20	350.52-505.45	452.31	42.85
	female	0	0	0	0
Tuktoyaktuk	male	14	304.80-487.68	399.14	72.54
	female	21	190.50-449.58	350.40	57.20
Delta Totals	male	70	304.80-505.45	430.34	47.79
	female	28	190.50-449.58	363.71	60.53

Table A3.5. Range, mean and standard deviation of total lengths from the 1983 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean (cm)	S.D. (cm)
West Whitefish	male	8	375.92-457.20	419.74	30.74
	female	5	363.22-429.26	388.49	24.46
Kendall Island	male	24	360.68-457.20	436.35	22.19
	female	1	396.24	396.24	0
East Whitefish	male	9	389.89-472.44	421.50	24.70
	female	1	360.68	360.68	0
Tuktoyaktuk	male	21	368.30-468.75	423.73	25.53
	female	7	335.60-372.50	358.06	11.99
Delta Totals	male	62	360.68-472.44	427.78	25.23
	female	14	335.60-429.26	371.84	22.69

Table A3.6. Range, mean and standard deviation of total lengths from the 1984 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean (cm)	S.D. (cm)
West Whitefish	male	10	365.76-495.30	442.47	32.43
	female	1	396.24	396.24	0
Kendall Island	male	21	309.88-477.52	430.47	36.60
	female	5	320.04-406.40	361.70	23.94
East Whitefish	male	20	403.86-487.68	425.13	20.76
	female	3	309.88-384.81	355.18	39.18
Tuktoyaktuk	male	35	349.25-477.52	429.93	30.44
	female	5	353.06-441.96	379.48	36.92
Delta Totals	male	86	309.88-495.30	430.40	30.29
	female	14	309.88-441.96	369.12	31.68

Table A3.7. Range, mean and standard deviation of total lengths from the 1985 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean (cm)	S.D. (cm)
West Whitefish	male	8	388-488	432	34
	female	3	295-366	342	41
Kendall Island	male	23	338-465	431	26
	female	2	376-381	378	4
East Whitefish	male	17	363-465	420	30
	female	5	335-368	367	39
Tuktoyaktuk	male	28	361-513	440	35
	female	16	333-434	374	28
Delta Totals	male	76	338-513	432	31
	female	26	295-434	367	28

Table A3.8. Range, mean and standard deviation of total lengths from the 1984 Mackenzie Delta beluga whale harvest. All lengths in cm.

Location	Sex	n	Range (cm)	Mean (cm)	S.D. (cm)
West Whitefish	male	7	396-467	429	27
	female	7	353-432	387	27
Kendall Island	male	8	386-493	427	32
	female	6	366-389	378	9
East Whitefish	male	19	366-448	412	23
	female	6	335-422	372	35
Tuktoyaktuk	male	25	343-470	425	30
	female	31	320-457	378	30
Delta Totals	male	59	343-493	422	28
	female	50	320-457	378	28

APPENDIX 4



Length frequencies of beluga harvested in the Mackenzie River Estuary:  
Annually, by area and cumulative, 1981-1985

## LENGTH FREQUENCY DISTRIBUTION 1981

FEMALES



MALES




## LENGTH FREQUENCY DISTRIBUTION 1981

west whitefish

FEMALES

MALES




## LENGTH FREQUENCY DISTRIBUTION 1981

kendall island

FEMALES

MALES




## LENGTH FREQUENCY DISTRIBUTION 1981

east whitefish

FEMALES

MALES




## LENGTH FREQUENCY DISTRIBUTION 1981

tuktoyaktuk

FEMALES

MALES




## LENGTH FREQUENCY DISTRIBUTION 1982

FEMALES



MALES






LENGTH FREQUENCY DISTRIBUTION 1982

west whitefish

FEMALES

MALES




LENGTH FREQUENCY DISTRIBUTION 1982

kendall island

FEMALES

MALES




LENGTH FREQUENCY DISTRIBUTION 1982

east whitefish

FEMALES

MALES





LENGTH FREQUENCY DISTRIBUTION 1982

tuktoyaktuk

FEMALES

MALES






LENGTH FREQUENCY DISTRIBUTION 1983

west whitefish

FEMALES

MALES




LENGTH FREQUENCY DISTRIBUTION 1983

kendall island

FEMALES

MALES




LENGTH FREQUENCY DISTRIBUTION 1983

east whitefish

FEMALES

MALES




LENGTH FREQUENCY DISTRIBUTION 1983


tuktoyaktuk





FEMALES

MALES



## LENGTH FREQUENCY DISTRIBUTION 1984











# LENGTH FREQUENCY DISTRIBUTION 1985

FEMALES




MALES





## LENGTH FREQUENCY DISTRIBUTION 1986






## LENGTH FREQUENCY DISTRIBUTION 1986

west whitefish

FEMALES

MALES




## LENGTH FREQUENCY DISTRIBUTION 1986

kendall island

FEMALES

MALES




## LENGTH FREQUENCY DISTRIBUTION 1986

east whitefish

FEMALES

MALES




## LENGTH FREQUENCY DISTRIBUTION 1986

tuktoyaktuk

FEMALES

MALES



Appendix 5. Number of teeth recorded from beluga harvested in the Canadian Beaufort Sea.

Whale Length cm	sex	number of teeth	upper			lower		
			right	left	total	right	left	total
335	M	32	8	8	16	8	8	16
349	M	37	10	9	19	9	9	18
389	M	34	9	8	17	9	8	17
396	M	40	10	10	20	10	10	20
403	M	35	9	9	18	8	9	17
410	M	35	8	9	17	9	9	18
410	M	28	6	7	13	8	7	15
411	M	27	7	7	14	7	6	13
411	M	28	8	8	16	6	6	12
414	M	34	9	9	18	8	8	16
414	M	38	10	10	20	9	9	18
415	M	36	9	9	18	9	9	18
417	M	28	7	7	14	7	7	14
421	M	36	9	9	18	9	9	18
422	M	35			18			17
426	M	38	11	9	20	9	9	18
426	M	34	9	9	18	8	8	16
429	M	31	8	8	16	7	8	15
430	M	33	8	8	16	8	9	17
435	M	39	10	11	21	9	9	18
457	M	32			18			14
472	M	37	10	10	20	9	8	17
477	M	38	11	9	20	9	8	17
327	F	31	6	7	13	9	9	18
360	F	39	10	10	20	10	9	19
411	F	32	9	9	18	7	7	14
?	F	30	8	8	16	7	7	14