



Scientific Excellence • Resource Protection & Conservation • Benefits for Canadians
Excellence scientifique • Protection et conservation des ressources • Bénéfices aux Canadiens

Life History Data on Alewife and Blueback Herring, Mactaquac Dam, 1982-88

B.M. Jessop and W.E. Anderson

Freshwater and Anadromous Division
Biological Sciences Branch
Department of Fisheries and Oceans
Halifax, Nova Scotia B3J 2S7

1991

**Canadian Data Report of Fisheries
and Aquatic Sciences No. 829**



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Data Report of Fisheries and Aquatic Sciences

Data reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of data reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was introduced with the publication of report number 161.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport statistique canadien des sciences halieutiques et aquatiques

Les rapports statistiques servent à classer et à archiver les compilations de données pour lesquelles il y a peu ou point d'analyse. Ces compilations auront d'ordinaire été préparées à l'appui d'autres publications ou rapports. Les sujets des rapports statistiques reflètent la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports statistiques ne sont pas destinés à une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans autorisation écrite préalable de l'établissement auteur. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 25 de cette série ont été publiés à titre de relevés statistiques, Services des pêches et de la mer. Les numéros 26 à 160 ont été publiés à titre de rapports statistiques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 161.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épousés seront fournis contre rétribution par des agents commerciaux.

Canadian Data Report of
Fisheries and Aquatic Sciences No. 829

1991



LIFE-HISTORY DATA ON ALEWIFE
AND BLUEBACK HERRING,
MACTAQUAC DAM, 1982-88

by

B.M. Jessop and W.E. Anderson

Freshwater and Anadromous Division
Biological Science Branch
Department of Fisheries and Oceans
Halifax, Nova Scotia
B3J 2S7

©Minister of Supply and Services Canada 1991
Cat. No. Fs97-13/0829 ISSN 0706-6465

Correct citation for this publication:

Jessop, B.M. and W.E. Anderson 1982-88. Life-history data on alewife and blueback herring, Mactaquac Dam, 1982-88. Can. Data Rep. Fish. Aquat. Sci. No. 829: 42 p.

CONTENTS

LIST OF TABLES	iv
ABSTRACT/RÉSUMÉ	vii
INTRODUCTION	1
METHODS	1
RESULTS AND DISCUSSION	1
FIGURE	2
ACKNOWLEDGEMENTS	42
REFERENCES	42

LIST OF TABLES

TABLE 1.	Mean fork length of alewife, by sex and age, Mactaquac Dam (1982-88)	3
TABLE 2.	Mean fork length of blueback herring, by sex and age, Mactaquac Dam (1982-88)	4
TABLE 3.	Spawning history of alewife, by sex and year, Mactaquac Dam (1982-88)	5
TABLE 4.	Spawning history of blueback herring, by sex and year, Mactaquac Dam (1982-88)	6
TABLE 5.	(a-g). Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam (1982-88)	7-12
TABLE 6.	(a-g). Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam (1982-88)	13-17
TABLE 7.	Age at first spawning (maturity), by sex and year, for alewife at Mactaquac Dam (1982-88)	18
TABLE 8.	Age at first spawning (maturity), by sex and year, for blueback herring at Mactaquac Dam (1982-88)	19
TABLE 9.	Length-frequency distribution of mature alewives, by sex and year, Mactaquac Dam (1982-88)	20
TABLE 10.	Length-frequency distribution of mature blueback herring, by sex and year, Mactaquac Dam (1982-88)	21
TABLE 11.	Mean weight of alewife, by sex and age, Mactaquac Dam (1982-88)	22
TABLE 12.	Mean weight of blueback herring, by sex and age, Mactaquac Dam (1982-88)	23

TABLE 13.	Mean fork lengths, by sample date and year, for alewife at Mactaquac Dam (sexes combined, 1982-88)	24
TABLE 14.	Mean fork lengths, by sample date and year, for blueback herring at Mactaquac Dam (sexes combined, 1982-88).....	25
TABLE 15.	(a-g). Length-frequency distribution of alewife, by age, sexes combined, Mactaquac Dam (1982-88)	26-30
TABLE 16.	(a-g). Length-frequency distribution of blueback herring, by age, sexes combined, Mactaquac Dam (species and sexes combined, 1982-88)	31-35
TABLE 17.	(a-g). Length-frequency distribution of gaspereau, by age, Mactaquac Dam (species and sexes combined, 1982-88)	36-41

ABSTRACT

Jessop, B.M. and W.E. Anderson. 1991. Life-history data on alewife and blueback herring, Mactaquac Dam, 1982-88. Can. Data Rep. Fish. Aquat. Sci. No. 829: 42p.

Life-history data were collected annually between 1982 and 1988 for the alewife and blueback herring migrating to the Mactaquac Dam, Saint John River, New Brunswick. This report tabulates the data, in various combinations, according to species, sex, length, weight, age, maturity and spawning history.

Key words: Life-history, alewife, blueback herring, gaspereau, anadromous, Mactaquac Dam, Saint John River, hydroelectric development.

RÉSUMÉ

Jessop, B.M. and W.E. Anderson. 1991. Life-history data on alewife and blueback herring, Mactaquac Dam, (1982-88.) Can. Data Rep. Fish. Aquat. Sci. No. 829: 42p.

Entre 1982 et 1988, on a recueilli chaque année des données relatives au cycle biologique du gaspareau et de l'aloise d'été remontant la rivière Saint-Jean jusqu'au barrage de Mactaquac (Nouveau-Brunswick). Ce rapport présente les données sous forme de tableaux, en les combinant de diverses façons, selon l'espèce, le sexe, la longueur, le poids, l'âge la maturité et l'historique du frai.

INTRODUCTION

Creation of a large headpond on the Saint John River, New Brunswick, by the completion of the Mactaquac Dam in 1967 led to the rapid development of major stocks of alewives (*Alosa pseudoharengus*) and blueback herring (*A. aestivalis*), collectively and colloquially termed gaspereau. In 1973, a study was begun of the life history and population dynamics of the gaspereau of the Saint John River, with the objectives of: (1) establishing whether alewife and blueback herring populations are homogeneous within the river or form distinct local populations, (2) evaluating the influence of the commercial fishery and hydroelectric development on the populations, and (3) providing a data base for the establishment and evaluation of management programs.

This report presents tabulations of the life-history data collected annually between 1982 and 1988. The data for 1973 and 1974 are given in Jessop (1977 a, b), and for 1975-81 in Jessop et. al. (1982).

Mactaquac Dam is the furthest downstream (148 km from the river mouth) of the three main-stem hydroelectric dams constructed on the Saint John River (Fig. 1). Its construction created a headpond of approximately 83.5 km², which extends about 104 km upriver. During low-flow periods, the dam is operated as a peak-load plant, and during high-flow periods (spring runoff and rainfall floods), as a base-load plant. Fish passage facilities include both a mechanical elevator and tank trucks, which in combination relay fish to release points upstream of the dam.

METHODS

During the run, random samples of 100 were collected twice weekly from the fish-lift. Each fish was measured for fork length to the nearest millimetre, weighed to the nearest gram and identified as to sex, state of maturity and species. Scale samples were collected for aging. Maturity was classified according to Nikolsky (1963). Species were identified according to color of the peritoneum (Bigelow and Schroeder 1953).

Whenever possible, scales were taken from the left side of each fish, just above the lateral line and behind the posterior edge of the dorsal fin. Eight to ten scales were placed on transparent plastic slides (Power 1964), and each sample was independently aged twice without reference to associated length or weight data. If readings differed, a third reading was made and an age assigned on the basis of majority agreement. A few fish were unageable, due either to a lack of scales, scales with regenerate centers or aging disagreements.

Age was determined for adult fish by counting the number of annuli and spawning marks and adding a year for the scale edge (Cating 1953; Marcy 1969). Fish under age four frequently exhibited "plus" growth, and judgement based upon capture date was used in deciding whether or not to add a year for the scale edge. Scale analysis included counting the number of transverse grooves in order to establish a criterion for locating annuli that were difficult to interpret.

The age at first spawning of mature, virgin (no previous spawns evident on scales) gaspereau was recorded as the age at capture during the spawning run. A small proportion of older, mature fish show checks, sometimes resembling spawning checks, between annuli laid previously at Age 2 and, occasionally, at Age 3. Such checks at Age 2 are presumed not to be spawning marks but evidence of freshwater entry, because virtually all (>99%) Age-2 fish sampled have been sexually immature and small numbers of immature fish (mostly alewives [>85%] and mostly Age 2, but with a few [<5%] Age 3) are frequently found in the lake system of the lower Saint John River, e.g., Washademoak and Grand lakes, during the latter part of the spawning run (mid to late June). Non-annular checks at Age 3 were considered as spawning marks if well defined erosion and little plus growth to the annulus were evident; otherwise they were considered freshwater checks. The effect on spawning history patterns of any errors in judgement should be minimal, given the small numbers of fish involved.

RESULTS AND DISCUSSION

The data have been tabulated in various combinations according to species, sex, length, weight, age, maturity and spawning history (Tables 1-17). Between tables, totals may not always agree due to incomplete data; i.e., unageable scales. Variance has been calculated using degrees of freedom (n-1) rather than n. The relationship between sample standard deviation (s) and the estimated population standard deviation (s)

$$is s = s \sqrt{\frac{n-1}{n}}$$

The calculation of standard deviation as the square root of s , as done here, consistently underestimates σ (parametric population standard deviation), particularly for sample sizes <30. The bias can be corrected by applying the correction factor given in Sokal and Rohlf (1981, p.53). No analysis has been presented of the data contained in this report; the analysis will be made in a subsequent technical report.

Conversion between fork and total lengths for alewives in the fork length interval from 130 mm to 320 mm and for blueback herring in the interval from 180 to 270 mm, respectively, can be obtained from Jessop et. al. (1982).




Figure 1. Lower Saint John River system, showing location of Mactaquac Dam.

Table 1. Mean fork length (mm) of alewife by sex, age and year, Mactaquac Dam, 1982 - 1988.

Table 2. Mean fork length (mm) of blueback herring by sex, age and year, Mactaquac Dam, 1982 - 1988.

Year	Age	Male				Female			
		N	Mean	Range	SD	N	Mean	Range	SD
1982	3	143	202.9	183-235	9.29	36	210.1	187-233	11.13
	4	97	232.4	202-261	12.25	93	238.2	207-268	12.05
	5	66	246.7	222-281	11.38	93	255.9	226-282	10.46
	6	40	251.6	230-267	8.40	68	263.7	237-279	7.41
	7	19	257.3	245-268	6.43	25	269.8	248-287	10.90
	8	28	263.6	246-292	10.76	26	276.8	262-299	9.31
	9	26	268.6	256-283	7.24	30	279.0	252-307	11.50
	10	13	266.5	253-284	8.79	9	287.0	278-299	7.30
	11	2	261.0	260-262	1.41	4	289.3	282-295	5.91
	12	1	265.0	265-265	-	1	302.0	302-302	-
1983	3	136	201.0	178-237	10.74	49	211.1	185-247	13.50
	4	120	220.4	190-255	12.88	107	231.8	191-265	10.76
	5	60	237.2	219-273	11.99	79	245.9	221-297	13.32
	6	13	251.5	225-278	14.81	25	260.9	240-292	13.41
	7	12	256.5	234-270	9.68	9	268.7	254-280	8.12
	8	6	262.5	259-271	4.51	8	273.8	256-281	7.91
	9	2	268.5	253-284	21.92	6	281.2	265-312	16.85
	10	7	266.1	254-277	6.96	0	-	-	-
	11	0	-	-	-	3	289.0	272-306	17.00
	12	1	288.0	288-288	-	1	293.0	293-293	-
1984	3	88	196.8	175-220	8.99	40	201.2	183-220	9.53
	4	196	219.1	195-251	7.26	208	226.8	208-255	7.74
	5	46	231.2	207-257	10.21	61	237.9	217-262	8.94
	6	13	243.6	231-271	12.18	13	252.9	240-280	12.77
	7	6	249.5	235-258	8.69	5	263.4	256-269	5.32
	8	0	-	-	-	5	270.0	242-302	22.44
	9	2	266.5	265-268	2.12	2	285.0	282-288	4.24
	10	1	260.0	260-260	-	1	280.0	280-280	-
	11	0	-	-	-	0	-	-	-
	12	0	-	-	-	1	302.0	302-302	-
1985	3	330	191.8	166-219	8.50	170	198.8	180-228	7.55
	4	130	217.2	186-255	10.89	177	226.7	195-250	9.77
	5	50	229.7	210-245	7.89	62	237.1	214-264	10.87
	6	11	238.7	216-252	10.17	11	249.8	235-270	9.61
	7	2	261.5	252-271	13.44	0	-	-	-
	8	0	-	-	-	2	280.0	280-280	0.00
1986	3	85	193.0	171-220	10.09	15	199.0	184-212	8.73
	4	138	213.2	190-247	9.82	223	221.3	198-246	8.60
	5	15	232.6	213-273	13.96	13	241.8	220-271	13.34
	6	4	246.3	240-253	5.32	2	247.5	235-260	17.68
1987	3	41	192.6	177-222	9.26	16	196.6	184-213	8.57
	4	249	210.4	188-237	9.56	278	215.6	197-242	8.44
	5	57	222.0	198-246	10.24	107	236.5	213-258	8.83
	6	3	243.0	235-257	12.17	5	251.0	242-262	9.67
	7	2	251.5	247-256	6.36	1	255.0	255-255	-
1988	3	121	187.8	173-202	6.22	20	193.5	185-208	6.30
	4	242	207.8	190-231	8.14	210	215.4	194-241	9.47
	5	150	217.9	197-240	8.92	171	230.2	198-260	9.15
	6	16	232.9	210-261	14.71	29	242.3	225-256	8.82

Table 3. Spawning history of alewife, by sex and year, Mactaquac Dam, 1982 - 1988.

No. previous spawnings	1982			1983			1984			1985			1986			1987			1988				
	N	%	Mean age																				
Male																							
0	163	51.3	4.7	117	31.0	4.6	151	43.5	4.6	107	31.2	4.3	196	50.7	4.1	167	44.1	4.3	106	31.9	4.5		
1	67	21.1	5.0	121	32.1	5.2	98	28.2	4.9	134	39.1	4.8	158	40.8	4.1	119	31.4	4.9	107	32.2	5.1		
2	43	13.5	5.7	86	22.8	5.7	62	17.9	5.7	72	21.0	5.5	26	6.7	5.6	86	22.7	5.2	86	25.9	5.8		
3	26	8.2	6.9	40	10.6	6.3	22	6.3	6.5	22	6.4	6.9	6	1.6	6.3	7	1.9	6.6	33	9.9	6.1		
4	9	2.8	8.0	7	1.9	7.4	10	2.9	7.5	6	1.8	7.2	1	0.3	9.0	0	-	-	0	-	-		
5	9	2.8	8.7	4	1.1	8.3	4	1.2	8.3	1	0.3	8.0	0	-	-	0	-	-	0	-	-		
6	0	-	-	1	0.3	9.0	0	-	-	1	0.3	10.0	0	-	-	0	-	-	0	-	-		
7	1	0.3	10.0	0	-	-	0	-	-	0	-	-	0	-	-	0	-	-	0	-	-		
8	0	-	-	1	0.3	12.0	0	-	-	0	-	-	0	-	-	0	-	-	0	-	-		
Totals	318			377			347			343			387			379			332				
Repeat spawners	48.7			69.0			56.5			68.8			49.3			55.9			68.1			61	
Female																							
0	122	38.7	4.9	98	26.9	4.7	148	38.2	4.8	71	26.1	4.8	163	54.5	4.0	162	49.2	4.3	119	37.5	4.6		
1	69	21.9	5.4	117	32.1	5.3	108	27.9	5.2	102	37.5	5.2	105	35.1	4.2	101	30.7	5.0	84	26.5	5.3		
2	46	14.6	5.8	81	22.2	5.9	76	19.6	5.8	56	20.6	5.8	25	8.4	5.5	60	18.2	5.2	87	27.4	5.9		
3	47	14.9	7.6	49	13.4	6.3	22	5.7	6.5	25	9.2	7.0	5	1.7	6.2	6	1.8	6.0	24	7.6	6.4		
4	18	5.7	8.2	8	2.2	8.0	25	6.5	7.4	11	4.0	7.6	1	0.3	8.0	0	-	-	3	1.0	7.0		
5	7	2.2	8.6	9	2.5	8.3	4	1.0	8.5	6	2.2	8.7	0	-	-	0	-	-	0	-	-		
6	2	0.6	9.0	2	0.5	9.0	1	0.3	9.0	0	-	-	0	-	-	0	-	-	0	-	-		
7	3	1.0	10.0	1	0.3	10.0	2	0.5	10.0	1	0.4	10.0	0	-	-	0	-	-	0	-	-		
8	1	0.3	11.0	0	-	-	1	0.3	11.0	0	-	-	0	-	-	0	-	-	0	-	-		
Totals	315			365			387			272			299			329			317				
Repeat spawners	61.2			73.2			61.8			73.9			45.5			50.8			62.5				
Sexes combined																							
Repeat spawners	55.0			71.0			59.3			71.1			47.7			53.5			65.3				

Table 4. Spawning history of blueback herring, by sex and year, Mactaquac Dam, 1982 - 1988.

No. previous spawnings	1982			1983			1984			1985			1986			1987		
	N	%	Mean age															
Male																		
0	194	44.6	3.3	200	56.0	3.4	180	51.1	3.5	374	71.2	3.1	155	64.1	3.5	164	46.6	3.8
1	88	20.2	4.4	101	28.3	4.3	128	36.4	4.2	124	23.6	4.3	74	30.6	4.1	145	41.2	4.1
2	53	12.2	5.5	23	6.4	5.3	28	8.0	5.2	21	4.0	5.1	11	4.6	5.2	41	11.7	5.1
3	18	4.1	6.4	11	3.1	6.6	9	2.6	6.2	6	1.1	6.3	2	0.8	6.0	1	0.3	7.0
4	19	4.4	7.4	8	2.2	7.3	5	1.4	7.4	0	-	-	0	-	-	1	0.3	7.0
5	27	6.2	8.2	4	1.1	8.0	0	-	-	0	-	-	0	-	-	0	-	-
6	22	5.1	9.1	5	1.4	9.6	2	0.6	9.5	0	-	-	0	-	-	0	-	-
7	11	2.5	10.0	4	1.1	10.0	0	-	-	0	-	-	0	-	-	0	-	-
8	2	0.5	11.0	1	0.3	12.0	0	-	-	0	-	-	0	-	-	0	-	-
9	1	0.2	12.0	0	-	-	0	-	-	0	-	-	0	-	-	0	-	-
Totals	435			357			352			525			242			352		
Repeat spawners	55.4			44.3			48.9			28.8			35.9			53.4		
Female																		
0	110	28.5	3.7	142	50.0	3.8	174	51.8	3.8	285	66.0	3.4	163	63.7	3.9	218	53.6	4.1
1	84	21.8	4.7	90	31.7	4.6	115	34.2	4.3	119	27.6	4.3	84	32.8	4.1	140	34.4	4.3
2	81	21.0	5.7	17	6.0	5.5	28	8.3	5.2	23	5.3	5.3	9	3.5	5.2	48	11.8	5.1
3	27	7.0	6.3	18	6.3	6.6	10	3.0	6.6	3	0.7	6.0	0	-	-	1	0.2	7.0
4	25	6.5	7.5	4	1.4	7.8	4	1.2	8.3	1	0.2	8.0	0	-	-	0	-	-
5	29	7.5	8.5	4	1.4	8.0	3	0.9	8.0	1	0.2	8.0	0	-	-	0	-	-
6	16	4.2	9.1	5	1.8	9.0	0	-	-	0	-	-	0	-	-	0	-	-
7	11	2.9	10.3	2	0.7	11.0	1	0.3	10.0	0	-	-	0	-	-	0	-	-
8	2	0.5	11.5	2	0.7	11.5	0	-	-	0	-	-	0	-	-	0	-	-
9	0	-	-	0	-	-	1	0.3	12.0	0	-	-	0	-	-	0	-	-
Totals	385			284			336			432			256			407		
Repeat spawners	71.5			51.6			48.2			34.0			36.3			46.4		
Sexes combined																		
Repeat spawners	62.9			46.6			48.5			31.1			36.1			49.7		

Table 5a. Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam, 1982.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	1	0.3	241.0	-	182.0	-
4	0	47	14.8	253.2	10.04	219.6	30.84
4	1	17	5.3	255.7	10.70	223.9	26.02
5	0	108	34.0	267.4	9.66	254.0	28.41
5	1	34	10.7	270.6	8.26	263.1	27.15
5	2	21	6.6	269.9	8.89	259.8	33.63
6	0	7	2.2	272.9	7.80	277.1	22.30
6	1	15	4.7	280.1	9.91	296.0	36.94
6	2	16	5.0	277.4	8.39	293.3	31.87
6	3	11	3.5	278.2	13.30	280.2	40.12
7	1	1	0.3	274.0	-	263.0	-
7	2	5	1.6	287.6	6.39	321.6	40.87
7	3	6	1.9	291.5	6.89	337.2	16.90
7	4	2	0.6	296.0	11.31	367.0	12.73
8	2	1	0.3	305.0	-	364.0	-
8	3	9	2.8	292.3	10.44	343.0	42.55
8	4	5	1.6	285.0	9.38	317.8	26.99
8	5	3	0.9	302.3	4.16	386.0	18.73
9	4	2	0.6	294.0	18.38	336.0	72.12
9	5	6	1.9	301.2	6.68	361.3	33.21
10	7	1	0.3	280.0	-	319.0	-
Female							
4	0	21	6.7	260.1	6.86	246.5	28.41
4	1	8	2.5	261.1	7.90	245.6	12.42
5	0	89	28.3	275.7	8.59	300.4	35.90
5	1	33	10.5	276.6	9.43	304.5	28.38
5	2	21	6.7	282.1	9.25	318.6	43.83
6	0	12	3.8	281.8	8.64	323.3	43.94
6	1	24	7.6	289.6	7.25	343.5	31.45
6	2	14	4.4	291.1	8.93	353.3	46.42
6	3	5	1.6	289.8	11.90	360.8	58.25
7	1	4	1.3	298.3	10.21	380.5	26.86
7	2	10	3.2	300.9	5.42	400.5	34.28
7	3	11	3.5	301.2	18.20	387.4	29.30
7	4	4	1.3	307.0	8.29	416.3	53.62
8	2	1	0.3	298.0	-	372.0	-
8	3	31	9.8	305.5	9.08	417.8	44.95
8	4	7	2.2	309.0	9.57	421.3	53.86
8	5	4	1.3	303.0	9.31	424.3	47.99
9	4	7	2.2	303.9	8.59	414.9	33.19
9	5	2	0.6	301.5	2.12	419.5	3.54
9	6	2	0.6	304.5	6.36	404.0	52.33
10	5	1	0.3	323.0	-	482.0	-
10	7	3	1.0	304.0	19.70	414.0	92.48
11	8	1	0.3	287.0	-	334.0	-

Table 5b. Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam, 1983.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	2	0.5	230.0	2.83	166.0	11.31
4	0	45	11.9	249.7	9.90	214.3	28.93
4	1	17	4.5	251.1	10.54	217.7	30.90
5	0	69	18.2	265.9	13.13	261.9	41.52
5	1	67	17.7	270.1	12.06	273.3	39.06
5	2	32	8.4	273.3	14.32	287.0	37.88
6	0	2	0.5	271.0	15.56	266.5	16.26
6	1	36	9.5	277.5	6.52	289.5	28.44
6	2	47	12.4	278.1	9.49	293.9	31.57
6	3	29	7.7	283.1	11.07	310.2	40.68
7	1	1	0.3	297.0	-	360.0	-
7	2	8	2.1	282.9	2.95	306.1	12.98
7	3	11	2.9	285.3	12.34	316.8	43.87
7	4	4	1.1	286.0	13.56	319.5	31.72
8	4	3	0.8	292.7	18.61	345.7	54.24
8	5	3	0.8	287.7	13.58	327.3	27.59
9	5	1	0.3	268.0	-	272.0	-
9	6	1	0.3	268.0	-	261.0	-
12	8	1	0.3	293.0	-	401.0	-
Female							
3	0	1	0.3	217.0	-	146.0	-
4	0	25	6.7	257.3	10.99	243.2	39.19
4	1	15	4.0	252.8	13.45	222.7	41.72
5	0	71	19.1	277.1	13.31	314.3	48.58
5	1	51	13.7	281.1	10.84	323.1	41.03
5	2	25	6.7	279.5	11.16	319.4	49.27
6	0	1	0.3	286.0	-	336.0	-
6	1	50	13.5	287.3	10.88	351.2	48.37
6	2	46	12.4	293.5	10.87	371.4	48.61
6	3	37	10.0	292.7	11.45	364.4	46.76
7	1	3	0.8	299.7	4.16	398.0	3.00
7	2	10	2.7	297.8	8.70	393.9	45.03
7	3	12	3.2	300.5	9.81	403.7	51.90
7	4	2	0.5	287.0	7.07	357.0	35.36
8	2	1	0.3	295.0	-	367.0	-
8	3	2	0.5	301.5	6.36	430.0	4.24
8	4	4	1.1	309.8	19.62	438.5	88.01
8	5	6	1.6	301.8	12.58	406.5	63.54
9	4	2	0.5	304.5	3.54	425.0	18.38
9	5	3	0.8	312.7	5.13	437.3	35.23
9	6	3	0.8	293.0	16.64	359.7	96.02
10	7	1	0.3	276.0	-	294.0	-

Table 5c. Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam, 1984.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	1	0.3	223.0	-	150.0	-
4	0	69	19.9	247.7	8.67	206.8	26.24
4	1	31	8.9	241.3	7.73	194.3	29.69
5	0	73	21.0	258.3	7.69	236.5	26.61
5	1	49	14.1	260.8	12.32	247.0	37.03
5	2	30	8.6	267.4	10.93	263.6	37.65
6	0	8	2.3	268.1	9.30	266.9	34.34
6	1	18	5.2	277.3	8.75	303.9	39.32
6	2	23	6.6	276.5	6.22	295.7	21.64
6	3	11	3.2	282.6	7.82	325.4	33.46
7	2	9	2.6	285.3	8.80	328.0	36.98
7	3	11	3.2	288.7	9.73	339.8	50.23
7	4	6	1.7	280.8	9.26	321.2	37.81
8	4	3	0.9	292.0	1.00	362.0	22.54
8	5	3	0.9	286.3	17.04	370.3	36.36
9	4	1	0.3	304.0	-	384.0	-
9	5	1	0.3	302.0	-	379.0	-
Female							
2	0	1	0.3	150.0	-	41.0	-
3	0	1	0.3	210.0	-	119.0	-
4	0	37	9.5	249.7	11.13	221.6	37.02
4	1	19	4.9	250.7	9.80	217.2	27.85
5	0	97	25.0	265.9	10.62	273.1	37.16
5	1	49	12.6	268.3	12.32	284.3	43.96
5	2	36	9.3	272.3	12.69	295.3	43.03
5	3	1	0.3	255.0	-	271.0	-
6	0	13	3.4	278.4	6.22	317.3	31.67
6	1	36	9.3	282.6	10.75	337.9	46.76
6	2	17	4.4	287.5	8.34	343.9	39.96
6	3	12	3.1	285.5	6.59	350.6	41.96
7	1	4	1.0	297.0	7.12	393.3	29.43
7	2	23	5.9	296.5	9.13	389.5	41.29
7	3	7	1.8	291.9	5.43	374.4	32.57
7	4	14	3.6	300.1	10.11	406.7	50.69
8	3	2	0.5	304.5	7.78	416.0	14.14
8	4	11	2.8	303.3	5.53	407.2	32.91
8	5	3	0.8	304.7	11.59	444.7	60.18
9	6	1	0.3	315.0	-	445.0	-
10	5	1	0.3	313.0	-	465.0	-
10	7	2	0.5	308.5	12.02	431.0	2.83
11	8	1	0.3	327.0	-	518.0	-

Table 5d. Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam, 1985.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	20	5.8	225.7	7.33	152.8	18.98
4	0	42	12.2	240.2	11.44	186.2	29.94
4	1	40	11.7	248.1	10.84	205.7	28.30
5	0	37	10.8	267.2	8.02	263.9	27.41
5	1	78	22.7	268.1	9.37	260.6	33.82
5	2	44	12.8	267.6	9.26	261.9	29.37
6	0	8	2.3	277.8	10.87	289.4	28.50
6	1	16	4.7	274.4	9.44	281.3	36.22
6	2	21	6.1	274.4	9.39	283.3	29.42
6	3	6	1.7	275.2	15.37	291.2	49.74
7	2	7	2.0	293.6	10.08	346.6	36.16
7	3	13	3.8	288.9	7.83	340.5	29.64
7	4	5	1.5	290.4	11.06	346.0	37.07
8	3	3	0.9	295.0	6.24	363.3	26.35
8	4	1	0.3	298.0	-	360.0	-
8	5	1	0.3	288.0	-	319.0	-
10	6	1	0.3	319.0	-	444.0	-
Female							
3	0	5	1.8	218.4	11.10	143.8	24.51
4	0	17	6.3	247.8	13.01	209.7	34.50
4	1	16	5.9	250.1	8.79	221.1	25.94
5	0	39	14.3	272.6	8.25	288.6	35.30
5	1	53	19.5	275.3	11.59	301.0	38.25
5	2	24	8.8	277.0	7.21	309.0	26.78
6	0	9	3.3	277.3	8.82	306.8	29.55
6	1	32	11.8	287.9	11.49	349.1	45.16
6	2	20	7.4	287.2	8.50	345.6	31.81
6	3	8	2.9	288.8	7.17	356.4	31.58
7	0	1	0.4	288.0	-	351.0	-
7	1	1	0.4	295.0	-	388.0	-
7	2	11	4.0	297.4	13.90	402.7	63.55
7	3	9	3.3	303.3	10.27	414.6	47.55
7	4	6	2.2	297.5	6.66	399.3	28.07
8	2	1	0.4	301.0	-	477.0	-
8	3	8	2.9	308.4	8.68	468.0	52.61
8	4	4	1.5	318.0	16.06	474.0	57.15
8	5	2	0.7	309.5	0.71	449.5	57.28
9	4	1	0.4	310.0	-	452.0	-
9	5	4	1.5	316.0	6.38	471.5	29.06
10	7	1	0.4	306.0	-	401.0	-

Table 5e. Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam, 1986.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	1	0.3	221.0	-	143.0	-
4	0	183	47.3	249.2	9.58	206.4	22.93
4	1	139	35.9	250.9	9.09	208.4	22.88
5	0	12	3.1	263.1	7.05	236.9	17.18
5	1	17	4.4	269.2	8.66	252.5	27.63
5	2	11	2.8	268.1	9.06	256.6	27.50
6	1	2	0.5	281.5	16.26	270.5	36.06
6	2	15	3.9	277.9	5.46	284.7	21.89
6	3	4	1.0	275.5	3.70	293.8	13.77
7	3	2	0.5	294.0	1.41	327.5	21.92
9	4	1	0.3	307.0	-	372.0	-
Female							
3	0	1	0.3	233.0	-	170.0	-
4	0	157	52.5	257.1	8.11	236.9	22.71
4	1	90	30.1	258.2	9.22	241.8	27.84
5	0	5	1.7	273.0	7.48	282.2	17.17
5	1	13	4.3	273.5	8.91	287.9	33.24
5	2	13	4.3	273.8	11.50	283.6	45.80
6	1	2	0.7	293.0	9.90	362.0	31.11
6	2	11	3.7	292.1	6.49	344.4	35.08
6	3	5	1.7	291.8	6.65	323.2	42.39
7	2	1	0.3	310.0	-	362.0	-
8	4	1	0.3	300.0	-	383.0	-

Table 5f. Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam, 1987.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	2	0.5	218.5	0.71	138.5	2.12
4	0	119	31.4	245.3	6.73	198.0	18.04
4	1	17	4.5	243.5	8.99	197.9	23.20
5	0	46	12.1	261.9	7.81	235.7	24.23
5	1	97	25.6	263.6	8.03	244.1	26.49
5	2	74	19.5	264.0	7.36	244.4	23.26
6	1	5	1.3	272.6	6.50	255.8	20.77
6	2	10	2.6	271.9	8.32	257.6	34.70
6	3	3	0.8	275.0	3.00	279.0	17.69
7	2	2	0.5	286.5	3.54	321.5	12.02
7	3	4	1.1	282.3	19.52	290.8	58.81
Female							
3	0	1	0.3	224.0	-	161.0	-
4	0	112	34.0	251.4	8.56	224.4	25.96
4	1	13	4.0	253.7	10.77	228.3	26.91
5	0	49	14.9	267.8	8.78	272.9	24.51
5	1	81	24.6	270.8	8.16	277.4	28.78
5	2	51	15.5	273.8	8.81	288.1	27.66
6	1	7	2.1	291.1	10.09	343.9	27.56
6	2	6	1.8	284.5	7.77	314.3	28.23
6	3	6	1.8	284.8	4.96	328.3	17.41
7	2	3	0.9	302.7	9.71	377.7	29.02

Table 5g. Mean fork length and weight of alewife, by age and sex, in relation to spawning history, Mactaquac Dam, 1988.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	1	0.3	227.0	-	154.0	-
4	0	56	16.9	238.4	7.57	187.6	19.53
4	1	12	3.6	241.2	6.48	190.0	16.41
5	0	48	14.5	255.9	7.30	233.3	26.07
5	1	76	22.9	258.7	6.22	233.3	22.00
5	2	21	6.3	260.2	7.31	240.7	23.04
6	0	1	0.3	285.0	-	295.0	-
6	1	19	5.7	270.6	5.30	266.4	22.05
6	2	61	18.4	270.8	10.35	262.6	24.90
6	3	29	8.7	267.8	7.24	261.5	27.33
7	2	4	1.2	277.5	5.45	288.0	14.17
7	3	4	1.2	288.0	13.22	291.0	35.54
Female							
3	0	1	0.3	219.0	-	150.0	-
4	0	44	13.9	243.3	7.50	206.3	21.49
4	1	7	2.2	246.1	5.34	215.1	19.42
5	0	73	23.0	263.5	7.63	266.7	26.10
5	1	42	13.2	265.2	6.79	270.2	23.83
5	2	14	4.4	261.8	8.77	250.8	30.50
6	0	1	0.3	290.0	-	345.0	-
6	1	35	11.0	278.5	7.29	312.4	32.59
6	2	69	21.8	279.8	8.05	311.8	33.68
6	3	15	4.7	276.9	8.48	311.9	25.37
7	2	4	1.3	289.3	5.32	328.3	14.01
7	3	9	2.8	287.8	8.88	346.8	30.39
7	4	3	0.9	284.7	12.86	329.7	41.63

Table 6a. Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam, 1982.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	143	32.9	202.9	9.29	101.2	15.38
4	0	46	10.6	231.2	13.44	154.9	32.70
4	1	51	11.7	233.5	11.09	160.8	26.40
5	0	5	1.1	254.8	15.80	203.8	38.23
5	1	35	8.0	246.3	12.01	192.2	26.61
5	2	26	6.0	245.5	9.27	195.7	34.29
6	1	2	0.5	256.0	7.07	217.0	2.83
6	2	27	6.2	251.3	8.81	203.2	26.34
6	3	11	2.5	251.6	7.99	208.5	31.34
7	3	7	1.6	258.6	6.16	226.7	19.14
7	4	12	2.8	256.5	6.72	212.5	31.37
8	4	6	1.4	265.3	17.00	247.2	58.62
8	5	22	5.1	263.1	8.88	238.2	36.24
9	4	1	0.2	276.0	-	263.0	-
9	5	5	1.1	266.0	7.78	247.6	29.74
9	6	20	4.6	268.9	7.17	247.5	29.44
10	6	2	0.5	264.5	0.71	228.5	41.72
10	7	11	2.5	266.9	9.58	244.2	19.19
11	8	2	0.5	261.0	1.41	236.0	15.56
12	9	1	0.2	265.0	-	249.0	-
Female							
3	0	36	9.4	210.1	11.12	116.8	19.58
4	0	67	17.4	236.8	12.30	171.2	31.13
4	1	26	6.8	242.0	10.70	187.3	24.87
5	0	7	1.8	249.4	10.95	208.6	24.47
5	1	54	14.0	257.1	9.69	228.9	29.01
5	2	32	8.3	255.4	11.36	219.6	30.45
6	1	4	1.0	264.5	3.51	241.5	27.36
6	2	45	11.7	263.6	7.89	247.5	27.79
6	3	19	4.9	263.7	7.05	250.8	30.20
7	2	4	1.0	270.0	8.83	238.0	34.13
7	3	7	1.8	273.3	8.08	271.3	25.15
7	4	14	3.6	268.1	12.71	256.3	43.50
8	3	1	0.3	275.0	-	327.0	-
8	4	10	2.6	279.1	9.87	302.4	49.94
8	5	15	3.9	275.3	9.26	277.3	35.44
9	4	1	0.3	265.0	-	282.0	-
9	5	14	3.6	278.8	11.09	297.4	39.25
9	6	15	3.9	280.1	12.01	306.7	48.27
10	6	1	0.3	298.0	-	414.0	-
10	7	8	2.1	285.6	6.44	308.3	36.59
11	7	3	0.8	287.3	5.51	287.3	36.90
11	8	1	0.3	295.0	-	335.0	-
12	8	1	0.3	302.0	-	377.0	-

Table 6b. Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam, 1983.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
2	0	1	0.3	154.0	-	37.0	-
3	0	137	38.4	201.0	10.70	97.2	18.74
4	0	50	14.0	218.1	15.17	132.4	34.16
4	1	70	19.6	222.1	10.77	138.3	23.13
5	0	12	3.4	236.4	10.09	175.9	35.94
5	1	30	8.4	237.5	12.86	173.7	29.94
5	2	17	4.8	235.1	9.13	174.9	26.00
6	1	1	0.3	271.0	-	274.0	-
6	2	6	1.7	251.2	11.50	212.5	32.61
6	3	5	1.4	242.8	11.30	203.4	31.75
7	3	6	1.7	256.7	12.55	222.3	36.24
7	4	6	1.7	256.3	6.98	221.0	21.82
8	4	2	0.6	261.5	3.54	233.0	5.66
8	5	4	1.1	263.0	5.35	243.5	5.97
9	6	2	0.6	268.5	21.92	252.5	65.76
10	6	3	0.8	265.0	1.73	247.0	10.39
10	7	4	1.1	267.0	9.63	247.3	19.02
12	8	1	0.3	288.0	-	278.0	-
Female							
1	0	2	0.7	98.0	2.83	10.5	0.71
2	0	1	0.4	144.0	-	29.0	-
3	0	49	17.3	211.1	13.50	114.1	21.95
4	0	67	23.6	231.1	12.04	159.5	28.20
4	1	40	14.1	232.9	8.20	166.4	21.16
5	0	21	7.4	246.4	12.82	197.4	34.61
5	1	46	16.2	243.1	8.95	192.3	28.92
5	2	9	3.2	245.6	13.95	198.4	36.37
6	0	2	0.7	242.0	1.41	184.0	-
6	1	4	1.4	246.8	4.72	208.3	15.71
6	2	8	2.8	262.5	7.58	242.0	23.00
6	3	9	3.2	263.3	7.45	241.6	27.14
7	3	8	2.8	267.8	8.17	263.5	24.10
7	4	1	0.4	276.0	-	246.0	-
8	3	1	0.4	280.0	-	313.0	-
8	4	3	1.1	267.0	9.64	281.3	27.10
8	5	4	1.4	277.3	2.99	310.0	17.98
9	6	5	1.8	275.0	8.34	282.4	51.31
11	7	2	0.7	297.5	12.02	344.0	89.10
11	8	1	0.4	272.0	-	240.0	-
12	8	1	0.4	293.0	-	321.0	-

Table 6c. Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam, 1984.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	88	25.0	196.8	8.99	90.6	11.74
4	0	88	25.0	220.5	7.38	129.6	16.70
4	1	108	30.7	217.9	6.96	126.1	17.17
5	0	4	1.1	241.8	11.81	182.0	26.94
5	1	20	5.7	232.0	8.45	161.6	21.35
5	2	22	6.3	228.7	10.50	150.1	22.78
6	2	6	1.7	244.7	15.27	193.2	31.90
6	3	7	2.0	242.7	10.01	180.3	23.95
7	3	2	0.6	253.5	0.71	207.5	31.82
7	4	4	1.1	247.5	10.47	190.0	29.02
9	4	1	0.3	268.0	-	241.0	-
9	6	1	0.3	265.0	-	264.0	-
10	6	1	0.3	260.0	-	303.0	-
Female							
3	0	40	11.9	201.2	9.52	96.2	13.37
4	0	122	36.3	227.1	7.67	145.3	17.77
4	1	85	25.3	226.2	7.86	143.7	19.03
4	2	1	0.3	228.0	-	162.0	-
5	0	12	3.6	238.3	11.00	172.1	31.84
5	1	28	8.3	238.3	7.56	179.6	21.65
5	2	21	6.3	237.1	9.75	176.0	29.59
6	1	2	0.6	259.0	22.63	225.5	77.07
6	2	6	1.8	249.0	6.51	208.5	22.16
6	3	5	1.5	255.0	16.17	218.2	51.50
7	3	4	1.2	264.3	5.74	233.8	5.91
7	4	1	0.3	260.0	-	257.0	-
8	3	1	0.3	302.0	-	328.0	-
8	4	1	0.3	265.0	-	290.0	-
8	5	3	0.9	261.0	19.00	246.3	36.86
9	4	2	0.6	285.0	4.24	300.5	45.96
10	7	1	0.3	280.0	-	308.0	-
12	9	1	0.3	302.0	-	330.0	-

Table 6d. Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam, 1985.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	332	63.2	191.8	8.48	81.3	12.44
4	0	38	7.2	218.8	12.29	129.7	25.33
4	1	92	17.5	216.6	10.26	126.3	21.26
5	0	3	0.6	226.7	12.70	142.7	18.01
5	1	29	5.5	229.7	8.77	150.2	24.84
5	2	18	3.4	230.1	5.65	158.5	14.63
6	0	1	0.2	252.0	-	212.0	-
6	1	3	0.6	235.0	5.00	164.7	12.50
6	2	3	0.6	242.3	9.07	191.7	31.82
6	3	4	0.8	235.5	13.18	170.8	31.02
7	3	2	0.4	261.5	13.44	224.0	24.04
Female							
3	0	179	41.4	198.8	7.62	90.0	11.18
4	0	94	21.8	227.2	10.00	146.2	22.69
4	1	84	19.4	225.9	9.66	145.2	23.67
5	0	12	2.8	240.9	9.28	177.8	28.46
5	1	33	7.6	235.2	12.00	164.2	31.88
5	2	17	3.9	237.9	9.18	171.3	26.00
6	1	2	0.5	247.5	10.61	205.0	31.11
6	2	6	1.4	249.2	12.42	203.7	37.23
6	3	3	0.7	252.7	1.53	227.0	16.46
8	4	1	0.2	280.0	-	313.0	-
8	5	1	0.2	280.0	-	313.0	-

Table 6e. Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam, 1986.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	85	35.1	193.0	10.09	84.2	13.24
4	0	70	28.9	214.9	9.97	118.7	21.99
4	1	68	28.1	211.4	9.38	113.3	17.52
5	1	6	2.5	229.8	9.62	147.2	28.65
5	2	9	3.7	234.4	16.55	159.1	42.11
6	2	2	0.8	246.5	9.19	172.0	33.94
6	3	2	0.8	246.0	0.00	176.0	5.66
Female							
3	0	18	7.0	198.3	8.31	91.7	11.53
4	0	144	56.3	221.9	8.86	132.3	18.80
4	1	79	30.9	220.2	8.02	128.5	17.06
5	0	1	0.4	237.0	-	172.0	-
5	1	5	2.0	245.4	8.20	184.2	17.34
5	2	7	2.7	239.9	17.08	177.7	44.31
6	2	2	0.8	247.5	17.68	199.5	31.82

Table 6f. Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam, 1987.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	41	11.6	192.6	9.26	82.1	16.40
4	0	120	34.1	210.7	9.38	109.3	18.25
4	1	129	36.6	210.1	9.73	110.2	16.87
5	0	3	0.9	236.0	7.21	155.0	14.73
5	1	16	4.5	223.6	8.25	136.5	17.77
5	2	38	10.8	220.2	10.39	127.5	21.33
6	2	3	0.9	243.0	12.17	180.3	26.27
7	3	1	0.3	256.0	-	227.0	-
7	4	1	0.3	247.0	-	183.0	-
Female							
3	0	16	3.9	196.6	8.57	87.4	12.86
4	0	172	42.3	215.3	8.60	119.6	16.23
4	1	106	26.0	216.0	8.15	122.0	15.40
5	0	30	7.4	238.7	9.89	171.1	23.51
5	1	33	8.1	235.1	8.52	163.7	21.69
5	2	44	10.8	236.0	8.17	164.1	21.93
6	1	1	0.2	262.0	-	203.0	-
6	2	4	1.0	248.3	8.62	193.5	31.64
7	3	1	0.2	255.0	-	216.0	-

Table 6g. Mean fork length and weight of blueback herring, by age and sex, in relation to spawning history, Mactaquac Dam, 1988.

Age (yr)	No. previous spawnings	N	%	Mean length (mm)	SD	Mean weight (g)	SD
Male							
3	0	121	22.9	187.8	6.22	75.8	9.14
4	0	132	25.0	207.8	9.32	107.6	18.41
4	1	110	20.8	207.8	6.48	108.7	14.03
5	0	6	1.1	221.8	10.53	131.3	15.85
5	1	49	9.3	220.0	7.16	129.9	14.94
5	2	95	18.0	216.5	9.42	123.3	18.98
6	0	1	0.2	249.0	-	199.0	-
6	2	6	1.1	239.8	16.44	177.8	46.47
6	3	9	1.7	226.6	11.10	142.4	28.48
Female							
3	0	20	4.6	193.5	6.30	83.9	8.66
4	0	156	36.2	215.0	9.59	123.0	20.59
4	1	55	12.8	216.2	9.26	124.2	19.88
5	0	39	9.0	231.6	7.47	162.6	22.09
5	1	71	16.5	230.7	9.09	159.6	27.56
5	2	61	14.2	228.8	10.09	153.9	22.38
6	1	5	1.2	248.6	6.80	213.6	20.01
6	2	8	1.9	241.1	8.90	180.9	27.92
6	3	16	3.7	240.9	8.92	178.6	25.12

Table 7. Age at first spawning (maturity) of alewife, by sex and year, Mactaquac Dam, 1982 - 1988.

Age at first spawning (yr)	1982				1983				1984				1985			
	Male		Female		Male		Female		Male		Female		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
3	56	17.6	48	15.2	93	24.5	98	26.4	89	25.7	95	24.5	118	34.4	63	23.3
4	114	35.9	88	27.9	171	45.1	135	36.4	149	42.9	117	30.2	154	44.9	104	38.5
5	139	43.7	162	51.4	112	29.6	132	36.4	101	29.1	158	40.7	63	18.4	91	33.7
6	9	2.8	17	5.4	3	0.8	3	0.8	8	2.3	17	4.4	8	2.3	11	4.1
7	0	-	0	-	0	-	0	-	0	-	0	-	0	-	1	0.4
Total	318		315		379		371		347		387		343		270	
Mean age	4.3		4.5		4.1		4.1		4.1		4.2		3.9		4.2	
SD	0.79		0.82		0.76		0.80		0.80		0.88		0.78		0.86	
<hr/>																
1986																
Age at first spawning (yr)	Male		Female													
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
	155	40.1	109	36.5	96	25.3	71	21.6	63	19.0	39	12.5				
3	217	56.1	182	60.9	232	61.2	200	60.8	197	59.3	161	51.4				
4	15	3.9	8	2.7	51	13.5	58	17.6	71	21.4	112	35.8				
5	0	-	0	-	0	-	0	-	1	0.3	1	0.3				
6	0	-	0	-	0	-	0	-	0	-	0	-				
7																
Total	387		299		379		329		332		313					
Mean age	3.6		3.7		3.9		4.0		4.0		4.2					
SD	0.56		0.53		0.61		0.63		0.65		0.66					
<hr/>																

Table 8. Age at first spawning (maturity) of blueback herring, by sex and year, Mactaquac Dam, 1982 - 1988.

Age at first spawning (yr)	1982				1983				1984				1985			
	Male		Female		Male		Female		Male		Female		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
3	299	68.7	166	43.1	247	69.4	121	43.1	232	65.9	160	47.6	445	85.1	275	65.2
4	128	29.4	202	52.5	96	27.0	132	47.0	115	32.7	159	47.3	71	13.6	133	31.5
5	8	1.8	17	4.4	13	3.6	26	9.2	5	1.4	17	5.1	6	1.1	14	3.3
6	0	-	0	-	0	-	2	0.7	0	-	0	-	1	0.2	0	-
Total	435		385		356		281		352		336		523		422	
Mean age	3.3		3.6		3.3		3.7		3.4		3.6		3.2		3.4	
SD	0.51		0.57		0.55		0.67		0.51		0.59		0.42		0.55	

Age at first spawning (yr)	1986				1987				1988			
	Male		Female		Male		Female		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%
3	164	67.8	101	39.9	209	59.4	168	41.3	335	63.3	152	35.4
4	78	32.2	151	59.7	140	39.8	209	51.3	187	35.4	234	54.4
5	0	-	1	0.4	3	0.9	30	7.4	6	1.1	44	10.2
6	0	-	0	-	0	-	0	-	1	0.2	0	-
Total	242		253		352		407		529		430	
Mean age	3.3		3.6		3.4		3.7		3.4		3.8	
SD	0.47		0.50		0.51		0.61		0.52		0.63	

Table 9. Length-frequency distribution of mature alewife, by sex and year, Mactaquac Dam, 1982 - 1988.

Fork length (mm)	1982				1983				1984				1985			
	Male		Female		Male		Female		Male		Female		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
200-209	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-
210-219	0	-	0	-	1	0.2	1	0.3	0	-	2	0.5	4	1.1	0	-
220-229	0	-	0	-	6	1.5	2	0.5	6	1.6	2	0.5	20	5.7	3	1.1
230-239	6	1.9	0	-	3	0.7	1	0.3	19	5.2	5	1.2	26	7.4	6	2.2
240-249	21	6.5	2	0.6	27	6.5	6	1.5	62	17.0	23	5.7	19	5.4	10	3.6
250-259	41	12.7	15	4.6	57	13.8	24	6.0	87	23.8	61	15.1	60	17.1	18	6.5
260-269	90	27.8	44	13.6	73	17.7	37	9.3	67	18.4	79	19.6	76	21.7	37	13.4
270-279	83	25.6	64	19.8	117	28.3	62	15.5	60	16.4	68	16.8	80	22.8	55	19.9
280-289	47	14.5	71	21.9	84	20.3	111	27.8	40	11.0	67	16.6	37	10.5	65	23.5
290-299	25	7.7	52	16.1	35	8.5	84	21.1	20	5.5	52	12.9	20	5.7	38	13.7
300-309	11	3.4	47	14.5	8	1.9	46	11.5	3	0.8	32	7.9	7	2.0	25	9.0
310-319	0	-	24	7.4	2	0.5	18	4.5	1	0.3	11	2.7	2	0.6	13	4.7
320-329	0	-	3	0.9	0	-	6	1.5	0	-	1	0.3	0	-	6	2.2
330-339	0	-	1	0.3	0	-	1	0.3	0	-	0	-	0	-	1	0.4
340-349	0	-	1	0.3	0	-	0	-	0	-	0	-	0	-	0	-
Total	324		324		413		399		365		404		351		277	

Fork length (mm)	1986				1987				1988			
	Male		Female		Male		Female		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%
200-209	0	-	0	-	0	-	0	-	0	-	0	-
210-219	1	0.3	0	-	2	0.5	0	-	1	0.3	1	0.3
220-229	10	2.6	1	0.3	3	0.8	1	0.3	9	2.6	2	0.6
230-239	29	7.4	4	1.3	31	8.0	11	3.3	21	6.1	10	3.0
240-249	91	23.2	31	10.2	73	18.9	39	11.5	50	14.4	28	8.5
250-259	153	39.0	107	35.2	90	23.3	67	19.8	79	22.8	49	14.8
260-269	66	16.8	98	32.2	123	31.9	83	24.6	106	30.6	71	21.5
270-279	28	7.1	33	10.9	55	14.3	79	23.4	61	17.6	87	26.3
280-289	9	2.3	12	4.0	8	2.1	46	13.6	16	4.6	62	18.7
290-299	4	1.0	13	4.3	0	-	8	2.4	2	0.6	20	6.0
300-309	1	0.3	4	1.3	1	0.3	2	0.6	1	0.3	1	0.3
310-319	0	-	1	0.3	0	-	2	0.6	0	-	0	-
320-329	0	-	0	-	0	-	0	-	1	0.3	0	-
330-339	0	-	0	-	0	-	0	-	0	-	0	-
340-349	0	-	0	-	0	-	0	-	0	-	0	-
Total	392		304		386		338		347		331	

Table 10. Length-frequency distribution of mature blueback herring, by sex and year, Mactaquac Dam, 1982 - 1988.

Fork length (mm)	1982				1983				1984				1985			
	Male		Female		Male		Female		Male		Female		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
160-169	0	-	0	-	0	-	0	-	0	-	0	-	1	0.2	0	-
170-179	0	-	0	-	1	0.3	0	-	1	0.3	0	-	13	2.5	0	-
180-189	5	1.1	1	0.3	8	2.1	1	0.3	19	5.4	3	0.9	110	20.9	13	3.0
190-199	54	11.9	2	0.5	66	17.4	8	2.7	35	9.9	13	3.8	160	30.4	89	20.8
200-209	53	11.7	15	3.8	65	17.1	25	8.3	39	11.0	22	6.4	57	10.8	63	14.8
210-219	39	8.6	19	4.8	52	13.7	15	5.0	101	28.5	34	9.9	75	14.3	43	10.1
220-229	35	7.7	14	3.5	61	16.1	43	14.2	92	26.0	107	31.1	50	9.5	83	19.4
230-239	41	9.0	40	10.1	43	11.3	60	19.9	43	12.2	99	28.8	43	8.2	79	18.5
240-249	73	16.0	50	12.7	37	9.7	63	20.9	8	2.3	31	9.0	12	2.3	38	8.9
250-259	70	15.4	71	18.0	19	5.0	33	10.9	12	3.4	14	4.1	4	0.8	14	3.3
260-269	53	11.7	86	21.8	15	4.0	24	8.0	3	0.9	10	2.9	0	-	2	0.5
270-279	25	5.5	50	12.7	10	2.6	13	4.3	1	0.3	2	0.6	1	0.2	1	0.2
280-289	5	1.1	34	8.6	3	0.8	10	3.3	0	-	7	2.0	0	-	2	0.5
290-299	2	0.4	11	2.8	0	-	5	1.7	0	-	0	-	0	-	0	-
300-309	0	-	2	0.5	0	-	1	0.3	0	-	2	0.6	0	-	0	-
310-319	0	-	0	-	0	-	1	0.3	0	-	0	-	0	-	0	-
Total	455		395		380		302		354		344		526		427	

Fork length (mm)	1986				1987				1988			
	Male		Female		Male		Female		Male		Female	
	N	%	N	%	N	%	N	%	N	%	N	%
160-169	0	-	0	-	0	-	0	-	0	-	0	-
170-179	8	3.3	0	-	1	0.3	0	-	7	1.2	0	-
180-189	22	9.0	2	0.8	19	5.4	4	1.0	66	11.7	6	1.3
190-199	32	13.1	7	2.8	46	13.0	10	2.4	78	13.8	19	4.2
200-209	76	31.2	16	6.3	91	25.6	72	17.1	141	25.0	54	11.9
210-219	53	21.7	93	36.5	111	31.3	132	31.4	160	28.4	102	22.5
220-229	31	12.7	81	31.8	61	17.2	83	19.7	77	13.7	120	26.5
230-239	14	5.7	42	16.5	21	5.9	71	16.9	27	4.8	101	22.3
240-249	6	2.5	10	3.9	3	0.9	36	8.6	7	1.2	37	8.2
250-259	1	0.4	2	0.8	2	0.6	10	2.4	0	-	12	2.7
260-269	0	-	1	0.4	0	-	3	0.7	1	0.2	2	0.4
270-279	1	0.4	1	0.4	0	-	0	-	0	-	0	-
280-289	0	-	0	-	0	-	0	-	0	-	0	-
290-299	0	-	0	-	0	-	0	-	0	-	0	-
300-309	0	-	0	-	0	-	0	-	0	-	0	-
310-319	0	-	0	-	0	-	0	-	0	-	0	-
Total	244		255		355		421		564		453	

Table 11. Mean weight (g) of alewife by sex, age and year, Mactaquac Dam, 1982 - 1988.

Table 12. Mean weight (g) of blueback herring by sex, age and year, Mactaquac Dam, 1982 - 1988.

Year	Age	Male				Female			
		N	Mean	Range	SD	N	Mean	Range	SD
1982	3	143	101.2	73-160	15.38	36	116.8	80-160	19.58
	4	97	158.0	92-244	29.55	93	175.7	107-286	30.26
	5	66	194.4	129-311	30.38	93	224.2	148-305	29.58
	6	40	205.3	147-274	26.98	68	248.1	192-322	28.12
	7	19	217.7	170-291	27.80	25	257.6	188-333	38.06
	8	28	240.1	173-337	40.89	26	288.9	220-386	42.57
	9	26	248.1	199-309	28.45	30	301.5	232-416	43.02
	10	13	241.8	199-277	22.06	9	320.0	269-414	49.13
	11	2	236.0	225-247	15.56	4	299.3	256-335	38.41
	12	1	249.0	249-249	-	1	377.0	377-377	-
1983	3	136	97.2	65-172	18.80	49	114.1	85-169	21.95
	4	120	135.9	78-245	28.27	107	162.1	79-242	25.91
	5	60	176.1	131-268	31.80	79	200.7	130-413	44.74
	6	13	218.9	161-279	38.12	25	244.2	184-407	54.02
	7	12	221.7	170-269	28.53	9	261.6	231-296	23.29
	8	6	240.0	229-249	7.56	8	299.6	253-327	24.06
	9	2	252.5	206-299	65.76	6	313.7	223-470	89.29
	10	7	247.1	229-273	14.73	0	-	-	-
	11	0	-	-	-	3	309.3	240-407	87.03
	12	1	278.0	278-278	-	1	321.0	321-321	-
1984	3	88	90.6	66-125	11.74	40	96.2	76-123	13.37
	4	196	127.7	82-204	17.01	208	144.7	99-206	18.27
	5	46	157.9	113-218	23.89	61	176.9	132-262	26.40
	6	13	186.2	155-254	27.48	13	214.9	171-293	40.34
	7	6	195.8	164-230	28.10	5	238.4	229-257	11.59
	8	0	-	-	-	5	271.4	218-328	45.14
	9	2	252.5	241-264	16.26	2	300.5	268-333	45.96
	10	1	303.0	303-303	-	1	308.0	308-308	-
	11	0	-	-	-	0	-	-	-
	12	0	-	-	-	1	330.0	330-330	-
1985	3	330	81.4	48-149	12.47	170	90.3	65-122	10.99
	4	130	127.3	78-215	22.48	177	145.9	87-203	22.97
	5	50	152.8	104-208	21.50	62	168.8	104-250	29.76
	6	11	178.6	125-212	27.71	11	210.3	161-259	30.98
	7	2	224.0	207-241	24.04	0	-	-	-
	8	0	-	-	-	2	313.0	313-313	0.00
1986	3	85	84.2	58-125	13.24	15	92.9	73-112	12.00
	4	138	116.1	81-196	20.03	223	130.9	88-199	18.25
	5	15	154.3	114-258	36.65	13	179.8	120-255	33.13
	6	4	174.0	148-196	20.00	2	199.5	177-222	31.82
1987	3	41	82.1	64-138	16.40	16	87.4	71-110	12.86
	4	249	109.8	73-170	17.52	278	120.5	84-166	15.94
	5	57	131.5	96-189	20.99	107	165.9	111-220	22.34
	6	3	180.3	160-210	26.27	5	195.4	171-239	27.73
	7	2	205.0	183-227	31.11	1	216.0	216-216	-
1988	3	121	75.8	55-98	9.14	20	83.9	72-100	8.66
	4	242	108.1	81-162	16.54	210	123.4	85-185	20.39
	5	150	125.8	88-164	17.85	171	158.2	104-258	24.71
	6	16	159.3	103-251	39.57	29	185.2	137-235	27.65

Table 13. Mean fork length (mm) of alewife, by sample date and year, Mactaquac Dam, 1982 - 1988 (sexes combined).

Table 14. Mean fork length (mm) of blueback herring, by sample date and year, Mactaquac Dam, 1982 - 1988 (sexes combined).

1982				1983				1984			
Date	N	Mean	SD	Date	N	Mean	SD	Date	N	Mean	SD
28 May	3	271.7	10.40	26 May	10	285.4	12.49	31 May	7	242.9	19.40
1 Jun	18	265.3	15.40	6 Jun	34	237.1	13.39	8 Jun	53	234.5	16.29
4 Jun	17	255.0	12.90	9 Jun	32	253.2	18.35	11 Jun	39	226.2	13.27
8 Jun	79	261.8	17.48	13 Jun	50	239.5	11.37	14 Jun	83	227.3	13.52
11 Jun	77	261.3	15.00	16 Jun	80	243.2	18.24	18 Jun	93	224.8	12.63
15 Jun	95	252.9	20.50	20 Jun	92	228.2	20.67	21 Jun	93	223.3	20.09
18 Jun	99	250.2	20.19	23 Jun	100	237.1	20.57	25 Jun	90	215.3	20.38
22 Jun	96	253.8	18.38	27 Jun	99	220.5	23.56	29 Jun	95	223.0	21.61
25 Jun	90	241.1	22.51	30 Jun	93	214.7	24.13	3 Jul	77	214.5	14.43
29 Jun	93	232.6	24.99	4 Jul	96	205.7	24.27	6 Jul	66	222.9	17.35
2 Jul	83	224.6	24.54								
6 Jul	100	210.7	19.39								
1985				1986				1987			
Date	N	Mean	SD	Date	N	Mean	SD	Date	N	Mean	SD
27 May	2	247.5	2.12	30 May	2	253.0	28.28	25 May	7	229.0	12.48
30 May	15	245.2	10.15	2 Jun	20	229.3	17.26	28 May	18	228.2	15.70
3 Jun	71	231.6	10.81	6 Jun	64	222.7	9.54	1 Jun	33	219.9	9.66
6 Jun	96	224.8	17.08	9 Jun	77	214.2	16.08	4 Jun	89	223.8	13.56
10 Jun	92	216.7	18.68	12 Jun	97	215.1	14.80	8 Jun	64	221.2	11.93
13 Jun	98	214.0	19.55	16 Jun	93	208.6	14.65	11 Jun	81	217.2	13.05
17 Jun	100	206.2	18.49	20 Jun	50	211.5	13.93	15 Jun	88	213.8	14.19
21 Jun	96	197.8	15.75	23 Jun	98	211.9	14.50	18 Jun	97	217.7	12.04
24 Jun	100	204.1	17.98					22 Jun	100	214.0	16.53
27 Jun	100	205.1	17.46					25 Jun	99	209.8	14.03
2 Jul	94	200.8	11.42					29 Jun	100	212.0	14.56
5 Jul	100	196.0	12.04								
1988											
				Date	N	Mean	SD				
				24 May	3	234.7	11.01				
				27 May	9	231.1	14.26				
				30 May	47	227.2	10.80				
				2 Jun	88	224.7	10.57				
				6 Jun	96	219.5	13.85				
				9 Jun	93	221.9	12.76				
				13 Jun	84	213.7	14.48				
				17 Jun	97	218.2	14.54				
				20 Jun	100	212.6	15.49				
				23 Jun	100	206.0	15.48				
				27 Jun	100	208.0	16.73				
				30 Jun	100	206.4	15.05				
				4 Jul	100	205.0	14.82				

Table 15a. Length-frequency distribution of alewife, by age, Mactaquac Dam, 1982
(sexes combined).

Fork length (mm)	Age										Total
	3	4	5	6	7	8	9	10	11		
230-239		5	1							6	
240-249	1	19	3							23	
250-259		35	20	1						56	
260-269		27	97	10						134	
270-279		7	107	25	1	2				142	
280-289			65	35	8	4	1	2	1	116	
290-299			11	25	14	19	5			74	
300-309			2	8	15	19	11			55	
310-319					4	15	2	1		22	
320-329						1		2		3	
330-339						1				1	
340-349					1					1	
Total	1	93	306	104	43	61	19	5	1	633	
Percent	0.2	14.7	48.3	16.4	6.8	9.6	3.0	0.8	0.2		

26

Table 15b. Length-frequency distribution of alewife, by age, Mactaquac Dam, 1983 (sexes combined).

Fork length (mm)	Age											Total
	3	4	5	6	7	8	9	10	11	12		
210-219	1	1										2
220-229	1	4	2									7
230-239	1	3										4
240-249		26	6									32
250-259		44	35	2								81
260-269		18	71	14	2			2				107
270-279		6	98	59	3	2			1			169
280-289			65	85	16	3	2					171
290-299			29	54	17	5					1	106
300-309			7	25	9	4	3					48
310-319			1	7	4	4	3					19
320-329			1	2								3
330-339						1						1
Total	3	102	315	248	51	19	10	1	0	1		750
Percent	0.4	13.6	42.0	33.1	6.8	2.5	1.3	0.1	0.0	0.1		

Table 15c. Length-frequency distribution of alewife, by age, Mactaquac Dam, 1984 (sexes combined).

Fork length (mm)	Age											Total
	2	3	4	5	6	7	8	9	10	11		
150-159	1											1
160-169												0
170-179												0
180-189												0
190-199												0
200-209												0
210-219	1	1										2
220-229	1	6	1									8
230-239		22	1									23
240-249		65	18									83
250-259		45	96	3								144
260-269		15	112	14								141
270-279		2	68	40	9	1						120
280-289			29	55	18	1						103
290-299			8	24	25	8						65
300-309			2	2	18	8	2	1	1			33
310-319					4	4	1	2				11
320-329									1			1
Total	1	2	156	335	138	74	22	3	3	1		735
Percent	0.1	0.3	21.2	45.6	18.8	10.1	3.0	0.4	0.4	0.1		

Table 15d. Length-frequency distribution of alewife, by age, Mactaquac Dam, 1985
 (sexes combined).

Fork length (mm)	Age										Total
	2	3	4	5	6	7	8	9	10		
140-149	1									1	
150-159										0	
160-169	1									1	
170-179	1									1	
180-189										0	
190-199										0	
200-209	3									3	
210-219	4									4	
220-229	12	11								23	
230-239	7	25								32	
240-249	26	2	1							29	
250-259	42	32	3							77	
260-269	11	86	14	1						112	
270-279		103	28	3						134	
280-289		43	42	13	2					100	
290-299		7	22	18	5					52	
300-309		2	10	11	6			1	30		
310-319				6	4	3	1		14		
320-329					1	2	2		5		
330-339						1			1		
Total	3	26	115	275	120	53	20	5	2	619	
Percent	0.5	4.2	18.6	44.4	19.4	8.6	3.2	0.8	0.3		

Table 15e. Length-frequency distribution of alewife, by age, Mactaquac Dam, 1986
(sexes combined).

Fork length (mm)	Age									Total
	2	3	4	5	6	7	8	9		
150-159	1								1	
160-169									0	
170-179									0	
180-189									0	
190-199									0	
200-209									0	
210-219	1	1							2	
220-229	1	10							11	
230-239	1	31							32	
240-249		121							121	
250-259		248	8						256	
260-269		135	26	1					162	
270-279		22	24	15					61	
280-289		1	12	8					21	
290-299			1	12	2				15	
300-309				3			1	1	5	
310-319					1				1	
Total	1	3	569	71	39	3	1	1	688	
Percent	0.1	0.4	82.7	10.3	5.7	0.4	0.1	0.1		

Table 15f. Length-frequency distribution of alewife, by age,
Mactaquac Dam, 1987 (sexes combined).

Fork length (mm)	Age					Total
	3	4	5	6	7	
210-219	2					2
220-229	1	3				4
230-239		41	1			42
240-249		103	9			112
250-259		89	66	1		156
260-269		23	174	3	1	201
270-279		2	113	13	1	129
280-289			33	14	3	50
290-299			2	5	1	8
300-309					2	2
310-319				1	1	2
Total	3	261	398	37	9	708
Percent	0.4	36.9	56.2	5.2	1.3	

Table 15g. Length-frequency distribution of alewife, by age,
Mactaquac Dam, 1988 (sexes combined).

Fork length (mm)	Age					Total
	3	4	5	6	7	
210-219	1	1				2
220-229	1	10				11
230-239		31				31
240-249	64	13	1			78
250-259	13	106	6			125
260-269		115	54			169
270-279		38	94	5		137
280-289		2	62	10		74
290-299			11	8		19
300-309			1	1		2
310-319				1		0
320-329				1		1
Total	2	119	274	230	24	649
Percent	0.3	18.3	42.2	35.4	3.7	

Table 16a. Length-frequency distribution of blueback herring, by age, Mactaquac Dam, 1982
(sexes combined).

Fork length (mm)	Age										Total
	3	4	5	6	7	8	9	10	11	12	
180-189	6										6
190-199	56										56
200-209	64	4									68
210-219	38	20									58
220-229	11	32	5								48
230-239	4	58	14	3							79
240-249	52	46	13	3	2						116
250-259	21	49	36	12	7	5	3				133
260-269	3	38	42	15	16	12	5	2	1		134
270-279		5	14	7	18	21	5				70
280-289		2		7	7	14	6	2			38
290-299					4	3	3	2			12
300-309						1			1		2
Total	179	190	159	108	44	54	56	22	6	2	820
Percent	21.8	23.2	19.4	13.2	5.4	6.6	6.8	2.7	0.7	0.2	

Table 16b. Length-frequency distribution of blueback herring, by age, Mactaquac Dam, 1983 (sexes combined).

Fork length (mm)	Age												Total
	1	2	3	4	5	6	7	8	9	10	11	12	
90-99	1												1
100-109	1												1
110-119													0
120-129													0
130-139													0
140-149		1											1
150-159		1											1
160-169													0
170-179			1										1
180-189			9										9
190-199		63	9										72
200-209		71	14										85
210-219		20	42	2									64
220-229		12	65	22	1								100
230-239		8	63	29			1						101
240-249		2	30	48	12	1							93
250-259			3	27	7	7		2		1		1	48
260-269				1	5	12	7	4	2	2	4		35
270-279					1	2	4	6	1	2	1		17
280-289					1	1	1	2	3		1		10
290-299												1	1
300-309											1		1
Total	2	2	186	227	135	35	21	14	7	7	3	2	641
Percent	0.3	0.3	29.0	35.4	21.1	5.5	3.3	2.2	1.1	1.1	0.5	0.3	

Table 16c. Length-frequency distribution of blueback herring, by age, Mactaquac Dam, 1984
 (sexes combined).

Fork length (mm)	Age											Total
	3	4	5	6	7	8	9	10	11	12		
170-179	1											1
180-189	22											22
190-199	47	1										48
200-209	44	15	1									60
210-219	12	117	6									135
220-229	2	178	18									198
230-239	85	48	8	1								142
240-249	6	24	5	1	1							37
250-259	2	9	9	5								25
260-269		1	1	4	2	2	1					11
270-279			2									2
280-289			1			1	2	1				5
290-299												0
300-309						1				1		2
Total	128	404	107	26	11	5	4	2	0	1		688
Percent	18.6	58.7	15.6	3.8	1.6	0.7	0.6	0.3	0.0	0.1		

Table 16d. Length-frequency distribution of blueback herring, by age, Mactaquac Dam, 1985 (sexes combined).

Fork length (mm)	Age						Total
	3	4	5	6	7	8	
160-169	1						1
170-179	14						14
180-189	123	1					124
190-199	248	7					255
200-209	95	27					122
210-219	30	80	9	1			120
220-229	1	102	29				132
230-239		73	44	5			122
240-249		16	21	9			46
250-259		2	8	6	1		17
260-269			1				1
270-279				1	1		2
280-289						2	2
Total	512	308	112	22	2	2	958
Percent	53.4	32.2	11.7	2.3	0.2	0.2	

Table 16e. Length-frequency distribution of blueback herring, by age, Mactaquac Dam, 1986 (sexes combined).

Fork length (mm)	Age				Total
	3	4	5	6	
170-179	8				8
180-189	25				25
190-199	35	5			40
200-209	28	64			92
210-219	6	135	2		143
220-229	1	105	6		112
230-239		46	9	1	56
240-249		6	7	3	16
250-259			2	1	3
260-269				1	1
270-279			2		2
Total	103	361	28	6	498
Percent	20.7	72.5	5.6	1.2	

Table 16f. Length-frequency distribution of blueback herring, by age, Mactaquac Dam, 1987 (sexes combined).

Fork length (mm)	Age					Total
	3	4	5	6	7	
170-179	1					1
180-189	18	5				23
190-199	26	29	1			56
200-209	8	147	6			161
210-219	3	216	21			240
220-229	1	111	27			139
230-239		17	70	2		89
240-249		2	31	3	1	37
250-259			8	1	2	11
260-269				2		2
Total	57	527	164	8	3	759
Percent	7.5	69.4	21.6	1.1	0.4	

Table 16g. Length-frequency distribution of blueback herring, by age, Mactaquac Dam, 1988 (sexes combined).

Fork length (mm)	Age				Total
	3	4	5	6	
170-179	7				7
180-189	72				72
190-199	54	40	2		96
200-209	8	158	25		191
210-219		169	74	4	247
220-229		67	108	2	177
230-239		18	85	15	118
240-249		1	23	15	39
250-259			3	8	11
260-269			1	1	2
Total	141	453	321	45	960
Percent	14.7	47.2	33.4	4.7	

Table 17a. Length-frequency distribution of gaspereau, by age, Mactaquac Dam, 1982
(species and sexes combined).

Fork length (mm)	Age											Total
	3	4	5	6	7	8	9	10	11	12		
180-189	6											6
190-199	56											56
200-209	64	4										68
210-219	38	20										58
220-229	11	32	5									48
230-239	4	63	15	3								85
240-249	1	71	49	13	3	2						139
250-259		56	69	37	12	7	5	3				189
260-269		30	135	52	15	16	12	5	2	1		268
270-279		7	112	39	8	20	21	5				212
280-289			67	35	15	11	15	8	3			154
290-299			11	25	14	23	8	3	2			86
300-309			2	8	15	19	12		1			57
310-319					4	15	2	1				22
320-329						1		2				3
330-339						1						1
340-349					1							1
Total	180	283	465	212	87	115	75	27	7	2		1453
Percent	12.4	19.5	32.0	14.6	6.0	7.9	5.2	1.9	0.5	0.1		

Table 17b. Length-frequency distribution of gaspereau, by age, Mactaquac Dam, 1983 (species and sexes combined).

Fork length (mm)	Age												Total
	1	2	3	4	5	6	7	8	9	10	11	12	
90-99	1												1
100-109	1												1
110-119													0
120-129													0
130-139													0
140-149		1											1
150-159		1											1
160-169													0
170-179			1										1
180-189			9										9
190-199		63	9										72
200-209		71	14										85
210-219		21	43	2									66
220-229		13	69	24	1								107
230-239		9	66	29			1						105
240-249		2	56	54	12	1							125
250-259		47	62	9	7		2		1		1		129
260-269		19	76	26	9		4		4		4		142
270-279		6	99	61	7		8		1		3		186
280-289			66	86	17		5		5			1	181
290-299			29	54	17		5					2	107
300-309				7	25		9		4			1	49
310-319				1	7		4		4				19
320-329				1	2				3				3
330-339								1					1
Total	2	2	189	329	450	283	72	33	17	8	3	3	1391
Percent	0.1	0.1	13.6	23.7	32.4	20.3	5.2	2.4	1.2	0.6	0.2	0.2	

Table 17c. Length-frequency distribution of gaspereau, by age, Mactaquac Dam, 1984 (species and sexes combined)

Fork length (mm)	Age												Total
	2	3	4	5	6	7	8	9	10	11	12		
150-159	1												1
160-169													0
170-179		1											1
180-189		22											22
190-199	47		1										48
200-209	44		15	1									60
210-219	13		118		6								137
220-229		3	184		19								206
230-239			107	49		8	1						165
240-249			71	42		5	1		1				120
250-259			47	105		12	5						169
260-269			15	113		15	4	2		2	1		152
270-279		2		68		42	9	1					122
280-289			29	56		18	2		2		1		108
290-299			8	24		25	8						65
300-309			2	2		18	9	2		1		1	35
310-319						4	4	1		2			11
320-329											1		1
Total	1	130	560	442	164	85	27	7	5	1	1		1423
Percent	0.1	9.1	39.4	31.1	11.5	6.0	1.9	0.5	0.4	0.1	0.1		

Table 17d. Length-frequency distribution of gaspereau, by age, Mactaquac Dam, 1985
(species and sexes combined).

Fork length (mm)	Age										Total
	2	3	4	5	6	7	8	9	10		
140-149	1									1	
150-159										0	
160-169	1	1								2	
170-179	1	14								15	
180-189		123	1							124	
190-199		248	7							255	
200-209		98	27							125	
210-219		34	80	9	1					124	
220-229		13	113	29						155	
230-239		7	98	44	5					154	
240-249		42	23	10						75	
250-259		44	40	9	1					94	
260-269		11	87	14	1					113	
270-279			103	29	4					136	
280-289			43	42	13	4				102	
290-299			7	22	18	5				52	
300-309			2	10	11	6		1		30	
310-319					6	4	3	1		14	
320-329					1	2	2			5	
330-339						1				1	
Total	3	538	423	387	142	55	22	5	2	1577	
Percent	0.2	34.1	26.8	24.5	9.0	3.5	1.4	0.3	0.1		

Table 17e. Length-frequency distribution of gaspereau, by age, Mactaquac Dam, 1986
 (species and sexes combined).

Fork length (mm)	Age									Total
	2	3	4	5	6	7	8	9		
150-159	1									1
160-169										0
170-179		8								8
180-189		25								25
190-199	35	5								40
200-209	28	64								92
210-219	7	136	2							145
220-229	2	115	6							123
230-239	1	77	9	1						88
240-249		127	7	3						137
250-259		248	10	1						259
260-269		135	26	2						163
270-279		22	26	15						63
280-289		1	12	8						21
290-299			1	12	2					15
300-309				3		1	1			5
310-319					1					1
Total	1	106	930	99	45	3	1	1	1186	
Percent	0.1	8.9	78.4	8.3	3.8	0.3	0.1	0.1		

Table 17f. Length-frequency distribution of gaspereau, by age, Mactaquac Dam, 1987 (species and sexes combined).

Fork length (mm)	Age					Total
	3	4	5	6	7	
170-179	1					1
180-189	18	5				23
190-199	26	29	1			56
200-209	8	147	6			161
210-219	5	216	21			242
220-229	2	114	27			143
230-239		58	71	2		131
240-249		105	40	3	1	149
250-259		89	74	2	2	167
260-269		23	174	5	1	203
270-279		2	113	13	1	129
280-289			33	14	3	50
290-299			2	5	1	8
300-309					2	2
310-319				1	1	2
Total	60	788	562	45	12	1467
Percent	4.1	53.7	38.3	3.1	0.8	

Table 17g. Length-frequency distribution of gaspereau, by age, Mactaquac Dam, 1988 (species and sexes combined).

Fork length (mm)	Age					Total
	3	4	5	6	7	
170-179	7					7
180-189	72					72
190-199	54	40	2			96
200-209	8	158	25			191
210-219	1	170	74	4		249
220-229	1	77	108	2		188
230-239		49	85	15		149
240-249		65	36	16		117
250-259		13	109	14		136
260-269			116	55		171
270-279			38	94	5	137
280-289			2	62	10	74
290-299				11	8	19
300-309				1	1	2
310-319						0
320-329				1		1
Total	143	572	595	275	24	1609
Percent	8.9	35.6	37.0	17.1	1.5	

ACKNOWLEDGEMENTS

The assistance of Mr. R. Price and the other members of the fish-lift crew at the Mactaquac Dam and of the numerous summer students who collected the sample data is gratefully acknowledged. C. Harvie and H. Stone helped in the preparation of this report.

REFERENCES

- Bigelow, H.B. and W.C. Schroeder. 1953. Fishes of the Gulf of Maine. U.S. Fish and Wildlife Serv., Fish. Bull. 53(74), 577 p.
- Cating, J.P. 1953. Determining age of Atlantic shad from their scales. U.S. Fish and Wildlife. Serv. Bull. 54:187-199.
- Jessop, B.M. 1977a. Life history data on the alewife and blueback herring of the Saint John River, New Brunswick, 1973. Dept. of Fisheries and the Environment Data Record Series No. MAR/D-77-2, 49 p.
- Jessop, B.M. 1977b. Life history data on the alewife and blueback herring of the Saint John River, New Brunswick, 1974. Dept. of Fisheries and the Environment Data Record Series No. MAR/D-77-3, 59 p.
- Jessop, B.M., A.H. Vromans and W.E. Anderson. 1982. Life-history data on alewife and blueback herring, Mactaquac Dam, 1975-81. Can. Data Rep. Fish. Aquat. Sci. No. 367, vii + 43p.
- MacLean, P., G.E. (Buck) Newsome and P.A. Dill. 1981. Discrimination by external features between alewife (*Alosa pseudoharengus*) and blueback herring (*A. aestivalis*). Can. J. Fish. Aquat. Sci. 38 (5): 544-546.
- Marcy, B.C. 1969. Age determinations from scales of *Alosa pseudoharengus* (Wilson) and *Alosa aestivalis* (Mitchill) in Connecticut waters. Trans. Amer. Fish. Soc. 98(4): 622-630.
- Nikolsky, G.V. 1963. The ecology of fishes. Academic Press, London and New York, 352 p.
- Power, G. 1964. A technique for preparing scale smears. Trans. Amer. Fish. Soc. 93: 201-202.
- Sokal, R.R. and F.J. Rohlf. 1981. Biometry. W.H. Freeman and Company, New York. 859p.