CHRONIC DISEASE INDICATOR FRAMEWORK QUICK STATS, 2013 EDITION

| INDICATOR GROUP | SPECIFIC INDICATOR MEASURE(S) | LATEST DATA | DATA SOURCE (YEAR) |
|--|---|-------------|-----------------------|
| SOCIAL AND ENVIRONME | ENTAL DETERMINANTS | | |
| Education | % of population with less than a high school education, population aged 20+ years | 14.4% | CCHS (2009-2010 |
| Income | % of population living below low-income cut-offs, after tax, all population | 9.0% | SLID (2010 |
| Employment | Average annual unemployment rate (% of labour force that was unemployed during reference period), population aged 15+ years | 7.5% | LFS (2011 |
| EARLY LIFE/CHILDHOOD I | RISK AND PROTECTIVE FACTORS | | |
| Birth Weight | % of live births with a low birth weight | 6.2% | CVS (2010 |
| Breastfeeding | % of women that report exclusive breastfeeding of their child for at least the first six months of life, women aged 15+ years | 25.9% | CCHS (2009–2010 |
| Exposure to Second-hand Smoke | % of households with children aged less than 12 years regularly exposed to environmental tobacco smoke at home | 3.9% | CTUMS (2011 |
| BEHAVIOURAL RISK AND | PROTECTIVE FACTORS | | |
| Smoking | % of population that reports being current smokers (daily and occasional), population aged 15+ years | 17.3% | CTUMS (2011 |
| | % of population that reports being current daily smokers, population aged 15+ years | 13.7% | CTUMS (2011 |
| Physical Activity | % of children and youth that attain at least 12,000 steps daily (measured), population aged 5 to 17 years | 7.0% | CANPLAY (2010–2011 |
| | % of population that reports being physically 'active' or 'moderately active' during their leisure-time, population aged 20+ years | 50.1% | CCHS (2009–2010 |
| Sedentary Behaviour | % of population that reports spending more than 14 hours per week watching television or using computers during leisure-time, population aged 12+ years | 50.5% | CCHS (2007–2008 |
| Healthy Eating | % of population that reports consuming fruit and vegetables at least 5 times/day, population aged 12+ years | 44.1% | CCHS (2009–2010 |
| Unhealthy Eating | % of population that reports drinking sugar-sweetened beverages daily, population aged 5 to 19 years | 27.2% | CHMS (2009–2011 |
| Alcohol Use | % of population that exceeds low risk alcohol drinking guidelines for chronic drinking, population aged 15+ years | 14.4% | CADUMS (2011 |
| Chronic Stress | % of population that reported life to be 'quite a bit' or 'extremely' stressful most days in the last 12 months, population aged 12+ years | 22.9% | CCHS (2009–2010 |
| RISK CONDITIONS | | | |
| Healthy Weight | % of children and youth that is obese (measured), children and youth aged 5 to 17 years | 11.7% | CHMS (2009–2011 |
| , , | % of population that is obese (measured), population aged 18+ years | 26.2% | CHMS (2009–2011 |
| Elevated Blood Glucose | % of population that has elevated blood glucose (measured), population aged 20+ years | 4.2% | CHMS (2009–2011 |
| Elevated Blood Pressure | % of population that has elevated blood pressure (measured), population aged 20+ years | 7.8% | CHMS (2009–2011 |
| Elevated Blood Cholesterol | % of population that has elevated blood cholesterol (TC/HDL-C) ratio (measured), population aged 20+ years | 17.3% | CHMS (2009–2011 |
| DISEASE PREVENTION PR | ACTICES (SECONDARY PREVENTION) | | |
| Contact with Health Care Professional | % of population that reported consulting a family physician or general practitioner at least once in the past 12 months, population aged 12+ years | 77.5% | CCHS (2009–2010 |
| | % of population that reported consulting a dentist, dental hygienist or orthodontist at least once in the past 12 months, population aged 12+ years | 66.3% | CCHS (2009–2010 |
| Disease Screening | % of women that reported having a mammogram at least once in the past five years, population aged 50 to 74 years | 81.9% | CCHS (2008 |
| | % of women that reported having at least one PAP smear test in the past three years, population aged 25 to 69 years | 80.1% | CCHS (2008 |
| | % of population that reported having at least one fecal occult blood test , colonoscopy and/ or sigmoidoscopy in the recommended time period, population aged 50 to 74 years | 39.3% | CCHS (2008 |
| Vaccination (Influenza) | % of population living with a chronic health condition that reported having a seasonal flu shot in the past 12 months, population aged 12+ years | 46.4% | CCHS (2009–2010 |
| HEALTH OUTCOMES/STAT | US | | |
| General Health | % of population that rates their health as 'very good' or 'excellent', population aged 12+ years | 60.2% | CCHS (2009–2010 |
| | % of population that rates their mental health as 'very good' or 'excellent', population aged 12+ years | 73.8% | CCHS (2009–2010 |
| | Life expectancy, at birth | 81.7 years | CCDSS (2006–2008 |
| | Life expectancy, at 65 years | 20.5 years | CCDSS (2006–2008 |
| | Health-adjusted life expectancy, at birth | 71.8 years | CCDSS (2006–2008 |
| | Health-adjusted life expectancy, at 65 years of age | 15.9 years | CCDSS (2006–2008 |
| Morbidity—Prevalence | % of population with at least one major chronic disease (cancer, diabetes, cardiovascular disease, chronic obstructive pulmonary disease), population aged 20+ years | 14.9% | CCHS (2009–2010 |
| | Prevalence of diabetes, children and youth aged 19 years or less | 0.3% | CCDSS (2008) |
| | Prevalence of diabetes, population aged 20+ years | 8.7% | CCDSS (2008) |
| | Prevalence of heart disease, population aged 20+ years | 5.4% | CCHS (2009-2010 |
| | Prevalence of stroke, population aged 20+ years | 1.2% | CCHS (2009-2010 |
| | Prevalence of asthma, children and youth aged 19 years or less | 15.2% | CCDSS (2008) |





| INDICATOR GROUP | SPECIFIC INDICATOR MEASURE(S) | LATEST DATA | DATA SOURCI (YEAR |
|----------------------|---|--------------------|----------------------|
| HEALTH OUTCOMES/STA | TUS | | |
| Morbidity—Prevalence | Prevalence of asthma, population aged 20+ years | 8.3% | CCDSS (2008) |
| | Prevalence of chronic obstructive pulmonary disease, population aged 35+ years | 8.7% | CCDSS (2008) |
| | Prevalence of arthritis, population aged 20+ years | 17.2% | CCHS (2009-2010 |
| | Prevalence of the use of health services for mental disorders, children and youth aged 19 years or less | 8.0% | CCDSS (2008 |
| | Prevalence of the use of health services for mental disorders, population aged 20+ years | 16.2% | CCDSS (2008 |
| | Prevalence of the use of health services for mood disorders and/or anxiety, children and youth aged 19 years or less | 3.3% | CCDSS (2008) |
| | Prevalence of the use of health services for mood disorders and/or anxiety, population aged 20+ years | 12.1% | CCDSS (2008) |
| | Number of individuals living with or beyond any cancer, that were diagnosed in a 5-year period, all population | 518,705 persons | CCR (2004–2008 |
| | Number of individuals living with or beyond prostate cancer, that were diagnosed in a 5-year period, all population | 105,179 persons | CCR (2004–2008 |
| | Number of individuals living with or beyond lung cancer, that were diagnosed in a 5-year period, all population | 29,780 persons | CCR (2004–2008 |
| | Number of individuals living with or beyond breast cancer, that were diagnosed in a 5-year period, all population | 90,677 persons | CCR (2004–2008 |
| | Number of individuals living with or beyond colorectal cancer, that were diagnosed in a 5-year period, all population | 67,173 persons | CCR (2004–2008 |
| Morbidity—Incidence | Incidence rate of diabetes, children and youth aged 19 years or less | 42.7 per 100,000 | CCDSS (2008 |
| | Incidence rate of diabetes, population aged 20+ years | 813.6 per 100,000 | CCDSS (200 |
| | Incidence rate of asthma, children and youth aged 19 years or less | 1097.8 per 100,000 | CCDSS (200 |
| | Incidence rate of asthma, population aged 20+ years | 392.1 per 100,000 | CCDSS (200 |
| | Incidence rate of chronic obstructive pulmonary disease, population aged 35+ years | 890.4 per 100,000 | CCDSS (200 |
| | Incidence rate of all cancers, all male population | 467.5 per 100,000 | CCR (200 |
| | Incidence rate of all cancers, all female population | 364.8 per 100,000 | CCR (200 |
| | Incidence rate of prostate cancer, all male population | 125.8 per 100,000 | CCR (200 |
| | Incidence rate of lung cancer, all male population | 69.0 per 100,000 | CCR (200 |
| | Incidence rate of lung cancer, all female population | 47.9 per 100,000 | CCR (200 |
| | Incidence rate of colorectal cancer, all male population | 60.8 per 100,000 | CCR (200 |
| | Incidence rate of colorectal cancer, all female population | 40.9 per 100,000 | CCR (200 |
| | Incidence rate of breast cancer, all female population | 98.8 per 100,000 | CCR (200 |
| Multimorbidity | % of population with multiple chronic diseases** (2+ of 10 chronic diseases), population aged 20+ years | 13.3% | CCHS (2009–201 |
| | % of population with multiple chronic diseases** (3+ of 10 chronic diseases), population aged 20+ years | 4.4% | CCHS (2009-201 |
| Disability | % of population that reports being limited in their activities 'sometimes' or 'often' due to disease/illness, population aged 12+ years | 28.4% | CCHS (2009–201 |
| Mortality | Mortality rate due to a major chronic disease (cardiovascular diseases, all cancers, chronic respiratory disease), total population | 458.0 per 100,000 | CVS (200 |
| | Mortality rate due to cardiovascular diseases, total population | 203.7 per 100,000 | CVS (200 |
| | Mortality rate due to cancer, total population | 210.9 per 100,000 | CVS (200 |
| | Mortality rate due to chronic respiratory diseases, total population | 43.5 per 100,000 | CVS (200 |
| | Mortality rate due to suicide, total population | 11.5 per 100,000 | CVS (200 |
| | All-cause mortality rate ratios among people with and without diabetes, population aged 20+ years | 4.5 rate ratio | CCDSS (200 |
| Premature Mortality | Potential years of life lost due to cancer | 1,504 per 100,000 | CVS (200 |
| | Potential years of life lost due to cardiovascular diseases | 755.4 per 100,000 | CVS (200 |
| | Potential years of life lost due to chronic respiratory diseases | 118.1 per 100,000 | CVS (200 |
| | Potential years of life lost due to suicide | 362.1 per 100,000 | CVS (200 |
| | Probability of dying (%) between ages 30 and 69 years from major chronic diseases (cardiovascular disease, cancer, chronic respiratory disease, diabetes) | 11.4% | CVS (200 |
| | Probability of dying (%) between ages 30 and 69 years from cardiovascular disease | 3.5% | CVS (200 |
| | Probability of dying (%) between ages 30 and 69 years from cancer | 7.1% | CVS (200 |
| | Probability of dying (%) between ages 30 and 69 years from chronic respiratory diseases | 0.7% | CVS (200 |
| | Probability of dying (%) between ages 30 and 69 years from diabetes | 0.5% | CVS (200 |

* CCHS 2009–2010 data exist for this indicator and are available for use when disaggregating by demographic and social markers.

** Multimorbidity: Chronic diseases included are heart disease, stroke, cancer, asthma, COPD, diabetes, arthritis, Alzheimer's or other dementia, mood disorder (depression), and anxiety

Acronyms of Data Sources: Canadian Alcohol and Other Drug Use Monitoring Survey (CADUMS), Canadian Cancer Registry (CCR), Canadian Chronic Disease Surveillance System (CCDSS), Canadian Community Health Survey (CCHS), Canadian Health Measures Survey (CHMS), Canadian Physical Activity Levels Among Youth (CANPLAY), Canadian Tobacco Use Monitoring Survey (CTUMS), Canadian Vitals Statistics (CVS), Labour Force Survey (LFS), Survey of Labour and Income Dynamics (SLID).

For questions or comments, please contact us at: Chronic.Publications.Chronique@phac-aspc.gc.ca

Suggested Citation: Centre for Chronic Disease Prevention, Public Health Agency of Canada (2013). Chronic Disease Indicator Framework, 2013 Edition.