

**Survey of Northern Abalone, *Haliotis kamtschatkana*, Populations in Southeast Barkley Sound, British Columbia, July 2000**

B.G. Lucas, A. Campbell, D. Brouwer, S. Servant,  
and N. Webb

Fisheries and Oceans Canada  
Science Branch, Pacific Region  
Pacific Biological Station  
Nanaimo, British Columbia  
V9R 5K6

2001

**Canadian Manuscript Report of  
Fisheries and Aquatic Sciences 2571**



Fisheries and Oceans  
Canada

Science

Pêches et Océans  
Canada

Sciences

**Canada**

## **Canadian Manuscript Report of Fisheries and Aquatic Sciences**

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 1426 - 1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

## **Rapport manuscrit canadien des sciences halieutiques et aquatiques**

Les rapports manuscrits contiennent des renseignements scientifiques et techniques que qui constituent une contribution aux connaissances actuelles, mais qui traitent de problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Manuscript Report of  
Fisheries and Aquatic Sciences 2571

2001

SURVEY OF NORTHERN ABALONE, *Haliotis kamtschatkana*,  
POPULATIONS IN SOUTHEAST BARKLEY SOUND,  
BRITISH COLUMBIA, JULY 2000

by

B.G. Lucas, A. Campbell, D. Brouwer,  
S. Servant<sup>1</sup> and N. Webb<sup>1</sup>

Fisheries and Oceans Canada  
Science Branch, Pacific Region  
Pacific Biological Station  
Nanaimo, British Columbia  
V9R 5K6

<sup>1</sup>Bamfield Marine Station  
Bamfield, B.C. V0R 1B0

© Her Majesty the Queen in right of Canada, 2001,  
as represented by the Minister of Fisheries and Oceans

Cat. No. Fs 97-4 /2571

ISSN 0706-6473

Correct citation for this publication:

Lucas, B.G., A. Campbell, D. Brouwer, S. Servant, and N. Webb. 2001. Survey of northern abalone, *Haliotis kamtschatkana*, populations in southeast Barkley Sound, British Columbia, July 2000. Can. Manusc. Rep. Fish. Aquat. Sci. 2571: 11 p.

## ABSTRACT

Lucas, B.G., A. Campbell, D. Brouwer, S. Servant, and N. Webb. 2001. Survey of northern abalone, *Haliotis kamtschatkana*, populations in southeast Barkley Sound, British Columbia, July 2000. Can. Manusc. Rep. Fish. Aquat. Sci. 2571: 11 p.

A survey was conducted to provide an estimate of population numbers of emergent northern abalone in areas known to have abalone present in southeast Barkley Sound, British Columbia, during July 5 to 7, 2000. Abalone shell lengths (SL) ranged from 38 to 119 mm. The estimated mean density for abalone of all sizes in all areas combined was  $0.10/\text{m}^2$ , while the estimated mean density for abalone 90 – 110 mm SL in all areas was  $0.04/\text{m}^2$ . The estimated mean total population number (and 90% confidence interval) of emergent abalone for all sizes in all areas combined was 28,485 (25,997 – 30,973). The mean total population number (and 90% confidence interval) of emergent abalone for all areas combined in the 90 – 110 mm SL size range was estimated to be 12,558 (11,275 – 13,842). The highest concentration of abalone was found on the east side of Edward King Island, where the mean density for all sizes was  $0.23/\text{m}^2$  and for abalone 90 – 110 mm SL was  $0.11/\text{m}^2$ . The estimated mean total population number (and 90% confidence interval) of emergent abalone for all sizes on the east side of Edward King Island was 20,624 (18,170 – 23,078), and for the 90 – 110 mm SL size range was 10,179 (8,206 – 12,153).

## RÉSUMÉ

Lucas, B.G., A. Campbell, D. Brouwer, S. Servant, and N. Webb. 2001. Survey of northern abalone, *Haliotis kamtschatkana*, populations in southeast Barkley Sound, British Columbia, July 2000. Can. Manusc. Rep. Fish. Aquat. Sci. 2571: 11 p.

Un relevé a été effectué du 5 au 7 juillet 2000 pour estimer l'effectif des populations d'ormeaux nordiques émergents dans les secteurs du sud-est de la baie Barkley (Colombie-Britannique) où leur présence est connue. La longueur des ormeaux échantillonnés variaient de 38 à 119 mm. La densité moyenne des ormeaux de toutes tailles échantillonnés dans tous les secteurs a été estimée à  $0,10/\text{m}^2$ , et celle des ormeaux de longueur allant de 90 à 110 mm, à  $0,04/\text{m}^2$ . Pour l'ensemble des secteurs, l'effectif total moyen de la population d'ormeaux émergents de toutes tailles a été estimé à 28 485 (intervalle de confiance à 90 % : 25 997 – 30 973), et celui des ormeaux de longueur allant de 90 à 110 mm, à 12 558 (intervalle de confiance à 90 % : 11 275 – 13 842). La plus forte concentration d'ormeaux a été trouvée du côté est de l'île Edward King, où la densité moyenne était de  $0,23/\text{m}^2$  pour les ormeaux de toutes tailles et de  $0,11/\text{m}^2$  pour les animaux faisant de 90 à 110 mm de longueur. Dans ce secteur, l'effectif total moyen de la population d'ormeaux émergents de toutes tailles a été estimé à 20 624 (intervalle de confiance à 90 % : 18 170 – 23 078), et celui des ormeaux de longueur allant de 90 à 110 mm, à 10 179 (8 206 – 12 153).

## INTRODUCTION

Northern abalone, *Haliotis kamtschatkana*, populations have declined in British Columbia (B.C.), despite complete closure of the fishery since 1990 for conservation purposes. Post-harvest surveys indicated that abalone densities were well below levels recommended to ensure sustainable populations (Breen 1986, Campbell et al. 1998). The future of *H. kamtschatkana* was sufficiently uncertain that in April 1999 this species was declared "Threatened" by COSEWIC (Committee on the Status of Endangered Wildlife in Canada).

Fisheries and Oceans Canada (DFO), in cooperation with coastal communities and other interested stakeholders, has begun an initiative to rehabilitate abalone stocks in B.C. Collection of broodstock will be required to implement the first step in this program, which is to develop aquaculture technology for this species.

To estimate the abundance of abalone stocks in areas of interest, an independent assessment using conventional survey methodology was required. Only emergent (visible) abalone were surveyed to provide a minimum estimate of mature adult population numbers; cryptic abalone were not surveyed in this census.

The objectives of this study were to determine densities, size frequencies, and population numbers of emergent northern abalone at four locations in Barkley Sound.

## MATERIALS AND METHODS

### FIELD METHODS

This survey was conducted during July 5 to 7, 2000, near Bamfield, B. C. (Figure 1). Four locations, Aguilar Point, Dixon Island, and east and west Edward King Island were selected, based on historical abundance of abalone and preliminary reconnaissance dives. The transect survey method (Cripps and Campbell 1998) was used for this study. Transects were randomly placed along the lengths of shoreline where populations were to be estimated. To avoid bias, these transect positions were determined in advance. The primary sampling unit was a "transect", made up of a cluster of secondary units, or quadrats. Each transect was 1 m wide and variable in length, depending on the slope of the substrate. Prior to entering the water, a lead line was laid perpendicular to shore from the boat in depths of approximately 12 m to 0 m from datum. The secondary sampling unit consisted of a 1 m x 1 m square quadrat that was placed along the transect, 1 m away, to avoid sampling areas disturbed by the lead line placement. Divers flipped the quadrat parallel to the transect line, from deep to shallow, and the number of "emergent" or "exposed" abalone, shell length (SL in mm) of each abalone, depth, substrate type, slope, and algal cover was recorded. Algal cover, sea urchins and starfish were removed from the quadrat to ensure all emergent abalone were detected. Boulders were not moved to search for cryptic abalone, since the majority of mature animals (i.e.,  $\geq 70$  mm SL) are exposed (Campbell 1996). On transects where the slope of the substrate would have resulted in many quadrats, only a proportion (e.g., every other quadrat) were surveyed.

## ANALYTICAL METHODS

All gauge depths were converted to depth (m) at chart datum. The survey results were analyzed according to Fisheries and Oceans Canada methodology (Campbell et al. 1999). To determine the size frequency distribution for a location, the number of abalone in each 5 mm SL size class was divided by the total number of abalone in each location.

The estimated mean density,  $\bar{d}$  (number/m<sup>2</sup>), of abalone was calculated as:

$$\bar{d} = \frac{\sum_t (d_t * L_t)}{\sum_t L_t} \quad (1)$$

The standard error of the mean density,  $s_d$ , was calculated as:

$$s_d = \sqrt{1 - \frac{n}{T}} * \sqrt{\frac{\sum_t (d_t * L_t - \bar{d} * L_t)^2}{n * (n-1) * \bar{L}^2}} \quad (2)$$

where  $n$  is the number of transects,  $d_t$  is the density at transect  $t$ ,  $L_t$  is the length of transect  $t$  (or area in square metres since each transect is one metre wide),  $\bar{L} = \frac{\sum_t L_t}{n}$  is the mean transect length (or mean area in square metres), and  $T$  is the total possible number of transects that can be sampled in the surveyed area. The expression  $\sqrt{1 - \frac{n}{T}}$  is approximately one, because the sample size  $n$  is small compared to  $T$ . This method accounts for the variable length of transects and for the variable proportion of quadrats surveyed along each transect.

To estimate the mean density (number/m<sup>2</sup>) for a specific size group ( $J$ ), (e.g., 90 – 110 mm SL) the value  $d_t$  was substituted with densities ( $P_{t,J}$ ) in equations 1 and 2. The mean density  $P_{t,J}$  (number/m<sup>2</sup>) of abalone of size group  $J$  in transect  $t$  was calculated as:

$$P_{t,J} = \frac{\left( N_t * \frac{\sum_{i \in J} m_{t,i}}{M_t} \right)}{S_t} \quad (3)$$

where  $J$  is a subset (e.g., 90 – 110 mm SL) of possible  $i$  values representing a range of shell lengths (SL <sub>$i$</sub> ),  $S_t$  is the number of quadrats surveyed for density estimates in transect  $t$ ,  $N_t$  is the

number of abalone counted in transect  $t$ ,  $M_t = \sum_i m_{t,i}$  the total number of abalone measured for size in transect  $t$ , and  $m_{t,i}$  is the number of abalone in size range  $J$  of transect  $t$ .

The approximate lower 90% confidence interval of the mean density (L90CI) was calculated as:

$$\text{L90CI} = \bar{d} - t_{\alpha/2} (s_d / \sqrt{n}) \quad (4)$$

where  $t_{\alpha/2}$  is the critical t-distribution value at degrees of freedom of  $n-1$ .

The total number of abalone (X) in each location was calculated as:

$$X = \bar{d} * A \quad \text{or} \quad X = \text{L90CI} * A \quad (5)$$

where L90CI is the lower 90% confidence interval of the mean density for all sizes or for a particular size group of abalone, and A is the estimated area ( $\text{m}^2$ ) of the surveyed location. The area of the location surveyed was calculated as the product of the mean transect length (since transects generally covered the breadth of suitable abalone habitat) and the length of the location as measured by the distance between the two furthest shoreline points used when generating random transects.

## RESULTS

King Edward Island and Aguilar Point were moderately exposed to wind and storms, with ground swells normal, while most of Dixon Island was well sheltered. The substrate consisted of bedrock, boulders, cobble and gravel, with some sand and shell (Table 1). The slope of the substrate (angle from horizontal) ranged from  $2^\circ$  –  $60^\circ$ . The canopy was mostly *Nereocystis* along with some *Macrocystis* and *Pterygophora*. The predominant understory species was *Desmarestia*, followed by *Laminaria*. An unknown filamentous algae and articulated corraline algae were the most common turf algae, and encrusting corraline algae were abundant as bottom cover.

The depths surveyed ranged from -1.5 to 14.6 m from datum (Table 2). One hundred eighteen abalone were counted in quadrats along the 33 transects surveyed (Tables 2, 3). Aguilar Point had 11 abalone along 5 transects, where the mean transect length was 62 m. Dixon Island had 1 abalone along 6 transects, where the mean transect length was 59 m. East Edward King Island had 77 abalone along 11 transects, where the mean transect length was 60 m. West Edward King Island had 29 abalone along 11 transects, where the mean transect length was 93 m. Eleven transects surveyed (33%) had no emergent abalone. The maximum density of emergent abalone on a transect was 0.47 abalone/ $\text{m}^2$ .

Sizes of abalone surveyed ranged from 38 to 119 mm SL (Figure 2). The mean size of emergent abalone for all areas was 88 mm SL. Sizes of abalone found at Aguilar Point ranged from 63 to 119 mm SL. The one abalone found at Dixon Island measured 110 mm SL. East



Edward King Island abalone ranged from 38 to 114 mm SL. At West Edward King Island, the sizes of abalone found ranged from 62 to 110 mm SL. The mean size of emergent abalone was 98 mm SL at Aguilar Point, and 87 mm SL at both East and West Edward King Island.

Estimated mean total density of emergent abalone of all sizes in all areas combined was  $0.10/\text{m}^2$  (Table 4). The mean population estimate was 28,485 emergent abalone in all the locations surveyed. The mean estimate of the number of abalone in the 90 – 110 mm SL size range for all areas was 12,558. The highest concentration of abalone was found on the east side of Edward King Island, where the mean density for all sizes was  $0.23/\text{m}^2$  and for abalone 90 – 110 mm SL was  $0.11/\text{m}^2$ . The mean total population number of emergent abalone in the 90 – 110 mm SL size range was estimated for East Edward King Island to be 10,179.

## DISCUSSION

The small mean size (< 100 mm SL), and the absence of large abalone (> 120 mm SL), in this survey suggest that some individuals are ‘surf’ abalone, whose growth is stunted by food or habitat limitations (Sloan and Breen 1988). The mean size is similar to the mean sizes (78 – 90 mm SL) measured at Edward King Island in 1984 (Emmett and Jamieson 1988), but smaller than the size range of abalone (51 – 146 mm SL) found in other areas of Barkley Sound in 1964 (Quayle 1971).

A conservative estimate of the total population numbers of emergent abalone of all sizes was provided by the lower 90% confidence interval (L90CI) for the mean, which was 654 abalone for Aguilar Point, 41 for Dixon Point, 18,170 for East Edward King Island, and 3,234 for West Edward King Island. The L90CI for mature emergent abalone 90 – 110 mm SL was 119 abalone for Aguilar Point, 0 for Dixon Point, 8,206 for east Edward King Island, and 923 for west Edward King Island.

This survey provides no evidence of recovery of abalone populations since the province-wide closure in 1990. The total abalone densities found during this survey at East Edward King Island were lower than those found in the same area ( $0.52/\text{m}^2$ ) in 1984 (Emmett and Johnstone 1985). All areas surveyed had low densities similar to other recent surveys in B.C. (Campbell et al. 1998; Lucas et al. 2000a, 2000b, 2001). Since these low densities are well below those recommended to ensure sustainable populations (Breen 1986), the removal of any abalone from these areas must be considered with caution.

## ACKNOWLEDGMENTS

We thank K. Fong and G. Parker for their assistance in organising this study; J. Harding, M. MacNab, and L. Verhegge for diving; and C. Hand for reviewing the manuscript.

## REFERENCES CITED

- Breen, P.A. 1986. Management of the British Columbia fishery for northern abalone (*Haliotis kamtschatkana*). Can. Spec. Publ. Fish. Aquat. Sci. 92: 300-312.
- Campbell, A. 1996. An evaluation of abalone surveys off southeast Queen Charlotte Islands. Can. Tech. Rep. Fish. Aquat. Sci. 2089: 111-121.
- Campbell, A., W. Hajas, and D. Bureau. 1999. Quota options for the red sea urchin fishery in British Columbia for fishing season 2000/2001. Can. Stock Assessment Secretariat Res. Doc. 99/201.
- Campbell, A., I. Winther, B. Adkins, D. Brouwer, and D. Miller. 1998. Survey of the northern abalone (*Haliotis kamtschatkana*) in the central coast of British Columbia, May 1997. Can. Stock Assessment Secretariat Res. Doc. 98/99.
- Cripps, K. and A. Campbell. 1998. Survey of abalone populations at Dallain Point and Higgins Pass, central coast of British Columbia, 1995-96. Can. Manuscr. Rep. Fish. Aquat. Sci. 2445: 31 p.
- Emmett, B. and G. S. Jamieson. 1988. An experimental transplant of northern abalone, *Haliotis kamtschatkana*, in Barkley Sound, British Columbia. Fish Bull. 77: 95-105.
- Emmett, B. and D. Johnstone. 1985. An experimental abalone transplant project in Barkley Sound, British Columbia. Final report for DFO contract 08SB.FP 597-4-0145.
- Lucas, B.G., D. Brouwer, and A. Campbell. 2000a. Survey of northern abalone, *Haliotis kamtschatkana*, populations at Malcolm Island and Cormorant Island, British Columbia, October 1999. Can. Manuscr. Rep. Fish. Aquat. Sci. 2520: 10 p.
- Lucas, B.G., D. Brouwer, and A. Campbell. 2001. Survey of northern abalone, *Haliotis kamtschatkana*, populations near Kitkatla, British Columbia, March 2000. Can. Manuscr. Rep. Fish. Aquat. Sci. 2572: 11 p.
- Lucas, B.G., A. Campbell, and D. Brouwer. 2000b. Survey of northern abalone, *Haliotis kamtschatkana*, populations in Lotbinière Bay, British Columbia, March 2000. Can. Manuscr. Rep. Fish. Aquat. Sci. 2545: 10 p.
- Quayle, D. B. 1971. Growth, morphometry and breeding in the British Columbia abalone (*Haliotis kamtschatkana* Jonas). Fish. Res. Board Can. Tech. Rep. 279: 84 p.
- Sloan, N. A. and P. A. Breen. 1988. Northern Abalone, *Haliotis kamtschatkana*, in British Columbia: fisheries and synopsis of life history information. Can. Spec. Publ. Fish. Aquat. Sci. 103: 46 p.

Table 1. Habitat summary for abalone transects surveyed in the Bamfield area, July 5 – 7, 2000. The most common substrates are listed (1 = first, 2 = second, 3 = third) where substrate code 1 = smooth bedrock, 2 = bedrock with crevices, 3 = boulders, 4 = cobble, 5 = gravel, 6 = pea gravel, 7 = sand, 8 = shell, and 9 = mud. The algae present are listed by growth characteristic for the most common (Sp 1) and second most common (Sp 2) species with the percentage of cover (%) for each, where AC = articulated coralline, AG = *Agarum*, AL = *Alaria*, CO = *Costaria*, DM = *Desmarestia*, EN = encrusting coralline, EG = *Egregia*, F = unknown foliose, GI = *Gigartina*, GR = sea grass, H = unknown hairy, LA = *Laminaria*, MA = *Macrocystis*, NT = *Nereocystis*, PL = *Pleurophyucus*, PT = *Pterygophora*, R = unknown red, and UL = *Ulva*.

Transect	Date	Algae																	
		Substrate			Slope degrees	Canopy			Understory				Turf				Bottom Cover		
		1	2	3		Sp 1	%	Sp 2	%	Sp 1	%	Sp 2	%	Sp 1	%	Sp 2	%	Sp	%
<b>Aguilar Point</b>																			
AP-9	July 7	3	4	7	10	NT	5				NT	5	LA	5	H	15		EN	40
AP-11	July 7	8	7	4	30	NT	1				DM	1	MA	1	H	50		EN	40
AP-16	July 7	3	4	7	10	NT	5				LA	5	NT	5	H	15		EN	30
AP-30	July 7	3	4	7	20						LA	10	NT	10	H	10		EN	30
AP-39	July 7	3	4	8	45	NT	1				CO	1						EN	50
<b>Dixon Island</b>																			
DX-4	July 7	4	5	8	8									UL	40	GI	15	EN	10
DX-9	July 7	7	4	3	20									DM	1			EN	60
DX-17	July 7	2	7	3	50				DM	4	AL	1						EN	50
DX-26	July 7	3	4	7	10				EG	5	MA	5		GI	5	UL	5	EN	30
DX-30	July 7	2	7	4	45									H	2	UL	2	EN	50
DX-34	July 7	7	4	2	25									H	2			EN	95
<b>East Edward King Island</b>																			
EEK-10	July 6	3			10	NT	30	PT	10	PT	5	LA	5	AC	5			EN	90
EEK-31	July 6	2			20	NT	5			DM	10	NT	5					EN	60
EEK-35	July 6	2	4	5	10	NT	5			NT	5	DM	5	F	5			EN	50
EEK-47	July 6	3	4	8	20	NT	8			AL	1	CO	1					EN	70
EEK-52	July 6	1	2		20	NT	1			LA	1			AC	3			EN	90
EEK-57	July 6	1	2	3	30	NT	5			NT	5	LA	5	AC	5			EN	70
EEK-76	July 6	2	4	5	20					DM	5			F	5			EN	70
EEK-82	July 6	1	2	3	30	NT	1			DM	5	LA	1	AC	2			EN	80
EEK-85	July 6	1	3		10					LA	10	NT	5					EN	60
EEK-86	July 6	2	3	4	15					LA	10	DM	5					EN	60
EEK-94	July 6	3	4	2	60	NT	1			LA	1							EN	80
<b>West Edward King Island</b>																			
WEK-11	July 5	3	4	6	15	NT	20			LA	20	NT	20	F	5			EN	40
WEK-25	July 5	1	2	3	45													EN	60
WEK-36	July 5	3	4	8	20	NT	2			DM	25	LA	1					EN	75
WEK-42	July 5	3	4	1	2	MA	5	NT	5	NT	5			AC	5	R	5	EN	90
WEK-43	July 5	4	3	1	3	MA	1	PT	1	DM	2			R	2			EN	80
WEK-45	July 5	1	2	7	5	NT	1			DM	5	EG	1	AC	5			EN	50
WEK-48	July 5	1	3	4	60	NT	30	MA	30	LA	10	AL	5	AC	2			EN	85
WEK-59	July 5	1	2	8	20	NT	20			DM	10	LA	1					EN	90
WEK-69	July 5	3	2	7	10	NT	1			NT	1			R	1			EN	80
WEK-89	July 5	4	3		3	NT	5			DM	5			AC	10				
WEK-97	July 5	1	3	4	50	NT	2			CO	2	NT	1	AC	5			EN	90

Table 2. Dive summary for abalone transects surveyed in the Bamfield area, July 5 – 7, 2000.

Transect	Date	Time			Depth (m)		Number of Quadrats	Transect Length (m)	Number of Abalone		Density (#/m <sup>2</sup> )	
		Start	Finish	Bottom	Min	Max			all sizes	90-110mm	all sizes	90-110mm
Aguilar Point												
AP-9	July 7	10:49	11:07	0:18	0.82	11.43	27	53	5	0	0.19	0.00
AP-11	July 7	10:05	10:30	0:25	0.67	10.30	29	57	3	3	0.10	0.10
AP-16	July 7	10:04	10:24	0:20	0.30	11.46	43	85	2	1	0.05	0.02
AP-30	July 7	9:15	9:40	0:25	-0.91	9.54	30	59	1	0	0.03	0.00
AP-39	July 7	9:20	9:45	0:25	0.06	10.21	28	55	0	0	0.00	0.00
Dixon Island												
DX-4	July 7	13:17	13:39	0:22	-0.46	10.67	18	101	0	0	0.00	0.00
DX-9	July 7	13:44	14:13	0:29	-0.12	10.18	34	67	0	0	0.00	0.00
DX-17	July 7	12:50	13:20	0:30	-0.34	11.43	31	61	0	0	0.00	0.00
DX-26	July 7	12:24	12:49	0:25	-0.15	10.00	31	61	1	0	0.03	0.00
DX-30	July 7	12:20	12:36	0:16	0.52	11.52	15	29	0	0	0.00	0.00
DX-34	July 7	10:55	11:16	0:21	0.24	11.16	18	35	0	0	0.00	0.00
East Edward King Island												
EEK-10	July 6	9:42	10:42	1:00	1.16	10.06	42	83	4	1	0.10	0.02
EEK-31	July 6	9:45	10:04	0:19	0.73	12.53	17	33	2	1	0.12	0.06
EEK-35	July 6	10:25	11:14	0:49	1.16	11.49	39	77	13	7	0.33	0.18
EEK-47	July 6	11:20	12:05	0:45	0.98	14.57	46	91	3	2	0.07	0.04
EEK-52	July 6	14:40	15:05	0:25	0.52	9.91	16	31	5	3	0.31	0.19
EEK-57	July 6	11:40	11:56	0:16	0.09	11.46	15	29	6	3	0.40	0.20
EEK-76	July 6	12:29	12:57	0:28	1.49	11.49	40	79	3	1	0.08	0.03
EEK-82	July 6	12:35	13:05	0:30	0.52	11.13	24	47	6	1	0.25	0.04
EEK-85	July 6	13:40	14:12	0:32	-0.76	9.33	38	75	9	6	0.24	0.16
EEK-86	July 6	14:25	15:15	0:50	-1.13	9.75	38	75	18	9	0.47	0.24
EEK-94	July 6	14:00	14:20	0:20	-0.55	11.55	20	39	8	4	0.40	0.20
West Edward King Island												
WEK-11	July 5	10:00	10:55	0:55	2.23	11.16	37	73	17	4	0.46	0.11
WEK-25	July 5	9:50	10:10	0:20	2.93	12.98	36	36	3	0	0.08	0.00
WEK-36	July 5	10:25	11:05	0:40	2.19	9.94	65	129	6	6	0.09	0.09
WEK-42	July 5	11:40	12:25	0:45	0.58	11.37	56	166	0	0	0.00	0.00
WEK-43	July 5	12:55	13:35	0:40	-0.15	9.39	47	139	0	0	0.00	0.00
WEK-45	July 5	13:03	13:45	0:42	-0.34	2.87	24	47	0	0	0.00	0.00
WEK-48	July 5	13:55	14:05	0:10	0.67	9.30	11	21	0	0	0.00	0.00
WEK-59	July 5	14:35	15:15	0:40	4.08	8.56	34	67	1	0	0.03	0.00
WEK-69	July 5	15:07	15:46	0:39	2.53	8.78	40	118	0	0	0.00	0.00
WEK-89	July 5	16:15	17:08	0:53	-1.52	7.32	38	185	1	0	0.03	0.00
WEK-97	July 5	16:15	16:30	0:15	0.88	9.14	19	37	1	0	0.05	0.00

Table 3. Summary statistics and area surveyed for transect survey of emergent abalone from the Bamfield area, July 5 – 7, 2000. Values in brackets are standard errors.

Details per transect	Aguilar Point	Dixon Island	E Edward King I	W Edward King I	All areas
Number of transects	5	6	11	11	33
Number of abalone	11	1	77	29	118
Mean transect length (m)	61.80 (5.89)	59.00 (10.51)	59.91 (7.22)	92.55 (17.21)	70.91 (6.89)
Mean number of quadrats	31.40 (2.94)	24.50 (3.41)	30.45 (3.61)	31.70 (2.18)	31.70 (2.18)
Mean depth (m)	6.14 (0.25)	4.86 (0.29)	6.08 (0.17)	5.36 (0.15)	5.64 (0.10)
Mean minutes/transect	22.60 (1.50)	23.83 (2.15)	34.00 (4.43)	36.27 (4.47)	31.18 (2.30)
Mean minutes/quadrat	0.74 (0.08)	1.01 (0.07)	1.14 (0.08)	1.03 (0.11)	1.02 (0.05)
Shoreline distance (m)	250	730	1500	1500	3980
Area surveyed (m <sup>2</sup> )	15,450	43,070	89,865	138,825	287,210

Table 4. Mean densities (number/m<sup>2</sup>) and populations estimates of emergent abalone from the Bamfield area, July 5 – 7, 2000. Values in brackets are 90% confidence intervals.

Details per location	Aguilar Point	Dixon Island	E. Edward King Is.	W. Edward King Is.	All areas
Mean density all sizes	0.07 (0.04 - 0.10)	0.01 (0.00 - 0.01)	0.23 (0.20 - 0.26)	0.06 (0.02 - 0.09)	0.10 (0.09 - 0.11)
Mean density 90 - 110 mm SL	0.03 (0.01 - 0.04)	0.00 (-0.01 - 0.01)	0.11 (0.09 - 0.14)	0.02 (0.01 - 0.03)	0.04 (0.07 - 0.05)
Total population all sizes	1,082 (654 - 1,509)	239 (41 - 438)	20,624 (18,170 - 23,078)	7,085 (3,234 - 12,376)	28,485 (25,997 - 30,973)
Population 90 - 110 mm SL	394 (119 - 668)	0 (-332 - 332)	10,179 (8,206 - 12,153)	2,700 (923 - 4,477)	12,558 (11,275 - 13,842)




Figure 1. Abalone survey locations in the Bamfield area, July 5 – 7, 2000. The inset shows the Barkley Sound area of Vancouver Island, British Columbia.

This page purposely left blank




Figure 2. Size frequencies of emergent abalone found in quadrats during dive surveys in the Bamfield area, July 5 – 7, 2000, are shown for A) all areas combined, B) Aguilar Point, C) Dixon Island, D) East Edward King Island, and F) West Edward King Island. Number of abalone (N) and mean shell length in mm are shown.