

**Life history parameters of Great Lakes populations of lake trout,  
lake whitefish, bloater, walleye, and yellow perch**

C. Chu and M.A. Koops

Great Lakes Laboratory for Fisheries and Aquatic Sciences  
867 Lakeshore Road  
Burlington, Ontario  
L7R 4A6

**2007**

**Canadian Manuscript Report of  
Fisheries and Aquatic Sciences 2811**



Fisheries and Oceans  
Canada

Pêches et Océans  
Canada

**Canada**

## **Canadian Manuscript Report of Fisheries and Aquatic Sciences**

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 901-1425 were issued as Manuscript Reports of the Fisheries Research Board of Canada. Numbers 1426-1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

## **Rapport Manuscrit Canadien des Sciences Halieutiques et Aquatiques**

Les rapports manuscrits contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui traitent de problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, e'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Manuscript Report of  
Fisheries and Aquatic Sciences 2811

2007

**Life history parameters of Great Lakes populations of lake trout, lake whitefish, bloater,  
walleye, and yellow perch**

by

C. Chu and M.A. Koops

Great Lakes Laboratory for Fisheries and Aquatic Sciences  
Bayfield Institute  
Fisheries and Oceans Canada  
867 Lakeshore Road  
Burlington, Ontario  
L7R 4A6  
E-mail: [KoopsM@dfo-mpo.gc.ca](mailto:KoopsM@dfo-mpo.gc.ca)

© Her Majesty the Queen in Right of Canada, 2007.  
Cat. No. Fs 97-4/2811E ISSN 0706-6473

Correct citation for this publication:

Chu, C. and M.A. Koops. 2007. Life history parameters of Great Lakes populations of lake trout, lake whitefish, bloater, walleye, and yellow perch. *Can. Manuscr. Rep. Fish. Aquat. Sci.* 2811: vi + 43 p.

## ABSTRACT

Chu, C. and M.A. Koops. 2007. Life history parameters of Great Lakes populations of lake trout, lake whitefish, bloater, walleye, and yellow perch. Can. Manusc. Rep. Fish. Aquat. Sci. 2811: vi + 43 p.

Life history parameters summarize the growth, maturity and mortality of populations. This report describes the data used to calculate the life history parameters of 12 lean and 6 siscowet lake trout (*Salvelinus namaycush*), 26 lake whitefish (*Coregonus clupeaformis*), 8 bloater (*Coregonus hoyi*), 9 walleye (*Sander vitreus*) and 12 yellow perch (*Perca flavescens*) populations in the Great Lakes. Interpopulation (among populations), intrapopulation (within populations through time) and virtual population analysis results are summarized in this report. These estimates of life history parameters can be used to inform fisheries management and ecological modelling.

## RÉSUMÉ

Chu, C. and M.A. Koops. 2007. Life history parameters of Great Lakes populations of lake trout, lake whitefish, bloater, walleye, and yellow perch. Can. Manusc. Rep. Fish. Aquat. Sci. 2811: vi + 43 p.

Les paramètres du cycle biologique rendent compte de la croissance, de la maturité et de la mortalité des populations. Le présent rapport décrit les données utilisées pour calculer les paramètres du cycle biologique de 12 populations de touladi maigre, 6 populations de siscowet (touladi gras), 26 populations de grand corégone, 8 populations de cisco de fumage, 9 populations de doré jaune et 12 populations de perchaude des Grands Lacs. Les résultats des analyses interpopulationnelles (entre populations) et intrapopulationnelles (à l'intérieur des populations, dans le temps) et des analyses séquentielles des populations sont présentés dans le rapport. Ces estimations des paramètres du cycle biologique peuvent être utiles pour la gestion des pêches et la modélisation écologique.

## TABLE OF CONTENTS

<b>ABSTRACT</b> .....	<b>iii</b>
<b>RÉSUMÉ</b> .....	<b>iii</b>
<b>LIST OF TABLES</b> .....	<b>v</b>
<b>LIST OF FIGURES</b> .....	<b>vi</b>
<b>INTRODUCTION</b> .....	<b>1</b>
<b>METHODS</b> .....	<b>1</b>
DATA SOURCES .....	1
EXAMINING SAMPLING BIAS IN THE DATA .....	2
LIFE HISTORY PARAMETERS .....	3
VIRTUAL POPULATION ANALYSES .....	3
<b>RESULTS</b> .....	<b>4</b>
LENGTH AT AGE ESTIMATES FOR YEAR-ROUND VERSUS FALL SAMPLING DATA .....	4
SAMPLING BIAS IN THE DATA .....	4
LIFE HISTORY PARAMETER ANALYSIS .....	4
VIRTUAL POPULATION ANALYSES .....	6
<b>DISCUSSION</b> .....	<b>6</b>
<b>ACKNOWLEDGEMENTS</b> .....	<b>7</b>
<b>REFERENCES</b> .....	<b>8</b>

## LIST OF TABLES

Table 1. Summary of the interpopulation data used to examine the life history parameters of fish populations in the Great Lakes Basin.....	10
Table 2. Summary of the intrapopulation (yearly) data used to examine the life history parameters of fish populations through time.....	12
Table 3. Life history parameters calculated from age, length and maturity data from each species and population.....	14
Table 4. Intrapopulation life history parameter values for lake whitefish (LWF), walleye (WALL) and yellow perch (YP) populations in Lake Erie.....	15
Table 5. Life history parameter values for lake whitefish (LWF), smallmouth bass (SMB), walleye (WALL) and yellow perch (YP) populations in Lake Ontario.....	18
Table 6. Life history parameter values for bloater (BLO), lean lake trout (LT), lake whitefish (LWF), walleye (WALL) and yellow perch (YP) populations in Lake Huron.....	19
Table 7. Life history parameter values for bloater (BLO), lake whitefish (LWF) and yellow perch (YP) populations in Lake Michigan.....	22
Table 8. Life history parameters for lake whitefish (LWF), lean lake trout (LT), siscowet lake trout (LT-SIS) and yellow perch (YP) populations in Lake Superior.....	24
Table 9. Life history parameter values of cohorts used in the virtual population analyses.....	29

## LIST OF FIGURES

Figure 1. Extent of (a) 12 lean lake trout and (b) 6 siscowet lake trout populations used to examine life history traits in the Great Lakes basin. ....	30
Figure 2. Extent of (a) 26 lake whitefish and (b) 8 bloater populations used to examine life history traits in the Great Lakes basin. ....	31
Figure 3. Extent of (a) 12 yellow perch and (b) 9 walleye populations used to examine life history traits in the Great Lakes basin. ....	32
Figure 4. Comparison of the length at age relationships produced using fish collected year-round and fish collected in the summer and/or fall for three populations, (a) lake whitefish in Lake Huron (b) yellow perch in Lake Erie, and (c) lake trout in Lake Michigan. ....	33
Figure 5. Comparison of the length at age relationships produced using yellow perch collected in the summer and fall within the same lake a) EYP41999, b) EYP42000 and c) EYP42001. .	34
Figure 6. Length at age for a) lake whitefish, b) walleye and c) yellow perch caught using different gillnet mesh sizes in Lake Erie (all populations). ....	35
Figure 7. Growth curves for a) lake whitefish, b) walleye and c) yellow perch in Lake Erie (all populations). ....	36
Figure 8. Growth curves for 5 populations of a) lake whitefish, b) yellow perch and c) walleye in Lake Erie. ....	37
Figure 9. Growth curves for 1 population of a) walleye, b) lake whitefish and c) yellow perch in Lake Ontario. ....	38
Figure 10. Growth curves for a) 3 bloater, b) 2 lean lake trout and c) 3 walleye populations in Lake Huron. ....	39
Figure 11. Growth curves for a) 7 lake whitefish and b) 3 yellow perch populations in Lake Huron. ....	40
Figure 12. Growth curves for a) 4 lake whitefish, b) 2 yellow perch and c) 5 bloater populations in Lake Michigan. ....	41
Figure 13. Growth curves for a) 10 lean lake trout and b) 6 siscowet lake trout populations in Lake Superior. ....	42
Figure 14. Growth curves for a) 9 lake whitefish population and b) 1 yellow perch population in Lake Superior. ....	43



## INTRODUCTION

Life history parameters summarize the growth, maturity and mortality of populations. Particular attention has been given to estimating mortality from life history parameters because mortality directly influences the population dynamics of individual species and affects fisheries management decisions (Beverton and Holt, 1959; Pauly, 1980; Hoenig, 1983, Roff, 1984; Charnov *et al.*, 1993). This report documents the data and methods used to calculate the life history parameters of several fish species and populations in the Great Lakes.

The objectives of this study were three-fold; the amalgamation of life history data for several fish species and populations within the Great Lakes into a single database, calculation of the life history parameters, and an initial examination of interpopulation (spatial, among population) differences, intrapopulation (temporal, within population) and cohort trends in the life history parameters.

## METHODS

### DATA SOURCES

Individual fish data consisting of the total length (cm), round weight (g), age (years), maturity (yes/no), sex (male, female or unknown), population code, population location and year sampled were gathered from several agencies throughout the Great Lakes (Table 1). Maturity data were not available for all populations. Fish that were not aged were not included in our analyses. In total, aged fish data were available for 41,394 lean and 8,815 siscowet lake trout (*Salvelinus namaycush*), 48,565 lake whitefish (*Coregonus clupeaformis*), 25,875 bloater (*Coregonus hoyi*), 18,505 walleye (*Sander vitreus*), and 73,304 yellow perch (*Perca flavescens*) in the Great Lakes basin (Table 1). Yearly data spanned the early 1980's to 2003 but were not consistently available for every year within that period (Table 2). The data were collected year-round which may produce underestimates of length at age and bias the length-based life history

parameter estimates as fish sampled in the spring will be smaller than the same fish sampled in the fall. To examine if year-round sampling underestimated length at age, year-round length at age data were compared using analysis of covariance (ANCOVA) to length at age data collected in the fall.

Locations of the populations for each species were compiled in ArcView 3.2 (ESRI, 1996) using population distinctions based on historical information, tracking and spawning data. Lean lake trout data exist for Lake Superior and Lake Huron however siscowet lake trout data were only available for populations in Lake Superior (Figure 1). Lake whitefish data were available from each Great Lake and bloater populations in lakes Michigan and Huron (Figure 2). Yellow perch data were available for all of the Great Lakes and data were available for walleye in lakes Huron, Erie and Ontario (Figure 3). Species data from Lake Ontario were not divided into different populations because population boundaries are not known for that lake (personal communication J. Dietrich, OMNR).

## **EXAMINING SAMPLING BIAS IN THE DATA**

Much of the data used in this study came from fish caught using gillnet gangs of different mesh sizes. This sampling method is size selective with smaller meshes excluding large individuals and larger meshes excluding small individuals. Therefore growth curves fitted to these data may underestimate growth in older fish while inflating the growth rates of younger fish (Beauchamp *et al.*, 2004; Taylor *et al.*, 2005). ANCOVA was used to determine if mesh sizes used in this study produced biased estimates of growth rates. Lake whitefish, yellow perch and walleye data from Lake Erie were chosen to test for bias in growth estimation as these fishes were sampled extensively using a wide range of mesh sizes, 32 mm – 152 mm.

We also wanted to determine if pooling data across all mesh sizes to estimate growth for the population could negate any potential biases from individual mesh sizes. One would expect biased samples to produce two different growth curves when young (age 0 and 1) fish and old (age 7 and older) fish were included and excluded from the analysis. To examine this, we tested for differences in the mean growth curves of fish aged 0 – 12 versus fish aged 2 – 7 for each species. Ages 2 – 7 was chosen since these ages are well represented in the data for each species.

### **LIFE HISTORY PARAMETERS**

Eight life history parameters (Table 3) were calculated for each population with age, length and maturity data. For populations without maturity data (Table 1), five life history parameters ( $k$ ,  $L_0$ ,  $t_0$ ,  $t_{max}$ ,  $Z$ ) were calculated. Asymptotic length, von Bertalanffy growth coefficient, age at length zero and length at age zero were calculated by fitting the length at age data to a von Bertalanffy growth function using Ford-Walford plots. Total mortality was calculated using the number of individuals caught at each age within each population and catch-curve analysis outlined in Hilborn and Walters (1992). Life history parameters were calculated for each lean lake trout, siscowet lake trout, lake whitefish, bloater, walleye and yellow perch population (interpopulation) and for the individual years within each population through time (intrapopulation).

### **VIRTUAL POPULATION ANALYSES**

Virtual population analyses (VPA) were also used in this study. This approach differs from the inter- and intra-population analyses because it estimates the life history parameters of individual cohorts rather than the mean of the populations or means of individual years. Populations with at least 10 consecutive years of data were used for these analyses. This

included 10 populations, EWAL1, EWAL2, EYP1, EYP2, HLWFON2, HYPON2, OLWF1, OYP1, SLTSISMI7 and SLTWI2 (Table 2).

## RESULTS

### LENGTH AT AGE ESTIMATES FOR YEAR-ROUND VERSUS FALL SAMPLING DATA

Comparison of the length at age estimates for year-round versus fall sampling data showed no consistent pattern among populations (Figure 4) or within populations through time (Figure 5). Therefore year-round data were used for our analyses.

### SAMPLING BIAS IN THE DATA

Growth curves for lake whitefish ( $F_{(13,43)}=7.052, p<0.000$ ) and yellow perch ( $F_{(13,96)}=2.829, p=0.002$ ) were significantly different with different mesh sizes (Figure 6). Mesh size had no effect on growth curves produced for walleye ( $F_{(13,119)}=1.729, p=0.063$ ; Figure 6). However, to calculate the life history parameters the entire sampled population is used, that is, data from all mesh sizes are pooled together. Comparison of the mean growth curves for the two age groups, ages 0 – 12 versus ages 2 – 7, were not significantly different for lake whitefish ( $F_{(1,15)}=0.122, p=0.732$ ), yellow perch ( $F_{(1,15)}=0.062, p=0.807$ ) or walleye ( $F_{(1,15)}=0.011, p=0.918$ ) (Figure 7). This suggests that although there is sampling bias for lake whitefish and yellow perch caught with different mesh sizes, the pooled data produce growth curves representative of the mean population growth. Sampling protocols were similar for species and populations throughout the Great Lakes therefore we conclude that the extensive datasets of pooled data produce accurate estimates of the growth rates exhibited by those populations.

### LIFE HISTORY PARAMETER ANALYSIS

The life history parameters calculated for the interpopulation and intrapopulation analyses are presented in Tables 4-8 with population codes corresponding to Figures 1-3. The

number of fish included in the interpopulation analyses (pooled years) does not always equal the sum of the number of fish included in the intrapopulation data (individual years) because years with less than 10 fish or only 2 age classes or only ascending catches (e.g. if fish age 1-3 were caught and there were more age 3 than age 1 or 2 fish, mortality estimates could not be calculated) were excluded from the intrapopulation analyses. Interpopulation (years of sampling data pooled for each population) results indicated that the von Bertalanffy growth coefficients ( $k$ ) and total mortality rates ( $Z$ ) varied the least among all of the populations (Tables 4-8). Asymptotic length ( $L_{\infty}$ ) and length at maturity ( $L_{\alpha}$ ) showed the greatest range among all of the species and populations.

Growth curves (averaged for all years with sampling data) are presented for each species and population in each Great Lake (Figure 8-12). In Lake Erie, the lake whitefish populations displayed similar growth curves with populations ELWF2 and ELWF3 having nearly identical growth curves. Four of the yellow perch populations had similar growth curves, the exception being the EYP1 population. The five walleye populations in Lake Erie had similar growth curves except for EWAL1 that showed lower growth than the other 4 populations from ages 1-9 (Figure 8). Lake Ontario was treated as a single population therefore no interpopulation comparisons of the growth curves could be made for that lake (Figure 9). In Lake Huron, bloater and lake trout in the main basin, HBLOON2 and HLTON2 grew faster at younger ages than populations in the other basins but had the shortest asymptotic size of the populations (Figure 10). Walleye in Lake Huron showed a similar growth pattern for all 3 populations. Lake whitefish and yellow perch in Lake Huron showed very different growth patterns among populations (Figure 11). The shape of the growth curves for lake whitefish populations in Lake Michigan were similar but the asymptotic sizes ranged from 45 cm to 65 cm (Figure 12). The

MYPMM2 populations had higher growth rates than the MYPMM1 population. Bloater populations in MM1-6 had similar growth patterns but bloater in MM6 showed higher growth rates than the other populations (Figure 12). Lake trout in Lake Superior waters off Ontario grew slower after age 5 than the other lake trout populations (Figure 13). Lake whitefish in MI7 displayed faster growth than the other 8 populations in Lake Superior. Only one yellow perch population was available for Lake Superior (Figure 14).

### **VIRTUAL POPULATION ANALYSES**

The life history parameters varied across all of the cohorts (Table 9). Lake whitefish showed the greatest variation in life history parameters with walleye showing the least. Yellow perch had the highest mortality rates with lake whitefish cohorts in Lake Ontario having the least. The  $t_{\max}$  and  $L_{\infty}$  parameters of the lake trout and siscowet lake trout were likely underestimated using the VPA approach because none of the populations have been studied long enough to follow an individual cohort through to their maximum life span, i.e. greater than 20 years.

### **DISCUSSION**

Our findings show that life history parameters vary among populations of the same species and within populations through time. These findings are consistent with recent publications documenting variation in life history parameters of fishes such as lake trout (Shuter *et al.*, 1998), walleye (Lester *et al.*, 2000) and yellow perch (Purchase *et al.*, 2005). The spatial and temporal variation in life history traits in the Great Lakes may be due to regional factors such as climate, and local factors such as lake size. Temperature can affect the growth of fishes and may directly or indirectly affect their mortality. Variation in early growth rate and female maximum size were positively related to lake surface area and water hardness, respectively, for

yellow perch populations in Ontario (Purchase *et al.*, 2005). Density-dependent growth also explained variance in intrapopulation growth parameters of bloater in Lake Michigan (Szalai *et al.*, 2003).

This report documents the life history parameters calculated for 65 populations of five fishes inhabiting the Great Lakes, providing estimates for future analyses to understand variation in life history traits. These estimates of life history parameters can be used to inform fisheries management and ecological modelling.

### **ACKNOWLEDGEMENTS**

This study was funded by the Great Lakes Fishery Commission. Many thanks to Tim Johnson, Lloyd Mohr, Bruce Morrison, Jim Bowlby, Ted Schaner, Jay Dietrich and Ken MacIntosh (all of the OMNR), Shawn Sitar (Michigan DNR), Michael Seider (Wisconsin DNR), Mark Ebener (Chippewa Ottawa Resource Authority) and David Clapp (Michigan DNR) for providing the data used in our analyses.

## REFERENCES

- Beauchamp, K.C., Collins, N.C. and Henderson, B.A. 2004. Covariation of growth and maturation of lake whitefish (*Coregonus clupeaformis*). *Journal of Great Lakes Research* 30: 451-360.
- Beverton, R.J.H. and Holt, S.J. 1959. A review of the lifespans and mortality rates of fish in nature, and their relation to growth and other physiological characteristics. In: Ciba Foundation Colloquia on Ageing. Vol. 5. The Lifespan of Animals. Edited by: G.E.W. Wolstenholme and M. O'Connor. J & A. Churchill Ltd., London.
- Charnov, E.L. 1993. Life history invariants: some explorations of symmetry in evolutionary ecology. Oxford University Press, Oxford.
- ESRI Inc. 1996. ArcView® GIS. ESRI Inc., Redlands, California.
- Hilborn, R. and Walters, C.J. 1992. Quantitative fisheries stock assessment: Choice, dynamics and uncertainty. Chapman Hall, New York.
- Hoenig, J.M. 1983. Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin* 82: 898-903
- Lassen, H. and Medley, P. 2000. Virtual population analysis. A practical manual for stock assessment. FAO Fisheries Technical Paper. No. 400. Rome, FAO. 129p.
- Lester, N.P., Shuter, B.J., Kushneriuk, R.S. and Marshall, T.R. 2000. Life history variation in Ontario walleye populations: implications for safe rates of fishing. Ontario Ministry of Natural Resources, Peterborough.
- Pauly, D. 1980. On the interrelationship between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. *Journal du Conseil* 39: 175-192.
- Purchase, C.F., Collins, N.C., Morgan, G.E. and Shuter, B.J. 2005. Predicting life history traits of yellow perch from environmental characteristics of lakes. *Transactions of the American Fisheries Society* 134: 1369-1381.
- Roff, D.A.. 1984. The evolution of life history parameters in teleosts. *Canadian Journal of Fisheries and Aquatic Sciences* 41: 55-66.
- Shuter, B.J., Jones, M.L., Korver, R.M. and Lester, N.P. 1998. A general, life history based model for regional management of fish stocks: the inland lake trout (*Salvelinus namaycush*) fisheries of Ontario. *Canadian Journal of Fisheries and Aquatic Sciences* 55: 2161-2177.
- Szalai, E.B., Fleischer, G.W. and Bence, J.R. 2003. Modeling time-varying growth using a generalized von Bertalanffy model with application to bloater (*Coregonus hoyi*) growth



dynamics in Lake Michigan. *Canadian Journal of Fisheries and Aquatic Sciences* 60: 55-66.

Taylor, N.G., Walters, C.J. and Martell, S. J.D. 2005. A new likelihood for simultaneously estimating von Bertalanffy growth parameters, gear selectivity, and natural and fishing mortality. *Canadian Journal of Fisheries and Aquatic Sciences* 62: 215-223.

Table 1. Summary of the interpopulation data used to examine the life history parameters of fish populations in the Great Lakes Basin. Population codes correspond to Figures 1-3. Lake trout, yellow perch and bloater were aged using scales, siscowet lake trout, lake whitefish and walleye with otoliths.

Species	Lake	Populations	Number of fish	Age	TL	RWT	SEX	MAT	Source
Lake trout lean	Huron	HLTON1	2988	X	X	X	X	X	Mohr, OMNR
		HLTON2	140						
	Superior	SLTWI1	3619	X	X	X	X	X*	Seider, WIDNR; Sitar, MDNR; Ebener, CORA; MacIntosh, OMNR
		SLTWI2	13967						
		SLTMI2	469						
		SLTMI3	1230						
		SLTMI4	3203						
		SLTMI5	2448						
		SLTMI6	1611						
SLTMI7	1293								
SLTON1	10426								
Lake trout siscowet	Superior	SLTSISMI2	493	X	X	X	X	X	Sitar, MDNR; Ebener, CORA
		SLTSISMI3	2255						
		SLTSISMI4	2254						
		SLTSISMI5	1900						
		SLTSISMI6	978						
		SLTSISMI7	935						
Lake whitefish	Erie	ELWF1	14	X	X	X	X	X	Johnson, OMNR
		ELWF2	279						
		ELWF3	233						
		ELWF4	107						
		ELWF7	113						
	Ontario	OLWF1	2381	X	X	X	X		Dietrich, OMNR
		HLWFON1	6527	X	X	X	X	X	
	Huron	HLWFON2	9114						Mohr, OMNR and Ebener, CORA
		HLWFON3	351						
		HLWFCORA5	593						
		HLWFCORA6	145						
		HLWFCORA8	213						
		HLWFCORA13	210						
	Michigan	MLWFCORA9	539	X	X	X	X	X	Ebener, CORA
		MLWFCORA10	1146						
		MLWFCORA12	52						
		MLWFCORA14	59						
	Superior	SLWFMI2	131	X	X	X	X	X*	Seider, WIDNR; Sitar, MDNR; Ebener, CORA; MacIntosh, OMNR
		SLWFMI3	840						
		SLWFMI4	28						
SLWFMI5		31							
SLWFMI6		257							
SLWFMI7		1663							
SLWFMI8		79							
SLWFWI2		41							
SLWFON1		23419							

Bloater	Michigan	MBLOMM1	1768	X	X	X	X	X	Clapp, MDNR
		MBLOMM3	5370						
		MBLOMM4	4037						
		MBLOMM5	2541						
		MBLOMM6	21						
		Huron	HBLOON1	5642	X	X	X	X	
	HBLOON2	5538							
	HBLOON3	958							
Walleye	Erie	EWAL1	8439	X	X	X	X	X	Johnson, OMNR
		EWAL2	3991						
		EWAL3	1236						
		EWAL4	799						
		EWAL7	132						
	Ontario	OWAL1	3647	X	X	X	X		Dietrich, OMNR
	Huron	HWALLON1	14	X	X	X	X	X	
		HWALLON2	208						
		HWALLON3	39						
Yellow perch	Erie	EYP1	11453	X	X	X	X	X	Johnson, OMNR
		EYP2	16946						
		EYP3	14622						
		EYP4	5648						
		EYP7	2201						
	Ontario	OYP1	4140	X	X	X	X		Dietrich, OMNR
	Huron	HYPON1	1929	X	X	X	X	X	
		HYPON2	11064						
		HYPON3	874						
	Superior	SYPON1	1590	X	X	X	X		MacIntosh, OMNR
	Michigan	MYPMM1	2202	X	X	X	X	X	
MYPMM2		635							

\* no maturity data for the OMNR and WIDNR datasets

Table 2. Summary of the intrapopulation (yearly) data used to examine the life history parameters of fish populations through time. Population codes correspond to Figures 1-3.

Species	Population	Number of years	Years
Lake trout – lean	HLTON1	2	1990, 2003
Lake trout – lean	HLTON2	2	1990, 2003
Lake trout – lean	SLTWI1	14	1987-95, 19997-00, 2002
Lake trout – lean	SLTWI2	21	1981-95, 1997-00, 2002-03
Lake trout – lean	SLTMI2	8	1995-96, 1998-2003
Lake trout – lean	SLTMI3	11	1977, 1981, 1987, 1995-96, 1998-2003
Lake trout – lean	SLTMI4	18	1971-72, 1975, 1977, 1980-83, 1987, 1995-2003
Lake trout – lean	SLTMI5	14	1975, 1980-83, 1987, 1995-96, 1998-2003
Lake trout – lean	SLTMI6	15	1975, 1980-83, 1987, 1995-2003
Lake trout – lean	SLTMI7	12	1981, 1983, 1987, 1995-2003
Lake trout – lean	SLTMI8	3	1975, 1981, 1982
Lake trout – lean	SLTON1	8	1988-95
Lake trout – siscowet	SLTSISMI2	8	1995-96, 1998-03
Lake trout – siscowet	SLTSISMI3	8	1995-96, 1998-03
Lake trout – siscowet	SLTSISMI4	9	1995-03
Lake trout – siscowet	SLTSISMI5	9	1995-03
Lake trout – siscowet	SLTSISMI6	10	1980, 1983, 1995-96, 1998-03
Lake trout – siscowet	SLTSISMI7	10	1995-04
Lake whitefish	ELWF2	9	1995-03
Lake whitefish	ELWF3	6	1997-03
Lake whitefish	ELWF4	5	1989-91, 1998, 2002
Lake whitefish	ELWF7	4	1991, 1993, 1994, 1998
Lake whitefish	HLWFCORA13	2	2001-02
Lake whitefish	HLWFCORA5	7	1983, 1985-86, 2000-03
Lake whitefish	HLWFCORA6	4	2000-03
Lake whitefish	HLWFCORA8	3	1985, 2001-02
Lake whitefish	HLWFON1	13	1981-82, 1985-92, 2001-03
Lake whitefish	HLWFON2	15	1981-92, 2001-03
Lake whitefish	HLWFON3	7	1987-92, 2001, 2003
Lake whitefish	MLWFCORA9	2	2001-02
Lake whitefish	MLWFCORA10	4	2000-03
Lake whitefish	MLWFCORA14	2	2002-03
Lake whitefish	SLWFMI3	3	1995, 1998, 2003
Lake whitefish	SLWFMI6	4	1995, 1998, 2002, 2003
Lake whitefish	SLWFMI7	7	1981-84, 2001-03
Lake whitefish	SLWFMI8	3	2001-03
Lake whitefish	SLWFWI2	2	1991-92
Lake whitefish	OLWF1	13	1992-04
Lake whitefish	SLWFON1	7	1989, 1991-1996
Bloater	MBLOMM1	5	1980-82, 86, 88
Bloater	MBLOMM3	7	1980-83, 85, 86, 88
Bloater	MBLOMM4	5	1980-82, 84, 86
Bloater	MBLOMM5	7	1980-1986
Bloater	HBLOON1	6	1989, 1991-92, 2001-03
Bloater	HBLOON2	9	1986-87, 1989-92, 2001-03
Bloater	HBLOON3	6	1986-90, 1992
Walleye	EWAL1	13	1990, 1992-03
Walleye	EWAL2	13	1990, 1992-03

---

Walleye	EWAL3	11	1989-92, 1994-95, 1997-98, 2001-03
Walleye	EWAL4	11	1989-92, 1994, 1998-03
Walleye	EWAL7	9	1991-92, 1994-95, 1998-03
Walleye	HWALLON2	9	1984, 1987-91, 2001-03
Walleye	OWALL1	13	1992-04
Yellow perch	EYP1	13	1990, 1992-03
Yellow perch	EYP2	13	1990, 1992-03
Yellow perch	EYP3	14	1989-95, 1997-03
Yellow perch	EYP4	13	1989-95, 1998-03
Yellow perch	EYP7	11	1991-95, 1998-03
Yellow perch	HYPON1	8	1981, 1983, 1985, 1987-91
Yellow perch	HYPON2	15	1981-92, 01-03
Yellow perch	HYPON3	4	1988, 2001-03
Yellow perch	OYP1	13	1992-04
Yellow perch	SYPON1	6	1991-96
Yellow perch	MYPMM1	4	1986, 1996-1998
Yellow perch	MYPMM2	4	1980-1983

---

Table 3. Life history parameters calculated from age, length and maturity data from each species and population.

Life history parameter	Description
$k$	von Bertalanffy growth coefficient ( $\text{yr}^{-1}$ )
$L_{\infty}$	asymptotic length (cm)
$t_0$	age at length 0 (yr)
$L_0$	length at age 0 (cm)
$\alpha$	age at maturity (yr)
$L_{\alpha}$	length at maturity (cm)
$Z$	total mortality rate ( $\text{yr}^{-1}$ )
$t_{\text{max}}$	maximum lifespan (yr)

Table 4. Intrapopulation life history parameter values for lake whitefish (LWF), walleye (WALL) and yellow perch (YP) populations in Lake Erie. Interpopulation (years pooled) parameters are highlighted with boxes. Population codes correspond to Figures 1-3.

Species	Population	Year	Fish	k	$t_0$	$L_\infty$	$L_0$	$t_{max}$	$\alpha$	$L_\alpha$	Z
LWF	E1		14	1.06	1.57	57.31	38.30	7	4	47.00	0.35
LWF	E2	1995	28	0.49	-0.57	46.91	16.58	4	3	32.40	0.23
LWF	E2	1996	13	1.03	0.56	54.71	23.52	7	3	37.70	0.16
LWF	E2	1997	60	0.77	1.82	62.61	12.53	10	3	41.40	0.22
LWF	E2	1998	22	1.26	1.96	56.01	21.47	8	2	36.00	0.13
LWF	E2	1999	43	0.81	0.96	54.11	32.94	10	3	42.10	0.18
LWF	E2	2000	47	0.68	2.54	63.01	6.40	12	2	33.30	0.14
LWF	E2	2002	17	0.64	0.98	56.11	32.10	9	3	42.70	0.24
LWF	E2	2003	29	0.50	0.96	58.81	6.26	8	4	46.20	0.14
LWF	E2		279	0.50	1.58	63.01	14.35	12	3	32.40	0.23
LWF	E3	1997	114	0.76	1.48	60.71	15.78	8	3	35.10	0.34
LWF	E3	1998	35	0.85	1.58	55.21	4.66	9	2	33.10	0.16
LWF	E3	1999	23	0.49	-0.96	54.71	35.56	7	3	42.80	0.21
LWF	E3	2000	14	0.61	0.15	49.41	22.13	5	3	35.10	0.20
LWF	E3	2002	14	0.30	-1.14	60.01	32.37	9	3	42.50	0.21
LWF	E3	2003	10	0.26	-2.29	53.51	26.84	5	4	40.20	0.13
LWF	E3		231	0.25	-1.60	60.71	16.43	9	2	32.50	0.25
LWF	E4	1989	10	1.45	0.42	49.11	20.18	5	4	48.30	0.35
LWF	E4	1990	13	2.79	1.10	51.51	-1.12	4	2	33.70	0.31
LWF	E4	1991	13	0.73	1.31	58.21	17.73	12	3	39.50	0.05
LWF	E4	1998	41	0.37	-0.42	60.21	30.49	8	2	30.10	0.10
LWF	E4	2002	10	0.56	-0.20	51.61	28.27	6	3	37.10	0.30
LWF	E4		107	0.31	-0.98	60.21	14.08	12	2	30.10	0.19
LWF	E7	1993	21	1.17	2.36	55.31	30.34	8	4	43.40	0.21
LWF	E7	1994	13	0.51	-0.64	54.21	62.79	7	3	50.10	0.23
LWF	E7	1998	47	0.60	0.81	54.91	15.64	7	3	41.30	0.33
LWF	E7		113	0.68	1.83	60.51	15.58	11	4	42.20	0.23
WALL	E1	1990	323	0.24	-1.93	65.31	19.27	7	2	37.80	0.27
WALL	E1	1992	554	0.55	1.03	67.41	12.17	10	2	33.30	0.23
WALL	E1	1993	268	0.61	1.07	66.21	17.26	10	3	42.50	0.23
WALL	E1	1994	872	0.31	-0.55	68.51	17.21	9	2	38.00	0.26
WALL	E1	1995	1444	0.51	0.59	63.51	14.54	10	3	43.60	0.28
WALL	E1	1996	722	0.25	-0.70	72.21	17.40	12	3	40.80	0.19
WALL	E1	1997	872	0.45	0.30	69.51	22.42	9	3	40.90	0.29
WALL	E1	1998	179	0.20	-1.30	75.51	11.10	11	2	35.10	0.17
WALL	E1	1999	575	0.47	0.45	66.41	18.13	11	3	41.10	0.25
WALL	E1	2000	689	0.16	-3.95	67.51	17.10	12	2	33.20	0.22
WALL	E1	2001	418	0.43	0.82	71.21	23.42	12	2	31.00	0.19
WALL	E1	2002	455	0.25	-2.66	60.01	23.23	10	2	36.30	0.23
WALL	E1	2003	1068	0.33	0.78	72.91	11.52	15	3	42.30	0.19
WALL	E1		8439	0.15	-2.61	75.51	14.93	15	3	36.30	0.26
WALL	E2	1990	266	0.34	-0.72	68.51	17.49	7	2	37.90	0.49
WALL	E2	1992	619	0.23	-2.25	67.21	20.64	9	2	36.70	0.24
WALL	E2	1993	310	0.18	-2.42	73.61	19.69	10	2	29.50	0.20
WALL	E2	1994	271	0.22	-1.79	72.61	18.53	10	2	41.00	0.21
WALL	E2	1995	773	0.19	-2.41	72.61	16.03	13	2	37.20	0.27
WALL	E2	1996	460	0.15	-3.46	71.41	20.56	10	2	34.50	0.21
WALL	E2	1997	377	0.40	-0.46	63.81	18.87	10	2	40.50	0.19
WALL	E2	1998	71	0.68	0.66	62.21	23.84	9	1	34.20	0.16
WALL	E2	1999	77	0.32	-0.75	70.71	19.45	8	2	40.00	0.09
WALL	E2	2000	319	0.27	-0.80	72.61	17.23	11	2	35.00	0.19
WALL	E2	2001	53	0.33	-2.26	59.51	25.78	8	2	40.20	0.17

WALL	E2	2002	160	0.29	-2.26	63.91	23.91	8	2	40.80	0.22
WALL	E2	2003	234	0.55	1.98	70.21	8.35	13	2	38.40	0.15
WALL	E2		3991	0.18	-2.38	73.61	16.98	13	2	29.50	0.26
WALL	E3	1989	333	0.68	1.68	71.01	10.34	10	2	38.20	0.29
WALL	E3	1990	68	0.55	1.98	70.21	8.35	13	2	38.40	0.15
WALL	E3	1991	82	0.33	-1.13	65.71	17.35	7	1	33.10	0.42
WALL	E3	1992	129	0.70	1.23	65.01	17.58	9	1	32.20	0.39
WALL	E3	1994	133	0.32	-0.59	69.81	16.91	10	2	34.20	0.42
WALL	E3	1995	146	0.17	-2.79	74.21	16.37	11	2	39.80	0.26
WALL	E3	1997	72	0.23	-1.04	75.61	13.02	12	3	44.20	0.10
WALL	E3	1998	26	0.40	0.40	72.71	35.46	11	2	38.80	0.08
WALL	E3	2001	24	0.19	-2.71	74.01	51.51	10	1	35.30	0.06
WALL	E3	2002	90	0.16	-4.34	74.51	22.73	14	2	40.50	0.05
WALL	E3	2003	61	1.33	1.63	62.51	5.91	8	2	41.30	0.19
WALL	E3		1236	0.19	-1.72	75.61	15.66	15	2	34.20	0.20
WALL	E4	1989	150	0.22	-2.90	67.21	23.87	9	3	41.50	0.24
WALL	E4	1990	48	0.32	-1.20	63.71	12.77	7	2	40.50	0.24
WALL	E4	1991	23	0.28	-1.21	66.81	37.87	7	3	47.10	0.10
WALL	E4	1992	12	0.45	-0.01	66.51	32.46	6	2	40.00	0.09
WALL	E4	1994	21	0.39	-0.11	70.71	25.61	8	3	46.40	0.39
WALL	E4	1998	25	0.51	1.99	69.91	12.83	13	3	46.20	0.10
WALL	E4	1999	89	0.16	-3.80	73.31	17.46	11	2	36.00	0.20
WALL	E4	2000	104	0.26	-0.60	71.81	22.68	13	2	40.00	0.18
WALL	E4	2001	117	0.15	-3.31	75.61	14.04	13	2	38.00	0.01
WALL	E4	2002	96	0.16	-2.93	74.11	16.44	14	2	42.00	0.12
WALL	E4	2003	102	0.22	-1.44	72.81	21.03	15	2	39.50	0.15
WALL	E4		799	0.14	-4.01	75.61	15.50	15	2	36.00	0.15
WALL	E7	1992	13	0.48	-0.69	61.31	12.11	7	1	42.40	0.11
WALL	E7	1994	14	0.65	0.84	65.71	15.60	8	3	48.40	0.07
WALL	E7	1995	16	1.01	1.57	68.21	14.67	9	1	32.70	0.08
WALL	E7	1998	17	0.96	1.89	67.21	-2.24	8	3	45.80	0.12
WALL	E7	1999	16	0.30	-1.90	70.01	31.09	12	2	40.00	0.06
WALL	E7	2001	13	0.40	0.33	69.31	51.51	10	2	41.60	0.06
WALL	E7	2003	16	0.29	-0.72	69.71	38.80	11	4	55.60	0.03
WALL	E7		132	0.17	-3.01	73.01	16.17	12	3	43.00	0.13
YP	E1	1990	508	0.27	-0.56	33.01	6.25	9	3	15.90	0.42
YP	E1	1992	799	0.38	0.65	34.81	0.67	8	1	12.70	0.40
YP	E1	1993	684	0.61	2.11	37.11	6.17	10	2	16.40	0.33
YP	E1	1994	1576	0.60	1.70	36.41	7.78	9	2	14.50	0.40
YP	E1	1995	1789	0.58	1.08	33.81	6.93	9	2	14.10	0.38
YP	E1	1996	845	0.65	1.84	34.71	7.77	10	2	13.90	0.36
YP	E1	1997	685	0.62	1.53	31.71	4.59	10	2	13.60	0.34
YP	E1	1998	622	0.34	0.20	33.71	1.60	7	1	12.90	0.49
YP	E1	1999	716	0.30	0.27	34.51	2.51	8	3	14.70	0.47
YP	E1	2000	773	0.33	-0.20	29.51	4.21	8	2	13.50	0.49
YP	E1	2001	831	0.26	-1.11	30.31	2.39	8	2	14.00	0.36
YP	E1	2002	759	0.20	-0.90	36.01	4.40	9	2	13.40	0.42
YP	E1	2003	866	0.17	-2.12	34.01	4.55	9	2	14.30	0.35
YP	E1		11453	0.20	-0.83	37.11	4.86	10	2	12.90	0.40
YP	E2	1990	282	0.90	1.53	34.41	4.62	7	2	17.20	0.20
YP	E2	1992	1267	0.28	-0.54	36.01	5.27	9	1	13.70	0.34
YP	E2	1993	734	0.25	-1.29	35.01	9.41	9	2	13.00	0.34
YP	E2	1994	1387	0.44	0.31	33.81	7.25	9	2	13.80	0.39
YP	E2	1995	1523	0.96	1.63	33.61	3.15	7	2	13.20	0.50
YP	E2	1996	1404	0.62	1.87	37.31	5.72	12	2	13.80	0.28
YP	E2	1997	1343	0.37	-0.20	30.51	5.98	6	2	12.30	0.69
YP	E2	1998	973	0.62	0.52	30.71	1.34	5	1	12.00	0.87
YP	E2	1999	1383	0.36	-0.86	28.41	4.59	5	1	11.50	1.29
YP	E2	2000	1609	0.25	-1.02	34.81	6.29	7	1	11.30	0.53
YP	E2	2001	1657	0.70	1.36	33.21	5.16	8	1	15.70	0.50



YP	E2	2002	1709	0.24	-1.86	33.51	10.67	7	1	12.40	0.62
YP	E2	2003	1675	0.21	-1.77	35.51	9.27	9	2	14.90	0.47
YP	E2		16964	0.53	1.98	37.31	5.18	12	1	11.00	0.42
YP	E3	1989	1389	0.21	-1.46	35.51	11.42	8	2	15.00	0.88
YP	E3	1990	228	0.32	-0.55	32.31	4.75	6	2	14.40	0.07
YP	E3	1991	763	0.32	-0.23	34.41	6.83	8	1	12.20	0.30
YP	E3	1992	853	0.30	-0.63	34.51	6.85	9	1	12.30	0.29
YP	E3	1993	89	0.96	-0.26	19.81	16.39	3			0.56
YP	E3	1994	606	0.25	-1.67	34.11	8.95	11	2	12.70	0.20
YP	E3	1995	890	0.64	1.48	35.41	6.50	9	2	13.70	0.35
YP	E3	1997	1163	0.27	-0.54	33.81	6.37	10	2	12.60	0.37
YP	E3	1998	957	0.53	0.77	33.21	5.88	7	1	12.00	0.55
YP	E3	1999	1718	0.26	-0.94	35.51	7.02	11	1	11.90	0.37
YP	E3	2000	1539	0.33	-0.05	34.51	5.37	8	1	11.50	0.62
YP	E3	2001	1447	0.32	-0.45	34.11	7.38	9	1	14.90	0.47
YP	E3	2002	1588	0.21	-1.80	35.71	10.40	8	1	11.60	0.44
YP	E3	2003	1391	0.24	-1.15	34.01	8.20	9	1	12.00	0.47
YP	E3		14621	0.20	-1.90	35.71	6.63	11	2	12.20	0.42
YP	E4	1989	424	0.28	-0.43	34.11	7.93	9	2	14.50	0.39
YP	E4	1990	255	0.48	1.37	35.71	1.42	9	3	16.00	0.38
YP	E4	1991	231	0.33	0.37	34.61	3.79	8	2	13.30	0.20
YP	E4	1992	103	0.37	0.59	34.61	5.71	9	3	17.30	0.21
YP	E4	1993	14	1.11	1.80	33.01	-70.35	9			0.12
YP	E4	1994	12	0.85	1.35	34.21	15.05	9	2	23.50	0.03
YP	E4	1995	34	2.08	1.11	25.51	-2.90	4	2	16.00	0.49
YP	E4	1998	169	0.59	2.61	37.81	6.00	12	2	12.90	0.19
YP	E4	1999	475	0.52	1.56	34.61	6.76	11	2	14.00	0.27
YP	E4	2000	756	0.23	-1.29	35.51	9.01	10	2	13.50	0.28
YP	E4	2001	979	0.24	-0.78	35.61	7.80	8	2	12.20	0.49
YP	E4	2002	974	0.61	1.33	33.61	7.30	9	2	14.70	0.49
YP	E4	2003	1222	0.28	-0.39	35.01	5.85	8	1	12.70	0.53
YP	E4		5648	0.41	1.71	37.81	5.11	12	2	12.20	0.40
YP	E7	1991	58	0.72	1.12	30.71	6.70	7	5	25.30	0.28
YP	E7	1992	15	0.36	-3.06	22.51	21.45	4	2	17.20	0.18
YP	E7	1993	64	1.16	0.73	22.51	10.15	5			0.42
YP	E7	1994	59	0.47	0.87	34.21	1.67	8	2	15.90	0.19
YP	E7	1995	133	0.75	1.31	32.71	6.80	11	1	14.80	0.19
YP	E7	1998	226	0.52	1.32	34.51	7.94	9	3	14.90	0.31
YP	E7	1999	238	0.79	1.69	31.71	5.34	9	3	14.40	0.27
YP	E7	2000	342	0.19	-2.08	35.31	12.40	8	2	14.60	0.39
YP	E7	2001	352	0.17	-2.92	33.31	10.31	8	2	16.00	0.46
YP	E7	2002	434	0.26	-0.85	33.01	7.74	8	2	17.10	0.41
YP	E7	2003	280	0.31	0.20	35.51	6.28	11	1	12.30	0.31
YP	E7		2201	0.25	-0.54	35.51	5.84	11	2	13.50	0.36

Table 5. Life history parameter values for lake whitefish (LWF), smallmouth bass (SMB), walleye (WALL) and yellow perch (YP) populations in Lake Ontario. Interpopulation (years pooled) parameters are highlighted with boxes. Maturity data were not available for these populations. Population codes correspond to Figures 1-3.

Species	Population	Year	Fish	k	$t_0$	$L_\infty$	$L_0$	$t_{max}$	$\alpha$	$L_\alpha$	Z
LWF	O1	1992	369	0.21	-2.19	64.71	10.96	16			0.17
LWF	O1	1993	439	0.14	-4.62	68.21	15.72	19			0.13
LWF	O1	1994	286	0.50	2.06	65.41	12.80	16			0.16
LWF	O1	1995	343	0.21	-1.37	70.31	9.15	17			0.14
LWF	O1	1996	123	0.52	2.04	61.61	9.18	11			0.23
LWF	O1	1997	120	0.44	2.74	63.71	7.30	17			0.11
LWF	O1	1998	141	0.49	2.19	61.01	5.55	11			0.33
LWF	O1	1999	94	0.22	-0.36	59.51	8.74	14			0.15
LWF	O1	2000	85	0.31	2.35	64.41	7.56	17			0.12
LWF	O1	2001	123	0.24	1.19	63.91	11.14	15			0.22
LWF	O1	2002	76	0.22	1.48	62.61	12.49	16			0.12
LWF	O1	2003	58	0.18	-0.01	63.21	10.12	17			0.19
LWF	O1	2004	124	0.10	-4.64	68.91	25.11	21			0.15
LWF	O1		2381	0.08	-6.38	70.31	10.36	21			0.14
WALL	O1	1992	455	0.22	0.20	77.71	11.68	15			0.26
WALL	O1	1993	168	0.23	-1.28	63.71	14.15	10			0.15
WALL	O1	1994	167	0.16	-2.04	74.61	11.45	15			0.14
WALL	O1	1995	159	0.35	2.00	79.31	11.59	14			0.14
WALL	O1	1996	328	0.09	-6.47	77.31	42.25	18			0.11
WALL	O1	1997	585	0.20	0.07	77.91	13.99	19			0.14
WALL	O1	1998	494	0.18	-1.23	76.71	13.24	20			0.11
WALL	O1	1999	373	0.08	-7.68	81.41	12.52	22			0.09
WALL	O1	2000	220	0.08	-11.15	76.21	19.32	22			0.09
WALL	O1	2001	175	0.11	-4.59	77.41	16.60	19			0.12
WALL	O1	2002	272	0.07	-10.35	80.01	17.41	21			0.06
WALL	O1	2003	353	0.07	-9.84	79.81	22.61	20			0.20
WALL	O1	2004	392	0.06	-13.18	80.11	22.86	22			0.14
WALL	O1		4141	0.07	-8.92	81.41	14.71	22			0.11
YP	O1	1992	232	0.19	-2.15	28.41	5.94	8			0.39
YP	O1	1993	378	0.19	-1.84	28.31	3.78	8			0.36
YP	O1	1994	292	1.03	2.49	27.61	-10.79	8			0.43
YP	O1	1995	216	0.15	-3.70	27.31	4.43	8			0.47
YP	O1	1996	208	0.09	-7.03	30.51	8.95	7			0.20
YP	O1	1997	281	0.22	-1.45	27.61	2.82	8			0.36
YP	O1	1998	387	0.27	-0.44	29.31	6.06	8			0.46
YP	O1	1999	494	0.57	1.73	28.61	2.52	10			0.33
YP	O1	2000	519	0.21	-1.29	30.01	1.73	8			0.48
YP	O1	2001	324	0.21	-3.14	25.31	4.89	8			0.35
YP	O1	2002	264	0.26	-1.42	26.41	2.03	6			0.43
YP	O1	2003	250	0.25	-2.32	25.41	8.54	7			0.42
YP	O1	2004	393	0.46	1.08	31.01	10.95	9			0.40
YP	O1		4238	0.18	-1.32	31.01	3.06	10			0.50*

Table 6. Life history parameter values for bloater (BLO), lean lake trout (LT), lake whitefish (LWF), walleye (WALL) and yellow perch (YP) populations in Lake Huron. Interpopulation (years pooled) parameters are highlighted with boxes. Maturity data were not available for all of the populations. Population codes correspond to Figures 1-3.

Species	Population	Year	Fish	k	t <sub>0</sub>	L <sub>∞</sub>	L <sub>0</sub>	t <sub>max</sub>	α	L <sub>α</sub>	Z
BLO	HON1	1989	1231	0.13	-2.68	41.41	1.74	12	2	20.20	0.49
BLO	HON1	1991	1028	0.10	-5.36	37.41	6.67	10	2	17.80	0.64
BLO	HON1	2001	569	0.12	-4.49	35.51	9.84	11	2	17.20	0.44
BLO	HON1	2002	686	0.15	-2.63	34.81	1.76	11	2	17.20	0.43
BLO	HON1	2003	986	0.11	-6.84	32.21	6.69	10	2	16.70	0.30
BLO	HON1		5642	0.11	-3.20	41.41	5.85	12	2	16.70	0.59
BLO	HON2	1986	231	0.29	-1.82	27.51	6.71	7	2	14.60	0.36
BLO	HON2	1987	151	0.24	-1.42	31.11	3.86	7	2	18.20	0.42
BLO	HON2	1989	1651	0.11	-4.58	36.31	7.66	9	2	17.40	0.60
BLO	HON2	1990	319	0.16	-4.19	32.01	7.54	9	2	17.60	0.48
BLO	HON2	1991	1341	0.13	-3.44	37.11	4.56	10	2	19.20	0.75
BLO	HON2	1992	977	0.09	-7.54	36.51	1.51	10	1	24.30	0.52
BLO	HON2	2001	268	0.25	-1.19	32.81	6.01	11	2	18.10	0.25
BLO	HON2	2002	163	0.51	0.85	29.71	2.26	10	2	16.50	0.19
BLO	HON2	2003	437	0.12	-6.97	31.51	6.47	11	3	19.10	0.35
BLO	HON2		5538	0.10	-5.04	37.11	6.41	11	2	14.60	0.49
BLO	HON3	1986	441	0.39	2.05	42.31	3.51	12	2	15.90	0.25
BLO	HON3	1987	178	0.19	-1.92	35.11	4.83	10	2	18.20	0.34
BLO	HON3	1988	121	0.61	1.13	32.91	2.81	8	2	16.70	0.58
BLO	HON3	1989	102	0.14	-7.83	30.91	31.46	7	2	28.40	0.40
BLO	HON3	1990	84	0.24	-0.73	32.71	6.56	8	2	13.80	0.27
BLO	HON3	1992	29	0.59	1.05	45.71	25.15	9	2	30.60	0.11
BLO	HON3		958	0.15	-0.76	45.71	3.25	12	2	13.80	0.35
LT	HON1	1990	1117	0.47	0.87	72.21	22.73	5	4	42.90	0.28
LT	HON1	2003	431	0.35	0.95	83.51	15.12	9	4	45.80	0.38
LT	HON1		2988	0.34	0.69	83.51	15.70	9	4	42.90	0.55*
LT	HON2	1990	15	0.77	1.55	70.51	25.42	6	5	65.40	0.17
LT	HON2	2003	113	0.45	1.38	72.01	26.46	9	5	54.30	0.22
LT	HON2		140	0.37	0.07	72.01	12.32	9	5	54.30	0.26
LWF	CORA13	2001	144	0.55	2.61	60.71	6.75	11	6	38.60	0.23
LWF	CORA13	2002	66	0.26	0.60	57.01	7.56	12	7	41.10	0.20
LWF	CORA13		210	0.24	0.58	60.71	8.67	12	7	41.10	0.22
LWF	CORA5	1983	270	0.50	2.43	65.01	9.57	10	4	43.40	0.46
LWF	CORA5	1985	56	1.08	5.95	70.81	36.13	12	6	45.00	0.36
LWF	CORA5	1986	45	0.32	0.82	64.51	8.54	9	4	43.80	0.17
LWF	CORA5	2000	30	0.78	3.20	60.91	-1.83	10	7	46.10	0.20
LWF	CORA5	2001	75	0.25	-0.18	52.01	6.92	9	7	43.50	0.42
LWF	CORA5	2002	82	0.18	-1.16	55.31	5.47	13	6	40.50	0.22
LWF	CORA5	2003	35	0.38	2.14	54.91	7.37	10	7	40.00	0.30
LWF	CORA5		593	0.32	2.04	70.81	22.79	13	5	36.50	0.38
LWF	CORA6	2000	15	0.21	-0.27	52.11	7.33	9	7	41.20	0.16
LWF	CORA6	2001	58	0.16	-1.79	54.81	9.01	11	7	40.20	0.17
LWF	CORA6	2002	34	0.20	-0.45	52.51	11.32	11	8	35.30	0.26
LWF	CORA6	2003	38	0.20	-0.37	55.51	8.19	10	6	42.60	0.18
LWF	CORA6		145	0.16	-1.46	55.51	8.11	11	7	40.20	0.32
LWF	CORA8	1985	69	0.26	1.08	70.31	16.32	11	5	45.50	0.47
LWF	CORA8	2001	107	0.27	0.32	57.71	8.00	10	6	40.70	0.42
LWF	CORA8	2002	37	0.34	2.80	64.61	11.91	11	6	40.20	0.09

LWF	CORAB		213	0.22	0.92	70.31	6.35	11	6	40.20	0.39
LWF	HON1	1981	52	1.77	1.47	59.51	11.67	5	4	50.60	0.71
LWF	HON1	1982	101	1.41	2.56	73.21	2.82	8	4	52.00	0.39
LWF	HON1	1985	261	0.52	2.13	78.51	15.33	15	5	53.40	0.17
LWF	HON1	1986	158	0.65	2.08	73.81	14.15	13	4	42.00	0.18
LWF	HON1	1987	196	0.40	0.12	68.21	17.20	9	4	49.90	0.45
LWF	HON1	1988	427	0.35	-0.06	69.51	16.84	9	5	52.40	0.45
LWF	HON1	1989	580	0.25	-1.12	71.61	18.42	11	4	47.40	0.43
LWF	HON1	1990	767	0.30	-0.35	68.91	20.29	9	4	49.30	0.35
LWF	HON1	1991	913	0.33	0.36	72.91	15.79	12	4	44.60	0.24
LWF	HON1	1992	860	0.47	1.45	75.51	9.53	12	5	51.70	0.30
LWF	HON1	2001	602	0.26	-0.76	65.01	14.44	8	6	53.00	0.37
LWF	HON1	2002	487	1.08	1.57	57.11	-2.07	9	9	57.10	0.37
LWF	HON1	2003	332	1.15	1.70	54.51	6.99	8	5	47.20	0.39
LWF	HON1		6530	0.35	1.48	78.51	12.74	15	4	18.00	0.30
LWF	HON2	1981	131	0.70	0.44	50.31	18.70	4	4	42.60	0.28
LWF	HON2	1982	154	0.90	2.04	68.71	7.79	10	4	44.20	0.23
LWF	HON2	1983	523	0.44	0.31	62.91	12.35	6	4	44.10	0.51
LWF	HON2	1984	338	0.58	0.39	55.71	11.74	5			0.49
LWF	HON2	1985	626	0.27	-1.34	60.61	18.31	7	3	35.00	0.81
LWF	HON2	1986	399	0.42	-0.26	61.01	13.48	6	4	44.80	0.34
LWF	HON2	1987	623	0.72	1.07	61.11	12.25	7	5	49.70	0.36
LWF	HON2	1988	1414	0.37	-0.31	62.61	20.06	7	4	45.10	0.43
LWF	HON2	1989	1238	0.82	1.75	62.31	11.47	10	4	42.00	0.38
LWF	HON2	1990	954	0.38	-0.21	59.41	17.42	8	4	40.40	0.43
LWF	HON2	1991	911	0.37	0.39	65.21	16.69	10	4	37.80	0.33
LWF	HON2	1992	778	0.30	-0.01	65.71	10.98	10	5	43.20	0.36
LWF	HON2	2001	216	0.21	-0.47	65.11	9.70	9	6	45.00	0.20
LWF	HON2	2002	263	0.17	-1.50	63.21	11.79	11	6	47.10	0.18
LWF	HON2	2003	547	0.21	-0.74	59.71	8.86	11	6	41.80	0.19
LWF	HON2		9115	0.12	-3.81	68.71	15.26	11	4	33.00	0.32
LWF	HON3	1988	96	0.53	1.92	73.41	14.78	16	4	42.40	0.11
LWF	HON3	1989	52	0.82	1.86	66.81	27.14	9	3	43.80	0.20
LWF	HON3	1990	71	0.66	1.37	62.31	13.43	12	3	37.20	0.16
LWF	HON3	1992	17	0.57	0.42	58.41	17.53	5	4	50.00	0.15
LWF	HON3	2001	83	0.16	-2.76	62.01	15.36	11	3	36.50	0.24
LWF	HON3	2003	21	0.89	2.57	53.11	11.98	10	6	45.00	0.12
LWF	HON3		351	0.37	2.44	73.41	11.97	16	4	41.30	0.17
WALL	HON1	1983	14	2.34	1.12	46.51	20.53	4	2	37.20	0.29
WALL	HON2	1984	11	1.25	1.81	57.31	43.48	7	4	51.80	0.17
WALL	HON2	1987	18	1.19	1.26	60.11	-3.20	7	2	37.30	0.21
WALL	HON2	1988	36	0.55	0.67	60.11	13.41	8	2	36.30	0.15
WALL	HON2	1989	23	0.17	-6.53	58.31	34.17	8	4	41.30	0.17
WALL	HON2	1990	25	0.09	-9.74	71.51	52.84	13	4	43.70	0.10
WALL	HON2	1991	33	0.14	-5.32	68.41	18.50	13	3	45.20	0.15
WALL	HON2	2001	17	0.23	-0.93	73.41	40.30	9	4	53.40	0.20
WALL	HON2	2002	18	0.94	3.11	67.71	-9.01	10	3	42.40	0.10
WALL	HON2	2003	19	0.71	2.56	70.21	16.68	11	4	43.50	0.15
WALL	HON2		208	0.11	-5.64	73.41	14.41	13	2	34.30	0.15
WALL	HON3	1998	39	0.67	4.88	78.71	8.30	13	5	60.90	0.35
YP	HON1	1981	230	1.12	2.09	28.91	8.23	7			0.42
YP	HON1	1983	73	1.08	2.31	27.71	-0.92	7	2	15.40	0.61
YP	HON1	1985	200	0.32	-0.71	31.01	14.44	6	6	25.60	0.23
YP	HON1	1987	144	0.53	2.02	35.51	9.64	13	4	18.30	0.18
YP	HON1	1988	323	0.28	-1.28	30.11	7.74	6	5	19.40	0.11
YP	HON1	1989	160	0.20	-1.88	33.01	10.69	7			0.30
YP	HON1	1990	54	0.33	-0.61	28.91	7.32	7	3	18.30	0.25
YP	HON1	1991	15	0.30	-0.57	32.41	13.01	4	2	17.20	0.30

YP	HON1		1930	0.50	2.06	35.51	2.55	13	4	17.70	0.31
YP	HON2	1981	13	0.96	3.16	35.31	7.33	9	3	19.80	0.17
YP	HON2	1982	213	0.75	1.33	34.41	5.84	8	1	13.80	0.35
YP	HON2	1983	204	0.49	0.28	32.51	8.66	9	2	15.10	0.31
YP	HON2	1984	465	0.49	0.06	34.41	6.57	10	2	14.10	0.32
YP	HON2	1985	755	0.19	-1.14	38.71	7.10	10	4	15.00	0.29
YP	HON2	1986	1220	0.69	1.79	34.71	9.13	10	2	14.40	0.40
YP	HON2	1987	1261	0.53	2.06	36.41	2.64	11	1	12.50	0.39
YP	HON2	1988	971	0.01	-6.94	32.31	7.45	11	2	11.60	0.30
YP	HON2	1989	1484	0.30	-0.26	36.71	7.63	12	1	11.10	0.35
YP	HON2	1990	1530	0.36	0.73	37.21	8.65	12	2	13.60	0.39
YP	HON2	1991	994	0.33	0.46	37.41	5.23	11	1	14.00	0.38
YP	HON2	1992	542	0.30	0.24	38.01	4.89	10	1	13.40	0.31
YP	HON2	2001	47	0.44	-0.44	26.31	4.05	4	1	13.30	0.28
YP	HON2	2002	987	0.62	1.43	35.01	3.72	10	2	13.50	0.41
YP	HON2	2003	378	0.10	-4.11	36.81	-6.29	6	1	14.30	1.23
YP	HON2		11064	0.14	-1.99	41.41	7.82	12	1	11.10	0.37
YP	HON3	1988	13	0.13	-6.06	28.61	22.28	7	3	22.20	0.45
YP	HON3	2001	551	0.31	-1.36	32.61	-0.82	6	0	17.80	0.67
YP	HON3	2002	193	0.45	0.30	31.21	8.68	5	3	16.80	0.70
YP	HON3	2003	95	0.35	0.33	30.61	9.40	7	3	13.50	0.45
YP	HON3		874	0.14	-4.86	32.61	4.50	8	0	17.80	0.52

\* mortality estimates for interpopulation are greater than intrapopulation because the sum of young fish (age 1-3) in each age class, steepened the catch curve used to calculate mortality.

Table 7. Life history parameter values for bloater (BLO), lake whitefish (LWF) and yellow perch (YP) populations in Lake Michigan. Interpopulation (years pooled) parameters are highlighted with boxes. Maturity data were not available for all of the populations. Population codes correspond to Figures 1-3.

Species	Population	Year	Fish	k	$t_0$	$L_\infty$	$L_0$	$t_{max}$	$\alpha$	$L_\alpha$	Z
BLO	MM1	1980	576	0.19	0.10	44.01	2.54	12	3	15.60	0.37
BLO	MM1	1981	351	0.68	1.40	33.31	4.43	8	3	15.60	0.46
BLO	MM1	1982	202	0.99	2.02	36.01	-2.55	9	4	18.40	0.13
BLO	MM1	1986	383	1.18	2.32	31.41	8.25	7	3	14.30	0.67
BLO	MM1	1988	256	0.38	-0.75	30.01	8.50	7	3	22.00	1.06
BLO	MM1		1768	0.18	-0.05	44.01	5.26	12	3	14.30	0.30
BLO	MM3	1980	2089	0.08	-4.39	49.01	7.04	12	2	15.50	0.34
BLO	MM3	1981	529	0.68	0.69	32.51	6.11	5	2	15.60	0.77
BLO	MM3	1982	487	1.01	1.70	33.21	4.78	7	3	16.50	0.74
BLO	MM3	1983	201	0.66	2.12	35.51	4.56	8	3	14.20	0.49
BLO	MM3	1985	74	3.26	0.99	19.51	-31.58	3			1.65
BLO	MM3	1986	1191	0.59	1.10	31.71	6.27	7	3	15.70	0.41
BLO	MM3	1988	582	0.87	3.31	37.11	-4.77	10	3	22.80	0.55
BLO	MM3		5370	0.09	-3.45	49.01	6.89	12	3	14.20	0.38
BLO	MM4	1980	404	0.15	-1.76	42.01	4.27	11	2	13.80	0.42
BLO	MM4	1981	1465	0.75	2.29	46.01	3.74	10	2	14.80	0.42
BLO	MM4	1982	478	0.24	-0.90	37.01	0.70	4	2	14.50	0.86
BLO	MM4	1984	1339	0.54	0.43	31.21	6.55	6	2	14.00	0.69
BLO	MM4	1986	199	0.61	1.44	32.51	7.08	10	4	22.70	0.38
BLO	MM4		4037	0.12	-2.24	46.01	7.08	11	2	13.80	0.38
BLO	MM5	1980	100	0.38	0.16	34.51	12.23	6	2	14.50	0.35
BLO	MM5	1981	452	0.30	-0.05	35.01	3.91	7	3	16.00	0.18
BLO	MM5	1982	296	0.28	-0.01	36.51	8.02	10	3	17.20	0.35
BLO	MM5	1983	143	0.24	-3.04	25.01	10.96	6	2	15.50	0.64
BLO	MM5	1984	154	0.38	-0.51	27.01	0.04	5	1	14.20	0.51
BLO	MM5	1985	1196	0.62	0.20	24.81	8.14	4	2	14.20	0.68
BLO	MM5	1986	200	0.45	0.03	31.01	-0.83	7	3	23.30	0.73
BLO	MM5		2541	0.27	-0.19	36.51	5.74	10	2	13.80	0.39
BLO	MM6		21	0.84	1.85	44.61	6.93	9	5	35.50	0.13
LWF	CORA10	2000	114	0.74	2.49	52.51	5.91	9	6	37.10	0.40
LWF	CORA10	2001	218	0.20	-1.74	53.31	9.48	13	6	35.30	0.25
LWF	CORA10	2002	148	0.45	3.06	63.71	3.46	12	6	35.30	0.27
LWF	CORA10	2003	51	0.16	-1.95	56.51	11.70	13	4	39.80	0.14
LWF	CORA10		531	0.12	-2.35	63.71	10.79	13	6	35.30	0.28
LWF	CORA12	1991	51	0.08	-11.51	67.31	9.08	12	4	44.20	0.32
LWF	CORA14	2002	31	0.24	-0.26	51.41	13.08	10	7	41.50	0.20
LWF	CORA14	2003	21	1.09	2.63	46.91	54.78	8	8	46.90	0.22
LWF	CORA14		52	0.25	-0.30	51.41	9.51	10	7	41.50	0.23
LWF	CORA9	2001	41	0.32	-0.11	56.51	17.01	7	4	46.10	0.70
LWF	CORA9	2002	12	0.93	3.29	49.61	11.33	11	8	46.80	0.11
LWF	CORA9		58	0.16	-3.55	56.51	19.89	11	4	46.10	0.24
YP	MM1	1986	276	0.47	1.12	25.01	1.42	5	2	10.00	0.39
YP	MM1	1996	1324	0.23	0.89	34.31	6.23	13	5	18.80	0.32
YP	MM1	1997	340	0.08	-5.98	34.81	7.62	14			0.17
YP	MM1	1998	257	0.08	-7.52	34.51	8.89	19			0.17
YP	MM1		2202	0.09	-4.60	34.81	7.10	19	3	10.30	0.24

YP	MM2	1980	114	0.52	0.67	25.51	-0.04	5	2	12.50	0.72
YP	MM2	1981	106	1.18	1.60	29.31	-0.31	6	1	10.70	0.59
YP	MM2	1982	276	0.22	-1.64	27.31	6.11	5	2	13.60	0.25
YP	MM2	1983	139	0.83	2.04	37.11	-0.78	8	4	18.70	0.18
YP	MM2		635	0.76	2.10	37.11	1.19	8	1	8.50	0.46

---

Table 8. Life history parameters for lake whitefish (LWF), lean lake trout (LT), siscowet lake trout (LT-SIS) and yellow perch (YP) populations in Lake Superior. Interpopulation (years pooled) parameters are highlighted with boxes. Maturity data were not available for all of the populations. Population codes correspond to Figures 1-3.

Species	Population	Year	Fish	k	t <sub>0</sub>	L <sub>∞</sub>	L <sub>0</sub>	t <sub>max</sub>	α	L <sub>α</sub>	Z
LT	MI2	1995	82	0.30	1.87	67.81	4.28	10			0.21
LT	MI2	1996	78	0.17	-0.50	69.81	-0.03	8			0.35
LT	MI2	1998	96	0.21	1.22	71.01	2.75	13	11	59.80	0.18
LT	MI2	1999	78	0.11	-2.88	69.31	26.16	9			0.30
LT	MI2	2000	87	0.31	0.81	59.91	12.09	9			0.51
LT	MI2	2001	33	0.91	3.96	71.11	9.71	11	11	71.10	0.20
LT	MI2	2002	72	0.20	0.48	73.51	9.47	12			0.11
LT	MI2	2003	80	0.23	1.50	70.71	4.00	12	11	62.60	0.17
LT	MI2		606	0.18	0.46	73.51	6.06	13	11	59.80	0.26
LT	MI3	1977	425	0.19	0.99	89.21	7.49	12			0.25
LT	MI3	1981	397	0.14	-2.11	85.11	0.91	12	12	71.10	0.22
LT	MI3	1987	39	0.35	2.79	75.41	16.46	11	9	61.70	0.39
LT	MI3	1995	63	0.32	2.02	64.01	-13.99	13			0.12
LT	MI3	1996	87	0.49	2.17	66.51	5.76	11	9	66.50	0.27
LT	MI3	1998	121	0.13	-2.17	61.51	8.06	15			0.15
LT	MI3	1999	105	0.06	-3.04	91.51	22.24	19	19	65.50	0.11
LT	MI3	2000	114	0.21	1.59	68.91	15.58	20	17	68.90	0.10
LT	MI3	2001	96	0.28	1.46	66.81	8.79	11	9	64.90	0.24
LT	MI3	2002	110	0.30	1.47	66.41	1.99	10			0.34
LT	MI3	2003	89	0.37	4.44	91.01	-12.04	24	24	91.00	0.07
LT	MI3		1672	0.11	-0.07	91.51	17.01	24	17	68.90	0.18
LT	MI4	1971	141	0.32	1.93	77.01	-23.17	10	9	56.40	0.31
LT	MI4	1972	151	0.11	-8.66	80.81	-30.09	11	6	64.00	0.31
LT	MI4	1975	111	0.15	-1.88	80.01	8.41	11	9	62.20	0.40
LT	MI4	1977	529	0.14	-1.64	87.11	9.30	13	9	61.00	0.56
LT	MI4	1980	220	0.15	-0.59	91.41	12.80	13	8	61.70	0.60
LT	MI4	1981	198	0.31	2.76	85.61	8.88	14	8	61.00	0.42
LT	MI4	1982	119	0.45	5.26	91.41	11.07	16	8	61.00	0.34
LT	MI4	1983	427	0.17	1.47	95.81	15.54	15	8	51.60	0.22
LT	MI4	1987	82	0.99	5.54	74.91	8.44	12	9	57.20	0.29
LT	MI4	1995	257	0.63	2.68	57.21	9.85	10			0.48
LT	MI4	1996	328	0.14	-1.33	68.61	12.45	13	11	63.20	0.32
LT	MI4	1997	366	0.13	-0.78	85.11	9.49	17	15	67.30	0.19
LT	MI4	1998	360	0.11	-2.20	69.61	28.38	17	14	66.50	0.18
LT	MI4	1999	322	0.12	-3.04	65.61	16.58	15			0.22
LT	MI4	2000	445	0.39	3.62	73.61	16.67	16	16	73.60	0.24
LT	MI4	2001	417	0.18	-1.32	61.81	7.00	10	10	55.60	0.52
LT	MI4	2002	279	0.17	0.11	72.91	11.17	12	12	60.60	0.35
LT	MI4	2003	215	0.10	-0.57	83.01	-0.57	12			0.22
LT	MI4		4967	0.10	-1.84	95.81	13.24	17	9	53.80	0.26
LT	MI5	1975	16	1.23	5.92	85.61	12.69	12	6	60.50	0.14
LT	MI5	1980	200	0.21	1.93	97.51	13.70	15	7	55.40	0.33
LT	MI5	1981	198	0.19	0.64	88.91	11.80	13	6	52.80	0.23
LT	MI5	1982	239	0.14	-1.16	89.41	17.52	17	7	54.10	0.18
LT	MI5	1983	265	0.38	4.70	92.71	22.97	15	8	55.90	0.25
LT	MI5	1987	50	0.96	6.11	75.71	11.08	12	9	58.40	0.29
LT	MI5	1995	180	0.67	4.23	73.21	-8.56	13	13	73.20	0.22
LT	MI5	1996	200	0.27	1.32	67.31	9.84	12	11	66.50	0.28
LT	MI5	1998	247	0.16	-0.99	64.61	10.57	13	13	59.50	0.22
LT	MI5	1999	277	0.26	2.16	68.81	14.48	15	11	40.60	0.22
LT	MI5	2000	206	0.29	5.14	90.31	-0.90	24	24	90.30	0.10
LT	MI5	2001	178	0.40	2.80	74.91	12.77	12	11	74.90	0.27
LT	MI5	2002	181	0.35	4.42	91.61	9.33	21	11	75.20	0.13
LT	MI5	2003	195	0.15	-0.68	77.81	10.51	15	10	61.90	0.19



LT	MI5		2653	0.11	-0.31	97.51	8.53	24	9	58.40	0.18
LT	MI6	1975	50	0.25	0.42	79.51	32.54	13	7	61.50	0.27
LT	MI6	1980	121	0.19	1.00	86.11	11.10	13	8	53.60	0.20
LT	MI6	1981	136	0.38	2.47	75.71	6.09	10	9	66.80	0.33
LT	MI6	1982	136	0.18	1.22	90.91	9.42	15	8	60.50	0.20
LT	MI6	1983	314	0.21	1.59	82.61	4.18	12	10	55.90	0.35
LT	MI6	1987	34	1.01	5.34	73.41	6.78	11	9	67.10	0.22
LT	MI6	1995	42	1.18	3.61	61.01	-4.97	9			0.40
LT	MI6	1996	51	0.20	0.75	73.21	11.61	10	9	62.00	0.29
LT	MI6	1997	224	0.14	-0.09	88.11	6.69	20	13	64.80	0.15
LT	MI6	1998	54	0.14	-3.17	57.81	62.65	9			0.28
LT	MI6	1999	169	0.09	-5.18	72.11	14.08	19	11	57.90	0.13
LT	MI6	2000	151	0.37	3.60	76.61	-2.89	15	11	56.20	0.20
LT	MI6	2001	46	0.29	0.72	62.51	-0.65	10	10	58.80	0.50
LT	MI6	2002	79	0.25	1.27	71.71	16.32	18	14	70.60	0.10
LT	MI6	2003	72	0.95	4.67	70.81	9.83	10			0.35
LT	MI6		1679	0.08	-4.29	90.91	14.41	20	10	48.80	0.22
LT	MI7	1981	222	0.19	0.22	83.81	20.32	13	8	63.50	0.23
LT	MI7	1983	171	0.13	-0.66	89.21	35.85	14	10	62.70	0.28
LT	MI7	1987	26	0.49	4.58	75.41	30.49	11	8	63.50	0.30
LT	MI7	1995	317	0.11	-3.63	76.21	16.04	15	11	59.40	0.21
LT	MI7	1996	54	0.32	0.96	59.91	8.48	9			0.65
LT	MI7	1997	179	0.12	-2.65	80.31	19.66	16	9	51.80	0.17
LT	MI7	1998	188	0.13	-2.09	72.81	15.22	15	9	53.00	0.22
LT	MI7	1999	289	0.09	-3.91	82.51	18.84	18	8	38.00	0.17
LT	MI7	2000	348	0.04	-19.51	79.61	16.26	26	10	54.20	0.10
LT	MI7	2001	187	0.24	0.86	78.31	27.04	15	9	51.70	0.27
LT	MI7	2002	199	0.24	1.94	75.71	8.70	14	9	54.50	0.27
LT	MI7	2003	162	0.09	-2.98	82.31	23.59	15	10	57.10	0.26
LT	MI7		2354	0.03	-22.96	89.21	16.66	26	2	58.80	0.17
LT	MI8	1975	66	1.05	5.26	80.81	90.77	12	8	69.10	0.27
LT	MI8	1981	49	0.76	3.83	82.31	-14.05	11	7	56.10	0.25
LT	MI8	1982	108	0.16	-0.93	87.61	30.67	12	10	71.40	0.19
LT	MI8		223	0.17	-0.35	87.61	16.06	12	10	71.40	0.28
LT	SON1	1988	2176	0.11	-1.67	84.01	4.56	14			0.45
LT	SON1	1989	1806	0.17	0.88	86.01	1.67	15			0.38
LT	SON1	1990	1590	0.08	-5.35	86.01	3.82	13			0.37
LT	SON1	1991	2106	0.13	0.30	91.01	10.22	14			0.41
LT	SON1	1992	366	0.11	-0.42	90.01	4.20	25			0.18
LT	SON1	1993	529	0.18	3.34	89.01	9.01	24			0.20
LT	SON1	1994	492	0.05	-10.34	92.01	24.26	30			0.14
LT	SON1	1995	1361	0.01	-89.56	96.01	34.72	30			0.14
LT	SON1		10429	0.02	-29.38	96.01	14.07	30			0.15
LT	WI1	1987	633	0.10	-4.29	80.31	4.95	15			0.23
LT	WI1	1988	208	0.27	2.36	82.81	20.56	16			0.23
LT	WI1	1989	344	0.33	4.23	85.31	8.64	17			0.19
LT	WI1	1990	264	0.35	4.30	79.51	12.76	18			0.23
LT	WI1	1991	338	0.18	-0.03	77.51	34.34	15			0.21
LT	WI1	1992	220	0.35	5.20	90.41	10.32	19			0.17
LT	WI1	1993	443	0.14	0.18	87.41	17.23	18			0.15
LT	WI1	1994	197	0.30	3.49	85.31	8.46	16			0.18
LT	WI1	1995	335	0.17	0.52	83.61	11.56	17			0.14
LT	WI1	1997	92	0.18	0.78	78.51	19.69	16			0.14
LT	WI1	1998	132	0.25	2.13	84.11	12.61	16			0.22
LT	WI1	1999	196	0.37	2.52	72.41	10.09	16			0.17
LT	WI1	2000	125	0.33	3.57	82.61	9.49	17			0.15
LT	WI1	2002	67	0.30	4.52	86.11	0.04	21			0.09
LT	WI1		3619	0.20	2.39	90.41	7.81	21			0.22
LT	WI2	1981	384	0.27	3.12	91.41	10.82	14			0.28


LT	WI2	1982	251	0.33	2.95	85.11	7.11	13			0.28
LT	WI2	1983	121	0.35	3.21	85.11	16.43	11			0.38
LT	WI2	1984	178	0.45	3.85	84.81	14.97	16			0.20
LT	WI2	1985	1153	0.14	-1.20	84.31	15.55	14			0.45
LT	WI2	1986	1975	0.10	-2.71	90.71	16.96	18			0.26
LT	WI2	1987	1543	0.13	0.03	95.81	8.67	15			0.25
LT	WI2	1988	1063	0.14	-0.61	87.41	12.94	19			0.20
LT	WI2	1989	1374	0.10	-1.49	100.31	9.77	17			0.27
LT	WI2	1990	783	0.07	-3.79	98.81	11.87	22			0.17
LT	WI2	1991	867	0.12	-1.24	87.11	11.99	18			0.21
LT	WI2	1992	560	0.13	-0.02	90.91	12.95	18			0.19
LT	WI2	1993	426	0.18	0.22	78.71	14.29	19			0.20
LT	WI2	1994	530	0.18	0.93	81.81	11.26	22			0.16
LT	WI2	1995	387	0.11	-2.22	85.61	12.49	17			0.17
LT	WI2	1997	511	0.19	3.83	91.21	11.02	26			0.12
LT	WI2	1998	442	0.15	-0.84	81.51	11.14	25			0.12
LT	WI2	1999	499	0.23	3.26	80.01	17.44	22			0.15
LT	WI2	2000	705	0.21	4.29	86.91	15.12	24			0.15
LT	WI2	2002	161	0.18	4.34	90.91	14.23	31			0.07
LT	WI2	2003	56	0.61	6.38	79.51	8.40	15	6	48.00	0.22
LT	WI2		13967	0.06	-5.58	100.31	12.70	31			0.18
LWF	MI2	1995	23	0.70	1.97	48.51	5.15	6	6	44.40	0.19
LWF	MI2		40	0.69	3.59	59.11	8.37	13	6	44.40	0.12
LWF	MI3	1995	184	0.20	0.65	63.01	-0.47	9	7	40.10	0.32
LWF	MI3	1998	70	0.62	3.74	56.31	7.82	12	7	41.60	0.19
LWF	MI3	2003	110	0.26	0.37	50.81	10.10	13	8	46.10	0.15
LWF	MI3		364	0.14	-1.35	63.01	12.17	13	7	40.10	0.24
LWF	MI4	1998	13	1.21	3.28	47.41	-6.42	6			1.00
LWF	MI4		13	1.21	3.28	47.41	-6.42	6			1.00
LWF	MI5	2002	17	0.27	-1.88	54.11	32.69	10	5	47.30	0.17
LWF	MI5		21	0.28	-1.21	54.11	31.42	10	6	43.90	0.20
LWF	MI6	1995	19	1.59	3.23	49.31	-2.25	7	7	49.30	0.36
LWF	MI6	1998	94	0.21	-0.60	50.41	18.29	9			0.34
LWF	MI6	2002	25	0.68	2.37	67.01	15.16	10	4	36.10	0.17
LWF	MI6	2003	61	0.58	1.83	47.61	25.25	10	9	47.60	0.22
LWF	MI6		199	0.10	-3.19	67.01	11.30	10			0.28
LWF	MI7	1981	178	0.23	0.13	75.41	19.09	14	4	45.00	0.20
LWF	MI7	1982	431	0.15	-2.33	80.01	15.80	16	4	47.80	0.21
LWF	MI7	1983	403	0.38	1.54	74.61	11.80	14	4	53.40	0.29
LWF	MI7	1984	417	0.20	-1.99	69.31	18.84	14	4	46.50	0.32
LWF	MI7	2001	79	0.65	2.71	63.51	11.84	13	5	41.30	0.14
LWF	MI7	2002	99	0.18	-1.15	67.51	7.06	9	5	39.50	0.24
LWF	MI7	2003	55	0.58	2.51	69.21	16.85	11	5	42.00	0.18
LWF	MI7		1662	0.15	-1.46	80.01	9.35	16	5	39.50	0.27
LWF	MI8	2001	57	0.21	-1.59	58.71	10.42	8	5	43.10	0.37
LWF	MI8	2002	13	0.53	0.66	44.01	-13.31	5			0.08
LWF	MI8		79	0.21	-1.35	58.71	8.94	8	6	40.70	0.22
LWF	WI2	1991	19	0.81	4.94	59.71	2.32	12			0.18
LWF	WI2		41	0.16	-0.37	71.61	-10.91	13			0.11
LWF	SON1	1989	2372	0.10	-3.09	79.01	11.79	16			0.36
LWF	SON1	1991	1989	0.10	-3.98	72.01	20.62	16			0.34
LWF	SON1	1992	5351	0.13	-1.80	76.01	7.64	19			0.37
LWF	SON1	1993	4421	0.05	-14.94	78.01	11.58	15			0.42
LWF	SON1	1994	3872	0.08	-5.89	79.01	7.75	16			0.42
LWF	SON1	1995	4042	0.05	-13.67	76.01	10.42	17			0.65
LWF	SON1	1996	1372	0.06	-10.37	72.01	18.85	18			0.20

LWF	SON1		23419	0.07	-6.37	79.01	13.49	19			0.32
LT-SIS	MI2	1995	20	0.47	6.26	65.31	7.99	17	17	65.30	0.14
LT-SIS	MI2	1996	82	0.13	2.80	77.01	4.62	22	16	55.90	0.09
LT-SIS	MI2	1998	48	0.29	8.14	77.21	3.41	23	16	57.50	0.15
LT-SIS	MI2	1999	42	0.29	5.93	69.01	6.78	19	14	56.30	0.14
LT-SIS	MI2	2000	66	0.19	5.66	70.41	4.39	25	20	59.50	0.07
LT-SIS	MI2	2001	67	0.09	-0.47	73.31	1.08	21	19	56.40	0.17
LT-SIS	MI2	2002	52	0.25	7.72	80.31	-2.84	29	19	57.50	0.05
LT-SIS	MI2	2003	105	0.07	-3.11	75.41	8.69	25	18	57.10	0.18
LT-SIS	MI2		482	0.14	5.06	80.31	2.13	29	19	54.60	0.09
LT-SIS	MI3	1995	213	0.12	-0.22	67.31	5.62	17	14	46.70	0.14
LT-SIS	MI3	1996	418	0.09	-0.86	75.71	4.91	19	15	52.30	0.23
LT-SIS	MI3	1998	216	0.22	6.43	73.21	3.78	26	17	51.70	0.10
LT-SIS	MI3	1999	163	0.21	6.86	78.01	-0.45	24	17	60.50	0.11
LT-SIS	MI3	2000	164	0.12	2.19	69.01	5.45	24	19	60.60	0.10
LT-SIS	MI3	2001	201	0.11	1.07	71.31	5.85	22	19	57.20	0.12
LT-SIS	MI3	2002	184	0.09	0.46	76.11	1.65	25	20	54.80	0.19
LT-SIS	MI3	2003	183	0.13	4.03	79.21	5.97	28	19	58.70	0.06
LT-SIS	MI3		1742	0.09	0.91	79.21	5.32	28	19	56.60	0.13
LT-SIS	MI4	1995	205	0.06	-4.62	75.41	12.66	20	13	50.00	0.24
LT-SIS	MI4	1996	362	0.10	-0.65	76.21	9.04	24	15	50.50	0.19
LT-SIS	MI4	1997	271	0.12	-1.32	76.71	11.01	25	17	54.60	0.18
LT-SIS	MI4	1998	227	0.08	-0.21	77.31	3.85	22	17	58.50	0.27
LT-SIS	MI4	1999	285	0.07	-0.72	80.51	3.06	22	18	55.20	0.12
LT-SIS	MI4	2000	172	0.08	0.27	77.01	4.07	24	16	60.20	0.07
LT-SIS	MI4	2001	214	0.14	3.72	77.81	2.78	22	18	61.70	0.10
LT-SIS	MI4	2002	234	0.09	0.18	77.51	2.49	25	19	56.50	0.06
LT-SIS	MI4	2003	201	0.08	-0.74	80.21	3.05	25	21	59.70	0.06
LT-SIS	MI4		2171	0.08	-0.81	80.51	4.88	25	17	47.80	0.10
LT-SIS	MI5	1995	122	0.09	-0.97	72.91	7.83	17	15	54.10	0.28
LT-SIS	MI5	1996	206	0.09	-0.86	76.71	4.61	18	15	49.80	0.40
LT-SIS	MI5	1997	502	0.14	2.69	76.51	11.17	25	15	51.30	0.23
LT-SIS	MI5	1998	189	0.03	-6.57	98.51	8.76	23	16	54.50	0.24
LT-SIS	MI5	1999	159	0.07	-2.23	77.81	6.05	23	16	56.40	0.19
LT-SIS	MI5	2000	164	0.11	-1.19	63.41	7.26	22	19	59.00	0.23
LT-SIS	MI5	2001	141	0.09	-0.95	73.81	5.86	22	15	58.20	0.10
LT-SIS	MI5	2002	161	0.08	0.44	82.91	8.24	22	16	52.90	0.42
LT-SIS	MI5	2003	192	0.17	5.88	79.51	1.50	26	18	57.00	0.15
LT-SIS	MI5		1836	0.04	-4.24	98.51	3.46	26	16	47.80	0.14
LT-SIS	MI6	1980	13	0.86	5.03	75.71	3.10	11	6	56.40	0.24
LT-SIS	MI6	1983	5	2.25	7.86	71.41	-2.05	11	8	55.90	0.15
LT-SIS	MI6	1995	88	0.11	-2.09	70.41	18.47	16	14	60.70	0.28
LT-SIS	MI6	1996	141	0.10	-2.63	69.61	14.56	20	15	57.90	0.19
LT-SIS	MI6	1998	124	0.10	-0.22	69.41	9.45	24	19	67.70	0.07
LT-SIS	MI6	1999	78	0.11	-0.40	65.01	11.08	22	15	46.50	0.14
LT-SIS	MI6	2000	111	0.12	3.46	74.41	4.45	28	20	51.40	0.05
LT-SIS	MI6	2001	153	0.18	1.95	69.01	26.11	19	16	56.60	0.18
LT-SIS	MI6	2002	137	0.08	-3.00	71.61	6.55	23	19	58.00	0.10
LT-SIS	MI6	2003	112	0.22	4.21	70.01	5.72	25	19	69.00	0.10
LT-SIS	MI6		963	0.07	-2.94	75.71	10.13	28	18	49.40	0.13
LT-SIS	MI7	1995	59	0.04	-23.76	64.81	11.56	28	12	57.90	0.03
LT-SIS	MI7	1996	82	0.02	-39.40	77.71	17.44	25	24	52.30	0.06
LT-SIS	MI7	1997	181	0.02	-68.58	68.61	20.61	28	12	51.30	0.10
LT-SIS	MI7	1998	226	0.03	-31.76	67.81	22.05	31	14	49.70	0.07
LT-SIS	MI7	1999	87	0.24	4.32	66.61	28.79	21	15	58.60	0.08
LT-SIS	MI7	2000	22	0.34	6.15	73.01	22.35	20	11	52.50	0.05
LT-SIS	MI7	2001	142	0.13	-1.73	67.51	11.27	19	9	45.40	0.14
LT-SIS	MI7	2002	86	0.06	-6.93	77.51	26.04	26	11	47.40	0.07
LT-SIS	MI7	2003	67	0.10	-7.25	66.01	22.34	31	11	51.60	0.05


LT-SIS	MI7	2004	12	0.20	5.54	72.41	-31.77	19	10	47.40	0.03
LT-SIS	MI7		964	0.02	-34.69	77.71	14.24	31	12	47.20	0.10
YP	SON1	1991	219	0.20	-3.79	30.01	3.51	6			1.07
YP	SON1	1992	241	0.12	-6.48	31.01	13.12	9			0.44
YP	SON1	1993	252	0.24	-0.22	37.01	2.07	11			0.32
YP	SON1	1994	288	0.15	-1.42	39.91	-1.45	11			0.31
YP	SON1	1995	433	0.07	-18.43	30.01	5.15	6			0.80
YP	SON1	1996	157	0.06	-20.01	30.01	9.29	9			0.27
YP	SON1		1590	0.13	-2.54	39.91	-1.36	11			0.37

Table 9. Life history parameter values of cohorts used in the virtual population analyses. Population codes correspond to Figures 1-3. The last two digits in the population code represent the start of each cohort for e.g. EWAL1C90 = Walleye population 1 from Lake Erie, cohort start 1990. Age represents the ages with data for each cohort.

Species	Population	Age	k	$t_0$	$L_\infty$	$L_0$	$t_{max}$	$\alpha$	$L_\alpha$	Z
WALL	EWAL1C90	0, 2-9, 11	0.18	-1.55	75.51	27.06	11	2	33.30	0.22
WALL	EWAL1C91	1-7, 9	0.59	2.08	72.91	6.16	12	3	39.60	0.23
WALL	EWAL1C92	0-8, 10-11	0.19	-3.20	67.51	16.70	11	2	38.00	0.13
WALL	EWAL2C89	1, 3-9, 11	0.58	1.85	69.21	22.88	11	3	39.10	0.21
WALL	EWAL2C90	0-8, 10, 13	0.44	0.39	71.31	20.90	13	2	36.70	0.23
WALL	EWAL2C91	1-6, 8-9	0.27	-1.01	70.71	19.26	9	2	29.50	0.26
YP	EYP1C91	2-9	0.31	-0.16	34.51	5.17	8	1	12.70	0.39
YP	EYP1C92	2-9	0.29	-0.92	31.41	8.21	8	2	14.50	0.39
YP	EYP1C93	2-9	0.25	-0.69	33.51	5.22	9	2	14.10	0.39
YP	EYP2C91	1-6	0.35	-1.00	30.81	7.80	6	1	13.70	0.54
YP	EYP2C93	1-5, 7-8	0.34	-0.04	34.81	4.75	8	2	13.20	0.44
LWF	HLWFON2C80	1-6	0.84	1.42	61.01	16.26	7	5	49.40	0.58
LWF	HLWFON2C81	1-7, 10	0.42	0.67	65.21	12.74	10	4	42.00	0.29
LWF	HLWFON2C82	1-8, 10	0.70	1.58	62.61	17.33	10	3	35.00	0.31
YP	HYPON2C81	1-10	0.58	1.94	35.71	4.13	10	1	13.80	0.39
LWF	OLWF1C91	1-13	0.12	-4.05	63.41	14.20	13			0.10
LWF	OLWF1C92	0-12	0.17	-1.85	61.21	8.13	12			0.10
LWF	OLWF1C93	1-11	0.21	-0.46	59.11	8.19	11			0.05
LWF	OLWF1C94	1-10	0.16	-1.11	63.61	6.27	10			0.14
YP	OYP1C91	1-8	0.23	-1.43	27.61	8.12	8			0.43
YP	OYP1C92	1-8	0.24	-0.84	29.31	5.02	8			0.36
YP	OYP1C94	1-7	0.18	-2.99	26.91	4.96	7			0.36
LT-SIS	SLTSISMI7C92	3-11	0.13	-2.83	61.91	10.27	11	9	45.40	0.31
LT	SLTWI2C85	1-10	0.21	-0.09	73.91	11.55	10			0.18




a)




b)

Figure 1. Extent of (a) 12 lean lake trout and (b) 6 siscowet lake trout populations used to examine life history traits in the Great Lakes basin. Black lines delineate population zones.



a)



b)

Figure 2. Extent of (a) 26 lake whitefish and (b) 8 bloater populations used to examine life history traits in the Great Lakes basin. Black lines delineate population zones.




Figure 3. Extent of (a) 12 yellow perch and (b) 9 walleye populations used to examine life history traits in the Great Lakes basin. Black lines delineate population zones.






Figure 4. Comparison of the length at age relationships produced using fish collected year-round and fish collected in the summer and/or fall for three populations, (a) lake whitefish in Lake Huron (HLWFCORA13) (b) yellow perch in Lake Erie (EYP4), and (c) lake trout in Lake Michigan (SLTMI5).




Figure 5. Comparison of the length at age relationships produced using yellow perch collected in the summer and fall within the same lake a) EYP41999, b) EYP42000 and c) EYP42001.




Figure 6. Length at age for a) lake whitefish, b) walleye and c) yellow perch caught using different gillnet mesh sizes in Lake Erie (all populations).




Figure 7. Growth curves for a) lake whitefish, b) walleye and c) yellow perch in Lake Erie (all populations), dashed lines represent the mean growth curve for all ages, the black line represents the growth curves for ages 2-7.




Figure 8. Growth curves for 5 populations of a) lake whitefish, b) yellow perch and c) walleye in Lake Erie. Population codes correspond to Figures 1-3.




Figure 9. Growth curves for 1 population of a) walleye, b) lake whitefish and c) yellow perch in Lake Ontario. Population codes correspond to Figures 1-3.




Figure 10. Growth curves for a) 3 bloater, b) 2 lean lake trout and c) 3 walleye populations in Lake Huron. Population codes correspond to Figures 1-3.




Figure 11. Growth curves for a) 7 lake whitefish and b) 3 yellow perch populations in Lake Huron. Population codes correspond to Figures 1-3.






Figure 12. Growth curves for a) 4 lake whitefish, b) 2 yellow perch and c) 5 bloater populations in Lake Michigan. Population codes correspond to Figures 1-3.




Figure 13. Growth curves for a) 10 lean lake trout and b) 6 siscowet lake trout populations in Lake Superior. Population codes correspond to Figures 1-3.




Figure 14. Growth curves for a) 9 lake whitefish population and b) 1 yellow perch population in Lake Superior. Population codes correspond to Figures 1-3.