

DESCRIPTIVE NOTES

The map area (NTS 87-144) lies within the Minto Inlier, a ~300 km long by 100–150 km wide belt of gently folded sedimentary and igneous rocks of early Neoproterozoic (late Tonian-early Cryogenian) age. The Neoproterozoic sedimentary strata belong to the Shaler Supergroup, an approximately 4 km thick succession of shallow marine carbonate rocks and evaporite rocks with interbedded terrigenous rocks that were mainly deposited in a shallow intracontinental epicritic sea, referred to as the Amundsen Basin (Rainbird et al., 1984; Rainbird et al., 1986a; Thorsen and Torsen, 1982; Young, 1981). The basin is considered to have formed within the supercontinent Rodinia and exposures of similar rocks in what are now the Mackenzie Mountains of the northern Cordillera suggest that it extended for more than 100 km to the southwest (Long et al., 2008; Rainbird et al., 1986a). The sedimentary succession is intercalated with mafic sills of the ca. 720 Ma Franklin gabbro complex (Heaman et al., 1982). The sills of variable thickness up to 100 m, but most are 20–50 m thick. In many cases, individual sills extend for 20 km or more along-strike with little significant change in thickness. Sills constitute anywhere from 10 to 50 per cent of the stratigraphic section. Sills of similar type and age also occur in the Coppeigne Horst, Brook Inlet and Duke of York Inlet to the south (Rainbird et al., 1986b; Shattuck et al., 2004) and several, geochemically similar intrusions and volcanic rocks associated with the Franklin event extend from Greenland to the western Yukon (Dewey et al., 2000; Heaman et al., 1992; Macdonald et al., 2010). The Shaler Supergroup in Minto Inlier is capped by a succession of flood basalts flows and interflow sedimentary rocks (Nauyasik Fm.) more than 1 km thick, which are the extensive equivalent of the sills (Bangor, 1976; Jefferson et al., 1985). Rare north-northeast-sloping dikes are interpreted to have intruded along syn-magmatic normal faults, to feed sills and possibly the flood basalts (Bedard et al., 2012). The mafic gabbro complex is identified in the lower, which have similarities in the different sills and dikes. The oldest sills and corresponding basal lavas are enriched in incompatible trace elements and may have alkali-enriched bases. Younger diatritic sills correspond to the major sheet flow units of the late succession. Basal strata of the Shaler Supergroup (Ran Group) are exposed only at the northeastern end of Minto Inlier, near Hadley Bay, where they unconformably overlie Paleoproterozoic sedimentary rocks, which, in turn, unconformably overlie Archean granitic rocks (Campbell, 1981; Rainbird et al., 1984). The irregular edge of Minto Inlier is defined by an erosional unconformity that separates the Neoproterozoic rocks from Lower Cambrian sandstone and siltstone that pass upward into a thick succession of mainly siliciclastic carbonate rocks, ranging in age from Cambrian to Devonian (Thorsen and Torsen, 1982). Structurally, the Minto Inlier is relatively simple, composed of the open, northeast-trending Holman Island syncline and a smaller Walker Bay anticline to the northwest. Dikes typically dip to more than 10° and there is generally no penetrative deformation or other apparent outcrop-scale fabric. The origin of the folding is unknown but it occurred after deposition of the early Neoproterozoic rocks and before uplift, erosion and deposition of overlying lower Cambrian siliciclastic rocks, which are not folded. All rocks are dissected by east-northeast to east-trending faults that form a horst and graben system with up to 200 m of relative stratigraphic separation on individual faults. The zone of faulting is about 100 km wide and stretches from the head of Minto Inlet in the west to Wynmatt Bay in the east and is spectacularly imaged as prominent lineaments on recently published aeromagnetic maps (e.g. Kias and Oleschak 2010).

NTS 87-144 is underlain by stratigraphic units from the middle to upper Kilian Formation, Kuujuaia Formation and Nauyasik Formation of the Shaler Supergroup. Together with diabase sills, the strata comprise the gently south-dipping northern limb of the Holman Island Syncline, whose axis lies along the southern edge of the map sheet. Exposures of the Kilian Formation (clastic-carbonate and tan carbonate members) are limited in the northern half of the map sheet area due to the nonconformable nature of the diabase sills that form a prominent southeast dip slope along the Kuujuaia River. The upper Kilian Formation, Kuujuaia Formation and basal Nauyasik Formation, are best exposed along a prominent cuesta that faces north-northeast along the south side of the Kuujuaia River. The Kilian generally is quite recessive but a good exposure of the tan carbonate member (nPK3) and upper evaporite member (nPK4) is located near UTM 566071E, 7894971N. In most places within the map area, the basal contact of the Kuujuaia with the upper evaporite member is poorly exposed, but at one locality (UTM 566103E, 7890758N), it is a well-developed erosional discontinuity overlain by a thin, discontinuous, chert-cobble conglomerate. The conglomerate, in turn, is overlain by a green, parallel-stratified volcanic (tuff) wedge, which passes upward into a dark green siltstone. That is interbedded with a cleaner (quartzite) sandstone, more typical of overlying cross-bedded quartzarenites of the Kuujuaia Formation. The underlying Kilian Formation strata at this location are well exposed (chert-cobble and tuff) and are (normally) red, suggesting that deposition of the overlying strata was accompanied by introduction of reducing hydrothermal fluids. This important outcrop shows that volcanism, normally associated with the overlying Nauyasik Formation, was already active at the beginning of Kuujuaia time. One of the thickest and best preserved sections of the Kuujuaia Formation is exposed near the eastern end of the cuesta at UTM 571406E, 789889N (see section 46-22 of Rainbird, 1992 and section 4 of Jefferson, 1980). Other good exposures of the Kuujuaia Formation are located at UTM 565403E, 789886N (section 86-10) and UTM 564148E, 7894971N (section 86-14). The Kuujuaia and Nauyasik formations are also well exposed along the flanks of two, north-flowing tributaries of the Kuujuaia River. The contact between the two formations generally is planar and conformable but there are numerous places where soft-sediment deformation features are preserved in the underlying sandstone indicating that the sand was un lithified at the time of the main eruption (see Rainbird, 1992). As well, irregular fragments of basalt have been observed within the sandstone. The most complete exposures of the Nauyasik Formation are located at approximately UTM 561945E, 7898591N and UTM 564916E, 7898591N and are described in detail by Williamson et al. (2013). Members nPK1 and nPK3 are composed of basaltic flows and associated interflow deposits that are regionally distributed, but the intervening volcanoclastic nPK2 member is discontinuous and of variable thickness. It thrives markedly from the central part of the map area toward the northwest and is absent in the western half of the map area. At least 5 diabase sills occur within the map area and generally are spaced at regular intervals within the host sedimentary rocks. The sills are of the type 2 (diabase) described in the legend. Several steep, northwest-sloping, mostly westside down, normal faults are spaced evenly across the map area and are evident as prominent linear topographic lows. At approximately UTM 564516E, 7894971N, the lineament coincides with a thin diabase dyke, suggesting that emplacement was coincident with faulting, a relationship that has been documented elsewhere in the region (Bedard et al., 2012). East-west block faulting, common in map areas to the north, is largely absent from NTS 87-144.

Abstract

NTS 87-144 is underlain by the middle to upper Kilian, Kuujuaia and Nauyasik formations of the Neoproterozoic Shaler Supergroup. Together with at least 5 diabase sills (type 2), spaced at regular intervals within the host sedimentary rocks, the strata comprise the gently south-dipping northern limb of the Holman Island Syncline. The northern half of the map area is dominated by a thick sill, which forms a prominent southeast dip slope down to the Kuujuaia River. The Kilian Formation is best exposed along a prominent cuesta that faces north-northeast along the south side of the Kuujuaia River. The upper two members of the Kilian Formation are sporadically exposed along the base of the cuesta. Cross-bedded quartzarenites of the Kuujuaia Formation overlie the cuestas. The Kuujuaia Formation is prominent along the cuestas, face along with conformably capping basalt flows of the Nauyasik Formation. Both are also well exposed along the flanks of two, north-flowing tributaries of the Kuujuaia River. Several steep, northwest-sloping, westside down normal faults are spaced evenly across the map area.

Résumé

Le feuillet NTS 87-144 expose des roches Néoproterozoïques des Formations du Kilian, Moyer et Supérieur, Kuujuaia et Nauyasik. Les roches sédimentaires appartiennent au Supergroupe de Shaler, les roches ignées par 5 filons-couches, diabasiques de l'événement Franklin. Les strates constituent le flanc nord du Synclinal Holman Island, et pendent doucement vers le sud. La partie nord du feuillet est dominée par une épaisse coulée de basalte, qui forme un banc qui va jusqu'à la rivière Kuujuaia. Les roches sédimentaires sont bien exposées le long d'une falaise qui s'élève au sud de cette rivière et aussi le long de ses tributaires du côté sud. Les deux membres supérieurs du Kilian sont exposés sporadiquement au pied de la falaise. Les arénites quartzifères à lit entrecroisé de la Formation de Kuujuaia sont en discordance sur les strates du Kilian. Le sommet de la montagne est formé par les lavas et volcanoclastes de la Formation de Nauyasik. Il y a plusieurs failles normales orientées nord-sud, avec pendage vers l'ouest.

Catalogue No. M185-1104-2012E-PDF
ISBN 978-1-105-21440-5
doi: 10.4095/293344

© Her Majesty the Queen in Right of Canada 2013

Cover illustration
Looking east at flood basalt-capped cuestas with Kuujuaia River and upper Kilian Formation and diabase sill in foreground, Victoria Island, Northwest Territories, 2011-07-13. Photograph by R. Rainbird 2013-204

National Topographic System reference and index to adjoining published Geological Survey of Canada maps

CGM 103	CGM 82
CGM 59	CGM 155
CGM 104	CGM 155

COVER ILLUSTRATION

Looking east at flood basalt-capped cuestas with Kuujuaia River and upper Kilian Formation and diabase sill in foreground, Victoria Island, Northwest Territories, 2011-07-13. Photograph by R. Rainbird 2013-204

COVER ILLUSTRATION

Looking east at flood basalt-capped cuestas with Kuujuaia River and upper Kilian Formation and diabase sill in foreground, Victoria Island, Northwest Territories, 2011-07-13. Photograph by R. Rainbird 2013-204

CANADIAN GEOSCIENCE MAP 104

GEOLOGY

TAKIYUAQATTAK

Victoria Island, Northwest Territories

1:50 000

Geological Survey of Canada

Geological Survey of Canada

CANADIAN GEOSCIENCE MAP 104

GEOLOGY

TAKIYUAQATTAK

Victoria Island, Northwest Territories

1:50 000

Geological Survey of Canada

Geological Survey of Canada

CANADIAN GEOSCIENCE MAP 104

GEOLOGY

TAKIYUAQATTAK

Victoria Island, Northwest Territories

1:50 000

Geological Survey of Canada

Geological Survey of Canada

CANADIAN GEOSCIENCE MAP 104

GEOLOGY

TAKIYUAQATTAK

Victoria Island, Northwest Territories

1:50 000

Geological Survey of Canada

Geological Survey of Canada

CENOZOIC

Quaternary sediments.

NEOPROTEROZOIC

Franklin intrusions: Typically massive, laterally extensive, diabasic sills with columnar jointing (~3–50 m thick, rarely up to 100 m). Some sills are composite with internal intrusive contacts. Two types: 1) An older, more primitive type is commonly layered, with microbasaltic lower and upper border zones and olivine-enriched basal cumulate (olivine gabbro). 2) Younger (based on cross cutting relationships), more evolved, diabasic sills showing enrichment in magnetite, ilmenite, quartz and alkali feldspar towards their cores, but are rarely layered. Some sills are porphyritic and contain 10–15% plagioclase-clinopyroxene phenocrysts and glomerocrysts up to 5 mm. Less common, 1–40 m wide dikes, irregular to very linear (generally oriented NNW). Commonly associated with fault breccias or drag folds in host metasediments. Dikes commonly connect to sills, some associated with calc-silicate contact metamorphic rocks (redish garnet rimmed by bright green vesicular), black Fe-calcite skarns, and minor sulphides.

Nauyasik Formation (nPN1–nPN4)

Sheet-flow member: Blue-green to orange-weathering, laterally extensive, subbasal basaltic flows, individual flows 15 to 50 m thick. Flow structure varies from columnar-entablature to a massive base with typically vesicular flow tops. Rare interflow scoria, spatter, fumarolic concretions, volcanic necks and pily to disseminated native copper. Maximum thickness of 200 m, limited by erosional preservation.

Lower recessive member: Massive to parallel-stratified volcanic-peggible volcanoclastic sandstone. Framework composed of volcanic, carbonate and sandstone clasts, matrix-rich in quartz and grains. Conglomerate subunit varies in thickness from 40 to 100 m; sandstone sub-unit from 5 to 10 m.

Lower member: Dark green to grey weathering, dominantly subbasal flows, varying from fine massive basalt to coarse sub-phicic basalt. Filled and hydrothermal breccias are common at unit base indicating emplacement flow-top, or discontinuous basaltic flows. Degree of vesicularity varies throughout. Thickness 40 to 70 m.

Shaler Supergroup (nPK1–nPK4)

Kuujuaia Formation: Two principal lithologies: coarse quartzarenites typified by stacked tabular cox-silt of simple and compound planar crossbedding and a less abundant fine-grained assemblage of interbedded fine sandstone, dolomite, siltstone and mudstone forming lenses up to 20 m thick. Rare basaltic pebbles, ~120 m thick.

Kilian Formation (nPK1–nPK4)

Upper Evaporite-Carbonate member: Base is dolomite and dolomite with 10–20% ripple cross-laminated gypsiferous silt. Bedding parallel to crosscutting salt-saturated veins and desiccation cracks common. Changes up-section from creamy grey to pinkish grey, reflecting increase in hematite relative to carbonate. Nodular siltstone more common in middle part of member. Upper consists mainly of parallel-laminated red dolomite, mudstone and siltstone, to vertical-bedded, buff to pink-weathering siltstone -no sulphate. Diagenetic redox horizons desiccation cracks, halite pseudomorphs and beeps structure are ubiquitous. Present only in the southwest domain of the Minto Inlier, ~80 m thick.

Tan Carbonate member: Tan to green-grey, flaggy weathering dolomite and limestone. Gradation between parallel-laminated tubes and flat to wavy and hummocky bedded siltstone. Luster-rich layers are generally plane parallel laminated with rare siltstone lenses (lat-vol ripple)? Bed bases typically scoured grading up to little-itch type. Interfingering clastic breccia commonly infilling swales and gutters. Black chert nodules throughout and stromatolites at several horizons. One destructive bedform, from the middle of the tan carbonate member, is laterally traceable from Ullahakook along the Kuujuaia River Valley to where it cuts across the Nauyasik plateau ~60 m thick.

Clastic-carbonate member: Variegated (red, green, grey and black) pe-stripe-laminated mudstone and siltstone, particularly at its base. Desiccation cracks common in mudstone and wavy basing ripple cross-lamination in coarse siltstone-fine sandstone interlayers. Wavy-flaser bedded and small-scale cross-bedded, 4 m thick, buff-weathering, fine-grained quartzarenite near top. Wavy bedded dolomite and laterally limited stromatolite interbeds are common and increase up-section, ~120 m thick.

Carbonate-evaporite member: Alternating, decimetre-scale subunits of evaporite and carbonate-dominated lithologies: evaporite: laminated mudstone and dolomite mudstone with interbedded nodular arthrite and laminated gypsum and arthrite, minor stromatolite dolomite. Carbonate lithologies: dolomite and minor limestone luster-rich rhythmic capped by arthrite-like laterally linked stromatolites, forming repetitive metre-scale cycles. Molar-tooth structure common.

REFERENCES

Bangor, W.R.A., 1976. The Nauyasik basalt, Victoria Island, District of Franklin, in Current Research, Part A: Geological Survey of Canada, Paper 76-1A, p. 347–352.

Bedard, J.H., Nasir, W.E., Nabelek, P., Wilpon, A., Hayak, M., Macdonald, W., Hayes, B., Stegwealdt, K., Haidt, T., Rainbird, R., Dewing, K., and Grant, E., 2012. Fault-mediated melt ascent in a Neoproterozoic continental flood basalt province: the Franklin sills, Victoria Island, Canada. Geological Society of America Bulletin, v. 124, p. 723–738. doi:10.1130/B3045011

Campbell, I.H.A., 1981. Geomorphology and tectono-depositional relationships of the Neoproterozoic rocks of the Hadley Bay area, northern Victoria Island, District of Franklin, in Current Research, Part A: Geological Survey of Canada, Paper 81-1A, p. 15–22.

Dewey, S.W., Halls, H.C., Davis, D.W., and Evans, D.A.D., 2009. Palaeomagnetism and U-Pb geochronology of Franklin dikes in high Arctic Canada and Greenland: Archaean age and palaeomagnetic pole containing about rotations in the Nares Strait region. Canadian Journal of Earth Sciences, v. 46, p. 989–1005.

Heaman, L.M., LeCheminant, A.N., and Rainbird, R.H., 1992. Nature and timing of Franklin igneous events, Canada: implications for a late Palaeozoic marine plate and the break-up of Laurentia. Earth and Planetary Science Letters, v. 109, p. 117–131.

Jefferson, C.W., 1985. Uppermost Shaler Group and its contact with the Nauyasik basalt, Victoria Island, District of Franklin, in Current Research, Part A: Geological Survey of Canada, Paper 85-1A, p. 203–214.

Jefferson, C.W., Nelson, W.E., Kishan, R.V., Rasmussen, J.L., and Spence, R.F., 1986. Geology and copper occurrences of the Nauyasik basalt, Victoria Island, District of Franklin, in Current Research, Part A: Geological Survey of Canada, Paper 86-1A, p. 203–214.

Kias, F. and Oleschak, D., 2010. First vertical derivative of the magnetic field, Minto Inlier Aeromagnetic Survey, Victoria Island, NTS 87-144, Northwest Territories. Geological Survey of Canada, Open File 6701, scale 1:100 000.

Long, D.G.F., Rainbird, R.H., Turner, E.C., and Macdonald, R.B., 2008. Early Neoproterozoic strata (Sequence B) in northern Victoria Island, Northwest Territories: a correlation with the Geological Atlas of the Northern Canadian Mainland Sedimentary Basin. Geological Survey of Canada, Open File 6701, scale 1:100 000.

Macdonald, P.A., Schmidt, M., D., Crowley, J.L., Rosta, C.F., Jones, D.S., Maked, A.C., Strauss, J.V., Cohen, P.A., Johnson, D., and Schrag, D. P., 2010. Calibrating the Cryogenian. Science, v. 327(5870), p. 1441–1443.

Rainbird, R.H., 1992. Anatomy of a large-scale tectonic platform from the Neoproterozoic Shaler Group, Victoria Island, N.W.T., Canada. Canadian Journal of Earth Sciences, v. 29, p. 2517–2530.

Rainbird, R.H., 1993. The sedimentary record of marine plate uplift preceding eruption of the Neoproterozoic Nauyasik flood basalt. Journal of Geology, v. 101, p. 305–318.

Rainbird, R.H., Jefferson, C.W., Hildebrand, R.S., and North, J.K., 1994. The Shaler Supergroup and revision of Neoproterozoic stratigraphy in the Amundsen Basin, Northwest Territories, in Current Research, Part A: Geological Survey of Canada, p. 61–70.

Rainbird, R.H., Jefferson, C.W., and Young, G.M., 1986a. The early Neoproterozoic sedimentary Succession of Victoria Island, Northwest Territories, in Current Research, Part A: Geological Survey of Canada, Open File 86-1A, p. 108–144.

Rainbird, R.H., LeCheminant, A.N., and Langer, G.L., 1986b. The Origin of the Minto Inlier related to the southern Victoria Island, District of Franklin, Northwest Territories, in Current Research 1986: Geological Survey of Canada, p. 125–134.

Shattuck, J.G., Dostal, J., and Kopp, J.D., 2014. Petrogenesis of the 723 Ma Connors sills, Amundsen basin, Arctic Canada: implications for the break-up of Rodinia. Precambrian Research, v. 125(4–5), p. 308–324.

Thorsen, R., and Torsen, E.T., 1982. Basalt, Victoria and Stefansson Islands, Arctic Archipelago. Geological Survey of Canada, Memoir 389.

Williamson, N., Cowens, B., Bedard, J.H., and Zagorovsk, A., 2013. Volcano-stratigraphy and significance of the southern side of the Nauyasik Formation flood basalts, Victoria Island, Northwest Territories, Current Research 2013-14, 15p, doi:10.4095/292708.

Young, G.M., 1981. The Amundsen Embayment, Northwest Territories: relevance to the upper Proterozoic (Proterozoic) of the Amundsen Basin of Canada, (ed.) J.H.A. Campbell. Geological Survey of Canada, Paper 81-10, p. 203–211.

Recommended citation

Rainbird, R.H., Bedard, J.H., and Williamson, N., 2013. Geology, Takiyuqaattak, Victoria Island, Northwest Territories. Geological Survey of Canada, Canadian Geoscience Map 104 (preliminary), scale 1:50 000, doi:10.4095/293344

CANADIAN GEOSCIENCE MAP 104

GEOLOGY

TAKIYUAQATTAK

Victoria Island, Northwest Territories

1:50 000

Geological Survey of Canada

Geological Survey of Canada

CANADIAN GEOSCIENCE MAP 104

GEOLOGY

TAKIYUAQATTAK

Victoria Island, Northwest Territories

1:50 000

Geological Survey of Canada

Geological Survey of Canada

ess.nrcan.gc.ca