Holocene earth materials and landforms REFERENCES Organic Deposits Peat bogs: fibric to humic organic matter; massive to stratified accumulations; Bednarski, J.M., 2003a. Betalamea Lake, Northwest Territories - Yukon Territory - British Columbia (NTS generally greater than 2 m thick; confined to topographic depressions or level This Surficial Geology Map of NTS 94-O/14 (Canadian Geoscience Map 120) is the product of collaboration between the Geological Survey of 95B/4); Geological Survey of Canada, Open File 4502, scale 1:50 000. Canada and the British Columbia Ministry of Energy, Mines and Natural Gas as part of the Geo-mapping for Energy and Minerals Program areas; underlain by poorly drained till, glaciolacustrine and other GEM-Energy Yukon Basins Project. The accompanying geodatabase includes field observation points and field photos, landform features as unconsolidated sediments; formed by decomposition of plant material in Bednarski, J.M., 2003b. Surficial geology of Fort Liard, Northwest Territories - British Columbia. Geological lines, and surficial geology unit polygons. The map and geodatabase are essential baseline geoscience information for a range of potential wetland areas: bogs with sporadically discontinuous permafrost and end-users including resource explorationists, geotechnical engineers, land-use managers, terrestrial ecologists, archaeologists Survey of Canada, Open File 1760, scale 1:50 000. thermokarst depressions potentially unstable if organic material is disturbed or geoscientists and communities in northern British Columbia. By providing new insight into the distribution and origins of surficial earth Bednarski, J.M., 2003c. Surficial geology of Lake Bovie, Northwest Territories - British Columbia. Geological Survey of Canada, Open File 1761, scale 1:50 000. materials, CGM 120 will help to reduce the economic costs and risks associated with the sustainable development of energy and mineral resources in NTS 94-O/14. Environmental impact assessments for new access roads, work camps, well pads, pipeline and power Fens: fibric organic matter; massive to stratified; generally greater than 2 m transmission line corridors, water storage and waste management systems and other infrastructure will benefit from the geoscience Bednarski, J.M., 2003d. Surficial geology of Celibeta Lake, Northwest Territories - British Columbia. Geological nformation presented here. By identifying areas prone to geological hazards (e.g., landslides, permafrost, flooding), CGM 120 will also help to thick; confined to topographic depressions, level areas and meltwater Survey of Canada, Open File 1754, scale 1:50 000. channels; underlain by poorly drained till, glaciolacustrine and other unconsolidated sediments; formed by decomposition of plant material in Bednarski, J.M., 2005a. Surficial Geology of Etsine Creek, British Columbia, Geological Survey of Canada, Open wetland areas; fens are prone to flooding following drainage damming by Terrain mapping and field-based benchmarking studies have led to a better understanding of the regional distribution of surficial deposits permafrost, landslides and other geomorphic processes in the NTS 94-O/14 map area (Huntley and Hickin, 2010; Huntley et al., 2011a-b) Bednarski, J.M., 2005b. Surficial Geology of Gote Creek, British Columbia, Geological Survey of Canada, Open Surficial earth materials and landforms were interpreted using a combination of stereo-pair air photos (BCB97010, 15BCB97015, 15BCB97029, 15BCB97075 and 15BCB97088 series), LANDSAT 7 satellite imagery (http://glovis.usgs.gov/ [URL 2011]) and Shuttle Radar Undifferentiated peat bogs and fens: humic to fibric organic matter; massive File 4846, scale 1:50 000. Topography Mission digital elevation models (http://dds.cr.usgs.gov/srtm/ [URL 2011]). The base map was generated from CANVEC shape files (http://geogratis.cgdi.gc.ca/geogratis/ [URL 2011]). Surficial geology polygons and landform line symbols were digitized using to stratified accumulations; generally greater than 2 m thick; confined to Clement, C., Kowall, R. Huntley, D. and Dalziel, R., 2004. Ecosystem units of the Sahtaneh area; Slocan Forest topographic depressions. level areas or channels; underlain by poorly drained commercially available computer software packages (Global Mapper, ArcMap and ArcGIS) and compared to published maps, reports and Products (Fort Nelson) Report, 39 pages and appendices till, glaciolacustrine and other unconsolidated sediments; formed by archived digital data (e.g., Stott and Taylor, 1968; Bednarski, 2003a-d; Clement et al., 2004; Bednarski, 2005a-b; Trommelen and Levson, 2008; Demchuk, 2010). The geodatabase accompanying this map conforms to the Science Language for the Data Management component decomposition of plant material in wetland areas; may contain sporadically Deblonde, C., Plouffe, A., Boisvert, E., Buller, G., Davenport, P., Everett, D., Huntley, D., Inglis, E.,Kerr, D., Moore, A., Paradis, S.J., Parent, M., Smith, R., St-Onge, D., and Weatherston, A., 2012. Science Language discontinuous permafrost and thermokarst depressions; potentially unstable if disturbed or removed during development. for an Integrated Geological Survey of Canada Data Model for Surficial Maps Version 1.1 Results of Fieldwork was undertaken in 2009 and 2010 to ground truth surficial geology polygons interpreted from air photos and satellite imagery, and to the properties of the propertGeological Survey of Canada Surficial Legend Review Committee; Geological Survey of Canada, Open File Alluvial deposits gather characteristics that could not be determined through remote predictive mapping. Earth materials were defined on the basis of facies and landform associations, texture, sorting, colour, sedimentary structures, degree of consolidation, and stratigraphic contact relationships at Alluvial terraced sediments: boulders, gravel, sand and silt; generally Demchuk, T., 2010. Surficial geology of the Komie Creek area (NTS 094P/05). British Columbia Ministry of field stations and remote observations from helicopters. The distribution of glacial and non-glacial landforms is depicted on the surficial massive to planar stratified; well to rapidly drained; greater than 2 m thick; may Energy, Mines and Petroleum Resources, Open File 2010-08; Geological Survey of Canada Open File 6568, geology map. Map units in the Legend are presented chronostratigraphically and include organic deposits, alluvial, colluvial, eolian, contain interbedded debris flows and buried organic material; underlain by outwash, till or bedrock; transported and deposited by modern rivers, streams Huntley. D.H. and Hickin, A.S., 2010. Surficial deposits, landforms, glacial history and potential for granular and creeks; subject to rare flooding; potential source of aggregate; land use aggregate and frac sand: Maxhamish Lake Map Area (NTS 94-O), British Columbia. Geological Survey of The distinctive landscape of NTS 94-O/14 is largely a product of underlying bedrock and geological structures, with ornamentation by the Late Wisconsinan Laurentide Ice Sheet. In the northwest, the plateau is underlain by conglomerate, sandstone, carbonaceous shale and coal activities may adversely affect stream courses and conditions, and impact fish Canada, Open File 6430, 17 pages. and wildlife resources Upper Cretaceous Wapiti Formation). Over much of the central map area, shale, siltstone and sandstone (Upper Cretaceous Kotaneelee Huntley, D., Hickin, A. and Chow, W., 2011a. Surficial geology, geomorphology, granular resource evaluation and geohazard assessment for the Maxhamish Lake map area (NTS 94-O), northeastern British Columbia; Formation) lie beneath surficial units. To the east, conglomerate, sandstone and carbonaceous shale (Dunvegan Formation) forms the Maxhamish Escarpment. Undifferentiated clastic bedrock (Lower Cretaceous Fort St. John Group) underlies the easternmost limit of the map Alluvial floodplain sediments: gravel, sand and silt; massive, trough cross-bedded, rippled-bedded, planar stratified; well to rapidly drained; greater Geological Survey of Canada, Open File 6883, 20 pages. than 2 m thick; underlain by till or bedrock; transported and deposited by Huntley, D.H., Hickin, A.S. and Ferri, F., 2011b. Provisional surficial geology, glacial history and paleogeographic modern rivers, streams and creeks; subject to seasonal flooding; land use Topography and drainage patterns were greatly modified during the phase of maximum ice cover (>18 ^{*}C ka BP or >21.4 calendar ka BP). reconstructions of the Toad River (NTS 94-N) and Maxhamish Lake map areas (NTS 94-O), British Columbia. activities may adversely affect stream courses and conditions, and impact fish olidated sediment thicknesses in excess of 2-5 m are observed in major valleys and it is suspected that similar drift thicknesses blanket Geoscience Reports 2011, BC Ministry of Energy, pages 37-55. bedrock (unit R) across the map area. Silt- and clay-rich Laurentide tills have low clast contents (<20%) of proximally derived Cretaceous iliciclastic sedimentary rocks and distal igneous and metamorphic clasts from the Canadian Shield, hundreds of kilometres to the northeast Huntley, D.H. and Sidwell, C.F., 2010. Application of the GEM surficial geology data model to resource evaluation Colluvial deposits Drumlin ridges up to several hundred metres in length suggest clay-rich tills (unit Ts) were deposited beneath active, rapidly flowing warmand geohazard assessment for the Maxhamish Lake map area (NTS 94-O), British Columbia. Geological based glacial ice (Huntley and Hickin, 2010; Huntley et al., 2011b). Streamlined till (unit Ts) is most pronounced south of the Petitot River and Survey of Canada, Open File 6553, 22 pages. Colluvial veneer: clast-supported diamictons and rubble; massive to stratified, west of the Maxhamish Escarpment where ice flowed down slope, and thick accumulations of till were deposited over soft bedrock and unconsolidated advance-phase sediments. Multiple directions and generations of iceflow are preserved in streamlined landforms across the poorly-sorted; well to rapidly drained; deposits less than 2 m thick; landslide Stott, D.F. and Taylor, G.C., 1968. Geology of Maxhamish Lake. Geological Survey of Canada, Map 2-1968, headscarps range from 300 m to 10.5 km; formed by the weathering and map area, that suggest converging and diverging ice streams prior to the onset of deglaciation. Iceflow from the northeast was deflected down-slope movement of earth materials by gravitational processes; bedrock northwest by ice flowing west toward the Liard River valley. Maxhamish Lake and numerous smaller basins were excavated by erosion and and unconsolidated debris on slopes above 10-15° with greater than 5 m relief Trommelen, M. and Levson, V.M., 2008. Quaternary stratigraphy of the Prophet River, northeastem British prone to mass-wasting; rock falls, topples, rock slides and debris flows occur Columbia; Canadian Journal of Earth Sciences, Vol. 45, pages 565-575. Declaciation began sometime after 18 ¹⁴C ka BP (or >21.4 calendar ka BP) and ended before 10 ¹⁴C ka BP (ca. 12 calendar ka BP) with the where shale, sandstone and carbonate strata is exposed close to the surface; retreating active Laurentide Ice Sheet, stagnant ice masses in lowlands, glaciofluvial outwash and landslide debris blocking and reordering retrogressive rotational debris slides, debris flows and slumps occur in regional drainage. The mapped distribution of moraine ridges (unit Tm) implies that ice margins receded to the east across the map shee glaciolacustrine sediments and outwash containing sporadically discontinuous ACKNOWLEDGMENTS (Huntley and Hickin, 2010). Some large end moraines are deformed and streamlined suggesting that receding lobes remained active during permafrost; where ground ice is found slope failure can occur on surfaces less retreat and occasionally rapidly advanced. Minor moraine ridges drape drumlins in cross-cutting patterns and are interpreted as crevassi fillings and squeeze moraines deposited shortly after drumlinization ended, or as ice retreated from the map area (Huntley et al., 2011b). than 5°; slope instability could present major problems for construction in Canadian Geoscience Map 120 is an output of the Geo-Mapping for Energy and Minerals Yukon Basins Project Hummocky till (unit Th) found with short segments of subareal-subglacial meltwater channels and eskers indicate that bodies of stagnant managed by Carl Ozyer and Larry Lane (GSC-Calgary). The assistance of Robert Cocking, Sean Eagles, Vic Dohar, Mike Sigouin, Scott Tweedy and Martin Legault (NRCAN Scientific Publishing Services) was greatly glacier ice remained in lowland areas west of the Maxhamish Escarpment. Eskers (unit GFr) are composed of hummocky till and glaciofluvial gravelly sand, and likely exploited pre-existing crevasse patterns beneath the retreating ice sheet or stagnant ice bodies (Huntley et al., 2011a; Late Pleistocene earth materials and landforms appreciated throughout the map-making process. A critical review of CGM 120 was provided by Ron DiLabio luntley et al., 2011b). Ås ice retreated from the map area, a proglacial lake system formed over the Liard Plateau. Proglacial lakes were linked by spillways that drained meltwater northward into the Mackenzie River basin. In the map area, glaciolacustrine deposits (unit GLb). glaciofluvial terraces (unit GFt), and meltwater channels incised into till and bedrock indicate that glacial lake levels fell stepwise through Glaciolacustrine blanket: silt and clay with subordinate sand, gravel and deglaciation, with stable elevations at approximately 420 m, 380 m and <300 m. Fine-grained glacial earth materials have been re-worked by diamicton; massive or rhythmically interbedded; slump structures and ropstones locally present; poor to moderately drained; generally greater than 2 m thick; kettle lakes and irregular topography underlain by bedrock, tills and Post-glaciation (10 ¹⁴C ka BP, or ca. 12 calendar ka BP to present), changes in regional base-level led to episodes of channel incision and aggradation, resulting in the formation of erosional alluvial terraces along most stream and river valleys. In the early Holocene, pulses of fluvial outwash; transported by and deposited from sediment-laden meltwater. terrace building followed initial valley incision by the Liard and other major rivers. Most streams and rivers have alluvial fans (unit Af) and subaqueous gravity flows and melting of ice in proglacial lakes; where terraces (unit At) <5 m above active floodplains (unit Ap) consisting of gravel overlain by silt and sand. Poorly drained clay-rich till on the sporadically discontinuous permafrost is, or was present, glaciolacustrine plateaux and glaciolacustrine sediments in lowland areas are covered by extensive postglacial peat deposits (unit Owb), fens (unit Owf) and sediments may be subject to thermokarst processes; slopes less than 5° are ndifferentiated wetlands (unit O). Discontinuous permafrost is sporadically encountered in glaciolacustrine and some peat deposits. potentially unstable and prone to landslides and debris flows. Charcoal, observed in dug pits on alluvial terraces, suggest forest fires may have contributed to periods of landslide activity on slopes and local fluvial aggradation. Landslides and colluviated deposits (units Cv, Cb) are common where bedrock outcrops form escarpments, and where Glaciofluvial deposits shale or fine-grained glacial deposits are exposed along steep cutbanks. Stream networks and wetlands draining plateau watersheds are disrupted by beaver activity and, to a lesser extent, by roads and infrastructure where they cross streams, rivers and organic deposits (Huntley Esker ridges: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamicton; generally massive to stratified, some slump structures; moderately to well-drained; greater than 2 m thick; range from 100 m to 8.8 km in length; in contact with, and overlying till units, outwash and La Carte géoscientifique du Canada 120 illustre la glaciolacustrine sediments; deposited by subglacial meltwater in contact with géologie des matériaux superficiels et les formes de glacial ice, potential source of groundwater and granular aggregate when terrain d'un territoire d'environ 790 km² couvert par le feuillet cartographique de Maxhamish Lake (SNRC 94material is gravel rich. O/14), dans le nord-est de la Colombie-Britannique. Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and Dans la région cartographique, le plateau de Liard se matrix-supported diamicton; generally massive to stratified, some slump termine à l'ouest contre l'escarpement de Maxhamish structures; moderately to well-drained; greater than 2 m thick; terrace scarps de direction nord. À l'ouest du lac Maxhamish, le range from 100 m to 8 km in length; in contact with, and overlying other till ruisseau Itssi et les affluents du ruisseau Sandv se units, outwash and glaciolacustrine sediments; deposited by meltwater déversent à l'ouest dans la rivière Liard. Le ruisseau confined to proglacial channels and spillways; potential source of groundwater d'Easum, Coule vers le nord-est pour se jeter dans la and granular aggregate when material is gravel rich. rivière Petitot. Le lac Maxhamish occupe un bassin en position centrale avec une petite zone d'alimentation et un émissaire se versant dans le ruisseau d'Easum. Le Hummocky till: sand and silt-rich diamictons; massive to stratified, matrixsocle rocheux est couvert de matériaux terrestres non and clast-supported: clast contents less than 20% and contain sub-rounded consolidés remontant au Pléistocène supérieur granitic erratic boulders with sources on the Canadian Shield: moderately to (Glaciation du Wisconsinien supérieur, de > 25 ka à well-drained; greater than 2 m thick; drapes till and other glacial deposits; env. 10 ka) ainsi que de matériaux non glaciaires de deposited by in situ melting from stagnant cold-based ice and modified by l'Holocène (d'env. 10 ka jusqu'à nos jours). Les dépôts meltwater; evidence for ice collapse includes slump structures, kettle lakes de till, de couleur verte sur la carte, sont généralement gravel rich; generally suitable for infrastructure placement. dépôts fluvioglaciaires, qui recèlent un potentiel en minéraux, en agrégats et en eau souterraine, sont Moraine ridges: sand, silt and clay-rich diamictons; massive figurés par la couleur orange. Les versants dérangés matrix-supported; clast contents less than 20% and contain sub-rounded par des glissements de terrain, des coulés de débris et granitic erratic boulders with sources on the Canadian Shield; moderately to des chutes de blocs sont représentés en brun et en well-drained; greater than 2 m thick; minor moraines less than 1 km long and 5 rose. Les dépôts glaciolacustres et organiques, qui m high; major moraines up to 12.5 km in length and 10 m high; ridges drape renferment sporadiquement du pergélisol discontinu bedrock and older glacial deposits; minor moraines include crevasse-fill ridges sont représentés en violet et en gris. Les dépôts and small recessional push moraines; major ridges features are large alluviaux sujets aux inondations, à l'érosion et à la recessional end moraines and ice-thrust ridges; generally suitable for infrastructure placement. Streamlined till: silt and clay-rich diamictons; massive, matrix-supported and compact; clast contents less than 20% and contain sub-rounded granitic erratic boulders with sources on the Canadian Shield; moderately well-drained; greater than 2 m thick mantling bedrock and older glacial deposits; drumlins and fluted till ridges typically under 1 km long but can exceed 9 km in length; generally less than 50 m wide and 20 m high; formed beneath the Laurentide Ice Sheet directly through lodgement, basal meltout, glacigenic deformation of sediment beneath rapidly-flowing warm-based ice; generally suitable for infrastructure placement. Pre-Quaternary earth materials and landforms Undifferentiated bedrock: conglomerate, sandstone, siltstone, shale and limestone; exposed in escarpments between 300 m and 80 km in length; slopes above 10-15° with greater than 5 m relief prone to rock falls, topples rock slides and debris flows; Paleozoic unconformably overlain by Mesozoic sedimentary rocks; limestone and clastic sedimentary rocks are a potential source of crushed granular aggregate. ---- Geological boundary (Confidence: approximate) • • • • • • • Major moraine ridge (end, interlobate, or unspecified) Other moraine ridge (DeGeer, minor lateral, recessional, rogen, washboard, other transverse or unspecified) <><><><> Esker ridge (sense: unknown or unspecified) © Her Majesty the Queen in Right of Canada 2013 Drumlin ridge Major meltwater channel scarp ######### Minor meltwater channel central axis (marginal, overflow, subglacial or unspecified; sense: known) Terrace scarp (environment: fluvial) Station location (ground observation or stratigraphic section) Recommended citation Huntley, D.H., Hickin, A.S., Chow, W., and Mirmohammadi, M., 2013. Surficial geology, Maxhamish Lake, British Columbia; Geological Survey of Canada, Canadian Geoscience Map 120 (preliminary), scale 1:50 000. doi:10.4095/292397 78 79 80 82 83 84 123°00' **Preliminary CANADIAN GEOSCIENCE MAP 120 Preliminary Preliminary Preliminary SURFICIAL GEOLOGY** Authors: D.H. Huntley, A.S. Hickin, W. Chow, and Geomatics by D.H. Huntley, W. Chow, and Map projection Universal Transverse Mercator, The Geological Survey of Canada welcomes zone 10. North American Datum 1983 M. Mirmohammadi M. Mirmohammadi corrections or additional information from users. **MAXHAMISH LAKE** Base map at the scale of 1:50 000 from Natural This publication is available for free download through Geology by D.H. Huntley and A.S. Hickin (2009-2010) Cartography by W. Chow Resources Canada, with modifications. GEOSCAN (http://geoscan.ess.nrcan.gc.ca/). **British Columbia** Geological compilation by D.H. Huntley (2009–2011) Initiative of the Geological Survey of Canada, Elevations in feet above mean sea level conducted under the auspices of the Yukon Basin reliminary publications in Magnetic declination 2013, 20°38'E. Project as part of Natural Resources Canada's Geothis series have not been

decreasing 22' annually.

CANADIAN GEOSCIENCE MAP 120

mapping for Energy and Minerals (GEM) program

DESCRIPTIVE NOTES

protect natural resources, infrastructure and communities vulnerable to climate change in Canada's north.

of the GEM Geological Map Flow process (cf. Huntley and Sidwell, 2010; Huntley et al., 2011a; Deblonde et al., 2012).

ice-thrusting as Laurentide ice and subglacial meltwater scoured and deformed older glacial deposits and bedrock.

eolian activity and discontinuous loess covers glacial lake and till deposits in some places.

APPROACH TO SURFICIAL GEOLOGY MAPPING

glaciolacustrine and glaciofluvial sediments, tills and areas of bedrock.

INFERRED GEOLOGICAL HISTORY

and Hickin, 2010; Huntley and Hickin, 2011a-b).

Canadian Geoscience Map 120 depicts the surficial

geology and landforms over some 790 km² covered by

the Maxhamish Lake map sheet (NTS 94-O/14) in

northeastern British Columbia. In the map area, the

western limit of the Liard Plateau terminates along the

north-trending Maxhamish Escarpment. West of

Maxhamish Lake, Itssi Creek and tributaries to Sandy

flows northeast into the Petitot River. Maxhamish Lake

occupies a centrally located basin with a small

catchment area and an outlet draining into d'Easum

Creek. Bedrock is mantled by unconsolidated earth

materials dating to the Late Pleistocene (Late

Wisconsinan Glaciation, > 25 ka to ca. 10 ka) and non-

glacial Holocene (ca. 10 ka to present). Deposits of till,

green on the map, are generally suitable for placement

of infrastructure. Glaciofluvial deposits with mineral,

aggregate, and groundwater potential are coloured

orange. Slopes disturbed by landslides, debris flows,

and rock falls appear brown and pink. Glaciolacustrine

permafrost are coloured purple and grey. Alluvial

deposits prone to flooding, erosion, and sedimentation

sédimentation apparaissent en jaune sur la carte.

Catalogue No. M183-1/120-2013E-PDF

Preliminary

Canadian

Geoscience Maps

Canada

ISBN 978-1-100-21791-8

doi:10.4095/292397

CGM 120 CGM 119

CGM 122 | CGM 125 | CGM 126

National Topographic System reference and index to adjoining published Geological Survey of Canada maps

appear yellow on the map.

Cover illustration

Canada

British Columbia

1:50 000

SURFICIAL GEOLOGY

MAXHAMISH LAKE

View north of moraine ridges and drumlins incised

Natural Resources Ressources naturelles

CANADIAN GEOSCIENCE MAP 120

du Canada

meltwater channels on the Liard Plateau near

Maxhamish Lake, northeast British Columbia.

Photograph by D.H. Huntley. 2013-098

Creek drain west into the Liard River, d'Easum Creek

area (Stott and Taylor, 1968).

INTRODUCTION

CANADIAN GEOSCIENCE MAP 120 SURFICIAL GEOLOGY MAXHAMISH LAKE British Columbia

scientifically edited.