

INTRODUCTION
 This Surficial Geology Map of NTS 94-06 (Canadian Geoscience Map 128) is the product of collaboration between the Geological Survey of Canada and the British Columbia Ministry of Energy, Mines, and Natural Gas as part of the Geoscience for Energy and Minerals Program (GEM-EM) Yukon Basins Project. The accompanying geodatabase includes field observation points and field photos, landform features as lines, and surficial geology unit polygons. The map and geodatabase are essential tools for geoscience information and potential end users including resource explorationists, geotechnical engineers, land-use managers, terrestrial ecologists, archaeologists, geologists, and consultants in northern British Columbia. By providing insight into the distribution and origin of surficial earth materials, CGM 128 will help to reduce the economic costs and risks associated with the sustainable development of energy and mineral resources in NTS 94-06. Environmental impact assessments for new access roads, work camps, well pads, pipelines and power transmission line corridors, water storage and waste management systems and other infrastructure will benefit from the geoscience information presented here. By identifying areas prone to geological hazards (e.g., mass wasting, permafrost, flooding), CGM 128 will also help to protect natural resources, infrastructure and communities vulnerable to climate change in Canada's north.

APPROACH TO SURFICIAL GEOLOGY MAPPING
 Terrain mapping and field-based geomorphology are critical to a better understanding of the regional distribution of surficial deposits, permafrost, landslides and other geomorphic processes in the NTS 94-06 map area (Huntley and Hickin, 2010; Huntley et al., 2011a-b). Surficial materials and landforms were classified using a combination of oblique air photos (8c99016, 15c299101, 15c299105, 15c299110, 15c299175 and 15c299185 photos), LANDSAT 7 satellite imagery (510-2600100000), AIRS (2011) and Shuttle Radar Topography Mission digital elevation models (<http://sftp2.cr.usgs.gov>), (RFL 2011). The base map was generated from CANVEC shape files (<http://geospatial.gc.ca/cegis/geoproducts>) (LRS, 2011). Surficial geology polygons and landform line symbols were digitized using commercially available computer software packages (Global Mapper, ArcMap and ArcGIS) and compared to published maps, reports and ancient digital data (e.g., Stett and Taylor, 1968; Stettmann, 2002a-d; Charnel et al., 2004; Beckwith, 2005a; Demchuk, 2010). The geodatabase accompanying this map conforms to the Science Language for the Data Management component of the GEM Geological Map Framework (cf. Huntley and Sedell, 2010; Huntley et al., 2011a; DeBlonde et al., 2012).

INFERRED GEOLOGICAL HISTORY
 This inferred geological history of NTS 94-06 is largely a product of underlying bedrock and geological structures, with ornamentation by the Late Wisconsinan Laurentide Ice Sheet. The Laird Plateau and Mauchamish Escarpment are underlain by conglomerate, sandstone and siltstone and metasediments of the Lower Cretaceous. Underlain by the Lower Cretaceous Fort St. John Group (Blatt and Taylor, 1963). Topography and drainage patterns were greatly modified by the Laurentide Ice Sheet (~15 ka) which covered the area from 15 ka to 10 ka. Thick basal tills and silty tills are observed in major valleys and it is suspected that similar drift thicknesses blanket bedrock (unit Tm) across the map area. In the northwest, the Laurentide Ice Sheet is suspected to have been a proglacial or marginal ice sheet. Several hundred kilometres west of the Laird Plateau and Mauchamish Escarpment, from the Yukon River to the Mackenzie River, hundreds of kilometres to the northeast. West of the Mauchamish Escarpment, from the Yukon River to the Mackenzie River, hundreds of kilometres to the northeast. West of the Mauchamish Escarpment, from the Yukon River to the Mackenzie River, hundreds of kilometres to the northeast. West of the Mauchamish Escarpment, from the Yukon River to the Mackenzie River, hundreds of kilometres to the northeast. West of the Mauchamish Escarpment, from the Yukon River to the Mackenzie River, hundreds of kilometres to the northeast.

ACKNOWLEDGMENTS
 Canadian Geoscience Map 128 is an output of the Geo-Mapping for Energy and Minerals Yukon Basins Project managed by Carl Cooper and Larry Lutz (GSC-CEM). The assistance of Robert Coakley, Sean Eagles, Vic Chouh, Mikal Simpson, Scott Trusley and Martin Legendt (NRCan Scientific Publishing Services) was greatly appreciated throughout the map-making process. A critical review of CGM 128 was provided by Isabelle Melinard (GSC-CEM).

Abstract
 Canadian Geoscience Map 128 depicts the surficial geology over some 750 km² covered by the Patry Lake map sheet (NTS 94-06) in northeastern British Columbia. The map (CGM 128) in the northeast of the southwest limit of the Laird Plateau, which terminates along the southwest to northeast-oriented Mauchamish Escarpment, and the southwest limit of the Fort Nelson Lowland. The plateau and lowland are incised by east- and south-draining tributaries to the Fort Nelson River. Bedrock is mantled by unconsolidated earth materials dating to the Late Pleistocene (Late Wisconsinan Glaciation) and Holocene (ca. 10 ka to present). Deposits of till, green on the map, are suitable for placement of infrastructure. Glaciofluvial and colluvial deposits with mineral, aggregate, and groundwater potential are coloured orange and buff. Slopes disturbed by landslides and debris flows appear brown. Glaciofluvial and organic deposits with sporadically discontinuous permafrost are coloured purple and grey. Alluvial deposits prone to flooding, erosion, and sedimentation appear yellow on the map.

| | | |
|---------|---------|---------|
| CGM 122 | CGM 125 | CGM 126 |
| CGM 109 | CGM 128 | CGM 127 |
| CGM 108 | CGM 107 | CGM 106 |

National Topographic System reference and index to adjoining published Geological Survey of Canada maps



Cover illustration: Organic deposits, moraines and drum ridges in the vicinity of Patry Lake and the Mauchamish Escarpment in northeast British Columbia, view south. Photograph by D.H. Huntley, 2013-106
 Catalogue No. M183-1125-2013E-PDF ISBN 978-1-105-21799-4 doi:10.4095/292405
 © Her Majesty the Queen in Right of Canada 2013

Natural Resources Canada Ressources naturelles du Canada

CANADIAN GEOSCIENCE MAP 128 SURFICIAL GEOLOGY PATRY LAKE British Columbia 1:50 000



Canadian Geoscience Maps
 Authors: D.H. Huntley, A.S. Hickin, W. Chow, and M. Mirzohammadi
 Geomatics by D.H. Huntley, W. Chow, and M. Mirzohammadi
 Cartography by W. Chow
 Initiative of the Geological Survey of Canada, conducted under the auspices of the Yukon Basin Project as part of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) program



SURFICIAL GEOLOGY PATRY LAKE British Columbia 1:50 000

Map projection: Universal Transverse Mercator, zone 10, North America Datum 1983
 Base map at the scale of 1:50 000 from Natural Resources Canada, with modifications.
 Elevations in feet above mean sea level
 Magnetic declination 2013, 20°27'E, decreasing 21' annually
 The Geological Survey of Canada welcomes corrections or additional information from users.
 This publication is available for free download through GEOCAN (<http://geocan.nrc.ca>).



- Holocene earth materials and landforms**
- Owb** Peat bogs: fibric to humic organic matter, massive to stratified accumulations, generally greater than 2 m thick; confined to topographic depressions or level areas, underlain by poorly drained till, glaciofluvial and other unconsolidated sediments, formed by decomposition of plant material in wetland areas; slopes with sporadically discontinuous permafrost and thermokarst depressions potentially unstable if organic material is disturbed or removed.
 - Owf** Fans: fibric organic matter, massive to stratified, generally greater than 2 m thick, confined to topographic depressions, level areas or channels; underlain by poorly drained till, glaciofluvial and other unconsolidated sediments, formed by decomposition of plant material in wetland areas; fans are prone to flooding following drainage damming by beaver activity.
 - O** Undifferentiated peat bogs and fans: humic to fibric organic matter, massive to stratified accumulations, generally greater than 2 m thick; confined to topographic depressions, level areas or channels; underlain by poorly drained till, glaciofluvial and other unconsolidated sediments, formed by decomposition of plant material in wetland areas; may contain sporadically discontinuous permafrost and thermokarst depressions, potentially unstable if disturbed or removed during development.
 - Af** Alluvial fan sediments: boulders, gravel, sand and silt; generally massive to planar stratified, well to rapidly drained, greater than 2 m thick; fan morphology with slopes up to 15°; may contain interbedded debris flows and banded organic material; transported and deposited by modern rivers, streams and creeks; subject to periodic flooding; potential source of aggregate.
 - AI** Alluvial terrace sediments: boulders, gravel, sand and silt, generally massive to planar stratified, well to rapidly drained, greater than 2 m thick; may contain interbedded debris flows and banded organic material; underlain by bedrock, silt and loess; transported and deposited by modern rivers, streams and creeks; subject to rare flooding; potential land use activities may adversely affect stream courses and conditions, and impact fish and wildlife resources.
 - Ap** Alluvial floodplain sediments: gravel, sand and silt; massive, tough crossbedded, rippled-bedded, planar stratified, well to rapidly drained, greater than 2 m thick, underlain by fill or bedrock; transported and deposited by modern rivers, streams and creeks; subject to seasonal flooding; land use activities may adversely affect stream courses and conditions, and impact fish and wildlife resources.
 - Cv** Colluvial veneer: clast-supported diamictons and rubble; massive to stratified, poorly-sorted, well to rapidly drained, deposits less than 2 m thick; landslide headscapes range from 300 m to 10 to 15 km, formed by the weathering and down-slope movement of earth materials by gravitational processes; bedrock and unconsolidated debris on slopes above 10 m relief greater than 5 m relief prone to mass-washing, rock falls, topples, rock slides and debris flows occur where shale, sandstone and carbonate strata is exposed close to the surface; retrogressive rotational debris slides, debris flows and slumps occur in glaciofluvial sediments and outwash containing sporadically discontinuous permafrost, where ground ice is found; slope failure can occur on surfaces less than 5°; slope instability could present major problems for construction in some areas.
 - Cb** Colluvial blanket: clast-supported diamictons and rubble; massive to stratified, poorly-sorted, well to rapidly drained, deposits greater than 2 m thick; landslide headscapes range from 300 m to 10 to 15 km, formed by the weathering and down-slope movement of earth materials by gravitational processes; bedrock and unconsolidated debris on slopes above 10 m relief greater than 5 m relief prone to mass-washing, rock falls, topples, rock slides and debris flows occur where shale, sandstone and carbonate strata is exposed close to the surface; retrogressive rotational debris slides, debris flows and slumps occur in glaciofluvial sediments and outwash containing sporadically discontinuous permafrost, where ground ice is found; slope failure can occur on surfaces less than 5°; slope instability could present major problems for construction in some areas.
- Late Pleistocene earth materials and landforms**
- Glaciofluvial deposits**
- GLB** Glaciofluvial blanket: silt and clay with subordinate sand, gravel and diamicton; massive to rhythmically interbedded, slump structures and diaposites locally present; poorly to moderately drained, generally greater than 2 m thick; kettle lakes and irregular topography underlain by bedrock, tills and outwash; transported and deposited from sediment-laden meltwater and subaqueous gravity flows and melting of ice in proglacial lakes, where sporadically discontinuous permafrost, or was present; glaciofluvial sediments may be subject to thermokarst processes, slopes less than 5° are potentially unstable and prone to landslides and debris flows.
 - GFl** Kames and hummocky outwash: boulders, cobbles, pebble-gravel, sand, silt and diamicton; generally massive to stratified, slopes moderate to well-drained, greater than 2 m thick, irregular hummocks and anticline topography; in contact with and overlain by tills, outwash and glaciofluvial sediments; deposited by rivers and streams flowing from, or in contact with glacial ice; potential source of groundwater and granular aggregate when material is gravel rich.
 - GFl** Outwash terraces: boulders, cobbles, pebble-gravel, sand, silt and matrix-supported diamicton; generally massive to stratified, some slump structures; moderately to well-drained; greater than 2 m thick, terrace scapes range from 100 m to 8 km in length; in contact with, and overlying other till units, outwash and glaciofluvial sediments; deposited by meltwater confined to proglacial channels and spillways; potential source of groundwater and granular aggregate when material is gravel rich.
 - Till deposits**
 - Th** Hummocky till: sand and silt-rich diamictons, massive to stratified, matrix- and clast-supported, clast contents less than 20% and contain sub-rounded granitic-erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick, minor moraines less than 1 km long and 5 m high, major moraines up to 12.8 km in length and 10 m high; ridges drape bedrock and older glacial deposits; minor moraines include crevasse-fill ridges and small recessional push moraines; major ridges feature are large recessional end moraines and ice-sheet ridges; generally suitable for infrastructure placement.
 - Tm** Moraine ridges: sand, silt and clay-rich diamictons; massive, matrix-supported, clast contents less than 20% and contain sub-rounded granitic-erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick, minor moraines less than 1 km long and 5 m high, major moraines up to 12.8 km in length and 10 m high; ridges drape bedrock and older glacial deposits; minor moraines include crevasse-fill ridges and small recessional push moraines; major ridges feature are large recessional end moraines and ice-sheet ridges; generally suitable for infrastructure placement.
 - Ts** Streamlined till: silt and clay-rich diamictons, massive, matrix-supported and compact, clast contents less than 20% and contain sub-rounded granitic-erratic boulders with sources on the Canadian Shield; moderately to well-drained; greater than 2 m thick, marling bedrock and older glacial deposits; deposited by in situ melting from stagnant cold-based ice and modified by meltwater; evidence for ice collapse includes slump structures, kettle lakes and irregular topography; potential source of aggregate when material is gravel rich; generally suitable for infrastructure placement.
 - Tv** Till veneer: sand, silt and clay-rich diamictons; massive, matrix-supported and compact, clast contents less than 20% and contain sub-rounded granitic-erratic boulders with sources on the Canadian Shield; moderately to well-drained; less than 2 m thick, draping bedrock and older glacial deposits; transported and deposited by the Laurentide Ice Sheet directly through lodgement, basal meltout, glacialic deformation beneath active, warm-based ice and in situ melting from stagnant cold-based ice; generally suitable for infrastructure placement.
- Pre-Quaternary earth materials and landforms**
- R** **Bedrock**
 Undifferentiated bedrock: conglomerate, sandstone, siltstone, shale and limestone, exposed in escarpments between 300 m and 80 km in length; slopes above 10° with greater than 5 m relief prone to rock falls, talus, rock slides and debris flows; Paleozoic unconformably overlain by Mesozoic sedimentary rocks; limestone and clastic sedimentary rocks are a potential source of crushed granular aggregate.

REFERENCES

Deblonde, J.M., 2003a. Batalemma Lake, Northwest Territories – Yukon Territory – British Columbia (NTS 9664); Geological Survey of Canada, Open File 6502, 1 map, scale 1:50,000.
 Deblonde, J.M., 2003b. Surficial geology of Fort Laird, Northwest Territories – British Columbia. Geological Survey of Canada, Open File 1704, scale 1:50,000.
 Deblonde, J.M., 2003c. Surficial geology of Lake Bovik, Northwest Territories – British Columbia. Geological Survey of Canada, Open File 1761, scale 1:50,000.
 Deblonde, J.M., 2004a. Surficial geology of Colwell Lake, Northwest Territories – British Columbia. Geological Survey of Canada, Open File 1754, scale 1:50,000.
 Deblonde, J.M., 2004b. Surficial geology of Etienne Creek, British Columbia, Geological Survey of Canada, Open File 4825, scale 1:50,000.
 Deblonde, J.M., 2004c. Surficial geology of Gato Creek, British Columbia, Geological Survey of Canada, Open File 4844, scale 1:50,000.
 Charnel, C., Kwiatk, R., Huntley, D. and Deblonde, R., 2004. Ecosystem units of the Sahtanen area. Sloon Forest Products (Fort Nelson) Report, 39 pages and appendices.
 Deblonde, C., Pfouffa, A., Boisvert, E., Bulter, G., Davernport, P., Everett, D., Huntley, D., Ingile, E., Korn, D., Moore, A., Paradis, S.J., Parnet, M., Smith, R., St-Onge, D., and Weatherston, A., 2012. Science Language for an Integrated Geological Survey of Canada. Data Model for Surficial Geology, Version 1.1. Results of Geological Survey of Canada Surficial Legend Review Committee. Geological Survey of Canada, Open File 7303, 227 pages.
 Demchuk, T., 2010. Surficial geology of the Komik Creek area (NTS 094905). British Columbia Ministry of Energy, Mines and Petroleum Resources, Open File 210-08, glacial history and potential for granular aggregate and fill; area: Mauchamish Lake Map Area (NTS 94-03), British Columbia, Geological Survey of Canada, Open File 6838, 17 pages.
 Huntley, D.H. and Hickin, A.S., 2010. Surficial deposits, landforms, glacial history and potential for granular aggregate and fill; area: Mauchamish Lake Map Area (NTS 94-03), British Columbia, Geological Survey of Canada, Open File 6838, 17 pages.
 Huntley, D., Hickin, A.S. and Chow, W., 2014. Surficial geology, geomorphology, granular resource evaluation and geohazard assessment for the Mauchamish Lake map area (NTS 94-03), northeastern British Columbia, Geological Survey of Canada, Open File 8883, 24 pages.
 Huntley, D.H., Hickin, A.S. and Ferris, J., 2015. Preliminary surficial geology, glacial history and paleogeographic reconstructions of the Toof River (NTS 94-04) and Mauchamish Lake map areas (NTS 94-03), British Columbia, Geoscience Reports 2015, IBC Ministry of Energy, pages 37-50.
 Huntley, D.H. and Sedell, C.F., 2010. Application of the GEM surficial geology data model to resource evaluation and geohazard assessment for the Mauchamish Lake map area (NTS 94-03), British Columbia, Geological Survey of Canada, Open File 6838, 22 pages.
 Stett, D.F. and Taylor, G.C., 1968. Geology of Mauchamish Lake, Geological Survey of Canada, Map 2-1968, scale 1:250,000.

Recommended citation
 Huntley, D.H., Hickin, A.S., Chow, W., and Mirzohammadi, M., 2013. Surficial geology, Patry Lake, British Columbia, Geological Survey of Canada, Canadian Geoscience Map 128 (preliminary), scale 1:50 000. doi:10.4095/292405

Preliminary publications in this series have not been scientifically edited.