

September 13 to September 26, 2015 (weeks 37 & 38)

Overall Summary

- Overall, there is low influenza activity in Canada; however, influenza activity and detections are increasing, especially in BC.
- In week 37, two influenza outbreaks were reported.
- In weeks 37 and 38, both paediatric and adult hospitalizations with influenza were reported.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

Are you a primary health care practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel for the 2015-16 influenza season? Contact us at FluWatch@phac-aspc.gc.ca

Influenza/Influenza-like Illness Activity (geographic spread)

In week 38, sporadic influenza activity was reported in the Western provinces and parts of Central Canada. The majority of influenza activity was reported in the BC and ON.

Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, Week 38




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available in mapping feature found in the [Weekly Influenza Reports](#).

Laboratory Confirmed Influenza Detections


Although the number of positive influenza tests increased over the two week period, the percent positive for influenza detections remains low (1.5%) (Figure 2).

Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2015-16



Most jurisdictions have reported only sporadic numbers of influenza detections in recent weeks. The majority of detections in Canada have been reported from BC, accounting for 77% of influenza detections in Canada in week 38. To date, 94% of influenza detections have been influenza A and the majority of those subtyped have been A(H3).

Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province, Canada, 2015-16



Note: Specimens from NT, YT, and NU are sent to reference laboratories in other provinces. Cumulative data includes updates to previous weeks.

Among cases with reported age, the largest proportion was in those ≥ 65 years of age (52%) (Table 1).

Table 1 – Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting¹, Canada, 2015-16

Age groups (years)	Weekly (September 13 to 26, 2015)					Cumulative (August 30, 2015 to September 26, 2015)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) ³		A Total	A(H1) pdm09	A(H3)	A (UnS) ³		Total	#
<5	1	0	1	0	0	3	0	3	0	0	3	5.2%
5-19	0	0	0	0	0	5	0	3	2	1	6	10.3%
20-44	1	0	0	1	0	5	0	1	4	0	5	8.6%
45-64	2	0	1	1	0	13	0	10	3	1	14	24.1%
65+	9	0	4	5	0	29	0	20	9	1	30	51.7%
Unknown	0	0	0	0	0	0	0	0	0	0	0	0.0%
Total	13	0	6	7	0	55	0	37	18	3	58	100.0%
Percentage²	100.0%	0.0%	46.2%	53.8%	0.0%	94.8%	0.0%	67.3%	32.7%	5.2%		

¹Table 1 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported.

²Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.


³UnS: untyped: The specimen was typed as influenza A, but no result for subtyping was available.

For additional data on other respiratory virus detections see the [Respiratory Virus Detections in Canada Report](#) on the Public Health Agency of Canada website.

Influenza-like Illness Consultation Rate

The national influenza-like-illness (ILI) consultation rate increased from 13.2 consultations per 1,000 patient visits in week 37 to 14.8 per 1,000 visits in week 38. In week 38, the highest ILI consultation rate was found in the 5-19 age group and the lowest was found in the ≥ 65 age group (Figure 4).

Figure 4 – Influenza-like illness (ILI) consultation rates by age group and week, Canada, 2015-16




Delays in the reporting of data may cause data to change retrospectively. In BC, AB, and SK, data is compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.

Influenza Outbreak Surveillance

In week 37, two new outbreaks of influenza were reported. One influenza A(H3) outbreak was reported in a long-term care facility (LTCF) and the other of unknown type was reported in an institutional or community setting (Figure 5). No outbreaks were reported in week 38.

Figure 5 – Overall number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2015-2016




¹All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of the report.

Sentinel Paediatric Hospital Influenza Surveillance


Paediatric Influenza Hospitalizations and Deaths (IMPACT)

In week 37, one laboratory-confirmed influenza-associated paediatric (≤ 16 years of age) hospitalization was reported by the Immunization Monitoring Program Active (IMPACT) network. The hospitalized case was aged 6-23 months and was due to influenza A(H3). This is the first hospitalization reported through IMPACT this season.

Figure 6 – Percentage of hospitalizations, ICU admissions and deaths with influenza reported by age-group, Canada, 2015-16, Paediatric hospitalizations (≤ 16 years of age, IMPACT)



**Figure 7 – Number of cases of influenza reported by sentinel hospital networks, by week, Canada, 2015-16
Paediatric hospitalizations (≤16 years of age, IMPACT)**




Note: The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated paediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Provincial/Territorial Influenza Hospitalizations and Deaths

Since the start of the 2015-16 season, nine laboratory-confirmed influenza-associated hospitalizations were reported from participating provinces and territories*. All were due to influenza A and the majority were patients ≥65 years of age. Two ICU admissions have also been reported.

* Note: Influenza-associated hospitalizations are not reported to PHAC by the following Provinces and Territory: BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by Saskatchewan. ICU admissions are not distinguished among hospital admissions reported from Ontario. Data may also include cases reported by the IMPACT networks. The number of new influenza-associated hospitalizations and deaths reported for the current week may include cases from Ontario that occurred in previous weeks, as a result of retrospective updates to the cumulative total. It is important to note that the hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.

Figure 8 – Percentage of hospitalizations, ICU admissions and deaths with influenza reported by age-group, Canada 2015-16



See additional data on [Reported Influenza Hospitalizations and Deaths in Canada: 2011-12 to 2015-16](#) on the Public Health Agency of Canada website.

Influenza Strain Characterizations

The National Microbiology Laboratory (NML) has not yet reported any influenza strain characterizations for the 2015-16 season.

Figure 9 – Influenza strain characterizations, Canada, 2015-16

The National Microbiology Laboratory (NML) has not yet conducted antigenic characterization of influenza viruses collected during the 2015-16 season.

The recommended components for the 2015-2016 northern hemisphere trivalent influenza vaccine include: an A/California/7/2009(H1N1)pdm09-like virus, an /Switzerland/9715293/2013(H3N2)-like virus, and a B/Phuket/3073/2013 -like virus (Yamagata lineage). For quadrivalent vaccines, the addition of a B/Brisbane/60/2008-like virus is recommended.

The NML receives a proportion of the number of influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Characterization data reflect the results of haemagglutination inhibition (HAI) testing compared to the reference influenza strains recommended by [WHO](#).

Antiviral Resistance

The NML has not yet reported antiviral resistance results for influenza viruses collected during the 2015-16 season (Table 2).

Table 2 – Antiviral resistance by influenza virus type and subtype, Canada, 2015-16

The National Microbiology Laboratory (NML) has not yet conducted antiviral resistance testing of influenza viruses collected during the 2015-16 season.

¹NA: Not Applicable

International Influenza Reports

[World Health Organization influenza update](#)

[World Health Organization FluNet](#)

[WHO Influenza at the human-animal interface](#)

[Centers for Disease Control and Prevention seasonal influenza report](#)

[European Centre for Disease Prevention and Control - epidemiological data](#)

[South Africa Influenza surveillance report](#)

[New Zealand Public Health Surveillance](#)

[Australia Influenza Report](#)

[Pan-American Health Organization Influenza Situation Report](#)

FluWatch Definitions for the 2015-2016 Season

Abbreviations: Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

Influenza-like-illness (ILI): Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI/Influenza outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

Hospitals and residential institutions: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Workplace: Greater than 10% absenteeism on any day which is most likely due to ILI.

Other settings: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.

Influenza/ILI Activity Levels

1 = No activity: no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

2 = Sporadic: sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

3 = Localized: (1) evidence of increased ILI* ;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

4 = Widespread: (1) evidence of increased ILI*;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

** More than just sporadic as determined by the provincial/territorial epidemiologist.*

† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.

We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.

This report is available on the Government of Canada Influenza webpage under [Weekly influenza reports](#).

Ce rapport est disponible dans les deux langues officielles.