

Carbon, Nitrogen and Sulfur Isotope Content in Bull Kelp (*Nereocystis luetkeana*) Collected Within and Adjacent to Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve and Haida Heritage Site, Haida Gwaii, British Columbia, Canada

H.L. Stewart, G.E. Piercey, and M. Nomura

Fisheries and Oceans Canada
Science Branch, Pacific Region
Centre for Aquaculture and Environmental Research
(CAER – West Vancouver Laboratory)
4160 Marine Drive
West Vancouver, BC V7V 1N6

2014

Canadian Data Report of Fisheries and Aquatic Sciences 1253



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Data Report of Fisheries and Aquatic Sciences

Data reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries management, technology and development, ocean sciences, and aquatic environments relevant to Canada.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written clearance from the issuing establishment. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Data Reports. The current series name was changed with report number 161.

Rapport statistique canadien des sciences halieutiques et aquatiques

Les rapports statistiques servent de base à la compilation des données de classement et d'archives pour lesquelles il y a peu ou point d'analyse. Cette compilation aura d'ordinaire été préparée pour appuyer d'autres publications ou rapports. Les sujets des rapports statistiques reflètent la vaste gamme des intérêts et politiques de Pêches et Océans Canada, notamment la gestion des pêches, la technologie et le développement, les sciences océaniques et l'environnement aquatique, au Canada.

Les rapports statistiques ne sont pas préparés pour une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans autorisation écrite préalable de l'établissement auteur. Le titre exact figure au haut du résumé de chaque rapport. Les rapports à l'industrie sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement d'origine dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 25 de cette série ont été publiés à titre de Records statistiques, Service des pêches et de la mer. Les numéros 26-160 ont été publiés à titre de Rapports statistiques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom de la série a été modifié à partir du numéro 161.

Canadian Data Report of
Fisheries and Aquatic Sciences 1253

2014

CARBON, NITROGEN AND SULFUR ISOTOPE CONTENT IN BULL KELP
(Nereocystis luetkeana) COLLECTED WITHIN AND ADJACENT TO
GWAII HAANAS NATIONAL PARK RESERVE, NATIONAL MARINE CONSERVATION
AREA RESERVE AND HAIDA HERITAGE SITE, HAIDA GWAII,
BRITISH COLUMBIA, CANADA

by

H.L. Stewart, G.E. Piercey, and M. Nomura

Fisheries and Oceans Canada
Science Branch, Pacific Region
Centre for Aquaculture and Environmental Research
(CAER – West Vancouver Laboratory)
4160 Marine Drive
West Vancouver, BC V7V 1N6

©Her Majesty the Queen in Right of Canada, 2014
Cat. No. Fs97-14/1253E ISSN 0706-6465

Correct citation for this publication:

Stewart, H.L., Piercey, G.E., and Nomura, M. 2014. Carbon, nitrogen and sulfur isotope content in bull kelp (*Nereocystis luetkeana*) collected within and adjacent to Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve and Haida Heritage Site, Haida Gwaii, British Columbia, Canada. Can. Data Rep. Fish. Aquat. Sci. 1253: iv + 11 p.

List of Figures

	Page
Figure 1. Locations of monitoring sites	4

List of Tables

	Page
Table 1. Names, locations and management policy of monitoring sites	5
Table 2. Carbon and nitrogen content in bull kelp tissue samples	6
Table 3. Sulfur content in bull kelp tissue samples	10

ABSTRACT

Stewart, H.L., Piercy, G.E., and Nomura, M. 2014. Carbon, nitrogen and sulfur isotope content in bull kelp (*Nereocystis luetkeana*) collected within and adjacent to Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve and Haida Heritage Site, Haida Gwaii, British Columbia, Canada. Can. Data Rep. Fish. Aquat. Sci. 1253: iv + 11 p.

Carbon, nitrogen and sulfur isotopes were analyzed from tissue samples of bull kelp (*Nereocystis luetkeana*) collected from 12 sites in Haida Gwaii, British Columbia. The data are expected to serve as a baseline for future nutrient cycling studies in the area, and may help with the evaluation of conservation and management policies for the Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve and rockfish conservation areas (RCAs).

RÉSUMÉ

Stewart, H. L., Piercy, G. E. et Nomura, M. 2014. Contenu en isotopes de soufre, d'azote et de carbone de la nereocystis de Lutke (*Nereocystis luetkeana*) recueillie à l'intérieur et à proximité de la Réserve de parc national, de la réserve d'aire marine nationale de conservation, et du site du patrimoine haïda Gwaii Haanas, Haida Gwaii (Colombie-Britannique) Canada. Rapp. stat. can. sci. halieut. aquat. 1253 : iv + 11 p.

Les auteurs ont analysé les isotopes de soufre, d'azote et de carbone dans les échantillons de tissus de la nereocystis de Lutke (*Nereocystis luetkeana*) recueillie à douze emplacements à Haida Gwaii, en Colombie-Britannique. Les données devraient servir de point de départ pour les prochaines études sur le cycle des éléments nutritifs dans la région et pourraient aider à l'évaluation des politiques de conservation et de gestion pour la Réserve de parc national et la réserve d'aire marine nationale de conservation Gwaii Haanas ainsi que les aires de conservation du sébaste (ACS).

INTRODUCTION

As part of the baseline ecosystem study conducted between 2009 and 2013 at the Gwaii Haanas National Marine Conservation Area Reserve (NMCAR) and associated rockfish conservation areas (RCAs) described by Trebilco et al. (2014), samples of bull kelp (*Nereocystis luetkeana*) collected in the summer of 2013 were taken and set aside for nutrient analysis.

Naturally occurring isotopes of carbon, nitrogen and sulfur tend to maintain their ratios as they cycle through a local ecosystem (Peterson and Fry 1987). Stable isotope ratios in kelp can vary according to various factors such as tidal gradient (Cooper and McCroy 1988), season and age (Frederiksen 2003, Dethier et al. 2013); however, ratios of multiple stable isotopes such as $^{13}\text{C}:\text{N}$ remain distinct for each species (Dethier et al. 2013). Thus it is possible to trace the flow of nutrients from primary producers such as *N. luetkeana* through the food chain (Hesslein et al. 1991, Ramshaw 2012, Fry 2013). Of particular interest, here on the British Columbia coast, may be the relationship between kelp, sea urchin and sea otters (Rodriguez 2003, Vanderklift et al. 2006, Ramshaw 2012).

METHODS

Study sites and sampling method

Bull kelp (*Nereocystis luetkeana*) samples were collected between July 15 and August 1 of 2013 from 12 monitoring sites as described by Trebilco et al. 2014 (Figure 1 and Table 1). A tissue sample from individual plants was cut from the blade approximately 15 cm above the bulb. These were placed in individually labeled Ziploc bags and frozen at -5°C in the field.

Nutrient analysis

In the laboratory, up to 10 samples from each site were dried for a minimum of 48 hours at 50 °C in a Sheldon Manufacturing Inc. Model 1305U oven. The samples were then ground in a mortar and pestle, and then ground further into a fine powder using a Rinn Crescent Wig-L-Bug MSD Amalgamator fitted with a stainless steel ball and polymeric vial. Five milligram aliquots were measured out using a Sartorius CPA225D analytical balance, and shipped to the University of California, Davis Stable Isotope Facility for carbon, nitrogen and sulfur isotope analyses.

Carbon and nitrogen were analyzed from a single sample using a PDZ Europa ANCA-GSL elemental analyzer interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd., Cheshire, UK). The results were presented relative to international standards V-PDB (Vienna PeeDee Belemnite) for carbon, and air for nitrogen. Sulfur was analyzed with an Elementar vario ISOTOPE cube interfaced to a SerCon 20-22 IRMS (Sercon Ltd., Cheshire, UK), and these results were presented relative to VCDT (Vienna Canyon Diablo Troilite) standards. For details see <http://stableisotopelab.ucdavis.edu/>.

REFERENCES

- Cooper, L.W. and McCroy, C.P. 1988. Stable carbon isotope ratio variations in marine macrophytes along intertidal gradients. *Oecol.* 77(2): 238-241.
- Dethier, M.N., Sosik, E., Galloway, A.W.E., Duggins, D.O., and Simenstad, C.A. 2013. Addressing assumptions: variation in stable isotopes and fatty acids of marine macrophytes can confound conclusions of food web studies. *Mar. Ecol. Prog. Ser.* 478: 1-14.
- Frederiksen, S. 2003. Food web studies in a Norwegian kelp forest based on stable isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) analysis. *Mar. Ecol. Prog. Ser.* 260: 71-81.
- Fry, B. 2013. Using stable CNS isotopes to evaluate estuarine fisheries condition and health. *Isotopes Environ. Health Stud.* 49(3): 295-304.
- Hesslein, R.H., Capel, M.J., Fox, D.E., and Hallard, K.A. 1991. Stable isotopes of sulfur, carbon, and nitrogen as indicators of trophic level and fish migration in the lower MacKenzie River basin, Canada. *Can. J. Fish. Aquat. Sci.* 48: 2258-2265.
- Peterson, B.J. and Fry, B. 1987. Stable isotopes in ecosystem studies. *Annu. Rev. Ecol. Evol. Syst.* 18:293-320.
- Ramshaw, B.C. 2012. Spatial and temporal variation in kelp-derived detritus and its dietary importance to consumers along the west coast of Vancouver Island, Canada. M.Sc. Thesis. University of British Columbia, Canada. 113 p.
- Rodríguez, S.R. 2003. Consumption of drift kelp by intertidal populations of the sea urchin *Tetrapygus niger* on the central Chilean coast: possible consequences at different ecological levels. *Mar. Ecol. Prog. Ser.* 251: 141-151.
- Trebilco, R., Demes, K.W., Lee, L.C., Keeling, B.E., Sloan, N.A., Stewart, H.L., and Salomon, A.K. 2014. Summary of baseline kelp forest surveys within and adjacent to Gwaii Haanas National Park Reserve, National Marine Conservation Area Reserve and Haida Heritage Site, Haida Gwaii, British Columbia, Canada. *Can. Data Rep. Fish. Aquat. Sci.* 1252: v + 25 p.
- Vanderklift, M.A., Kendrick, G.A., and Smit, A.J. 2006. Differences in trophic position among sympatric sea urchin species. *Estuar. Coast. Mar. Sci.* 66: 291-297.

ACKNOWLEDGEMENTS

We thank Jenn Burt, Ryan Cloutier, Matt Drake, Nick Duprey, Jim Hayward, Margot Hessing-Lewis, Sharon Jeffery, Britt Keeling, Joanne Lessard, Steve Romaine, Robert Russ, Leandre Vigneault, Taimen Lee Vigneault, Eric White, and Mark Wunsch for assistance with field data collection. We also thank Paula Anglada, Dominique Bureau, Seaton Taylor and DFO's Shellfish Group for logistical support and equipment loans. Ship time on a Canadian Coast Guard research fleet vessel provided by the Department of Fisheries and Oceans provided in-kind support for this work. We thank the Captains (Kent Reid and Simon Dockerill) and crew of the Canadian Coast Guard Ship Vector for their help facilitating fieldwork.

Figure 1. Locations of monitoring sites. Black points indicate monitoring sites. The Gwaii Haanas National Marine Conservation Area Reserve (NMCAR) and Rockfish Conservation Areas (RCAs) are indicated.




Table 1. Names, locations and management policy of monitoring sites.

RCA = Rockfish Conservation Area, NMCAR = National Marine Conservation Area Reserve.

Area	Management policy	Site	Latitude	Longitude
Louise	Commercial Red Urchin Fishery Closure	Skedans	52.9258° N	131.6175° W
		Vertical	52.8983° N	131.6259° W
		Nelson	52.8757° N	131.6719° W
Lyell	Commercial Red Sea Urchin Fishery Closure, RCA, NMCAR	Fuller	52.7102° N	131.4430° W
		Lyell SE	52.6427° N	131.4391° W
		Murchison	52.6157° N	131.4346° W
Kunghit E	NMCAR	Benjamin	52.2128° N	130.9963° W
		Koya	52.1799° N	131.0128° W
		Moorehead	52.1442° N	131.0467° W
Kunghit W	RCA, NMCAR	Louscoone	52.1291° N	131.2283° W
		Fanny	52.1191° N	131.1776° W
		Arnold	52.0987° N	131.1257° W

Table 2. Carbon and nitrogen content in bull kelp tissue samples. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ are ratios calculated relative to V-PDB (Vienna PeeDee Belemnite) and air, respectively. Sample mean and standard error are shown.

Area	Site	n	Sample (mg)	C (ug)	N (ug)	% C	% N	C:N ratio	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$
Louise	Skedans 1		4.74	1775.59	163.03	37.5	3.44	10.9	-17.74	6.69
	Skedans 2		5.38	2023.61	213.04	37.6	3.96	9.5	-19.47	6.28
	Skedans 3		5.01	1759.06	163.03	35.1	3.25	10.8	-17.30	6.49
	Skedans 4		5.54	1940.93	170.11	35.0	3.07	11.4	-16.14	8.31
	Skedans 5		5.40	1924.40	195.80	35.6	3.63	9.8	-17.54	6.91
	Skedans 6		5.41	1858.26	173.64	34.3	3.21	10.7	-18.85	6.40
	Skedans 7		4.82	1709.46	155.91	35.5	3.23	11.0	-16.54	7.39
	Skedans 8		5.12	1709.46	151.14	33.4	2.95	11.3	-16.59	6.86
	Skedans 9		5.42	1940.93	191.16	35.8	3.53	10.2	-17.23	6.63
	Skedans 10		5.22	1825.19	163.03	35.0	3.12	11.2	-17.95	6.77
	$\bar{x} \pm \text{SE}$	10	5.21 ± 0.09	1846.69 ± 34.33	173.99 ± 6.25	35.48 ± 0.41	3.34 ± 0.10	10.67 ± 0.20	-17.53 ± 0.33	6.87 ± 0.19
	Vertical 1		5.40	1940.93	183.02	35.9	3.39	10.6	-18.73	7.17
	Vertical 2		5.36	1907.86	186.51	35.6	3.48	10.2	-19.32	5.88
	Vertical 3		4.96	1808.66	159.47	36.5	3.22	11.3	-18.34	8.60
	Vertical 4		5.22	1602.00	139.16	30.7	2.67	11.5	-17.87	8.91
	Vertical 5		5.05	1436.68	119.91	28.4	2.37	12.0	-18.63	8.25
	Vertical 6		5.06	1621.84	145.16	32.1	2.87	11.2	-17.97	8.92
	Vertical 7		5.48	1542.48	148.75	28.1	2.71	10.4	-19.20	7.29
	Vertical 8		5.20	1603.65	148.75	30.8	2.86	10.8	-19.18	7.15
	Vertical 9		5.23	1431.72	118.44	27.4	2.26	12.1	-18.45	8.57
	$\bar{x} \pm \text{SE}$	9	5.22 ± 0.06	1655.09 ± 63.03	149.91 ± 7.96	31.73 ± 1.18	2.87 ± 0.14	11.12 ± 0.22	-18.63 ± 0.18	7.86 ± 0.35
	Nelson 1		5.24	1775.59	143.96	33.9	2.75	12.3	-17.99	8.34
	Nelson 2		5.13	1676.39	124.65	32.7	2.43	13.4	-16.95	8.69
	Nelson 3		4.84	1692.93	142.76	35.0	2.95	11.9	-18.86	8.22
	Nelson 4		5.14	1759.06	147.56	34.2	2.87	11.9	-20.27	6.92
	Nelson 5		5.44	1874.80	145.16	34.5	2.67	12.9	-16.56	8.88
	Nelson 6		5.10	1568.93	123.56	30.8	2.42	12.7	-17.40	8.61
	Nelson 7		4.90	1606.96	123.07	32.8	2.51	13.1	-17.05	8.79
	Nelson 8		5.24	1725.99	133.13	32.9	2.54	13.0	-16.75	9.02
	Nelson 9		4.87	1592.08	140.36	32.7	2.88	11.3	-15.82	8.96
	Nelson 10		5.18	1449.90	112.33	28.0	2.17	12.9	-18.37	8.65
	$\bar{x} \pm \text{SE}$	10	5.11 ± 0.06	1672.26 ± 38.50	133.65 ± 3.81	32.74 ± 0.65	2.62 ± 0.08	12.54 ± 0.21	-17.60 ± 0.41	8.51 ± 0.19

Table 2 (Continued). Carbon and nitrogen content in bull kelp tissue samples.

Area	Site	n	Sample (mg)	C (ug)	N (ug)	% C	% N	C:N ratio	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$
Fuller	Fuller 1		4.96	1709.46	143.96	34.5	2.90	11.9	-18.92	7.55
	Fuller 2		4.75	1562.32	128.29	32.9	2.70	12.2	-17.41	7.77
	Fuller 3		4.72	1621.84	136.75	34.4	2.90	11.9	-18.36	6.73
	Fuller 4		5.44	1709.46	137.95	31.4	2.54	12.4	-19.33	6.75
	Fuller 5		5.39	1792.13	153.53	33.2	2.85	11.7	-19.10	6.69
	Fuller 6		5.55	1656.55	173.64	29.8	3.13	9.5	-21.82	5.51
	Fuller 7		5.00	1327.57	111.23	26.6	2.22	11.9	-18.00	7.52
	Fuller 8		4.87	1461.48	121.98	30.0	2.50	12.0	-18.80	7.20
	Fuller 9		5.48	1595.38	134.34	29.1	2.45	11.9	-19.94	6.77
	Fuller 10		5.14	1598.69	148.75	31.1	2.89	10.7	-19.86	5.42
Lyell	$\bar{x} \pm \text{SE}$	10	5.13 ± 0.10	1603.49 ± 42.07	139.04 ± 5.51	31.30 ± 0.79	2.71 ± 0.09	11.61 ± 0.27	-19.15 ± 0.39	6.79 ± 0.25
	Lyell SE 1		5.40	1759.06	153.53	32.6	2.84	11.5	-15.81	7.63
	Lyell SE 2		5.02	1725.99	146.36	34.4	2.92	11.8	-16.80	7.17
	Lyell SE 3		5.01	1825.19	166.58	36.4	3.32	11.0	-16.38	7.67
	Lyell SE 4		5.31	1759.06	146.36	33.1	2.76	12.0	-16.67	7.84
	Lyell SE 5		5.25	1742.53	152.33	33.2	2.90	11.4	-16.97	7.77
	Lyell SE 6		4.73	1759.06	154.72	37.2	3.27	11.4	-14.76	8.12
	Lyell SE 7		5.31	1841.73	148.75	34.7	2.80	12.4	-15.27	7.93
	Lyell SE 8		5.18	1709.46	140.36	33.0	2.71	12.2	-14.89	7.99
	Lyell SE 9		5.41	1759.06	147.56	32.5	2.73	11.9	-14.57	8.29
Murchison	Lyell SE 10		5.51	1858.26	166.58	33.7	3.02	11.2	-17.66	7.42
	$\bar{x} \pm \text{SE}$	10	5.21 ± 0.07	1773.94 ± 15.87	152.31 ± 2.71	34.08 ± 0.51	2.93 ± 0.07	11.67 ± 0.15	-15.98 ± 0.34	7.78 ± 0.10
	Murchison 1		5.12	1626.79	129.50	31.8	2.53	12.6	-17.11	7.85
	Murchison 2		5.38	1775.59	139.16	33.0	2.59	12.8	-16.31	6.75
	Murchison 3		5.28	1841.73	155.91	34.9	2.95	11.8	-17.54	6.41
	Murchison 4		5.18	1841.73	148.75	35.6	2.87	12.4	-17.30	7.69
	Murchison 5		5.31	1725.99	145.16	32.5	2.73	11.9	-18.43	7.71
	Murchison 6		5.42	1907.86	168.94	35.2	3.12	11.3	-18.47	7.15
	Murchison 7		5.24	1611.92	140.36	30.8	2.68	11.5	-20.00	7.51
	Murchison 8		4.72	1552.40	128.29	32.9	2.72	12.1	-16.72	6.54
7	Murchison 9		5.12	1725.99	142.76	33.7	2.79	12.1	-16.99	6.38
	Murchison 10		5.17	1742.53	152.33	33.7	2.95	11.4	-19.75	5.92
7	$\bar{x} \pm \text{SE}$	10	5.19 ± 0.06	1735.25 ± 35.70	145.12 ± 3.86	33.40 ± 0.48	2.79 ± 0.06	11.98 ± 0.16	-17.86 ± 0.40	6.99 ± 0.21

Table 2 (Continued). Carbon and nitrogen content in bull kelp tissue samples.

Area	Site	n	Sample (mg)	C (ug)	N (ug)	% C	% N	C:N ratio	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$
Kunghit E	Benjamin 1		5.38	1924.40	157.10	35.8	2.92	12.2	-19.89	5.79
	Benjamin 2		5.36	2023.61	224.42	37.8	4.19	9.0	-22.47	5.51
	Benjamin 3		5.40	1858.26	210.76	34.4	3.90	8.8	-23.58	5.28
	Benjamin 4		5.09	1891.33	191.16	37.2	3.76	9.9	-24.07	4.91
	Benjamin 5		4.97	1825.19	179.51	36.7	3.61	10.2	-22.00	5.78
	Benjamin 6		5.31	1907.86	183.02	35.9	3.45	10.4	-21.05	5.38
	Benjamin 7		5.36	1990.54	168.94	37.1	3.15	11.8	-19.52	4.18
	Benjamin 8		5.40	1957.47	200.42	36.2	3.71	9.8	-21.98	5.51
	Benjamin 9		4.74	1656.55	141.56	34.9	2.99	11.7	-18.70	4.71
	Benjamin 10		5.19	1742.53	141.56	33.6	2.73	12.3	-19.58	5.19
$\bar{x} \pm \text{SE}$		10	5.22 ± 0.07	1877.77 ± 35.59	179.84 ± 8.86	35.97 ± 0.42	3.44 ± 0.15	10.61 ± 0.41	-21.29 ± 0.58	5.22 ± 0.16
Moorehead	Koya 1		5.52	2023.61	217.61	36.7	3.94	9.3	-22.52	5.47
	Koya 2		4.83	1623.49	171.29	33.6	3.55	9.5	-23.39	6.05
	Koya 3		5.03	1676.39	146.36	33.3	2.91	11.5	-22.05	5.72
	Koya 4		5.05	1725.99	170.11	34.2	3.37	10.1	-21.92	5.96
	Koya 5		5.42	1907.86	185.35	35.2	3.42	10.3	-21.98	5.07
	Koya 6		4.64	1383.78	142.76	29.8	3.08	9.7	-22.57	6.60
	Koya 7		5.15	1709.46	153.53	33.2	2.98	11.1	-21.47	5.27
	Koya 8		5.46	1676.39	165.39	30.7	3.03	10.1	-21.58	6.41
	$\bar{x} \pm \text{SE}$	8	5.14 ± 0.11	1715.87 ± 67.30	169.05 ± 8.55	33.34 ± 0.79	3.28 ± 0.12	10.20 ± 0.27	-22.19 ± 0.22	5.82 ± 0.19
	Moorehead 1		5.35	2007.07	181.85	37.5	3.40	11.0	-18.09	5.99
Kunghit E	Moorehead 2		4.93	1825.19	177.17	37.0	3.59	10.3	-19.86	6.17
	Moorehead 3		5.34	1924.40	177.17	36.0	3.32	10.9	-18.26	6.40
	Moorehead 4		5.43	2007.07	188.84	37.0	3.48	10.6	-19.36	3.62
	Moorehead 5		5.24	1552.40	137.95	29.6	2.63	11.3	-17.75	5.36
	Moorehead 6		5.21	1573.89	139.16	30.2	2.67	11.3	-16.10	6.31
	Moorehead 7		5.39	1588.77	142.76	29.5	2.65	11.1	-18.78	5.99
	Moorehead 8		5.40	1578.85	139.16	29.2	2.58	11.3	-16.43	6.16
	Moorehead 9		5.02	1387.08	125.62	27.6	2.50	11.0	-17.09	6.22
	Moorehead 10		5.02	1418.49	128.29	28.3	2.56	11.1	-16.99	5.84
	$\bar{x} \pm \text{SE}$	10	5.23 ± 0.06	1686.32 ± 74.14	153.80 ± 7.71	32.20 ± 1.30	2.94 ± 0.14	11.00 ± 0.10	-17.87 ± 0.39	5.81 ± 0.26

Table 2 (Continued). Carbon and nitrogen content in bull kelp tissue samples.

Area	Site	n	Sample (mg)	C (ug)	N (ug)	% C	% N	C:N ratio	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$
Kunghit W	Louscoone 1		5.51	1808.66	154.72	32.83	2.81	11.69	-19.79	5.19
	Louscoone 2		5.15	1676.39	180.68	32.55	3.51	9.28	-23.08	5.95
	Louscoone 3		4.71	1479.66	130.71	31.42	2.78	11.32	-19.53	5.84
	Louscoone 4		4.90	1497.84	149.95	30.57	3.06	9.99	-21.00	6.59
	Louscoone 5		4.75	1421.80	136.75	29.93	2.88	10.40	-21.17	4.88
	Louscoone 6		5.45	1709.46	149.95	31.37	2.75	11.40	-20.45	5.21
	$\bar{x} \pm \text{SE}$	6	5.08 ± 0.14	1598.97 ± 62.72	150.46 ± 7.09	31.44 ± 0.45	2.96 ± 0.12	10.68 ± 0.39	-20.84 ± 0.52	5.61 ± 0.26
	Fanny 1		4.96	1426.76	123.31	28.77	2.49	11.57	-16.59	8.25
	Fanny 2		5.54	1676.39	158.29	30.26	2.86	10.59	-19.18	7.52
	Fanny 3		5.24	1527.60	134.34	29.15	2.56	11.37	-16.57	8.11
	Fanny 4		5.23	1358.98	116.49	25.98	2.23	11.67	-15.96	8.80
	Fanny 5		4.91	1299.47	108.78	26.47	2.22	11.95	-18.65	8.00
	Fanny 6		5.45	1375.51	111.97	25.24	2.05	12.29	-16.21	8.00
	Fanny 7		5.27	1527.60	146.36	28.99	2.78	10.44	-18.97	7.82
	Fanny 8		5.59	1676.39	147.56	29.99	2.64	11.36	-18.27	7.76
	Fanny 9		5.48	1373.86	114.05	25.07	2.08	12.05	-16.80	7.88
	Fanny 10		4.95	1651.59	154.72	33.37	3.13	10.67	-20.59	6.28
	$\bar{x} \pm \text{SE}$	10	5.26 ± 0.08	1489.42 ± 45.03	131.58 ± 6.00	28.33 ± 0.83	2.50 ± 0.11	11.39 ± 0.20	-17.78 ± 0.49	7.84 ± 0.20
Arnold	Arnold 1		4.75	1692.93	139.16	35.6	2.93	12.2	-19.69	4.08
	Arnold 2		5.12	1891.33	157.10	36.9	3.07	12.0	-19.42	5.28
	Arnold 3		4.85	1676.39	134.34	34.6	2.77	12.5	-18.97	6.22
	Arnold 4		5.51	1858.26	158.29	33.7	2.87	11.7	-19.48	5.67
	Arnold 5		5.23	1709.46	136.75	32.7	2.61	12.5	-18.91	6.92
	Arnold 6		5.11	1742.53	142.76	34.1	2.79	12.2	-20.31	5.55
	Arnold 7		4.96	1405.27	121.12	28.3	2.44	11.6	-21.07	6.27
	Arnold 8		5.42	1507.76	134.34	27.8	2.48	11.2	-18.70	7.37
	Arnold 9		5.33	1742.53	152.33	32.7	2.86	11.4	-19.26	5.21
	Arnold 10		5.06	1522.64	128.29	30.1	2.54	11.9	-19.75	6.58
	$\bar{x} \pm \text{SE}$	10	5.13 ± 0.08	1674.91 ± 48.87	140.45 ± 3.88	32.66 ± 0.96	2.74 ± 0.07	11.93 ± 0.14	-19.56 ± 0.22	5.92 ± 0.30

Table 3. Sulfur content in bull kelp tissue samples. $\delta^{34}\text{S}$ was calculated relative to the VCDT (Vienna Canyon Diablo Troilite) standard. Sample mean and standard error are shown.

Area	Site	n	Sample (mg)	S (μg)	%o S	$\delta^{34}\text{S}$ vs. VCDT
Louise	Skedans 1		2.38	22.80	9.58	21.51
	Skedans 2		2.51	31.72	12.64	21.99
	Skedans 3		2.13	22.33	10.49	21.11
	$\bar{x} \pm \text{SE}$	3	2.34 ± 0.11	25.62 ± 3.05	10.90 ± 0.91	21.54 ± 0.35
	Vertical 1		1.55	21.98	14.18	21.36
	Vertical 2		2.13	23.30	10.94	21.32
	Vertical 3		2.52	33.26	13.20	21.83
	$\bar{x} \pm \text{SE}$	3	2.07 ± 0.28	26.18 ± 3.56	12.77 ± 0.96	21.51 ± 0.16
	Nelson 1		2.51	31.42	12.52	21.67
	Nelson 2		2.47	27.93	11.31	21.83
	Nelson 3		1.57	19.41	12.36	21.63
	$\bar{x} \pm \text{SE}$	3	2.18 ± 0.31	26.25 ± 3.57	12.06 ± 0.38	21.71 ± 0.06
Lyell	Fuller 1		2.09	21.53	10.30	20.27
	Fuller 2		2.39	27.99	11.71	20.37
	Fuller 3		2.35	23.41	9.96	20.29
	$\bar{x} \pm \text{SE}$	3	2.28 ± 0.09	24.31 ± 1.92	10.66 ± 0.54	20.31 ± 0.03
	Lyell SE 1		1.95	21.20	10.87	20.48
	Lyell SE 2		2.45	28.53	11.64	21.45
	Lyell SE 3		2.49	30.07	12.08	21.58
	$\bar{x} \pm \text{SE}$	3	2.30 ± 0.17	26.60 ± 2.74	11.53 ± 0.35	21.17 ± 0.35
	Murchison 1		1.86	20.53	11.04	21.35
	Murchison 2		2.14	22.85	10.68	21.65
	Murchison 3		2.15	25.63	11.92	21.54
	$\bar{x} \pm \text{SE}$	3	2.05 ± 0.10	23.00 ± 1.47	11.21 ± 0.37	21.51 ± 0.09

Table 3 (Continued). Sulfur content in bull kelp tissue samples.

Area	Site	n	Sample (mg)	S (µg)	%o S	$\delta^{34}\text{S}$ vs. VCDT
Kunghit E	Benjamin 1		1.68	17.28	10.29	21.06
	Benjamin 2		2.51	27.51	10.96	21.29
	Benjamin 3		2.47	29.17	11.81	21.26
	$\bar{x} \pm \text{SE}$	3	2.22 ± 0.27	24.65 ± 3.72	11.02 ± 0.44	21.20 ± 0.07
	Koya 1		2.2	20.73	9.42	19.94
	Koya 2		2.51	29.49	11.75	20.45
	Koya 3		2.42	32.16	13.29	21.18
	$\bar{x} \pm \text{SE}$	3	2.38 ± 0.09	27.46 ± 3.45	11.49 ± 1.12	20.52 ± 0.36
	Moorehead 1		2.46	21.14	8.59	21.51
	Moorehead 2		2.25	21.27	9.45	21.22
	Moorehead 3		2.24	21.05	9.40	21.60
	$\bar{x} \pm \text{SE}$	3	2.32 ± 0.07	21.15 ± 0.07	9.15 ± 0.28	21.44 ± 0.11
Kunghit W	Louscoone 1		2.64	26.40	10.00	20.25
	Louscoone 2		2.53	26.48	10.47	20.16
	Louscoone 3		2.00	21.96	10.98	19.83
	$\bar{x} \pm \text{SE}$	3	2.39 ± 0.20	24.95 ± 1.49	10.48 ± 0.28	20.08 ± 0.13
	Fanny 1		2.44	23.46	9.61	18.92
	Fanny 2		2.13	20.50	9.63	19.68
	Fanny 3		1.95	18.73	9.61	19.62
	$\bar{x} \pm \text{SE}$	3	2.17 ± 0.14	20.90 ± 1.38	9.61 ± 0.01	19.40 ± 0.24
	Arnold 1		1.56	13.97	8.95	21.50
	Arnold 2		2.15	19.93	9.27	21.55
	Arnold 3		1.86	18.50	9.94	21.47
	$\bar{x} \pm \text{SE}$	3	1.86 ± 0.17	17.47 ± 1.80	9.39 ± 0.29	21.51 ± 0.02