

August 27 to September 9, 2017 (Weeks 35-36)


## Overall Summary

- Influenza activity remains at interseasonal levels across the country, with a few regions reporting sporadic or localized activity.
- In weeks 35-36, the majority of influenza detections continued to be A(H3N2). The percentage of laboratory tests positive for influenza is higher for this time of year compared to previous seasons.
- This is the first FluWatch report of the 2017-18 season. The next FluWatch report will be published on September 29<sup>th</sup> and weekly reporting will resume on October 13<sup>th</sup>, 2017.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

## Influenza/Influenza-like Illness (ILI) Activity (geographic spread)

In weeks 35-36, influenza or influenza-like illness activity levels remained at low levels and the majority of regions reported no activity. In week 36, one region in Alberta reported localized activity, and 11 regions in British Columbia (3), Alberta (3), Ontario (4), and Quebec (1) reported sporadic activity.


**Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, 2017-18, Week 36**



## Laboratory-Confirmed Influenza Detections

In weeks 35-36, the number of tests positive for influenza remained at interseasonal levels. The percentage of tests positive was low but increased slightly from 1.9 to 2.4% in weeks 35 and 36, which is higher for this time of year than was observed during the previous seven seasons. The number of detections of influenza A has increased during this two-week period, while influenza B detections have remained low. The majority of influenza A viruses subtyped were A(H3N2). For data on other respiratory virus detections, see the [Respiratory Virus Detections in Canada Report](#).


**Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2017-18, weeks 35 to 36**



The shaded area indicates weeks where the positivity rate was at least 5% and a minimum of 15 positive tests were observed, signalling the period of [seasonal influenza activity](#).

To date this season, 73 laboratory-confirmed influenza detections have been reported, of which 90% have been influenza A. Influenza A(H3N2) has been the most common subtype detected this season, representing 87% of influenza A detections. Detections from BC and AB represent 74% of the cases reported in this 2-week period. For more detailed weekly and cumulative influenza data, see the text descriptions for Figures 2 and 3 or the [Respiratory Virus Detections in Canada Report](#).

**Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province/territory, Canada, 2017-18, weeks 35 to 36**



To date this season, detailed information on age and type/subtype has been received for more than 45 laboratory-confirmed influenza cases (Table 1). Based on the limited data to date, among influenza cases with reported age and type/subtype information, more than half of cases were in adults aged 65 years and older.

**Table 1 – Cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting, Canada, 2017-18, weeks 35 to 36**

Age groups (years)	Cumulative (August 27 to September 9, 2017)						
	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) <sup>1</sup>	Total	#	%
0-4	<5	<5	0	0	<5	5	11%
5-19	<5	<5	<5	0	<5	5	11%
20-44	8	0	8	0	0	8	17%
45-64	<5	0	<5	0	0	<5	4%
65+	27	<5	23	<5	0	27	57%
<b>Total</b>	>40	6	34	<5	<5	>45	100%


<sup>1</sup>UnS: unsubtyped: The specimen was typed as influenza A, but no result for subtyping was available;  
 x - Suppressed to prevent residual disclosure

## Syndromic / Influenza-like Illness Surveillance

### Healthcare Practitioners Sentinel Syndromic Surveillance

In week 36, 0.8% of visits to healthcare professionals were due to influenza-like illness.

**Figure 4 – Percentage of visits for ILI reported by sentinels by report week, Canada, 2017-18, weeks 35 to 36**  
 Number of Sentinels Reporting in Week 36: 84



The shaded area represents the maximum and minimum percentage of visits for ILI reported by week from seasons 2012-13 to 2016-17

### Participatory Syndromic Surveillance


FluWatchers is a participatory ILI surveillance system that relies on weekly voluntary submissions of syndromic information from the Canadians across Canada.

The 2017-18 FluWatchers reporting season will begin October 1<sup>st</sup>, 2017 (week 40).

## Influenza Outbreak Surveillance

In weeks 35-36, one new laboratory-confirmed influenza outbreak was reported in a long-term care facility.

**Figure 6 – Number of new outbreaks of laboratory-confirmed influenza by report week, Canada, 2017-18, weeks 35 to 36**



## Severe Outcomes Influenza Surveillance

### Provincial/Territorial Influenza Hospitalizations and Deaths

In weeks 35-36, the number of weekly influenza-associated hospitalizations reported by participating provinces and territories<sup>1</sup> remained at low levels with fewer than five hospitalizations reported during this 2-week period.

<sup>1</sup>Influenza-associated hospitalizations are reported by NL, PE, NS, NB, MB, AB, YT and NT. Only hospitalizations that require intensive medical care are reported by SK.

### Pediatric Influenza Hospitalizations and Deaths

To date this season, less than five laboratory-confirmed influenza-associated pediatric ( $\leq 16$  years of age) hospitalizations were reported from the Immunization Monitoring Program Active (IMPACT) network.

## Influenza Strain Characterizations

The National Microbiology Laboratory has not yet reported influenza strain characterization results for influenza viruses collected during the 2017-18 season.

## Antiviral Resistance

The National Microbiology Laboratory has not yet reported antiviral resistance results for influenza viruses collected during the 2017-18 season.

## Provincial and International Influenza Reports

- Alberta – [Influenza Surveillance Report](#)
- British Columbia – [Influenza Surveillance](#)
- Manitoba – [Manitoba – Seasonal Influenza Reports](#)
- New Brunswick – [Influenza Surveillance Reports](#)
- Newfoundland and Labrador – [Surveillance and Disease Reports](#)
- Nova Scotia – [Respiratory Watch Report](#)
- Ontario – [Respiratory Pathogen Bulletin](#)
- Prince Edward Island – [Influenza Summary](#)
- Saskatchewan – [Influenza Reports](#)
- Québec – [Flash Grippe](#)
- Australia – [Influenza Surveillance Report](#)
- European Centre for Disease Prevention and Control – [Surveillance reports and disease data on seasonal influenza](#)
- New Zealand – [Influenza Weekly Update](#)
- Public Health England – [Weekly national flu reports](#)
- Pan-American Health Organization – [Influenza Situation Report](#)
- United States Centres for Disease Control and Prevention – [Weekly Influenza Surveillance Report](#)
- World Health Organization – [Influenza update](#)
- World Health Organization – [FluNet](#)

## FluWatch Surveillance for the 2017-2018 Season – Notes and Definitions

The FluWatch report is compiled from a number of data sources. Surveillance information contained in this report is a reflection of the surveillance data available to FluWatch at the time of production. Delays in reporting of data may cause data to change retrospectively

### Influenza/Influenza-like Illness (ILI) Activity

Influenza/ILI activity levels, as represented on the map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, primary care consultations for ILI and reported outbreaks. ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls, and the determination of an increase is based on the assessment of the provincial/territorial epidemiologist. Maps from previous weeks, including any retrospective updates, are available in the mapping feature found in the [Weekly Influenza Reports](#).

### Influenza/ILI Activity Level definitions

- 1 = No activity:** no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported
- 2 = Sporadic:** sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†
- 3 = Localized:** (1) evidence of increased ILI\* ; (2) lab confirmed influenza detection(s); (3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region**†
- 4 = Widespread:** (1) evidence of increased ILI\*; (2) lab confirmed influenza detection(s);(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region**†;

\* More than just sporadic as determined by the provincial/territorial epidemiologist.

†Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist

### Laboratory-Confirmed Influenza Detections

Provincial, regional and some hospital laboratories report the weekly number of tests and detections of influenza and other respiratory viruses. Provincial public health laboratories submit demographic information for cases of influenza. This case-level data represents a subset of influenza detections reported through aggregate reporting. Specimens from NT, YT, and NU are sent to reference laboratories in the provinces for testing.

### Syndromic/Influenza-like Illness Surveillance

FluWatch maintains a network of primary care practitioners who report the weekly proportion of ILI cases seen in their practice. Independent sentinel networks in BC, AB, and SK compile their data for reporting to FluWatch. Not all sentinel physicians report every week.

**Definition of Influenza-like-illness (ILI):** Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5 years of age, gastrointestinal symptoms may also be present. In patients under 5 or 65 years and older, fever may not be prominent.

### Influenza Outbreak Surveillance

Outbreaks of influenza or ILI are reported from all provinces and territories, according to the definitions below. However, reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions. All provinces and territories with the exception of NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals.

#### **Outbreak definitions:**

**Schools:** Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI.

**Hospitals and residential institutions:** two or more cases of ILI within a seven-day period, including at least one laboratory-confirmed case of influenza. Residential institutions include but are not limited to long-term care facilities (LTCF) and prisons.

**Workplace:** Greater than 10% absenteeism on any day which is most likely due to ILI.

**Other settings:** two or more cases of ILI within a seven-day period, including at least one laboratory-confirmed case of influenza; i.e. closed communities.

### **Serious Outcome Influenza Surveillance**

#### **Provincial/Territorial Influenza Hospitalizations and Deaths**

Influenza-associated hospitalizations and deaths are reported by 8 Provincial and Territorial Ministries of Health (excluding BC, NU, ON and QC). The hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting. Only hospitalizations that require intensive medical care are reported by SK.

Due to changes in participating provinces and territories, comparisons to previous years should be done with caution.

#### **Pediatric Influenza Hospitalizations and Deaths**

The Immunization Monitoring Program Active (IMPACT) network reports the weekly number of hospitalizations with influenza among children admitted to one of the 12 participating paediatric hospitals in 8 provinces. These represent a subset of all influenza-associated pediatric hospitalizations in Canada.

#### **Influenza Strain Characterizations and Antiviral Resistance**

Provincial public health laboratories send a subset of influenza virus isolates to the National Microbiology Laboratory for strain characterization and antiviral resistance. These represent a subset of all influenza detections in Canada and the proportion of isolates of each type and subtype is not necessarily representative of circulating viruses.

Antigenic strain characterization data reflect the results of hemagglutination inhibition (HI) testing compared to the reference influenza strains recommended by [WHO](#). Genetic strain characterization data are based on analysis of the sequence of the viral hemagglutinin (HA) gene.

Antiviral resistance testing is conducted by phenotypic and genotypic methods on influenza virus isolates submitted to the National Microbiology Laboratory. All isolates are tested for oseltamivir and zanamivir and a subset are tested for resistance to amantadine.

**Abbreviations:** Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

This [report](#) is available on the Government of Canada Influenza webpage.

Ce [rapport](#) est disponible dans les deux langues officielles.

*We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.*