



WATER

SEDIMENTS

SHORELINES

BIOLOGICAL RESOURCES

USES

2nd edition

NORTHERN GANNET

A Sentinel Species for the Gulf

Background

The Northern Gannet is a seabird whose expansive feeding grounds and a diet rich in fish can provide information on the abundance of its prey and the degree of contamination of the

ecosystem. That's why it was chosen as a sentinel species (or "bioindicator") of the state of the Gulf of St. Lawrence. This bird can sometimes travel over 200 km in search of mackerel, herring, capelin and sand lance. There are six Northern



Photo: Jean-François Rill, Canadian Wildlife Service

Figure 1. Location of Northern Gannet colonies



Gannet colonies in eastern North America, including three along the east coast of Newfoundland and three others in the Quebec portion of the Gulf of St. Lawrence (Figure 1). The populations of the eastern tip of Anticosti Island, Bird Rocks in the Magdalen Islands and Bonaventure Island together make up 75% of the North American population.


The Bonaventure Island population of Northern Gannet grew by about 3% annually from the turn of the century until the mid-1960s, when its numbers fell by almost 25% in ten years, at the same time that the Newfoundland colonies were remaining stable. Scientists discovered that high concentrations of residual organochlorine substances detected in gannet eggs, including dieldrin and DDT, were responsible for the low reproductive rate observed during

this period. At high concentrations, these toxic substances interfere with the synthesis of calcium carbonate, the main component of eggshells; thin eggshells are even more vulnerable to impact. This reproductive problem, combined with the fact that the species produces only one egg per clutch, led to a spectacular drop in hatching success rates and the consequent reduction, equally dramatic, in the number of breeding adults. Large-scale spraying of DDT in the 1960s to control insect infestations in New Brunswick and the Gaspé region largely explains the high concentrations detected in some species in the Gulf of St. Lawrence.

Overview of the Situation

The population growth of Bonaventure Island gannets remained relatively steady between 1887 and 1966, when the number of pairs grew from 1 500 to 21 215. It was only in 1984 that numbers returned to near-1966 levels, after having reached a low of 16 400 pairs in 1976 (Figure 2). The colony has continued

Figure 2. The Bonaventure Island gannet population since the turn of the century



to grow ever since, expanding from 24 125 pairs in 1989 to a population of 53 635 pairs in 2004.

Figure 3 (A and B) provides a good indication of reductions in DDT concentrations in eggs, expressed here as a percentage relative to the 1968 level, which coincides with the sudden jump in the net productivity rate and in the



Photo: Nathalie Brunelle, © Québec en images, CCDMD


Photo: Pierre Brousseau, Canadian Wildlife Service

number of pairs, after a 5- to 7-year lag (corresponding with gannet breeding age). With the reproductive success rate remaining above 67%, the population continues to grow. Unless the gannets

colonize new sites, however, scientists are of the opinion that population growth will stabilize within the next few years, when they will have occupied almost all the available habitat.

Figure 3. Population, DDE contamination rate in eggs and breeding success of Bonaventure Island gannets from 1966 to 2004

A. Population trends



B. Relationship between DDE contamination rate in gannet eggs and breeding success




Photo: Nathalie Brunelle, © Québec en images, CCDMD

KEY VARIABLES

A net reproductive rate (or breeding success, expressed as the percentage of fledglings relative to the number of eggs laid) of greater than 67% and the maintenance or growth of the population indicate that the Gulf of St. Lawrence Northern Gannet is thriving.

Outlook

By monitoring the number of nesting pairs, hatching success and reproductive success rates, as well as observing nesting habitat, scientists can readily determine the state of health of a population and identify the pressures that may be acting on its dynamics. The pursuit of the five-year monitoring




Photo: Nathalie Brunelle, © Québec en images, CCDMD

plan will provide timely information on changes affecting the structure or abundance of the gannet population, in order to then identify which pressures are being brought to bear on the species, whether in the St. Lawrence system or in its wintering grounds in the Gulf of Mexico.

To Know More

CHAPDELAINE, G. 1995. "The Northern Gannet," pp. 224–227, in J. Gauthier and Y. Aubry (eds.), *The Breeding Birds of Québec: Atlas of the Breeding Birds of Southern Québec*. Association québécoise des groupes d'ornithologues, Quebec Society for the Protection of Birds and Canadian Wildlife Service, Environment Canada – Quebec Region. Montreal.

CHAPDELAINE, G., P. LAPORTE, and D. N. NETTLESHIP. 1987. Population, productivity and DDT contamination trends of Northern Gannet (*Sula bassanus*) at Bonaventure Island, Quebec, 1976–1984. *Canadian Journal of Zoology* 65: 2922–2926.

NETTLESHIP, D. N. and G. CHAPEDELAINE. 1988. Population size and status of the Northern Gannet *Sula bassanus* in North America, 1984. *Journal of Field Ornithology* 59: 120–127.

RAIL, J.-F., G. CHAPDELAINE, P. BROUSSEAU, and J.-P. L. SAVARD. 1996. *Utilisation des oiseaux marins comme bioindicateurs de l'écosystème du Saint-Laurent*. Technical Report Series No. 254, ii + 113 pp. Environment Canada – Quebec Region, Canadian Wildlife Service. Sainte-Foy.

Prepared by: Gilles Chapdelaine and Jean-François Rail
Environmental Stewardship Branch
Environment Canada

State of the St. Lawrence Monitoring Program

Six government partners — Environment Canada, Fisheries and Oceans Canada, the Canadian Space Agency, Parks Canada Agency, the Ministère du Développement durable, de l'Environnement et des Parcs du Québec, the Ministère des Ressources naturelles et de la Faune du Québec — and Stratégies Saint-Laurent, a nongovernmental organization that works actively with riverside communities, are pooling their expertise and efforts to provide Canadians with information on

the state of the St. Lawrence and long-term trends affecting it.

To this end, environmental indicators have been developed on the basis of data collected as part of each organization's ongoing environmental monitoring activities. These activities cover the main components of the environment, namely water, sediments, biological resources, uses and shorelines.

For more information on the State of the St. Lawrence Monitoring Program, please visit our Web site at <www.planstlaurent.qc.ca> or contact our offices at the following address:

St. Lawrence Plan
Coordination Office
1141 Route de l'Eglise
P.O. Box 10 100
Sainte-Foy, Quebec
G1V 4H5
Tel: (418) 648-3444

Published by Authority of the Minister of the Environment
© Her Majesty the Queen in Right of Canada, 2002, 2nd edition 2005
Published by Authority of the Ministère du Développement durable,
de l'Environnement et des Parcs du Québec
© Gouvernement du Québec, 2002, 2nd edition 2005
Catalogue No.: En153-114/3-2005E-PDF
ISBN 0-662-42141-8
Envirodoq: ENV/2005/0269A

Legal deposit – National Library of Canada, 2005
Aussi disponible en français sous le titre: *Le fou de Bassan – Une espèce sentinelle du golfe*