



# TUBERCULOSIS IN CANADA

## 2010 PRE-RELEASE



Public Health  
Agency of Canada

Agence de la santé  
publique du Canada

Canada

Our mission is to promote and protect the health of Canadians through leadership, partnership, innovation and action in public health.  
- Public Health Agency of Canada

Également disponible en français sous le titre :  
La tuberculose au Canada 2010 – Prédifffusion

To obtain a copy of the report, send your request to:  
Centre for Communicable Diseases and Infection Control  
Public Health Agency of Canada  
Room 2410, 100 Eglantine Driveway, Health Canada Building  
A.L. 0602C, Tunney's Pasture  
Ottawa, ON K1A 0K9  
E-mail: [ccdic-clmti@phac-aspc.gc.ca](mailto:ccdic-clmti@phac-aspc.gc.ca)

This publication can be made available in alternative formats upon request.

This report was prepared by:  
Victor Gallant, MA  
Chris Archibald, MDCM, MHSc, FRCPC  
Annie-Claude Bourgeois, MSc  
Kathryn Dawson, BSc  
Jessica Halverson, MPH, MSW  
Derek Scholten, MSc  
Ania Zycki, MA

Suggested citation: Public Health Agency of Canada. Tuberculosis in Canada 2010 – Pre-release. Ottawa (Canada): Minister of Public Works and Government Services Canada; 2012

#### **ACKNOWLEDGEMENT**

The authors would like to acknowledge the provincial and territorial tuberculosis programs and their teams for their contribution to and ongoing participation in the Canadian Tuberculosis Reporting System.

© Her Majesty the Queen in Right of Canada, represented by the Minister of Health, 2012

Cat.: No.HP37-5/1-2010E-PDF

# INTRODUCTION

To facilitate an earlier release of *Tuberculosis in Canada, 2010*, the following seven tables have been extracted from the Canadian Tuberculosis Reporting System (CTBRS) for the year 2010. Data included in this pre-release report represent the total number of active tuberculosis cases reported to the Public Health Agency of Canada as of August 31, 2011. These data should be considered provisional until publication of the full report, *Tuberculosis in Canada 2010*.

---

## HIGHLIGHTS

For 2010 there were 1,577 new active and re-treatment TB cases reported to the CTBRS, with a corresponding incidence rate of 4.6 per 100,000 population. This represents an unprecedented national low since collection of TB data began in Canada in 1924. However, reported TB incidence rates varied by jurisdiction, and in particular there was a disproportionately high number of cases reported in Nunavut. As a result of an ongoing outbreak in Nunavut, the territory reported 101 active TB cases for 2010, almost twice the number as reported in 2009, resulting in a reported incidence rate of 304.0 per 100,000 population (Table 1).

From the preliminary reports, all provinces and territories reported at least one case of active TB disease in 2010. Newfoundland and Labrador reported a drop in the number of reported cases from 22 in 2009 to 8 in 2010. For the remaining provinces and territories with the exception of Nunavut, the number of reported cases in 2010 was stable relative to the number of cases reported in 2009 (Table 1).

Individuals between the ages of 25 and 34 years made up the largest number of reported cases in 2010, accounting for 18% of the total. However, the corresponding reported incidence rate of 6.0 per 100,000 population for this age group was surpassed by the rate of 9.6 per 100,000 reported for those greater than 74 years of age (Table 2).

Across the provinces and territories, patterns generally follow the national trend with most cases reported among individuals between 25 and 44 years of age. However, there were a couple of deviations from this trend. Almost 50% of the cases reported in 2010 from Saskatchewan were among individuals between 0 and 24 years of age, whereas this age group comprised only 18% of overall cases across Canada. In British Columbia, 21% of reported cases were 75 years of age and older, whereas this proportion was only 14% for all cases reported in Canada (Table 3).

In 2010, 66% of all reported TB cases were among foreign-born individuals, 21% among Canadian-born Aboriginal Peoples and 12% of cases were among Canadian-born non-Aboriginal people. In Alberta, British Columbia, Ontario and Quebec, the majority of cases were reported among

foreign-born individuals. In Manitoba, Saskatchewan and the North (which includes Northwest Territories, Nunavut and Yukon), the majority of cases were among Canadian-born Aboriginal peoples. In the Atlantic provinces, the small number of reported cases were mostly distributed between foreign-born and Canadian-born non-Aboriginal peoples; Newfoundland and Labrador, however did report cases among Aboriginal peoples (Table 4).

For 2010, pulmonary TB continued to be the main disease site, representing 64% of all reported cases (Table 5). In all provinces and territories the majority of cases were diagnosed with pulmonary disease. In Saskatchewan and the Northwest Territories, a greater proportion of cases were diagnosed with primary disease compared to other jurisdictions (Table 6).

Finally, there were a total of 64 cases of primary TB reported in 2010; 67% of these were among Canadian-born Aboriginal peoples. Of the 197 peripheral lymph node TB cases, 89% were diagnosed in foreign-born individuals (Table 7).

Table 1.

Reported new active and re-treatment tuberculosis cases and incidence rate per 100,000 - Canada and provinces/territories: 2000-2010

Year of diagnosis	Province/territory														
	CANADA	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	Y.T.	N.W.T.	Nvt.	
2000	Cases	1,724	10	2	3	10	318	700	98	104	133	286	3	10	47
	Rate	5.6	1.9	1.5	0.3	1.3	4.3	6.0	8.5	10.3	4.4	7.1	9.9	24.7	170.9
2001	Cases	1,772	19	3	8	10	261	699	115	114	116	379	0	8	40
	Rate	5.7	3.6	2.2	0.9	1.3	3.5	5.9	10.0	11.4	3.8	9.3	0.0	19.6	142.2
2002	Cases	1,667	9	1	9	11	288	716	98	89	128	287	0	4	27
	Rate	5.3	1.7	0.7	1.0	1.5	3.9	5.9	8.5	8.9	4.1	7.0	0.0	9.6	93.7
2003	Cases	1,631	7	3	6	12	257	693	127	91	110	305	1	12	7
	Rate	5.2	1.3	2.2	0.6	1.6	3.4	5.7	10.9	9.1	3.5	7.4	3.2	28.2	23.9
2004	Cases	1,613	7	1	8	10	219	700	144	70	109	299	4	10	32
	Rate	5.0	1.4	0.7	0.9	1.3	2.9	5.6	12.3	7.0	3.4	7.2	12.7	23.1	107.2
2005	Cases	1,641	9	1	7	6	255	643	114	139	146	265	3	8	45
	Rate	5.1	1.7	0.7	0.7	0.8	3.4	5.1	9.7	14.0	4.4	6.3	9.4	18.4	148.4
2006	Cases	1,654	12	0	10	2	228	673	134	87	131	320	3	6	48
	Rate	5.1	2.4	0.0	1.1	0.3	3.0	5.3	11.3	8.8	3.8	7.5	9.3	13.9	155.8
2007	Cases	1,576	7	0	7	5	229	680	103	106	112	278	3	15	31
	Rate	4.8	1.4	0.0	0.7	0.7	3.0	5.3	8.6	10.6	3.2	6.5	9.2	34.4	99.1
2008	Cases	1,643	8	0	4	5	240	603	141	95	167	300	8	13	59
	Rate	4.9	1.6	0.0	0.4	0.7	3.1	4.7	11.7	9.4	4.7	6.8	24.2	29.7	186.7
2009	Cases	1,659	22	1	8	11	196	629	156	93	176	294	4	12	57
	Rate	4.9	4.3	0.7	0.9	1.5	2.5	4.8	12.8	9.0	4.8	6.6	11.9	27.5	177.1
2010*	Cases	1,577	8	1	10	10	208	633	132	82	134	241	6	11	101
	Rate	4.6	1.6	0.7	1.1	1.3	2.6	4.8	10.7	7.8	3.6	5.3	17.4	25.1	304.0

\* Cases and rates for 2010 are provisional until publication of the Tuberculosis in Canada – 2010 Annual Report. These numbers will likely change.

**Table 2.**

Reported new active and re-treatment tuberculosis cases and incidence rate per 100,000 by age group - Canada: 2000 – 2010

Year of diagnosis	TOTAL	Age group												
		< 1	01-04	05-14	15-24	25-34	35-44	45-54	55-64	65-74	75+	Age Unk.		
2000	Cases	1,724	17	50	44	207	316	279	208	160	204	239	0	
	Rate	5.6	5.0	3.4	1.1	5.0	7.3	5.3	4.8	5.7	9.5	14.0	-	
2001	Cases	1,772	11	33	70	180	325	289	207	182	219	256	0	
	Rate	5.7	3.3	2.3	1.7	4.3	7.5	5.5	4.6	6.2	10.1	14.5	-	
2002	Cases	1,667	11	43	45	211	314	264	202	162	199	216	0	
	Rate	5.3	3.4	3.1	1.1	4.9	7.3	5.0	4.4	5.2	9.2	11.9	-	
2003	Cases	1,631	7	34	41	198	332	277	207	154	178	203	0	
	Rate	5.2	2.1	2.5	1.0	4.6	7.7	5.3	4.4	4.7	8.1	10.8	-	
2004	Cases	1,613	6	33	45	198	324	272	198	167	177	193	0	
	Rate	5.0	1.8	2.4	1.1	4.6	7.5	5.3	4.1	4.9	8.0	10.0	-	
2005	Cases	1,641	10	38	71	254	279	278	212	143	168	188	0	
	Rate	5.1	2.9	2.8	1.8	5.8	6.4	5.4	4.3	4.0	7.5	9.5	-	
2006	Cases	1,654	10	46	51	261	253	287	201	158	168	219	0	
	Rate	5.1	2.9	3.3	1.3	5.8	5.8	5.7	4.0	4.3	7.4	10.7	-	
2007	Cases	1,576	12	33	53	200	254	285	209	160	152	218	0	
	Rate	4.8	3.3	2.4	1.4	4.4	5.7	5.7	4.0	4.2	6.5	10.4	-	
2008	Cases	1,643	8	30	52	205	297	279	232	165	172	203	0	
	Rate	4.9	2.2	2.1	1.4	4.5	6.6	5.7	4.4	4.2	7.2	9.4	-	
2009	Cases	1,659	11	33	46	232	297	295	235	177	142	191	0	
	Rate	4.9	2.9	2.3	1.2	5.1	6.4	6.2	4.4	4.3	5.7	8.7	-	
2010*	Cases	1,577	9	29	39	200	284	268	210	175	147	216	0	
	Rate	4.6	2.4	1.9	1.0	4.4	6.0	5.7	3.9	4.1	5.7	9.6	-	

\* Cases and rates for 2010 are provisional until publication of the Tuberculosis in Canada – 2010 Annual Report. These numbers will likely change.

Table 3.

Reported new active and re-treatment tuberculosis cases and incidence rate per 100,000 by age group–Canada and provinces/territories: 2010

Age group	CANADA										Province/territory						
	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	Y.T.	N.W.T.	Nvt.				
< 1	Cases	9	0	0	0	0	0	0	0	2	0	0	7				
	Rate	2.4	0.0	0.0	0.0	0.0	0.0	0.0	0.0	4.5	0.0	0.0	875.0				
01-04	Cases	29	0	0	0	6	4	12	1	0	0	0	4				
	Rate	1.9	0.0	0.0	0.0	1.8	6.5	22.1	0.5	0.0	0.0	0.0	140.8				
05-14	Cases	39	0	0	0	3	11	8	4	3	0	0	2				
	Rate	1.0	0.0	0.0	0.0	0.4	7.1	6.2	0.9	0.6	0.0	0.0	29.3				
15-24	Cases	200	0	1	1	24	17	19	18	15	0	3	28				
	Rate	4.4	0.0	0.8	1.1	2.4	9.5	12.5	3.4	2.5	0.0	41.0	439.6				
25-34	Cases	284	2	2	2	38	23	13	26	39	0	2	25				
	Rate	6.0	3.4	1.8	2.2	3.5	14.0	9.2	4.3	6.3	0.0	26.5	470.0				
35-44	Cases	268	1	1	2	34	30	9	31	27	1	2	15				
	Rate	5.7	1.4	0.8	2.0	3.2	18.9	7.2	5.7	4.3	19.0	32.3	336.9				
45-54	Cases	210	1	2	1	30	16	8	15	34	3	0	7				
	Rate	3.9	1.2	1.3	0.8	2.3	8.7	5.2	2.6	4.7	47.4	0.0	197.0				
55-64	Cases	175	1	0	2	27	11	4	12	39	1	2	7				
	Rate	4.1	1.3	0.0	1.8	2.6	7.5	3.3	3.0	6.6	21.0	46.8	342.5				
65-74	Cases	147	2	0	1	20	10	5	11	30	1	1	5				
	Rate	5.7	4.4	0.0	1.5	3.0	11.6	6.7	5.1	8.3	49.7	68.0	672.0				
75+	Cases	216	1	0	4	26	10	4	16	52	0	1	1				
	Rate	9.6	3.1	0.0	1.8	4.7	11.6	5.1	8.8	16.4	0.0	111.7	373.1				
Total*	Cases	1,577	8	1	10	208	132	82	134	241	6	11	101				
	Rate	4.6	1.6	0.7	1.1	2.6	10.7	7.8	3.6	5.3	17.4	25.1	304.0				

\* Cases and rates for 2010 are provisional until publication of the Tuberculosis in Canada – 2010 Annual Report. These numbers will likely change.

Table 4

Reported new active and re-treatment tuberculosis cases by origin – Canada and provinces/territories: 2010

Origin	CANADA										Province/territory						
	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	North						
North American Indian	Cases	188	0	0	0	3	75	52	11	25	16						
	Rate	22.2	0.0	0.0	0.0	3.7	60.9	44.4	9.2	16.5	69.4						
Status (registered) Indian	Cases	184		0		3	75	52	9	24	16						
	Rate	21.8		0.0		4.0	53.4	37.5	8.0	18.1	59.4						
Status (registered) Indian - On reserve	Cases	106		0		0	52	34	6	12	0						
	Rate	21.5		0.0		0.0	58.3	46.7	8.1	17.4	0.0						
Status (registered) Indian - Off reserve	Cases	72		0		0	23	18	3	12	16						
	Rate	20.4		0.0		0.0	45.0	27.4	7.8	18.9	154.3						
Status (registered) Indian - Unknown	Cases	6		0		3	0	0	0	0	0						
	Rate	-		-		-	-	-	-	-	-						
Non-status Indian	Cases	4	0	0	0	0	0	0	2	1	0						
	Rate	-	-	-	-	-	-	-	-	-	-						
Inuit	Cases	116	3	0	0	12	2	0	0	0	99						
	Rate	198.6	56.2	0.0	0.0	98.8	104.1	0.0	0.0	0.0	281.3						
Métis	Cases	26	1	0	0	0	1	18	4	2	0						
	Rate	7.5	15.4	0.0	0.0	0.0	1.5	34.1	4.9	4.0	0.0						
<b>Total Aboriginal</b>	Cases	330	4	0	0	15	76	70	15	27	115						
	Rate	26.4	16.7	0.0	0.0	13.5	39.8	41.1	7.4	13.4	181.0						
Non-Aboriginal	Cases	183	2	5	5	50	13	5	10	39	2						
	Rate	0.7	0.4	0.6	0.7	0.7	0.6	0.6	0.4	1.3	5.1						
<b>Total Canadian Born</b>	Cases	513	6	5	5	65	89	75	25	66	117						
	Rates	1.9	1.2	0.6	0.7	1.0	0.7	7.7	0.8	2.1	113.6						
Africa, High HIV Prevalence	Cases	90	0	1	1	18	41	1	19	4	0						
	Rate	37.4	0.0	55.2	72.3	46.1	35.1	24.3	56.5	11.8	0.0						
Africa, Low HIV Prevalence	Cases	28	0	0	1	10	9	0	5	0	0						
	Rate	21.5	0.0	0.0	137.7	13.7	22.2	0.0	70.9	0.0	0.0						
American Region - Latin American and Caribbean Countries	Cases	64	0	0	0	25	33	0	2	4	0						
	Rate	7.0	0.0	0.0	0.0	11.3	6.2	0.0	3.0	6.7	0.0						



Table 4 cont'd

Origin	CANADA										Province/territory				
	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	North				
Established Market Economies and Central Europe	Cases	51	0	1	0	14	27	0	4	5	0				
	Rate	1.9	0.0	2.7	0.0	4.1	1.9	0.0	1.5	1.0	0.0				
Eastern Europe	Cases	18	0	0	1	5	10	0	1	1	0				
	Rate	4.9	0.0	0.0	97.8	6.4	5.0	0.0	3.5	2.7	0.0				
Eastern Mediterranean	Cases	90	0	2	0	14	58	2	11	2	0				
	Rate	11.8	0.0	22.7	0.0	7.9	13.4	22.4	17.8	3.2	0.0				
South-East Asia	Cases	249	1	0	0	14	154	5	20	54	0				
	Rate	30.3	83.2	0.0	0.0	29.8	30.1	34.9	28.6	32.0	0.0				
Western Pacific Region	Cases	429	1	1	2	33	221	18	46	101	1				
	Rate	22.8	42.2	12.0	28.7	24.5	25.8	29.8	21.4	17.7	48.8				
Unknown	Cases	19	0	0	0	2	4	10	1	2	0				
	Rate														
<b>Total Foreign-Born</b>	Cases	1,038	2	5	5	135	557	43	109	173	1				
	Rate	13.3	14.3	9.1	12.4	12.1	13.6	20.7	14.7	12.2	11.7				
Unknown	Cases	26	0	0	0	8	16	0	0	2	0				
	Cases	1,577	8	10	10	208	633	132	134	241	118				
<b>Total Cases*</b>	Rate	4.6	1.6	0.7	1.3	2.6	4.8	10.7	3.6	5.3	105.8				

NOTE: Provincial/territorial data are suppressed and reported in aggregate form if the P/T population denominator for the cell is less than 40. For the Atlantic Provinces (New Brunswick, Newfoundland and Labrador, Nova Scotia, and Prince Edward Island), the population figures for the Status-Indian on and off reserve are available in aggregate only. The indicated rates for these groups are for the entire Atlantic region.

\* Cases and rates for 2010 are provisional until publication of the Tuberculosis in Canada – 2010 Annual Report. These numbers will likely change.

**Table 5.**

Reported new active and re-treatment tuberculosis cases and incidence rate per 100,000 by main diagnostic site - Canada: 2000-2010

Main diagnostic site	Year of diagnosis											
	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010**	
Respiratory	Cases	99	121	88	79	94	106	91	64	59	62	64
	Rate	0.3	0.4	0.3	0.2	0.3	0.3	0.3	0.2	0.2	0.2	0.2
Pulmonary†	Cases	1,085	1,145	1,038	974	948	972	1,079	1,023	1,121	1,108	1,016
	Rate	3.5	3.7	3.3	3.1	3.0	3.0	3.3	3.1	3.4	3.3	3.0
Other respiratory‡	Cases	63	51	55	64	97	115	92	97	84	103	108
	Rate	0.2	0.2	0.2	0.2	0.3	0.4	0.3	0.3	0.3	0.3	0.3
Military	Cases	13	7	12	17	20	18	19	16	28	21	19
	Rate	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.1	0.1	0.1
CNS	Cases	16	17	20	26	19	20	18	22	12	14	23
	Rate	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.1
Nonrespiratory	Cases	258	234	240	249	251	245	210	205	183	209	197
	Rate	0.8	0.8	0.8	0.8	0.8	0.8	0.6	0.6	0.5	0.6	0.6
Other §	Cases	161	163	180	170	181	165	144	149	156	142	150
	Rate	0.5	0.5	0.6	0.5	0.6	0.5	0.4	0.5	0.5	0.4	0.4
Unknown	Cases	29	34	34	52	3	0	1	0	0	0	0
	Rate	0.1	0.1	0.1	0.2	0.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>TOTAL</b>	Cases	1,724	1,772	1,667	1,631	1,613	1,641	1,654	1,576	1,643	1,659	1,577
	Rate	5.6	5.7	5.3	5.2	5.0	5.1	5.1	4.8	4.9	4.9	4.6

\* Primary includes primary respiratory tuberculosis and tuberculous pleurisy in primary progressive tuberculosis, (ICD-9 codes 010.0-010.9; ICD-10 A15.7 and A16.7).

† Pulmonary includes tuberculosis of the lungs and conducting airways which includes tuberculous fibrosis of the lung, tuberculous bronchiectasis, tuberculous pneumonia, tuberculous pneumothorax, isolated tracheal or bronchial tuberculosis and tuberculous laryngitis; (ICD-9 codes 011-011.9, 012.2, 012.3; ICD-10 codes A15.0-A15.3, A15.5, A15.9, A16.0-A16.2, A16.4, A16.9). Included within the pulmonary counts are cases, regardless of ICD site designation (except for primary diagnosis) for which smear results and culture results are positive for sputum, bronchial wash and GI wash.

‡ Other respiratory includes tuberculous pleurisy (non-primary) and tuberculosis of: intrathoracic lymph nodes, mediastinum, nasopharynx, nose (septum), and sinus (any nasal) (ICD-9 codes: 012.0, 012.1 and 012.8; ICD-10 codes: A15.4, A15.6, A15.8, A16.3, A16.5, A16.8).

§ Other includes tuberculosis of intestines, peritoneum and mesenteric glands, bones and joints, genitourinary system, skin, eye, ear, thyroid, adrenal and spleen.

\*\* Cases and rates for 2010 are provisional until publication of the Tuberculosis in Canada – 2010 Annual Report. These numbers will likely change.

**Table 6:**

Reported new active and re-treatment tuberculosis cases and incidence rate per 100,000 by main diagnostic site – Canada and provinces/territories: 2010

Main diagnostic site	Province/territory												
	Canada	N.L.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	Y.T.	N.W.T.
Cases	64	0	0	1	2	2	9	8	18	5	3	0	4
Rate	0.2	0.0	0.0	0.1	0.3	0.0	0.1	0.6	1.7	0.1	0.1	0.0	9.1
Cases	1,016	7	0	4	6	144	392	85	48	74	172	6	4
Rate	3.0	1.4	0.0	0.4	0.8	1.8	3.0	6.9	4.6	2.0	3.8	17.4	9.1
Cases	108	0	1	1	0	13	41	9	8	12	12	0	0
Rate	0.3	0.0	0.7	0.1	0.0	0.2	0.3	0.7	0.8	0.3	0.3	0.0	0.0
Cases	19	0	0	1	0	4	6	5	0	1	1	0	1
Rate	0.1	0.0	0.0	0.1	0.0	0.1	0.0	0.4	0.0	0.0	0.0	0.0	2.3
Cases	23	0	0	0	0	4	12	1	1	0	3	0	1
Rate	0.1	0.0	0.0	0.0	0.0	0.1	0.1	0.1	0.1	0.0	0.1	0.0	2.3
Cases	197	0	0	1	1	19	97	13	6	26	34	0	0
Rate	0.6	0.0	0.0	0.1	0.1	0.2	0.7	1.1	0.6	0.7	0.8	0.0	0.0
Cases	150	1	0	2	1	22	76	11	1	16	16	0	1
Rate	0.4	0.2	0.0	0.2	0.1	0.3	0.6	0.9	0.1	0.4	0.4	0.0	2.3
Cases	0	0	0	0	0	0	0	0	0	0	0	0	0
Rate	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Cases	1,577	8	1	10	10	208	633	132	82	134	241	6	11
Rate	4.6	1.6	0.7	1.1	1.3	2.6	4.8	10.7	7.8	3.6	5.3	17.4	25.1

\* Primary includes primary respiratory tuberculosis and tuberculous pleurisy in primary progressive tuberculosis, (ICD-9 codes 010.0-010.9; ICD-10 A15.7 and A16.7).

† Pulmonary includes tuberculosis of the lungs and conducting airways which includes tuberculous fibrosis of the lung, tuberculous bronchiectasis, tuberculous pneumonia, tuberculous pneumothorax, isolated tracheal or bronchial tuberculosis and tuberculous laryngitis; (ICD-9 codes 011-011.9, 012.2, 012.3; ICD-10 codes A15.0-A15.3, A15.5, A15.9, A16.0-A16.2, A16.4, A16.9). Included within the pulmonary counts are cases, regardless of ICD site designation (except for primary diagnosis) for which smear results and culture results are positive for sputum, bronchial wash and GI wash.

‡ Other respiratory includes tuberculous pleurisy (non-primary) and tuberculosis of: intrathoracic lymph nodes, mediastinum, nasopharynx, nose (septum), and sinus (any nasal) (ICD-9 codes: 012.0, 012.1 and 012.8; ICD-10 codes: A15.4, A15.6, A15.8, A16.3, A16.5 A16.8).

§ Other includes tuberculosis of intestines, peritoneum and mesenteric glands, bones and joints, genitourinary system, skin, eye, ear, thyroid, adrenal and spleen.

\*\* Cases and rates for 2010 are provisional until publication of the Tuberculosis in Canada – 2010 Annual Report. These numbers will likely change.

**Table 7.**

Reported new active and re-treatment tuberculosis cases by origin and main diagnostic site - Canada: 2010

Origin	Total	Main diagnostic site					
		Primary*	Respiratory	Other Respiratory†	Miliary	CNS	Nonrespiratory
		Pulmonary‡				Peripheral lymph node	Unknown
North American Indian	188	117	17	4	2	8	0
Status (registered) Indian	184	114	17	4	2	8	0
Status (registered) Indian - On reserve	106	62	12	1	0	7	0
Status (registered) Indian - Off reserve	72	47	5	3	1	1	0
Status (registered) Indian - Unknown	6	5	0	0	1	0	0
Non-Status	4	3	0	0	0	0	0
Inuit	116	86	12	1	1	0	0
Métis	26	18	1	0	1	0	0
Total Aboriginal	330	221	30	5	4	8	0
Non-aboriginal	183	125	13	5	2	10	0
Total	513	346	43	10	6	18	0
Africa, High HIV Prevalence	90	54	4	0	3	19	0
Africa, Low HIV Prevalence	28	19	1	0	1	4	0
American Region - Latin American and Caribbean Countries	64	42	5	0	1	10	0
Eastern Europe	18	14	1	1	0	1	0
Established Market Economies and Central Europe	51	38	10	1	0	1	0
Eastern Mediterranean	90	50	6	1	0	17	0
South-East Asia	249	155	17	1	5	44	0
Western Pacific Region	429	271	18	4	7	76	0
Unknown	19	12	2	0	0	3	0
Total	1,038	655	64	8	17	175	0
Unknown Birthplace	26	15	1	1	0	4	0
<b>TOTAL **</b>	<b>1,577</b>	<b>1016</b>	<b>108</b>	<b>19</b>	<b>23</b>	<b>197</b>	<b>0</b>

\* Primary includes primary respiratory tuberculosis and tuberculous pleurisy in primary progressive tuberculosis, (ICD-9 codes 010.0-010.9; ICD-10 A15.7 and A16.7).

† Pulmonary includes tuberculosis of the lungs and conducting airways which includes tuberculous fibrosis of the lung, tuberculous bronchiectasis, tuberculous pneumonia, tuberculous pneumothorax, isolated tracheal or bronchial tuberculosis and tuberculous laryngitis; (ICD-9 codes 011-011.9, 012.2, 012.3; ICD-10 codes A15.0-A15.3, A15.5, A15.9, A16.0-A16.2, A16.4, A16.9). Included within the pulmonary counts are cases, regardless of ICD site designation (except for primary diagnosis) for which smear results and culture results are positive for sputum, bronchial wash and GI wash.

‡ Other respiratory includes tuberculous pleurisy (non-primary) and tuberculosis of: intrathoracic lymph nodes, mediastinum, nasopharynx, nose (septum), and sinus (any nasal) (ICD-9 codes: 012.0, 012.1 and 012.8; ICD-10 codes: A15.4, A15.6, A15.8, A16.3, A16.5 A16.8).

§ Other includes tuberculosis of intestines, peritoneum and mesenteric glands, bones and joints, genitourinary system, skin, eye, ear, thyroid, adrenal and spleen.

\*\* Cases and rates for 2010 are provisional until publication of the Tuberculosis in Canada – 2010 Annual Report. These numbers will likely change.