

**West coast of Newfoundland capelin (*Mallotus villosus* M.) and Atlantic herring (*Clupea harengus harengus* L.) larval survey, part 5: Description of the data collected in partnership with the industry (Barry Group) in July 2007**

F. Grégoire, W. Barry, J. Barry, C. Lévesque, J.-L. Beaulieu, and M.-H. Gendron

Department of Fisheries and Oceans  
Science Branch  
Maurice Lamontagne Institute  
P.O. Box 1000, 850 Route de la Mer  
Mont-Joli (Québec)  
G5H 3Z4

2008

**Canadian Data Report of Fisheries  
and Aquatic Sciences 1205**



Pêches  
et Océans

Fisheries  
and Oceans

**Canada**

## **Canadian Data Report of Fisheries and Aquatic Sciences**

Data reports provide a medium for filing and archiving data compilations where little or no analysis is included. Such compilations commonly will have been prepared in support of other journal publications or reports. The subject matter of data reports reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are indexed in the data base *Aquatic Sciences and Fisheries Abstracts*.

Numbers 1-25 in this series were issued as Fisheries and Marine Service Data Records. Numbers 26-160 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Data Reports. The current series name was introduced with the publication of report number 161.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

## **Rapport statistique canadien des sciences halieutiques et aquatiques**

Les rapports statistiques servent à classer et à archiver les compilations de données pour lesquelles il y a peu ou point d'analyse. Ces compilations auront d'ordinaire été préparées à l'appui d'autres publications ou rapports. Les sujets des rapports statistiques reflètent la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports statistiques ne sont pas destinés à une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans autorisation écrite préalable de l'établissement d'origine. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont indexés dans la base de données *Aquatic Sciences and Fisheries Abstracts*.

Les numéros 1 à 25 de cette série ont été publiés à titre de relevés statistiques, Services des pêches et de la mer. Les numéros 26 à 160 ont été publiés à titre de rapports statistiques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 161.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement d'origine dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Data Report of  
Fisheries and Aquatic Sciences 1205

2008

**WEST COAST OF NEWFOUNDLAND CAPELIN (*Mallotus villosus* M.)  
AND ATLANTIC HERRING (*Clupea harengus harengus* L.)  
LARVAL SURVEY, PART 5: DESCRIPTION OF THE DATA COLLECTED IN  
PARTNERSHIP WITH THE INDUSTRY (BARRY GROUP) IN JULY 2007**

François Grégoire<sup>1</sup>, William Barry<sup>2</sup>, Jim Barry<sup>2</sup>, Charline Lévesque<sup>1</sup>,  
Jean-Louis Beaulieu<sup>1</sup>, and Marie-Hélène Gendron<sup>1</sup>

<sup>1</sup>Department of Fisheries and Oceans  
Science Branch  
Maurice Lamontagne Institute  
P.O. Box 1000, 850 Route de la Mer  
Mont-Joli (Québec)  
G5H 3Z4

---

<sup>2</sup>Barry Group, 415 Griffin Drive, Corner Brook, Newfoundland, A2H 3E9

© Her Majesty the Queen in Right of Canada, 2008  
Cat. No. Fs 97-13/1205E ISSN 1488-5395

This publication should be cited as follows:

Grégoire, F., W. Barry, J. Barry, C. Lévesque, J.-L. Beaulieu, and M.-H. Gendron. 2008. West coast of Newfoundland capelin (*Mallotus villosus* M.) and Atlantic herring (*Clupea harengus harengus* L.) larval survey, part 5: Description of the data collected in partnership with the industry (Barry Group) in July 2007. Can. Data Rep. Fish. Aquat. Sci. 1205: vi + 31 pp.

**TABLE OF CONTENTS**

LIST OF TABLES .....iv

LIST OF FIGURES .....iv

LIST OF APPENDICES ..... v

ABSTRACT / RÉSUMÉ .....vi

INTRODUCTION ..... 1

MATERIAL AND METHODS ..... 1

RESULTS ..... 2

ACKNOWLEDGEMENTS ..... 2

REFERENCES ..... 2

## LIST OF TABLES

Table 1.	Description of the stations and tows completed during the capelin and Atlantic herring larval survey of July 2007 .....	4
Table 2.	Mean water temperature (°C) by depth interval (m) for the stations sampled during the capelin and Atlantic herring larval survey of July 2007 .....	5
Table 3.	Mean water salinity by depth interval (m) for the stations sampled during the capelin and Atlantic herring larval survey of July 2007 .....	7
Table 4.	Descriptive statistics of the length measurements (mm) of capelin (A), Atlantic herring (B), and Atlantic mackerel (C) larvae sampled during the July 2007 survey .....	9

## LIST OF FIGURES

Figure 1.	Map of the west coast of Newfoundland showing the study area and the other locations mentioned in the document.....	12
Figure 2.	Map of the 46-station sampling grid of the capelin and Atlantic herring larval survey of July 2007 .....	13
Figure 3.	Mean water temperature (°C) by depth intervals of 2 m from 0 m to 20 m and intervals of 5 m between 20 m and 55 m .....	14
Figure 4.	Length (mm) frequency distributions of capelin, Atlantic herring, and Atlantic mackerel larvae sampled during the July 2007 survey .....	16

**LIST OF APPENDICES**

Appendix 1. Temperature profiles for stations sampled during the capelin and Atlantic herring larval survey of July 2007 on the west coast of Newfoundland .....	17
Appendix 2. Volume of the plankton sample (mL) and number of eggs sorted and identified from samples collected during the capelin and Atlantic herring larval survey of July 2007 on the west coast of Newfoundland .....	26
Appendix 3. Number of larvae sorted and identified from samples collected during the capelin and Atlantic herring larval survey of July 2007 on the west coast of Newfoundland .....	29

**ABSTRACT**

Grégoire, F., W. Barry, J. Barry, C. Lévesque, J.-L. Beaulieu, and M.-H. Gendron. 2008. West coast of Newfoundland capelin (*Mallotus villosus* M.) and Atlantic herring (*Clupea harengus harengus* L.) larval survey, part 5: Description of the data collected in partnership with the industry (Barry Group) in July 2007. Can. Data Rep. Fish. Aquat. Sci. 1205: vi + 31 pp.

In partnership with the Barry Group in Corner Brook, a larval survey was conducted on the west coast of Newfoundland from July 18 to 21, 2007. Eggs and larvae of several of species of fish were found at all sampled stations. The two most abundant groups of eggs were CYT (cunner [*Tautogolabrus adspersus*] and yellowtail flounder [*Limanda ferruginea*]) and CHW (cod [*Gadus morhua*], haddock [*Melanogrammus aeglefinus*], and witch flounder [*Glyptocephalus cynoglossus*]). Mackerel eggs (*Scomber scombrus*) were found at most of the stations. Of the thirteen species of larvae identified, the most abundant were those of cunner, righteye flounder (Pleuronectidae), mackerel, fourbeard rockling (*Enchelyopus cimbrius*), and capelin (*Mallotus villosus*). Depth-based profiles of water temperature and salinity were made for each station. All the biological and oceanographic data collected during the survey were compiled and are presented in the tables, figures, and appendices found in this document. These data will be analyzed and published in a forthcoming report.

**RÉSUMÉ**

Grégoire, F., W. Barry, J. Barry, C. Lévesque, J.-L. Beaulieu et M.-H. Gendron. 2008. West coast of Newfoundland capelin (*Mallotus villosus* M.) and Atlantic herring (*Clupea harengus harengus* L.) larval survey, part 5: Description of the data collected in partnership with the industry (Barry Group) in July 2007. Can. Data Rep. Fish. Aquat. Sci. 1205: vi + 31 pp.

En partenariat avec le Groupe Barry de Corner Brook, un relevé larvaire a été réalisé sur la côte ouest de Terre-Neuve entre le 18 et le 21 juillet 2007. Des œufs et des larves de plusieurs espèces de poissons ont été retrouvés à toutes les stations échantillonnées. Les deux plus abondants groupes d'œufs ont été CYT (tanche-tautogue [*Tautogolabrus adspersus*] et limande à queue jaune [*Limanda ferruginea*]) et CHW (morue [*Gadus morhua*], aiglefin [*Melanogrammus aeglefinus*] et plie grise [*Glyptocephalus cynoglossus*]). Des œufs de maquereau (*Scomber scombrus*) ont été retrouvés à la plupart des stations. Des treize espèces de larves identifiées, les plus importantes ont été celles de la tanche-tautogue, de la plie (Pleuronectidae), du maquereau, de la motelle à quatre barbillons (*Enchelyopus cimbrius*) et du capelan (*Mallotus villosus*). Des profils de la température et de la salinité de l'eau en fonction de la profondeur ont été réalisés à chaque station. Toutes les données biologiques et océanographiques recueillies lors du relevé ont été compilées et sont présentées dans les tableaux, figures et annexes du présent document. Ces données seront analysées et publiées dans un rapport à venir.



## INTRODUCTION

In collaboration with the industry, a plankton survey was conducted on the west coast of Newfoundland in July 2004 and 2005. The objective was to collect basic information on the distribution and abundance of capelin and herring larvae in an active fishing area located close to the coast, between Bonne Bay and Port au Port Bay (Figure 1). This area of the west coast of Newfoundland had never before been covered by this type of survey. Eggs from three species and three groups of species were identified as well as the larvae from 16 to 20 species (Grégoire et al. 2005, 2006a). The average abundances of capelin and herring larvae were estimated, and various biodiversity measurements indicated that the larval community structure in the sampled area was characterized by the occurrence of a large number of scarce species and by commercial species concentrated in specific locations (Grégoire et al. 2006b).

A third plankton survey was conducted in July 2007, again with the financial and logistic support of the Barry Group. While the objective was the same, the sampling area was extended to include St. George's Bay (Figure 1).

The present report compiles and documents the biological and oceanographic data collected during the 2007 survey. The detailed analysis of these data will be presented in a forthcoming report.

## MATERIAL AND METHODS

Plankton sampling was made by the *Ocean Provider*, a large seiner from the west coast of Newfoundland. The sampling procedures were the same as those performed in 2004 and 2005: At each station, plankton was collected using a 61 cm bongo sampler equipped with two nets lined with 333  $\mu\text{m}$  mesh (Posgay and Marak, 1980). A General Oceanics flowmeter was placed at the opening of each net to measure the volume of water filtered. The 10-min tows were carried out following a saw-tooth pattern (Hempel, 1973) between the surface and a maximum depth of 50 m, or to within 5 m of the bottom at shallower stations. Once on deck, the nets were raised and rinsed with seawater. The plankton sample from the first net was preserved in a diluted buffered formaldehyde solution (4-5%) (Hunter, 1985) while that of the second net was placed in a concentrated ethanol solution (95%). A CTD instrument (Sea-Bird Electronics Inc.) was installed on the bongo sampler to obtain a profile of water temperature and salinity in the water column.

Plankton sorting was performed at the Fisheries and Oceans laboratory in Mont-Joli, Québec, during the fall of 2007. Each sample was split using the Van Guelpen beaker technique (Van Guelpen et al. 1982). The criteria used to identify fish eggs and larvae were mainly drawn from Fritzsche (1978), Elliott and Jimenez (1981), and Fahay (1983). Total lengths (Hardy, 1978) of capelin, herring, and mackerel larvae preserved in the formaldehyde solution were measured to within 0.01 mm. The larvae preserved in ethanol will be used in growth studies based on otolith microstructure. The results of the sorting were input to data files and validated.

## RESULTS

The survey took place between 18 and 21 July 2007. A total of 45 stations were sampled in the study area (Figure 2). Station 22 was not sampled due to the lack of time and a malfunction of the CTD probe occurred at stations 5 and 28. Positions and depths station are presented in Table 1 along with the duration of each tow and the volume of water filtered. Temperature profiles by station are shown in Appendix 1. Mean temperatures and salinities were calculated by depth intervals of 2 m from 0 m to 20 m and by intervals of 5 m between 20 m and 65 m (Tables 2 and 3). Between 0 m and 12 m, warmer water temperatures were found in St. George's Bay and inshore Port au Port Bay and Bay of Islands (Figure 3). Starting from approximately 12–14 m of depth, water temperatures fell quickly at the stations located in the western part of St. George's Bay and for inshore stations in Port au Port Bay and Bay of Islands.

Fish eggs were found at each station (Appendix 2), the most abundant groups of which were CYT and CHW. The CYT group includes eggs of cunner (*Tautoglabrus adspersus*) and yellowtail flounder (*Limanda ferruginea*) and the CHW group cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), and witch flounder (*Glyptocephalus cynoglossus*). Mackerel eggs (*Scomber scombrus*) were found at most of the stations. Mackerel egg stages were identified based on criteria presented in Girard (2000).

The larvae of 13 fish species were also found during the survey (Appendix 3). The most abundant were those of cunner, righteye flounder (Pleuronectidae), mackerel, fourbeard rockling (*Enchelyopus cimbrius*), and capelin (*Mallotus villosus*). The mean lengths of capelin and herring larvae were 6.49 mm (n=484) (Table 4A) and 14.08 mm (n=86) (Table 4B), respectively. The mean length of mackerel larvae was 3.66 mm (n=714) (Table 4C). Two main modes characterize the length frequency distribution of mackerel larvae (Figure 4) compared to one main mode for capelin and herring larvae.

## ACKNOWLEDGEMENTS

A very sincere thanks to captain Bill McCarthy and to all the members of the *Ocean Provider* crew (Frank, Ron, and Peter) for their contribution to the success of this survey. We also want to thank Laure Devine and Alain Fréchet for their reviews of this report.

## REFERENCES

- Elliott, E. M., and D. Jimenez. 1981. Laboratory manual for the identification of ichthyoplankton from the Beverly - Salem Harbor area. Dept. of Fisheries, Wildlife and Recreational Vehicles. Massachusetts. 230 pp.
- Fahay, M. P. 1983. Guide to the early stages of marine fishes occurring in the western North Atlantic Ocean, Cape Hatteras to the southern Scotian Shelf. J. Northwest Atl. Fish. Sci. 4: 1-423.

- Fritzsche, R. A. 1978. Development of fishes of the Mid-Atlantic Bight: an atlas of egg, larval and juvenile stages. Vol. V. Chaetodontidae through Ophidiidae. U.S. Fish Wildl. Serv., Biol. Serv. Program FWS/OBS-78/12. 340 pp.
- Girard, L. 2000. Identification of mackerel (*Scomber scombrus* L.) eggs sampled during abundance surveys in the southern Gulf of St. Lawrence. In *The Atlantic mackerel (Scomber scombrus L.) of NAFO subareas 2 to 6, Chapter 4. Edited by F. Grégoire.* CSAS Res. Doc. 2000/021, pp. 119-137.
- Grégoire, F., W. Barry, J. Barry, L. Lefebvre, C. Lévesque, and J. Hudon. 2005. West coast of Newfoundland capelin (*Mallotus villosus* M.) and Atlantic herring (*Clupea harengus harengus* L.) larval survey, part 1: Description of the data collected in partnership with the industry (Barry Group) in July 2004. Can. Data Rep. Fish. Aquat. Sci. 1168 (Part1): vi + 32 pp.
- Grégoire, F., W. Barry, J. Barry, L. Lefebvre, C. Lévesque, and J. Hudon. 2006a. West coast of Newfoundland capelin (*Mallotus villosus* M.) and Atlantic herring (*Clupea harengus harengus* L.) larval survey, part 3: Description of the data collected in partnership with the industry (Barry Group) in July 2005. Can. Data Rep. Fish. Aquat. Sci. 1168 (Part3): vi + 32 pp.
- Grégoire, F., W. Barry, and J. Barry. 2006b. West coast of Newfoundland capelin (*Mallotus villosus* M.) and Atlantic herring (*Clupea harengus harengus* L.) larval survey, part 2: Abundance estimates and marine community analyses of the data collected in partnership with the industry (Barry Group) in July 2004. Can. Tech. Rep. Fish. Aquat. Sci. 2650 (Part2): viii + 50 pp.
- Hardy, J. D. 1978. Development of fishes of the Mid-Atlantic Bight: an atlas of egg, larval and juvenile stages. Vol. II. Anguillidae through Syngnathidae. U.S. Fish Wildl. Serv., Biol. Serv. Program FWS/OBS-78/12. 458 pp.
- Hempel, G. 1973. Fish egg and larval surveys. FAO Fisheries Technical Paper No. 122. 82 pp.
- Hunter, J. R. 1985. Preservation of northern anchovy in formaldehyde solution. In *An egg production method for estimating spawning biomass of pelagic fish: Application to the northern anchovy, *Eugraulis mordax*.* Edited by R. Lasker. NOAA Technical Report NMFS 36, pp. 63-66.
- Posgay, J. A., and R. R. Marak. 1980. The MARMAP Bongo zooplankton samplers. J. Northwest Atl. Fish. Sci. 1: 91-99.
- Van Guelpen, L., D. F. Markle, and D. J. Duggan. 1982. An evaluation of accuracy, precision, and speed of several zooplankton sub-sampling techniques. J. Cons. Int. Explor. Mer. 40: 226-236.

Table 1. Description of the stations and tows completed during the capelin and Atlantic herring larval survey of July 2007.

STATION NUMBER	DATE (yyyy-mm-dd)	TIME (hh:mm) (NDT)	LATITUDE LONGITUDE		DEPTH		TOW DURATION (mm:ss)	VOLUME OF WATER FILTERED (m <sup>3</sup> )
			°N (degrees minutes)	°W (degrees minutes)	Station bottom (m)	Maximum sampled (m)		
1	2007-07-21	08:18	49° 09'	58° 12'	111	48	11:09	286
2	2007-07-21	05:49	49° 12'	58° 12'	160	63	12:34	279
3	2007-07-20	18:28	49° 27'	58° 12'	43	34	10:23	249
4	2007-07-20	17:39	49° 33'	58° 12'	61	56	11:08	277
5*	2007-07-18	09:49	49° 06'	58° 18'	51		11:00	203
6	2007-07-21	06:32	49° 11'	58° 18'	135	61	11:56	292
7	2007-07-21	07:13	49° 15'	58° 18'	80	62	10:04	238
8	2007-07-20	19:35	49° 20'	58° 18'	37	30	10:26	251
9	2007-07-20	16:38	49° 27'	58° 18'	56	48	10:57	254
10	2007-07-18	10:28	49° 11'	58° 24'	64	52	10:28	207
11	2007-07-20	14:47	49° 15'	58° 24'	40	33	10:57	277
12	2007-07-20	15:32	49° 20'	58° 24'	47	33	10:34	262
13	2007-07-20	12:51	49° 03'	58° 30'	48	48	11:51	286
14	2007-07-18	11:19	49° 07'	58° 30'	88	51	11:19	216
15	2007-07-20	14:01	49° 12'	58° 30'	65	52	10:27	256
16	2007-07-20	11:17	48° 51'	58° 36'	42	33	10:48	268
17	2007-07-18	14:20	48° 57'	58° 36'	35	24	10:50	242
18	2007-07-18	13:22	49° 03'	58° 36'	44	33	09:58	208
19	2007-07-18	12:37	49° 07'	58° 36'	70	57	12:04	210
20	2007-07-20	09:23	48° 39'	58° 42'	34	28	09:56	245
21	2007-07-20	10:18	48° 45'	58° 42'	27	17	10:03	261
22**			48° 51'	58° 42'				
23	2007-07-18	15:05	48° 57'	58° 42'	31	16	10:07	248
24	2007-07-20	08:37	48° 36'	58° 48'	21	16	10:24	255
25	2007-07-20	08:02	48° 39'	58° 48'	23	17	10:21	262
26	2007-07-20	07:23	48° 42'	58° 48'	16	11	10:01	248
27	2007-07-18	17:15	48° 48'	58° 48'	29	16	10:03	249
28*	2007-07-18	16:18	48° 51'	58° 48'	43		10:31	231
29	2007-07-20	06:37	48° 39'	58° 54'	15	10	10:24	242
30	2007-07-18	18:00	48° 45'	58° 54'	29	19	09:56	245
31	2007-07-18	19:19	48° 39'	59° 06'	43	28	11:10	258
32	2007-07-19	18:35	48° 30'	59° 18'	59	44	12:26	312
33	2007-07-19	06:00	48° 28'	59° 06'	68	47	10:23	244
34	2007-07-19	16:01	48° 28'	58° 54'	63	50	10:14	259
35	2007-07-19	14:47	48° 30'	58° 42'	44	35	11:08	269
36	2007-07-19	14:12	48° 30'	58° 36'	48	33	10:54	253
37	2007-07-19	13:38	48° 27'	58° 36'	89	55	10:14	241
38	2007-07-19	17:33	48° 24'	59° 12'	87	53	10:10	235
39	2007-07-19	06:55	48° 24'	59° 00'	91	50	11:20	256
40	2007-07-19	12:04	48° 24'	58° 48'	44	37	09:51	248
41	2007-07-19	12:49	48° 24'	58° 42'	47	41	10:34	256
42	2007-07-19	07:59	48° 18'	59° 06'	99	52	12:03	263
43	2007-07-19	10:29	48° 18'	58° 54'	104	52	11:47	266
44	2007-07-19	11:11	48° 18'	58° 48'	42	35	10:33	255
45	2007-07-19	09:02	48° 12'	59° 00'	85	54	10:30	249
46	2007-07-19	09:43	48° 13'	58° 54'	34	26	10:43	257

\* Malfunction of the CTD probe

\*\* Not sampled

Table 2. Mean water temperature (°C) by depth interval (m) for the stations sampled during the capelin and Atlantic herring larval survey of July 2007.

DEPTH (m)	STATION																						
	1	2	3	4	5*	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22**	23
0-2	16.00	16.90	15.73	14.43		14.82	14.96	16.01	14.49	14.57	14.73	14.78	15.20	13.71	14.51	14.90	14.33	13.85	14.30	15.51	15.69		14.42
2-4	15.96	16.86	15.18	14.38		14.73	14.92	14.85	14.43	14.43	14.23	14.69	14.82	13.62	14.39	14.62	14.22	13.76	14.29	15.46	15.42		14.20
4-6	15.56	16.08	14.64	14.21		14.74	14.84	14.55	14.20	14.29	14.03	14.54	14.52	13.55	14.22	14.43	14.12	13.67	14.24	15.43	15.16		14.03
6-8	14.84	14.90	14.59	14.14		14.70	14.70	14.52	14.02	14.06	13.92	14.44	14.28	13.47	14.11	14.27	13.97	13.56	14.19	15.24	14.94		13.86
8-10	14.29	14.66	14.56	14.12		14.58	14.52	14.51	13.93	13.87	13.79	14.32	14.10	13.37	14.01	14.23	13.90	13.49	14.10	14.75	14.76		13.69
10-12	14.18	14.24	14.51	13.99		14.43	14.33	14.50	13.85	13.69	13.70	13.98	14.01	13.24	13.95	14.19	13.87	13.39	13.79	13.20	14.65		13.65
12-14	14.00	13.75	14.38	13.71		14.22	14.20	14.45	13.73	13.61	13.55	13.78	13.94	13.17	13.80	14.12	13.83	13.12	13.30	11.47	14.48		13.62
14-16	13.82	13.63	13.92	13.38		14.07	14.14	14.35	13.47	13.56	13.42	13.65	13.90	13.14	13.70	13.98	13.81	12.40	13.20	10.69	14.09		13.57
16-18	13.63	13.44	13.55	13.15		13.94	14.03	14.03	13.22	13.53	13.35	13.24	13.83	12.98	13.54	13.74	13.78	11.77	13.08	9.96	13.77		
18-20	13.43	13.27	13.12	12.75		13.78	13.60	13.57	12.99	13.50	13.29	12.56	13.70	12.78	13.23	13.67	13.68	10.98	12.93	9.14			
20-25	12.82	12.53	12.35	10.06		13.39	12.38	12.34	11.69	13.47	12.60	11.19	13.51	12.27	12.27	13.10	13.23	8.58	12.03	7.16			
25-30	11.40	11.26	10.01	5.67		11.36	10.62	8.97	8.66	12.88	9.06	6.95	11.42	10.88	9.72	10.26		6.54	7.93	5.66			
30-35	8.48	9.80	8.21	4.03		7.71	7.49	7.74	4.29	11.17	6.08	5.20	6.22	6.36	4.42	5.12		5.57	5.22				
35-40	4.96	5.24		3.06		5.02	3.93		2.56	6.70			3.73	3.84	2.79					3.20			
40-45	2.98	3.33		2.45		3.26	1.82		1.80	4.37			2.39	2.40	1.79					2.39			
45-50	2.33	1.99		1.38		1.96	1.11		1.44	3.57			2.05	1.80	1.22					1.68			
50-55		1.21		0.27		0.85	0.53			2.95			1.52	0.74						1.01			
55-60		0.59		0.11		0.39	0.06													0.61			
60-65		0.34				0.42	-0.15																

Table 2. (Continued).

DEPTH (m)	STATION																							
	24	25	26	27	28*	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	
0-2	16.10	15.57	15.72	13.97		15.17	13.96	13.62	14.84	15.53	16.07	16.52	17.93	15.70	15.85	15.14	15.37	15.88	15.58	15.27	15.12	15.31	15.27	
2-4	16.08	15.46	15.62	13.95		15.01	13.94	13.59	14.44	15.51	15.97	16.30	17.16	15.19	15.72	15.10	15.28	15.75	15.56	15.25	15.08	15.29	15.24	
4-6	15.95	15.38	15.47	13.94		14.92	13.93	13.57	14.26	15.17	15.93	15.93	16.65	15.01	15.67	15.08	15.22	15.60	15.48	15.18	15.02	14.79	15.19	
6-8	15.72	15.28	15.22	13.93		14.87	13.92	13.55	14.10	14.45	15.79	15.25	16.13	14.87	15.53	14.96	15.10	15.45	15.17	14.98	14.99	14.39	15.15	
8-10	15.23	14.91	15.07	13.90		14.62	13.90	13.50	13.95	14.11	15.55	14.79	15.28	14.76	15.08	14.37	14.90	15.21	14.17	14.58	14.92	14.08	15.07	
10-12	13.55	14.44	14.95	13.89		14.02	13.89	13.46	13.85	13.93	15.04	14.44	14.83	14.56	14.03	13.74	14.64	14.90	12.19	14.23	14.77	13.55	14.84	
12-14	11.25	13.83		13.88			13.87	13.42	13.69	13.79	14.35	14.26	14.56	14.42	13.08	12.37	14.32	14.60	9.20	14.15	14.33	13.26	14.36	
14-16	10.64	13.13		13.85			13.87	13.40	13.17	13.59	14.01	14.10	14.16	14.31	11.43	9.95	13.93	14.47	8.00	14.13	13.98	12.85	14.30	
16-18		12.47		13.83			13.85	13.39	12.98	12.98	13.60	13.91	13.94	14.11	10.32	8.45	13.76	14.30	6.94	14.03	13.78	11.37	14.16	
18-20							13.84	13.39	12.84	11.26	13.16	13.66	13.74	13.90	8.77	7.63	13.56	14.06	6.49	13.76	13.66	9.41	13.85	
20-25									13.36	12.42	9.23	10.84	11.40	12.42	12.73	7.28	6.09	11.35	11.04	5.43	12.34	12.67	7.33	11.44
25-30									11.82	11.31	6.68	7.98	5.67	7.62	9.44	5.36	5.08	7.60	5.27	3.95	6.50	7.87	5.42	7.14
30-35										7.73	4.06	6.24	2.09	3.32	3.60	3.84	4.33	4.78	1.78	2.86	4.59	2.96	3.51	
35-40										6.29	2.85	3.37			1.90	3.01	3.36	3.61	0.83	1.89	2.79	2.34	1.97	
40-45										4.44	1.79	2.10			0.91	2.42	2.07		0.77	1.20	1.50		1.15	
45-50											1.38	1.04			0.51	1.85	1.20			0.72	0.80		0.96	
50-55															0.44	1.15	0.89			0.49	0.33		0.48	
55-60																								
60-65																								

\* Malfunction of the CTD probe

\*\* Not sampled

Table 3. Mean water salinity by depth interval (m) for the stations sampled during the capelin and Atlantic herring larval survey of July 2007.

DEPTH (m)	STATION																							
	1	2	3	4	5*	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22**	23	
0-2	28.15	28.10	30.11	31.37		31.21	31.12	29.59	31.32	31.75	30.98	31.45	31.21	31.60	31.44	31.33	31.56	31.69	31.66	31.26	31.29			31.59
2-4	28.80	28.66	30.94	31.50		31.18	31.06	31.28	31.34	31.63	31.41	31.53	31.40	31.62	31.52	31.43	31.53	31.65	31.62	31.33	31.35			31.58
4-6	30.22	29.88	31.62	31.63		31.18	31.13	31.62	31.56	31.60	31.56	31.63	31.51	31.61	31.58	31.47	31.54	31.65	31.57	31.35	31.41			31.59
6-8	31.12	30.89	31.66	31.66		31.20	31.24	31.63	31.66	31.68	31.62	31.65	31.56	31.61	31.60	31.52	31.58	31.63	31.58	31.36	31.46			31.60
8-10	31.46	31.47	31.66	31.65		31.27	31.39	31.65	31.66	31.69	31.63	31.66	31.57	31.62	31.61	31.52	31.59	31.64	31.59	31.39	31.48			31.60
10-12	31.55	31.60	31.65	31.65		31.37	31.50	31.65	31.66	31.72	31.63	31.65	31.57	31.62	31.62	31.52	31.60	31.64	31.61	31.52	31.50			31.60
12-14	31.54	31.58	31.66	31.65		31.49	31.57	31.65	31.66	31.69	31.64	31.66	31.58	31.63	31.62	31.52	31.61	31.56	31.61	31.62	31.51			31.60
14-16	31.56	31.58	31.66	31.66		31.54	31.58	31.65	31.65	31.68	31.65	31.67	31.59	31.61	31.63	31.53	31.61	31.66	31.65	31.64	31.52			31.60
16-18	31.54	31.58	31.69	31.66		31.54	31.59	31.66	31.66	31.67	31.67	31.68	31.59	31.62	31.64	31.56	31.62	31.74	31.65	31.68	31.56			
18-20	31.56	31.60	31.67	31.67		31.55	31.61	31.67	31.67	31.68	31.67	31.70	31.59	31.61	31.63	31.58	31.63	31.76	31.57	31.72				
20-25	31.57	31.58	31.68	31.79		31.59	31.52	31.69	31.71	31.66	31.66	31.74	31.60	31.67	31.68	31.62	31.71	32.05	31.62	31.80				
25-30	31.57	31.57	31.71	31.97		31.61	31.60	31.83	31.86	31.63	31.76	31.92	31.73	31.75	31.82	31.77		32.15	32.05	31.89				
30-35	31.67	31.66	31.82	32.01		31.74	31.74	31.83	32.03	31.75	31.96	31.99	31.97	32.30	32.10	32.15		32.45	32.19					
35-40	31.88	31.88		32.04		31.92	31.96		32.11	32.17				32.13	32.37	32.16				32.30				
40-45	32.00	32.01		32.07		32.02	32.07		32.13	32.33				32.21	32.36	32.18				32.33				
45-50	32.09	32.08		32.19		32.08	32.08		32.15	32.19				32.23	32.29	32.21				32.42				
50-55		32.12		32.27		32.11	32.10			32.21					32.37	32.23				32.32				
55-60		32.14		32.24		32.14	32.14													32.70				
60-65		32.17				32.25	32.15																	

Table 3. (Continued).

DEPTH (m)	STATION																							
	24	25	26	27	28*	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	
0-2	31.13	31.33	31.32	31.56		31.25	31.61	31.53	31.34	31.36	31.42	31.05	29.93	31.11	31.30	31.33	31.41	31.42	31.33	31.33	31.44	31.35	31.33	
2-4	31.32	31.36	31.38	31.56		31.40	31.59	31.49	31.34	31.37	31.40	31.11	30.49	31.18	31.37	31.36	31.41	31.38	31.31	31.33	31.43	31.32	31.32	
4-6	31.34	31.38	31.41	31.57		31.43	31.59	31.49	31.33	31.42	31.41	31.20	30.88	31.22	31.38	31.36	31.41	31.38	31.32	31.34	31.44	31.34	31.32	
6-8	31.36	31.39	31.40	31.57		31.44	31.59	31.49	31.31	31.46	31.41	31.31	31.03	31.25	31.40	31.36	31.44	31.38	31.37	31.39	31.44	31.37	31.31	
8-10	31.38	31.41	31.40	31.57		31.47	31.59	31.49	31.31	31.50	31.41	31.39	31.23	31.25	31.39	31.39	31.42	31.40	31.39	31.47	31.44	31.37	31.30	
10-12	31.45	31.42	31.43	31.57		31.48	31.59	31.49	31.30	31.52	31.43	31.47	31.38	31.28	31.44	31.44	31.43	31.41	31.54	31.50	31.45	31.38	31.30	
12-14	31.58	31.46		31.57			31.59	31.49	31.31	31.53	31.47	31.52	31.46	31.39	31.49	31.57	31.45	31.43	31.76	31.50	31.46	31.42	31.27	
14-16	31.68	31.52		31.59			31.59	31.49	31.29	31.55	31.53	31.53	31.50	31.49	31.70	31.78	31.47	31.45	31.90	31.50	31.49	31.50	31.31	
16-18		31.60		31.61			31.60	31.49	31.31	31.61	31.54	31.52	31.49	31.50	31.70	31.88	31.48	31.47	31.92	31.51	31.50	31.63	31.44	
18-20							31.62	31.49	31.34	31.65	31.58	31.52	31.50	31.51	31.85	31.95	31.52	31.48	31.96	31.53	31.51	31.80	31.50	
20-25									31.50	31.45	31.83	31.68	31.57	31.57	31.51	31.89	32.00	31.69	31.65	32.00	31.61	31.60	31.90	31.71
25-30									31.57	31.61	31.97	31.77	32.00	31.88	31.72	32.00	32.01	31.94	32.04	32.06	31.99	31.91	31.99	32.15
30-35										31.89	32.14	31.93	32.26	32.19	32.14	32.09	32.06	32.11	32.26	32.12	32.07	32.32	32.14	
35-40										31.96	32.24	32.11			32.20	32.14	32.14	32.16	32.30	32.21	32.20	32.80	32.26	
40-45										32.11	32.32	32.26			32.23	32.20	32.24		32.23	32.25	32.29		32.39	
45-50											32.41	32.38			32.21	32.26	32.32			32.33	32.38		32.42	
50-55															32.18	32.32	32.49			32.38	32.46		32.43	
55-60																								
60-65																								

∞

\* Malfunction of the CTD probe

\*\* Not sampled



Table 4. Descriptive statistics of the length measurements (mm) of capelin (A), Atlantic herring (B), and Atlantic mackerel (C) larvae sampled during the July 2007 survey.

(A) Capelin

Station	Mean	Std. Dev.	Minimum	Maximum	Range	n
1	7.47	1.56	5.09	9.78	4.69	13
2	5.01	1.25	3.12	7.60	4.48	19
3	5.41	0.74	4.27	6.29	2.02	10
5	6.21	1.56	3.34	11.20	7.86	41
6	6.50	1.88	4.41	10.00	5.59	10
7	6.90	2.01	4.63	11.20	6.57	14
8	5.65	0.20	5.53	5.88	0.35	3
9	8.72	---	---	---	---	1
10	6.63	2.04	4.88	9.47	4.59	5
11	6.23	1.16	4.91	8.11	3.20	10
13	5.64	0.70	4.21	6.41	2.20	12
15	6.51	1.40	4.89	7.41	2.52	3
16	5.46	0.60	4.33	6.46	2.13	9
17	4.96	0.43	4.65	5.26	0.61	2
20	6.63	2.22	5.06	8.20	3.14	2
21	5.04	0.55	4.37	6.33	1.96	16
30	9.58	2.03	8.12	11.90	3.78	3
31	8.52	3.23	4.87	11.00	6.13	3
32	7.68	0.81	7.12	8.60	1.48	3
33	6.81	1.59	5.17	9.92	4.75	9
34	9.17	2.57	5.69	12.10	6.41	10
35	7.41	1.87	3.52	9.94	6.42	11
36	7.14	1.61	5.35	10.20	4.85	19
37	8.33	1.67	5.75	12.00	6.25	50
39	9.61	1.32	7.89	11.50	3.61	5
40	4.70	0.30	4.28	5.55	1.27	18
41	7.99	2.07	4.47	11.30	6.83	50
42	8.70	---	---	---	---	1
43	5.04	0.63	3.92	5.95	2.03	15
44	5.80	0.64	4.05	6.84	2.79	50
45	7.83	2.00	5.42	12.20	6.78	17
46	4.30	0.39	3.17	5.21	2.04	50
<b>TOTAL</b>	6.49	2.00	3.12	12.20	9.08	484

Table 4. (Continued).

**(B) Atlantic herring**

<b>Station</b>	<b>Mean</b>	<b>Std. Dev.</b>	<b>Minimum</b>	<b>Maximum</b>	<b>Range</b>	<b>n</b>
1	12.61	3.13	9.23	17.50	8.27	5
3	15.30	---	---	---	---	1
7	10.70	---	---	---	---	1
8	9.30	---	---	---	---	1
10	9.80	---	---	---	---	1
11	13.51	4.23	8.62	16.10	7.48	3
12	14.10	---	---	---	---	1
13	16.00	---	---	---	---	1
15	10.90	---	---	---	---	1
16	20.60	---	---	---	---	1
17	12.80	---	---	---	---	1
20	18.70	---	---	---	---	1
21	15.23	4.09	11.50	19.60	8.10	3
23	14.65	1.91	13.30	16.00	2.70	2
27	12.37	0.31	12.10	12.70	0.60	3
28	12.80	0.69	12.00	13.20	1.20	3
30	18.70	---	---	---	---	1
31	16.04	1.90	13.20	19.00	5.80	8
32	17.13	2.15	14.90	19.20	4.30	3
33	12.23	1.01	11.30	13.30	2.00	3
35	14.35	1.75	12.00	18.40	6.40	15
36	13.73	0.98	12.60	14.30	1.70	3
37	13.21	2.71	8.31	15.60	7.29	9
38	14.97	1.40	13.60	16.40	2.80	3
41	15.33	1.88	13.30	17.80	4.50	4
42	12.52	1.54	9.85	14.10	4.25	7
44	14.10	---	---	---	---	1
<b>TOTAL</b>	14.08	2.59	8.31	20.60	12.29	86

Table 4. (Continued).

## (C) Atlantic mackerel

Station	Mean	Std. Dev.	Minimum	Maximum	Range	n
1	2.58	0.46	1.89	3.08	1.19	7
2	2.93	0.57	2.09	3.55	1.46	5
3	3.19	0.51	2.39	4.53	2.14	19
5	2.49	0.15	2.28	2.60	0.32	4
6	2.82	0.51	1.95	3.80	1.85	9
7	3.23	0.44	2.85	3.76	0.91	4
8	2.85	0.74	2.03	4.58	2.55	10
9	2.81	0.36	2.24	3.13	0.89	5
10	3.75	---	---	---	---	1
11	3.05	0.31	2.80	3.50	0.70	4
12	2.89	0.34	2.35	3.58	1.23	16
13	3.43	1.55	2.27	6.17	3.90	8
15	4.01	1.30	2.86	5.96	3.10	10
16	5.15	1.74	3.00	8.03	5.03	8
17	7.94	3.20	5.67	10.20	4.53	2
19	3.32	0.70	2.58	3.98	1.40	3
21	2.82	0.71	1.63	5.78	4.15	31
23	2.77	0.26	2.47	2.97	0.50	3
25	2.66	0.42	2.36	2.95	0.59	2
26	2.34	0.42	1.75	3.14	1.39	11
27	4.36	1.57	1.97	9.95	7.98	25
30	4.90	1.33	2.72	7.09	4.37	8
31	2.99	0.82	2.25	4.99	2.74	9
32	3.59	1.22	2.31	6.81	4.50	36
33	3.37	1.16	2.10	8.13	6.03	50
34	4.21	1.13	2.86	6.58	3.72	31
35	2.93	0.74	1.68	5.88	4.20	50
36	3.83	1.02	2.36	6.01	3.65	50
37	3.76	0.99	2.65	6.68	4.03	49
38	4.18	1.47	2.46	7.63	5.17	34
39	4.01	1.28	2.36	6.85	4.49	33
40	5.54	1.81	2.62	9.46	6.84	21
41	3.55	1.15	2.25	8.92	6.67	50
42	4.82	1.49	2.10	7.13	5.03	25
43	4.74	---	---	---	---	1
44	4.58	1.51	2.52	6.68	4.16	13
45	3.47	1.52	2.07	11.40	9.33	41
46	3.32	0.55	2.51	4.84	2.33	26
<b>TOTAL</b>	3.66	1.33	1.63	11.40	9.77	714




Figure 1. Map of the west coast of Newfoundland showing the study area and the other locations mentioned in the document.




Figure 2. Map of the 46-station sampling grid of the capelin and Atlantic herring larval survey of July 2007.



Figure 3. Mean water temperature (°C) by depth intervals of 2 m from 0 m to 20 m and intervals of 5 m between 20 m and 55 m.




Figure 3. (Continued).






















Figure 4. Length (mm) frequency distributions of capelin, Atlantic herring, and Atlantic mackerel larvae sampled during the July 2007 survey.















Appendix 1. Temperature profiles for stations sampled during the capelin and Atlantic herring larval survey of July 2007 on the west coast of Newfoundland.
































Appendix 2. Volume of the plankton sample (mL) and number of eggs sorted and identified from samples collected during the capelin and Atlantic herring larval survey of July 2007 on the west coast of Newfoundland. Egg codes are as follows: H4B = hake (*Urophycis* sp.), fourbeard rockling (*Enchelyopus cimbrius*), and butterfish (*Peprilus triacanthus*); CHW = cod (*Gadus morhua*), haddock (*Melanogrammus aeglefinus*), and witch flounder (*Glyptocephalus cynoglossus*); CYT = cunner (*Tautoglabrus adspersus*) and yellowtail flounder (*Limanda ferruginea*).

Descriptions and pictures of mackerel eggs are found in Girard (2000).

Station	Sample volume (mL)	Eggs (n)									
		Mackerel					H4B	CHW	CYT	American plaice ( <i>Hippoglossoides platessoides</i> )	Windowpane ( <i>Scophthalmus aquosus</i> )
		Stage 1	Stage 2	Stage 3	Stage 4	Stage 5					
1	155	1	1	5	4	0	6	30	1608	2	0
2	140	0	1	2	1	1	4	30	160	2	0
3	90	10	3	2	0	0	16	10	416	0	0
4	110	0	0	0	0	0	0	222	0	0	0
5	90	2	5	15	9	0	24	28	1864	0	0
6	110	5	0	4	3	0	2	160	496	1	0
7	140	2	0	0	2	0	4	10	146	1	0
8	80	10	3	0	1	1	2	30	704	0	0
9	170	3	0	1	1	0	2	280	26	0	0
10	75	0	0	0	1	0	0	78	370	1	0
11	70	2	0	0	2	1	4	44	112	0	0
12	150	6	0	0	1	1	4	116	36	2	0
13	90	4	2	5	0	1	0	56	2976	0	0
14	50	0	0	0	0	0	0	90	4	0	0
15	60	2	0	3	2	0	1	304	448	1	0
16	75	2	2	10	4	0	16	32	4592	0	0
17	30	1	0	0	2	0	0	38	768	0	0
18	40	0	0	0	0	0	0	196	0	0	0
19	100	0	0	0	1	0	0	134	4	0	1
20	60	5	5	10	6	0	216	8	1360	0	0
21	80	3	1	15	9	2	64	20	2664	0	24
22*											
23	15	0	0	0	0	0	0	25	22	0	0
24	110	7	7	9	8	0	224	0	96	0	0
25	80	17	3	16	24	0	264	2	152	0	0
26	90	1	2	11	17	1	128	1	0	0	64
27	50	0	0	0	0	0	0	104	904	0	0
28	40	0	0	0	0	0	0	66	54	0	0

Station	Sample volume (mL)	Eggs (n)										
		Mackerel					H4B	CHW	CYT	American plaice	Windowpane	
		Stage 1	Stage 2	Stage 3	Stage 4	Stage 5				( <i>Hippoglossoides platessoides</i> )	( <i>Scophthalmus aquosus</i> )	
29	160	43	24	3	2	3	1024	0	5824		0	704
30	85	2	0	2	0	0	10	100	8640		2	0
31	90	18	0	0	0	0	64	86	10176		0	0
32	170	10	2	3	0	0	0	180	424		0	0
33	230	17	3	10	6	0	12	262	210		2	0
34	210	12	14	60	9	0	0	364	10		0	0
35	215	9	8	20	36	0	8	47	920		0	8
36	190	25	4	15	66	0	58	26	316		0	0
37	150	23	8	8	5	0	22	66	76		3	0
38	240	11	1	3	6	0	0	206	0		2	0
39	290	13	2	1	0	0	0	118	2		0	0
40	210	8	3	6	7	0	0	220	28		2	0
41	150	57	31	11	13	1	0	134	1280		0	8
42	230	5	1	0	0	0	4	186	12		0	0
43	80	10	2	1	1	0	1	105	145		0	0
44	140	58	18	9	4	7	8	132	4032		1	0
45	280	2	0	0	1	0	0	2	132		0	0
46	20	9	0	0	0	0	5	40	111		1	0

\* Not sampled

Appendix 3. Number of larvae sorted and identified from samples collected during the capelin and Atlantic herring larval survey of July 2007 on the west coast of Newfoundland. Scientific names are as follows: mackerel (*Scomber scombrus*), sand lance (*Ammodytes* sp.), herring (*Clupea harengus harengus*), cod (*Gadus morhua*), snailfish (*Liparis* sp.), capelin (*Mallotus villosus*), redfish (*Sebastes* sp.), arctic shanny (*Stichaeus punctatus*), radiated shanny (*Ulvaria subbifurcata*), fourbeard rockling (*Enchelyopus cimbrius*), righteye flounder (Pleuronectidae), cunner (*Tautoglabrus adspersus*), and windowpane flounder (*Scophthalmus aquosus*). Also included are counts of broken and other unidentified larvae.

Station	Mackerel	Sand Lance	Herring	Cod	Snailfish	Capelin	Redfish	Arctic shanny	Radiated shanny	Fourbeard rockling	Righteye flounder	Cunner	Windowpane flounder	Broken larvae	Other larvae not identified
1	7	0	6	5	4	15	5	0	8	6	32	1096	0	0	2
2	6	0	0	6	2	19	6	0	0	6	20	312	0	0	2
3	20	0	1	7	1	10	0	0	0	0	168	600	0	0	0
4	0	0	0	22	14	0	0	0	2	0	52	0	0	0	4
5	4	0	0	8	0	44	1	0	0	0	8	304	0	72	0
6	9	0	0	8	0	13	0	2	4	6	28	54	4	0	2
7	4	0	1	3	0	14	1	0	0	4	36	180	2	0	1
8	8	0	0	2	0	3	0	0	0	0	36	132	0	0	0
9	5	0	0	5	2	1	2	0	0	0	50	0	0	0	0
10	1	0	1	4	2	5	0	0	6	4	24	42	0	0	0
11	4	0	3	8	0	10	0	0	3	4	20	106	0	0	0
12	16	0	1	17	0	0	0	0	0	0	254	24	0	0	0
13	8	0	1	4	4	12	0	0	6	0	42	108	2	0	7
14	0	0	0	18	8	0	0	0	0	0	10	6	0	0	0
15	10	0	1	4	1	3	0	0	8	4	16	70	0	0	0
16	8	0	1	0	0	10	0	0	0	10	54	96	0	0	2
17	2	0	1	1	0	2	1	0	7	0	12	20	0	0	0
18	0	0	0	20	0	0	2	0	0	0	32	0	0	0	0
19	3	0	0	2	2	0	0	0	7	1	17	18	0	0	0
20	0	0	1	0	0	2	0	0	1	21	6	42	1	0	0
21	35	0	3	0	0	16	0	0	4	62	58	1408	4	0	5
22*															0
23	5	3	0	0	0	0	2	0	1	0	4	25	0	0	0
24	0	1	0	0	0	0	0	0	1	84	0	156	0	0	0
25	4	0	0	0	0	0	0	0	0	184	0	600	4	0	0
26	13	0	0	0	0	0	0	0	2	104	6	240	6	0	3
27	46	4	6	4	0	0	0	0	86	0	26	66	0	0	0
28	0	2	6	2	0	0	0	0	14	0	44	22	0	0	0

Station	Mackerel	Sand Lance	Herring	Cod	Snailfish	Capelin	Redfish	Arctic shanny	Radiated shanny	Fourbeard rockling	Righteye flounder	Cunner	Windowpane flounder	Broken larvae	Other larvae not identified
29	0	0	0	0	0	0	0	0	0	16	2	46	0	0	0
30	8	0	1	2	0	3	1	0	36	4	90	32	0	0	0
31	9	0	8	3	0	3	8	0	32	0	34	16	0	0	1
32	38	5	3	0	2	4	14	0	2	2	10	26	0	0	0
33	82	0	3	12	0	9	0	0	6	6	156	58	0	0	1
34	31	0	0	5	2	11	0	0	2	0	70	88	0	0	0
35	200	0	0	4	1	12	0	12	0	14	108	186	2	1	3
36	199	0	5	5	0	20	0	0	2	8	20	140	6	0	1
37	81	0	9	7	0	56	0	0	0	1	82	46	4	0	1
38	39	0	3	5	1	0	2	1	0	0	66	108	0	0	4
39	36	0	0	4	2	5	2	0	2	2	64	62	2	8	0
40	21	0	0	0	0	21	0	0	0	1	60	56	0	0	0
41	123	0	4	2	4	71	0	0	0	2	72	126	0	0	0
42	25	0	7	6	0	1	0	0	4	0	92	0	0	0	2
43	1	0	0	1	7	16	0	0	5	1	16	4	0	0	0
44	13	0	1	1	0	70	1	0	8	0	114	22	0	0	2
45	38	1	0	11	0	17	0	0	0	2	162	16	0	0	0
46	27	0	0	1	0	69	0	0	8	1	19	37	0	0	1

\* Not sampled