

The Canadian Albacore Tuna Catch and Effort Relational Database

M. Stocker, H. Stiff, W. Shaw, and A.W. Argue

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, BC
V9T 6N7

2007

Canadian Technical Report of Fisheries and Aquatic Sciences 2701



Fisheries
and Oceans

Pêches
et Océans

Canada

Canadian Technical Report of Fisheries and Aquatic Sciences

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport technique canadien des sciences halieutiques et aquatiques

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 456 de cette série ont été publiés à titre de rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Technical Report of
Fisheries and Aquatic Sciences 2701

2007

THE CANADIAN ALBACORE TUNA
CATCH AND EFFORT
RELATIONAL DATABASE

by

M. Stocker¹, H. Stiff², W. Shaw³, and A. W. Argue⁴

¹Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
3190 Hammond Bay Road
Nanaimo, B.C. V9T 6N7

²1960 Murray Road
Gabriola, B.C. V0R 1X7

³Fisheries and Oceans Canada
Fisheries Management Branch
South Coast Division
3225 Stephenson Point Road
Nanaimo, B.C. V9T 1K3

⁴Argus Bio-Resources Limited
4050 Lockehaven Dr
Victoria, B.C. V8N 4J5

© Her Majesty the Queen in Right of Canada, 2007
Cat. No. Fs97-6/2701E ISSN 0706-6457

Correct citation for this publication:

Stocker, M., Stiff, H., Shaw, W., and Argue, A.W. 2007. *The Canadian Albacore Tuna Catch and Effort Relational Database*. Can. Tech. Rep. Fish. Aquat. Sci. 2701: vi + 76 p.

TABLE OF CONTENTS

LIST OF FIGURES.....	IV
LIST OF TABLES.....	V
LIST OF APPENDICES.....	V
ABSTRACT	VI
RESUME.....	VI
1. INTRODUCTION	1
2. BACKGROUND.....	2
2.1. DESCRIPTION OF THE FISHERY	2
2.2. FISHERY DATA MANAGEMENT	2
3. DATABASE MANAGEMENT SYSTEM.....	4
3.1. DATA SOURCES	4
3.1.1. <i>Logbooks</i>	5
3.1.2. <i>Sales Slips</i>	5
3.1.3. <i>Hail Data</i>	6
3.1.4. <i>Trans-shipment Data</i>	6
3.1.5. <i>Unreported Vessels Data</i>	7
3.1.6. <i>Supplemental Data</i>	7
3.2. DATABASE DEVELOPMENT	7
3.2.1. <i>Conceptual Data Model</i>	8
3.2.1.1. <i>Catch and Effort Sources</i>	8
3.2.1.2. <i>Calculating Catch and Effort for Unique Vessels</i>	9
3.2.1.3. <i>Reported Catch and Effort Undefined By Area</i>	10
3.2.1.4. <i>Effort and Catch per Effort</i>	10
3.2.1.5. <i>Expanding the Unique Vessels Fishing</i>	10
3.2.1.6. <i>Expanding the Catch and Effort</i>	11
3.2.1.7. <i>Summarizing Vessels, Catch and Effort</i>	12
3.2.2. <i>Physical Data Model</i>	12
3.2.2.1 <i>System Requirements</i>	13
3.2.2.2. <i>File Organization</i>	13
3.2.2.3. <i>Database Data Structures</i>	13
3.2.2.4. <i>Data Relationships</i>	14
3.2.2.5. <i>Codebase Objects</i>	15
3.3. DATA ENTRY, EDITING, VALIDATION AND QUALITY CONTROL	15
3.3.1. <i>Triplogs</i>	15
3.3.2. <i>Saleslips</i>	16
3.3.3. <i>Hails</i>	16
3.3.4. <i>Data Reconciliation</i>	16
4. DATA ANALYSIS.....	17
4.1. <i>Reported Catch and Effort</i>	17
4.2. <i>Expanded Catch and Effort</i>	17
4.3. <i>Reporting Details</i>	18
4.4. <i>Catch and Effort Spatial Analysis for Export to GIS</i>	19
5. DISCUSSION.....	20
5.1. POST-SEASON AND IN-SEASON ANALYSES	20
5.2. COMPARISON OF ESTIMATES	20
5.3. ADDRESSING POTENTIAL SOURCES OF ERROR.....	20

7. LITERATURE CITED	23
8. FIGURES	25
9. TABLES	50
10. APPENDICES	59

LIST OF FIGURES

Figure 1. Albacore tuna catch and effort relational database diagram.....	26
Figure 2. Triplog data entry form.....	27
Figure 3. Saleslip data entry form.....	28
Figure 4. Food and Agricultural Organization (FAO) fishing areas.....	29
Figure 5. DFO Albacore Tuna Fishing Areas within the FAO fishing areas.....	29
Figure 6. Venn diagram demonstrating the intersection of fisher logbook (triplogs) and landed catch (saleslip) data subsets.....	30
Figure 7. Expanded Venn diagram demonstrating the intersection of fisher logbook (triplogs) and landed catch (saleslip), and hail data subsets.....	30
Figure 8. Reported and expanded albacore catch estimates, 1995-2005.....	31
Figure 9. Triplog data analytical flow model: (a) pseudo-code; (b) database objects.....	32
Figure 10. Saleslips analytical flow model: (a) pseudo-code; (b) database objects.....	34
Figure 11. Unique vessels analytical flow model: (a) pseudo-code; (b) database objects.....	36
Figure 12. Hail data analytical flow model: (a) pseudo-code; (b) database objects.....	38
Figure 13. Catch and effort analytical flow model (saleslips-based approach): (a) pseudo-code; (b) database objects.....	40
Figure 14. Catch and effort analytical flow model (triplog-based approach): (a) pseudo-code; (b) database objects.....	44
Figure 15. Analytical flow model for catch, effort, and CPUE by year-vessel-trip-date at latitude and longitude coordinates, for export to geographical information systems (GIS): (a) pseudo-code; (b) database objects.....	46
Figure 16. Reported and expanded albacore catch as a function of percent of vessels reporting.....	48
Figure 17. Saleslip trip catch weight as a function of triplog trip catch weight, 1995-2005.....	49
Figure 18. Comparison of post-season catch estimates based on triplog-based approach ($T+Y$) and saleslip-based approach ($S+X$).....	49

LIST OF TABLES

Table 1. Total reported catch, supplemental catch, and percent supplemental of total catch, 1995-2005.....	50
Table 2. DFO Albacore Tuna Fishing Areas and associated FAO areas.....	50
Table 3. Annual record of unique vessels fishing and percent of vessels submitting triplogs in Canadian albacore tuna fishing fleet.....	51
Table 4. Example catch and effort summary report ($T+Y$ method).....	52
Table 5. Example catch and effort summary report ($S+X$ method).....	53
Table 6. Logbook data form used on albacore tuna fishery vessels, 2006.....	54
Table 7. Saleslip data form used in the albacore tuna fishery.....	55
Table 8. Example catch and effort summary report for the month of July ($T+Y$ method).....	56
Table 9. Correlation and regression analysis of saleslip landings as a function of triplog landings, by year, vessel, and trip.....	57

LIST OF APPENDICES

Appendix 1. Historical Canadian albacore tuna jig-troll catch (tonnes), 1939-1994.....	59
Appendix 2. Canadian albacore tuna jig-troll catch summary reports, 1995-2005.....	60
Appendix 3. Albacore Relational Database – Data Table Field Properties and Data Structure.....	71
Appendix 4. Albacore Relational Database – Lookup Table Field Properties and Data Structure.....	75

ABSTRACT

Stocker, M., Stiff, H., Shaw, W., and Argue, A.W. 2007. *The Canadian Albacore Tuna Catch and Effort Relational Database*. Can. Tech. Rep. Fish. Aquat. Sci. 2701: vi + 76 p.

The *Canadian Albacore Tuna Catch and Effort Relational Database Management System* was developed by Fisheries and Oceans Canada to address the issues of tracking albacore catch and effort data from fishing logbooks and sales slips landings from the Canadian troll fleet operating in the Pacific Ocean. This document provides an overview of the structure and function of the database. The description includes a *conceptual data model*, which defines the logical relationship of fields and tables, and a *physical data model*, which describes the hardware/software implementation of the conceptual model. The description includes an outline of the data compilation, formulation, and summary procedures used to convert raw fishery data into an expanded catch and effort estimate at geospatial coordinates. Two analytical approaches to catch and effort estimation are presented which depend on the relative availability of saleslip and logbook data sources. Limitations and potential sources of error for each method are discussed.

RESUMÉ

Stocker, M., Stiff, H., Shaw, W., and Argue, A.W. 2007. *The Canadian Albacore Tuna Catch and Effort Relational Database*. Can. Tech. Rep. Fish. Aquat. Sci. 2701: vi + 76 p.

Pêches et Océans Canada a mis sur pied le Système canadien de gestion de base de données relationnelles sur les prises et l'effort de pêche du germon (Canadian Albacore Tuna Catch and Effort Relational Database Management System) dans le but de régler les problèmes liés à la traçabilité des données sur les prises et l'effort de pêche du germon provenant des journaux de bord des ligneurs canadiens pêchant dans le Pacifique et des bordereaux d'achat des débarquements. Nous faisons dans le présent document un survol de la structure et de la fonction de la base de données. La description inclut un *modèle conceptuel de données*, qui définit la relation logique des champs et des tableaux, et un *modèle de données réelles*, qui décrit la mise en œuvre logicielle et matérielle du modèle conceptuel. La description inclut également un survol des procédures de compilation, de présentation et de synthèse des données utilisées pour convertir les données brutes sur la pêche en des estimations pondérées des prises et de l'effort selon des coordonnées géospatiales. Nous présentons deux approches analytiques d'estimation des prises et de l'effort, qui reposent sur la disponibilité relative de données de bordereaux d'achat et de journal de bord, et nous établissons les limites et les sources potentielles d'erreur pour chacune.

1. INTRODUCTION

This document describes the catch data management processes for the Canadian albacore troll jig fishery in the Pacific ocean.

Canada has a long history of fishing for North Pacific albacore tuna (*Thunnus alalunga*), one of five species of highly migratory tunas that support some of the most lucrative fisheries in the world (Joseph et al. 1988). The first recorded albacore landings by British Columbia jig-troll vessels were in 1939 (Anonymous 1917-1950). These early statistics were incomplete compared to the catch statistics since 1995, the period for which the database described in this report was constructed.

The mid-1990s was a time of increased international concerns over growing, unregulated fisheries for highly migratory species (HMS) in the Pacific Ocean. Renegotiation of the *Inter-American Tropical Tuna Convention* for waters east of 150°E longitude was underway as was negotiation of a new western Pacific convention for HMS fisheries west of 150°E longitude.

Canada is committed to providing detailed catch and effort statistics, logbook data, and fishing vessel information, as required under various international agreements. The Convention for the Conservation of Highly Migratory Fish Stocks in the Western and Central Pacific was negotiated to conserve and manage highly migratory species in the central and western Pacific Ocean. Canada ratified the Convention on November 1, 2005, becoming the 22nd member of the Western and Central Pacific Fisheries Commission (WCPFC). The most important aspect of the WCPFC for Canada is the management of the northern albacore tuna fishery. The Canadian fishery occurs within the WCPFC and the adjacent Inter-American Tropical Tuna Commission (IATTC) Convention Areas and in the US and Canadian exclusive economic zones (EEZ). The IATTC, established by international convention in 1950, is responsible for the conservation and management of fisheries for tunas and other species taken by tuna-fishing vessels in the eastern Pacific Ocean. Currently Canada is a Cooperating Non-Party in the IATTC.

The *Canadian Albacore Tuna Catch and Effort Relational Database* is the instrument used to comply with the obligations of data provision to these international bodies. In 1998, DFO began construction of the relational database that would be used to compile albacore catch and effort from sales slips, log books, and hail data. The idea was to combine these three data sources in order to account for all Canadian catches from high seas areas, the EEZ off British Columbia, and each of the US Pacific northwest states.

Section 2 of this report provides background information on the history of the Canadian albacore tuna fishery and data collection. In Section 3, we describe the *Canadian Albacore Tuna Catch and Effort Relational Database*, designed to provide annual catch estimates for the Canadian albacore troll fishery based on sales slips, log books, and hail information. The description includes a conceptual data model, which outlines the logical relationship of fields and tables, and a physical data model, which describes the hardware/software implementation of the conceptual model. Section 4 includes an outline of the data compilation, formulation, and summary procedures used to convert raw fishery data into an expanded catch and effort estimate at geospatial coordinates, and summarized by fishing area. Section 5 provides a discussion of the potential sources of

error in data sources and compilation, and the implications for using the database for in-season and post-season analyses.

2. BACKGROUND

2.1. Description of the Fishery

The Canadian fishery for albacore tuna commenced as a coastal fishery in the waters off British Columbia (B.C.). It is now comprised of two fleet types, smaller vessels fishing coastal B.C. and USA waters, and larger vessels fishing on the high seas of the north and south Pacific Ocean. Geographical designations for DFO and associated UN FAO (United Nations Food and Agricultural Organization) fishing areas are listed in Table 1.

The coastal fleet operates within the Canadian EEZ and the USA EEZ in accordance with fishing and port access privileges under the treaty. Vessels in this fleet, mostly 10.67 m to 18.29 m in length (35 to 60 feet), concentrate their fishing effort primarily from the southern Oregon coast to the northern tip of Vancouver Island (Stocker and Shaw 2005). Fishing activity is dependent on price, ocean and weather conditions, albacore availability, the strength of other fisheries, particularly the salmon fishery, and fuel costs (Argue et al. 1999). Effort in the coastal fishery normally peaks in September, after the salmon season for trollers has wound down. Catch from the coastal fleet is sold into both the canned and the blast bled frozen tuna markets (Stocker and Shaw 2005).

The Canadian high seas fleet is comprised of larger troll vessels, mostly greater than 60 feet in length, with crews typically of two to four fishers. These vessels typically remain at sea for trips of several months. Many of these vessels are equipped with larger freezers and operate primarily from west of the International Dateline to the Canadian EEZ in the north Pacific. Some offshore vessels trans-ship their catch to carrier vessels at sea in order to continue fishing operations on migrating schools of tuna. Offshore vessel catches are sold primarily into the blast bled frozen sashimi market. The north Pacific fishery operates primarily in May - October each year when albacore are abundant offshore and in coastal waters. The south Pacific fishery lasts from December - March (Stocker and Shaw 2005).

Two to five vessels operate in the south Pacific, 20 to 30 vessels fish in waters outside the Canadian and USA EEZ to as far west as 170°E in the north Pacific and 130 to 179 vessels troll the waters of the USA EEZ, for a total of up to 230 vessels in the coastal waters. In total, there are about 350 unique Canadian vessels that have participated in the albacore fishery in at least one year since 1995.

Catches since 1996 by the Canadian fleet in the north Pacific albacore troll fishery have ranged from 3,591 tonnes in 1996 to 7,842 tonnes in 2004, with an average catch of 4,403 tonnes. Between 100 and 400 tonnes of this catch are taken in the south Pacific. Canadian caught albacore is worth up to \$28 million per year in landed value.

2.2. Fishery Data Management

Prior to 1951, fish landings were developed from statistics on the amount of products produced by Canadian processing companies (Appendix 1). Fisheries and Oceans Canada (DFO) calculated landings by converting products to landed weights using

industry standard conversion factors (Argue and Shepard 2005). DFO attempted to assign landings to catch areas with varying success; and there were instances of products (and landings) being double counted (A.W. Argue, unpublished data). Canadian caught albacore sold to buyers in United States ports were not captured by the product-based system.

The pre-1951 product and landed weight data were published annually by the Canada Dominion Bureau of Statistics. Between 1939 and 1950 Canadian trollers landed between zero (1942) and 1,012 tonnes (1949) of albacore in Canada (Anonymous 1917-1950; Ware and Yamanaka 1991:Table 1).

In 1951, DFO implemented the multiple sales slips system in order to provide more accurate and timely estimates of catch and effort. Fish buyers were responsible for completing sales slips at the time fish were first sold. They recorded landings at the point of sale in weight and value; the statistical area of catch was noted. Buyers, fishers and/or processing companies were required to return sales slips to DFO personnel who compiled and published catch data by month (week for numbers of salmon) and area in annual reports (Anonymous 1952-1996). This was a definite improvement over the product-based system, but still did not capture albacore landings in US ports (only a handful of US buyers sent copies of their landing records to DFO), nor did it fully capture direct sales of albacore to the public.

Sales slip records of Canadian albacore catch between 1951 and 1994 ranged from zero (1954 and 1955) to 3,921 tonnes (1972) (Ware and Yamanaka 1991; Stocker 2005:Table 1). For 1970 to 1976, these catches included a ten percent upward adjustment of sales slip amounts based on results from a logbook program between 1972 and 1976 (Bourque and Humphreys 1973; Bourque 1974, 1975; Lockner 1977a, 1977b). This was a period during which there was high abundance of albacore off the British Columbia coast as far north as Moresby Island in the Queen Charlotte Islands.

The Canadian catch increased again starting in 1994 (1,998 tonnes), reflecting a shift in trolling effort due to a severe downturn in the salmon troll fishery. By the late-1990s, high prices for blast frozen albacore, the predominant product from the Canadian fleet, and increased restrictions on salmon fishing turned more fishers to albacore fishing. Canadians were then consistently harvesting 2,500 to 4,000 tonnes of albacore per year, much of it from the United States Exclusive Economic Zone (EEZ), as allowed under the *1981 Canada/US Albacore Treaty* between Canada and the US governing reciprocal EEZ fishing and port access for vessels jig-trolling albacore (Shaw and Argue 2000).

There was growing concern that the sales slip system was not accurately capturing the full albacore harvest by Canadian vessels. Starting in the late 1980s, two to five Canadian jig-trollers annually fished south Pacific albacore well below the equator, and by the late 1990s some 40 to 60 Canadian vessels fished albacore on the high seas in the north Pacific as far west as the dateline (Argue et al. 1999; Argue and Shaw 2000). By the late 1990s, the combined coastal and high seas Canadian albacore fleet numbered between 150 and 200 vessels each year, producing a catch worth in excess of \$20 million in landed value. Much of this harvest was landed in continental US ports or in far off ports such as Papeete in French Polynesia, and Pago Pago in American Samoa. Sales slip

records from Canadian buyers by this point substantially underestimated the total Canadian catch.

When the Canadian fishery, and to a lesser degree the US fishery, intensified in the mid-1990s, Canada and the US recognized that improvements were needed to their catch and effort statistics which, under the *1981 Canada/US Albacore Treaty*, are required to be exchanged annually, a practice that had not been followed until the mid-1990s.

The database described herein is a major component of Canada's response to the need for improved annual catch statistics.

3. DATABASE MANAGEMENT SYSTEM

The *Canadian Albacore Tuna Catch and Effort Relational Database* is a database management system (DBMS) developed in Microsoft Access[®] by Fisheries and Oceans Canada (DFO) to relate catch and effort data by geographical area for the albacore tuna fleet from various data sources, 1995-present.

3.1. Data Sources

There are six sources of catch and effort information utilized in the database.

As a condition of licence all albacore tuna fleet vessel masters are required to:

1. Notify Canadian and USA authorities of their fishing activity by hailing out with their intention to start fishing, notifying of changing zones or canceling of trips and hailing in when fishing activity has ceased. The hail data information is used in the albacore database to estimate total vessels fishing.
2. Complete harvest logbooks at sea to be reported in hard copy to DFO. These constitute the triplog or logbook information, which give the best estimate of catch in numbers by geographical location and date.
3. Keep accurate catch records by way of fish slips to be submitted to DFO. Fish slips or saleslips supply the most accurate catch data in terms of total weight by fishing trip.

The other sources of annual catch and effort data required to complete the total annual estimates include:

4. The total annual tonnages of trans-shipments (t) of albacore that have not been identified in logbook or saleslip information, if any. These data are included in the total annual catch.
5. The total annual unreporting vessel counts, which refer to reliable estimates of vessels fishing that were not submitting catch data to DFO. These data, if any, are used in conjunction with reporting vessels to estimate the total vessel effort.
6. Documented vessel-specific annual corrections to fisher catch records that are supplemental to triplog or saleslip records. These data are used in both catch and effort calculations.

The relevant data tables and data fields for these sources are depicted in the Relationships table (Figure 1), and described below.

3.1.1. Logbooks

The principle source of catch and effort information is obtained from vessel trip logbooks (“triplogs”), which provide daily catch and effort at the highest temporal and geo-spatial resolution. All Canadian vessels must carry logbooks while fishing for highly migratory fish species, in any waters. Logbook entries include daily catch and by-catch (pieces), daily effort (hours fished, number of jigs), daily position (latitude/longitude coordinates), average weight of fish caught, and sea surface temperature. Daily estimates of mean weight are derived onboard by fishers using length-weight conversion tables (Clemens 1961).

All logbook data are recorded in the albacore database in three linked tables (Figure 1, Figure 2, Appendix Table 3.1-3.3). Vessel trip metadata (vessel captain, crew size, offload port, etc.) are captured in the *Triplog* table. Originally, fishers were encouraged to record catch data multiple times per day. Thus catch date (in table *CatchData*) was separated from actual catch at location (in table *CatchDataSets*) to permit multiple data records per date, and this structure persists to this day even though fishers are not required to record information at that level of temporal detail.

Key fields used by the database to track logbook catch and effort include:

1. *CFV#* - this refers to the Vessel Registration Number (VRN), which uniquely identifies each vessel.
2. *Trip#* - sequential trip number assigned by vessel master to identify a particular outing, from hail out to hail in and catch landed.
3. *Year* – fishing year (calendar year for northern hemisphere fleet, November to March for southern hemisphere fleet).
4. *Date* – date of fishing activity.
5. *Latitude* and *Latitude Hemisphere* – latitudinal position.
6. *Longitude* and *Longitude Hemisphere* – longitudinal position.
7. *NumberOfFish* – catch (pcs) for that date and geographical position.
8. *AvgWtOfFish* – mean weight (lbs) for the catch for that date and geographical position.

Daily total catch weight may be derived from the product of the number of fish caught and the average weight of fish.

3.1.2. Sales Slips

A secondary source of vessel catch and effort information is available from landings records (“saleslips”, or “fishslips”) from fish processing plants. These data are incorporated into the DFO *Catch Statistics* data management system, and made available post-season to managers, scientists, and the public. Saleslip data provide albacore catch weight estimates by vessel, trip, landing date, and albacore size class. These catch data are considered to be the most accurate estimates of albacore landings, though they do not

fully account for *international sales*, domestic *public sales* or *take-home* totals, and thus underestimate total landings by an unknown factor.

All saleslip data are recorded in the albacore database in two linked tables (Figure 1, Figure 3, Appendix Table 3.4 and 3.5). Vessel trip metadata (vessel captain, saleslip number, landing date, offload location, etc.) are captured in the *FishSlipHeader* table. Multiple landing weights may be recorded per fish slip, according to size, condition of fish, price per pound, etc., so the metadata table is linked in a one-to-many relationship to the *FishSlipData* table, where catch weights are recorded.

Key fields used by the database to track saleslip catch and effort include:

1. *CFV#* - this refers to the Vessel Registration Number (VRN), which uniquely identifies each vessel.
2. *Slip#* - unique saleslip number for landed catch.
3. *Year* and *Date* – landed catch offload date.
4. *AreaOfCatch* – geographical area where most of the fish were caught.
5. *TripNumber* – sequential trip number assigned by vessel master to identify a particular outing, from hail out to hail in and catch landed. This is not included in the saleslip information, but is added to the table later through the triplog/saleslip reconciliation process (see Section 3.3.4).
6. *SpeciesCode* – used to ensure that only albacore data are included in the analysis.
7. *Weight* – landed catch in pounds (lbs).

3.1.3. Hail Data

The hail reporting information includes vessel registration number (VRN), home port, operator/captain, and radio call sign (Stocker and Shaw 2005). Data relevant to the albacore database are stored in table *VesselFishingYears* (Figure 1, Appendix Table 3.6).

Key fields used by the database to track vessel effort include:

1. *VesselID* – uniquely identifies the vessel.
2. *CalledIn* – confirms vessel fishing activity.

3.1.4. Trans-shipment Data

Data in the *Transshipment* table consist of records of total annual albacore weights (t) offloaded en-route to or from the fishing grounds, that are not already included in logbook or saleslip data sources (Figure 1, Appendix Table 3.8). These data may come from various sources (fishers, processors, etc.) but are only included for documented transfers.

Key fields used by the database to incorporate trans-shipment weights include:

1. *Year* – tuna fishing year of trans-shipment.
2. *FAOArea* – geographical area (FAO designation) of catch.
3. *Tonnage* – Total weight in metric tonnes (t) of trans-shipment.

3.1.5. Unreported Vessels Data

The total annual unreporting vessel counts are used in conjunction with reporting vessels to estimate the total vessel effort. Key fields used by the database to incorporate trans-shipment weights include:

1. *Year* – tuna fishing year.
2. *Hemisphere* – north or south hemisphere location of vessels.
3. *UnreportedVessels* – number of vessels.

3.1.6. Supplemental Data

Retrospective corrections to fisher catch records that are supplemental to triplog or saleslip datasets are recorded in the *SupplementalCatch* table (Table 1). Key fields used by the database to incorporate trans-shipment weights include:

1. *Year* – tuna fishing year.
2. *VRN* – vessel registration number.
3. *AddToUSCatch* – weight to be included for US area catch (lbs).
4. *AddToTotalCatch* – weight to be included for total catch (lbs).

3.2. Database Development

The albacore database was developed to provide the best estimate of total annual (monthly summaries also available) catch and effort by vessel and area using disparate, and potentially duplicative, data sources. In addition, source data may be missing, incomplete, or erroneous, and, where required, must be estimated in a reasonable manner.

Post-season analyses of landings may be constrained by the unavailability of saleslip data, which are generally not accessible before March of the following year. Until saleslip data become available, albacore catch and effort estimates must be wholly based on triplog information.

The albacore relational database design rules were developed to generate estimates for catch and effort based on the above contingencies. For example, a vessel may hypothetically submit triplogs representing catch data for areas 1 and 2, while saleslips may provide landings for that vessel from areas 2 and 3 only; the best estimates of catch (weight) will be derived from triplogs for area 1, and saleslips for areas 2 and 3. The best estimates of effort (days fished or boat-days) and area fished (location) will be derived from triplogs for areas 1 and 2, and from saleslips for area 3.

The logical assembly of albacore tuna landings data is described in the *Conceptual Data Model* section of this document. The analytical techniques described in the *Physical Data Model* section are formulated to address the shortcomings in the data for the estimation of catch and effort by area.

3.2.1. Conceptual Data Model

A *conceptual data model* outlines the logical relationship of datasets, fields, and/or tables that resolve the database requirements. The solution to the principle data requirements of the albacore tuna database can be represented diagrammatically in a series of Venn diagrams based on set theory, described in this section.

Standard database query techniques can be used to classify the catch and effort data according to the Venn diagrams described below, based on the intersection of the key variables: *year*, *vessel registration number* (VRN), *trip*, and *fishing area*. The analytical processes and mathematical formulations applied to the data are described in the *Physical Data Model* section.

3.2.1.1. Catch and Effort Sources

Albacore catch estimates are based on trip logbook records and/or saleslip landings, recorded in *pounds* (lb) and converted to *metric tonnes* (t). Size classes, where distinguished on saleslip records, are pooled.

Effort is measured in *days fished*. Any date with a non-zero number of hours of fishing activity by a vessel, recorded in triplogs, constitutes a unit of effort. Non-fishing days (due to travel, inclement weather, mechanical break-down, etc.) are omitted from effort analyses.

Saleslip records distinguish between *length of trip* (days at sea, including travel to and from the fishing grounds) and *days fished*. Where corresponding triplog data is missing, saleslip *days fished* is used to define trip fishing effort.

Catch and effort are organized and summarized by “DFO fishing area”. These fishing areas comprise the EEZs of Canada (BC) and the US (Washington, Oregon and California), and offshore waters inside and outside of the convention¹ area (Figures 4 and 5, Table 2). The data can be further summarized at the FAO² statistical area level.

Triplog catch weights for a given year-vessel-trip are derived from the daily tally of fish multiplied by the mean daily weight of albacore caught. If no mean daily weight is available, the weighted trip mean weight, or (if absent) the weighted annual area mean weight is used, where area is derived from the conversion of the daily triplog record of latitude and longitude coordinates into designated fishing areas. Albacore catch weights are then summed by vessel-trip-area to get total catch and effort by area.

Saleslip catches for a year-vessel-trip are calculated as the sum of albacore weights across size classes as recorded on the processing plant saleslip. Effort is recorded as total days fished on the trip. Since only one fishing area is assigned per tuna fish saleslip, and detailed date-specific area of catch is not recorded, catch and effort may be aggregated from multiple fishing areas on a saleslip record.

The assignment of a single fishing area to a saleslip catch record may occur at the processing plant via inquiry with the vessel owner/operator, by DFO personnel based on interview data at a later date, or by the database administrator during data compilation,

¹ Convention on the Conservation and Management of Highly Migratory Fish Stocks in the Western and Central Pacific Ocean.

² United Nations Food and Agricultural Organization.

based on a tally of days fished in each area, where matching triplogs exist. Clearly, saleslip records by themselves may constitute a large source of error in estimates of catch by fishing area, since the designation of trip fishing location is decided based on a simple majority (i.e., more than 50%) of days fished in an area.

In the case where saleslips and triplogs are matched by year-vessel-trip, however, saleslip total trip weights can be apportioned to individual trip dates and fishing areas based on the distribution of catch (pieces) in the logbook records. This method utilizes the most accurate weight data (from saleslips) in conjunction with the high spatial and temporal resolution of triplog data to provide the best estimate of total albacore catch by date and location.

3.2.1.2. Calculating Catch and Effort for Unique Vessels

Let A represent the total annual landed catch (weight) of albacore tuna from Canadian vessels.

Let T represent the subset of albacore catch weight data, available from triplogs, for a combination of *key variables*, such as *year*, *vessel*, *trip*, and *geographical fishing area*.

Let S represent the subset of similar data available from saleslips.

Let Z symbolize the intersection of T and S , which represents catch data where records exist from both triplog and saleslip sources with the same key variables ($Z = T \cap S$). Then X is the subset of triplog catch data for which no saleslip records exist for the key variables, and Y is the subset of saleslip catch data for which no triplog records exist. Therefore, segments $X + Y$ represent the unique contributions of catch and effort from triplog and saleslips sources, respectively. The intersection of triplog and saleslip datasets may be illustrated in a Venn diagram (Figure 6).

The set of records for total annual catch of albacore from Canadian fishing vessels can be described as:

$$A = X + Y + Z, \text{ where } Z = T \cap S \quad (1)$$

In conceptual terms, circle T grows as triplog data are received at the end of the tuna fishing season. Before saleslip data become available post-season, circle S (and thus segment Z) are non-existent. At this preliminary stage, an estimate of the total albacore catch can be generated using triplog data only, using appropriate expansion factors based on known licensing information concerning overall fleet size.

As saleslip data are incorporated into the database, circle S grows, and segment Z expands. As the overlap of circles S and T increases, $X \rightarrow 0$, $Y \rightarrow 0$, and $Z \rightarrow T \approx S$. In other words, when triplog submission compliance is close to 100% and all saleslip data have been retrieved from DFO Catch Statistics, the *unique* contributions of triplog (X) and saleslip (Y) data go to zero, and there is a high level of redundancy between triplog data and saleslip data ($T \approx S$). This overlap can be used to cross-check and verify the data.

However, due to minor and often irreconcilable differences, the post-verification catch data for Z may not be equivalent between the two sources, nor may it be clear which value is the most accurate for a given set of key variables. For example, triplog daily weight estimates are based on length measurements and may therefore be biased by

length-weight conversion errors, while saleslip trip weight totals may underestimate trip catch where *dock sales* to the public have occurred. Thus, two estimates for total annual catch weight may be derived depending on whether saleslip or triplog weights are selected where matches occur on key variables. Procedures that eliminate the possibility of erroneous duplication of catch and effort are required regardless of which estimation method is employed.

The *triplogs-based* approach provides the preferred estimate when saleslip data are unavailable or significantly incomplete. For this method, trip catch (lbs) for segment *Z* is derived from the sum (across key variables year-vessel-trip) of the product of triplog daily catch (pcs) and average daily fish weight. Matching saleslip catch estimates are ignored. This estimate therefore utilizes triplog data as the best estimator where both triplog and saleslip data exist for a given year-vessel-trip, with the end result that:

$$A = T + Y, \text{ if } Z \text{ is based on triplog data } (T). \quad (2)$$

The *saleslip-based* approach provides the preferred estimate after saleslip data have been incorporated into the database. This method utilizes saleslip landing records as the best estimator of catch weight where both triplog and saleslip data exist for a given vessel-year-trip, such that:

$$A = S + X, \text{ if } Z \text{ is based on saleslip data } (S). \quad (3)$$

3.2.1.3. Reported Catch and Effort Undefined By Area

A special case of reported catch data that is undefined by fishing area arises from unmatched triplogs (*X*) or saleslips (*Y*) for which fishing area cannot be determined. The distribution of catch and effort (vessel fishing days) associated with these records can, however, be assumed to follow the geographic distribution of the overall fleet, and can therefore be distributed to area in proportion to the known catch and vessel distribution, respectively. Catch and effort from *undefined areas* can then be added to the catch and effort from *defined areas* to generate *total reported catch by area*.

3.2.1.4. Effort and Catch per Effort

Once the unique catch (lbs) and effort (days fished) information is obtained based on reported catch records, using either the *triplog-* or *saleslip-based* approach above, *effort* and *catch per unit effort* indices can be derived at the individual vessel, fishing area, and fleet-wide annual levels.

$$\text{Days per Vessel} = \text{Total Fishing Days} / \text{Total Vessels} \quad (4)$$

$$\text{Catch per Vessel} = \text{Total Catch} / \text{Total Vessels} \quad (5)$$

$$\text{Catch per Day} = \text{Total Catch} / \text{Total Days} \quad (6)$$

These indices can be used to adjust the reported catch to account for non-reporting vessels, as described in the *Expanding the Catch and Effort* section below.

3.2.1.5. Expanding the Unique Vessels Fishing

The conceptual model above relies on information at the year-vessel-area level, and thus must be expanded to incorporate the occurrence of vessels fishing in unknown areas. The

source of such unknowns may be either vessels fishing, as indicated by hail data, for which triplogs or landings records are not available, or “ghost” vessels (observed but unidentified vessels that cannot be accounted for by hail data, vessel registration, etc.) from interview reports. To accommodate these possibilities, Circle H is added to the Venn diagram to represent the unique set of all Canadian vessels fishing: hailing, reporting, non-reporting, and unidentified “ghost” vessels.

The intersection of triplog, saleslip, and hail datasets may be illustrated in a Venn diagram (Figure 7).

Because most of the vessels operating in the albacore tuna fishery also submit triplogs (T) or saleslips (S), the only component of circle H that is of interest is the small proportion of vessels (less than 10% in recent years, see *Other Vessels* in Table 3) that did not submit either triplogs or saleslips (segment Q).

The difficulty with Q is that, because these vessels have not submitted catch information, it is not possible to know exactly where the Q vessels fished, or how many fish they caught. However, it may be assumed that the catch and effort of these vessels follow the geographic distribution of the catch and effort of the overall fleet. Thus, the annual Q vessels can be distributed across fishing areas in proportion to the distribution of reported fishing vessels, and added to the reported vessels in each area to yield an expanded estimate of vessels by area (equation 7).

Let A_V represent the total number of vessels in the Canadian tuna fleet, and let Q_V represent the total annual estimate of non-reporting vessels. Let A_{Va} represent the total number of *unique* Canadian tuna vessels fishing in area a , based on triplog (X_{Va}), saleslip (S_{Va}), and non-reporting (Q_{Va} , i.e., hail-ins, “ghost” vessels, etc.) information sources.

The proportion of reporting vessels fishing in area a can be derived from:

$$P_{Va} = (S_{Va} + X_{Va}) / (A_V - Q_V) \quad (7)$$

The number of non-reporting vessels fishing in area a can then be calculated as:

$$Q_{Va} = P_{Va} * Q_V \quad (8)$$

The expanded number of vessels fishing in area a can then be expressed as:

$$A_{Va} = S_{Va} + X_{Va} + Q_{Va} \quad (9)$$

Note that, because all the vessels reporting via either triplog or saleslip sources can be uniquely identified, A_{Va} is also equivalent to the sum of T_{Va} , Y_{Va} , and Q_{Va} , since $S_{Va} + X_{Va} = T_{Va} + Y_{Va}$.

3.2.1.6. Expanding the Catch and Effort

Once the expanded number of vessels fishing in an area is calculated, reported catch and effort indices such as the *days per vessel* and *catch per day* statistics derived in the *Effort and Catch Per Effort* section above can be used to expand the area catch to provide the best estimate of total albacore catch.

Let D_{Va} represent the reported *days per vessel* in area a . Let C_{Da} represent the reported *catch per day* in area a . Let A_{Va} represent the expanded number of vessels in area a (equation 9), and let A_{Ca} represent the expanded albacore catch in area a to be estimated.

To account for the non-reporting fishing effort in area a , the expanded number of days fished D_{Vae} must first be computed as the product of the expanded vessels A_{Va} and the reported days per vessel D_{Va} :

$$D_{Vae} = A_{Va} * D_{Va} \quad (10)$$

The expanded albacore catch A_{Cae} for area a can then be estimated from the expanded days D_{Vae} and the reported catch per day C_{Da} in that area:

$$A_{Cae} = D_{Vae} * C_{Da} \quad (11)$$

Substituting, the expanded catch estimate can be expressed as the product of the expanded vessel count x days/vessel x catch/day for area a :

$$A_{Cae} = A_{Va} * D_{Va} * C_{Da} \quad (12)$$

3.2.1.7. Summarizing Vessels, Catch and Effort

Catch (lbs) and *effort* (fishing days), whether reported or expanded, are additive across DFO-area, FAO-area, and hemispheric geographic levels (Tables 4-5, D, E, K, L).

Miscellaneous catch estimates from other sources that are aggregated at the FAO level, such as *trans-shipment* data and *supplemental catch* data, can be combined with the expanded catch to generate a grand total annual albacore catch estimate (Tables 4-5, N).

Let A_{Ce} represent the annual grand total (expanded) albacore catch (t). It can be derived from the sum of albacore catch across DFO (or FAO) areas (A_{Cae}), plus the sum of FAO area trans-shipsments (T_a) and/or supplemental catch data (S_a):

$$A_{Ce} = \Sigma (A_{Va} * D_{Va} * C_{Da}) + \Sigma (T_a) + \Sigma (S_a) \quad (13)$$

Let A_{De} represent the annual grand total (expanded) albacore fishery effort (days fished). It can be derived from the sum of boat-days (days fished) across DFO (or FAO) areas (D_{Vae}):

$$A_{De} = \Sigma (A_{Va} * D_{Va}) \quad (14)$$

Unlike catch and effort indices, the number of *unique vessels fishing* can only be determined at particular aggregate levels, as they are not additive across aggregate levels. Since a vessel cannot fish in two places at one time, but may fish in two places at different times, the derivation of unique vessels fishing depends on the specific spatial and temporal scales selected. For example, the entire Canadian tuna fleet comprises the set of unique vessels fishing in the Pacific Ocean in a given fishing season. However, over the course of the season, a vessel may fish in multiple FAO areas, contributing to the unique vessels fishing in each of those FAO areas. It would, however, be incorrect to add the unique vessel contributions from different FAO areas together, because they would no longer be unique at the larger aggregate level, and would therefore overestimate fleet size.

3.2.2. Physical Data Model

The physical data model identifies the DBMS file structure that houses the database and codebase elements, and describes the hardware and software requirements to operate the

DBMS. It also includes a definition of the data structure (data tables, fields, relationships) and an outline of the functional and analytical interface developed to allow users to analyze and interpret the albacore catch and effort data.

3.2.2.1 System Requirements

The *Canadian Albacore Tuna Catch and Effort Relational DBMS* was developed using Microsoft Access[®] software as a stand-alone application on Windows[®] (IBM[®] PC-compatible) computers within the DFO network. Hardware requirements include a colour monitor with a minimum resolution of 1028 x 760 pixels, and 30 Mb of free disk space.

Software requirements include MS-Access[®] 97 (SR-1) for Windows 98[®], or later.

3.2.2.2. File Organization

The *Canadian Albacore Tuna Catch and Effort Relational Database Management System* comprises two MS-Access[®] database components:

- Albacore.mdb – the *codebase*, containing source code, queries, reports, lookup tables, and forms comprising the functional and analytical components of the user interface with the database. This file is often referred to as the “front-end”.
- AlbacoreData.mdb – the *database* proper, which contains the database tables for triplog, saleslip, hail and vessel data, and is often referred to as the “back-end” file.

The “back-end” database tables are programmatically linked to the “front-end” codebase upon execution, such that the data in the “back-end” database are available to forms, reports, procedures, and queries in the codebase.

3.2.2.3. Database Data Structures

The “back-end” database objects consist of data tables only, and comprise the *triplog* tables, the *saleslip* tables, and the *hail* data table, all linked together via the *vessel information* table. See *Data Relationships* below for more information on logical table relations.

The *vessel information* table contains vessel-specific information (vessel name, size, owner, etc.) uniquely identified by vessel registration number (VRN), for all tuna fleet vessels registered with DFO. Vessel information is updated on an annual basis from DFO Licensing.

Triplog data are recorded by albacore tuna fishers on a daily basis during fishing trips in official *Albacore Tuna Log Books*, published by the Canadian Highly Migratory Species Foundation³, and purchased by fishers as part of the license agreement. Data recorded on a daily basis include date, total hours fished, latitude and longitude, number of jigs used, water temperature, species, number of fish, and average weight of fish caught. By-catch species and numbers are also recorded. Trip specific information recorded on the logbook sheets include vessel name and registration number, trip number, captain, crew size, gear, days fished, offload date, offload port, offload weight and pieces, and buyer (Table 6).

³ Canadian Highly Migratory Species Foundation (CHMSF), 4829 Maplegrove Street, Victoria, B.C. Canada V8Y 3B9.

The *triplog* component of the database is comprised of three tables:

1. *Triplog* header table, housing trip-specific details (gear, start date, offload date, crew size, etc.) for unique combinations of year, vessel and trip number;
2. *CatchData* table, housing date-specific details (hours fished);
3. *CatchDataSets* table, housing catch, effort, location, and by-catch data for each catch event within a date.

Saleslip data are obtained from buyers by DFO for data entry of saleslip number, offload date, vessel name and registration number, price per pound and total weight by species, size class, and/or quality/condition code. Other meta-data recorded are skipper, area fished (if available), length of trip (days) and number of days fished, and buyer (Table 7).

The *sales slip* component of the database is comprised of two tables:

1. *FishslipHeader* table, housing catch landings details (e.g., date, gear, sales slip number, buyer, etc.) for unique combinations of year, vessel and trip number;
2. *FishslipData* table, housing species-specific details categorized by size-class (e.g., weight (lbs), condition).

The *hail data* table stores the vessel identification (Vessel ID) and date of hail for each vessel catalogued by the Coast Guard authority as actively fishing, on an annual basis.

A *supplemental catch* table was incorporated into the database to address discrepancies in DFO albacore catch data with respect to vessel owner's documented catch for the years 1995-2002. These supplemental catch data are based on substantiated submissions made by vessel owner/operators for the purposes of *qualification ranking* for the *limited vessel entry program*⁴ initiated in 2005. The percent supplemental catch of the total annual catch is listed in Table 1.

3.2.2.4. Data Relationships

The primary data tables of the albacore database are related to each other via the *Vessel Information* table (Figure 1).

The data tables are *normalized*⁵ to maximize data storage efficiency, maintain data integrity, and reduce data redundancy. For example, the saleslip data are housed in two tables: *FishSlipHeader*, containing the meta-data for each saleslip (vessel VRN, landing date, saleslip number, etc.), which is linked in a one-to-many relationship with table *FishSlipData*, which contains the catch (weight) by species data. Similarly, table *Triplog* contains vessel logbook meta-data for each tuna fishing trip (e.g., vessel VRN, trip

⁴ The *limited vessel entry* program was implemented in accordance with international agreements to reduce annual albacore tuna fishing effort in U.S. waters beginning in the 2005 season. Vessels eligible for fishing in U.S. waters were determined from a weighted rank analysis of annual albacore landings from U.S. waters from 1995-2002.

⁵ Database *normalization* refers to the process of designing the database to minimize redundancy while maximizing the flexibility required to provide the necessary management reports and to support *ad hoc* requests which have not been pre-defined. Where practical, the database is normalized to the 3rd *normal form*, meaning that tables are organized such that: [1] redundant data and calculated fields are eliminated; [2] secondary attributes are uniquely identified by the full primary key attributes; and [3] secondary attributes that depend on other secondary attributes are eliminated.

number, gear type, etc.), which is linked to daily catch event meta-data (date, hours fished) in table *CatchData*, which is itself linked to catch event data (location, number of fish caught, average weight, etc.) in table *CatchDataSets*.

These data structures are further detailed (e.g., field names, data types, primary keys, validation rules, etc.) in Appendix 3.

3.2.2.5. Codebase Objects

The “front-end” codebase objects consist of the queries, forms, reports, “look-up” tables and visual basic source code comprising the “user-interface” that enables the user to view, analyze, and interpret the data. The codebase is where all data analysis occurs. The user-interface is loaded automatically upon start-up of the codebase, and consists of a main form that provides links to sub-forms for the purposes of data entry, error-checking, data analysis, and report generation. Import and export utilities provide a means of communicating with external data sources.

Forms and reports are primarily based on queries, which ultimately reference the data in the linked database tables and look-up tables. The sequential flow that links tables and queries in an analytical sequence to arrive at the annual reported and expanded catch and effort estimates is described in the *Codebase Analysis* section below.

Look-up tables reside in the codebase and pertain to reference information, such as species codes, gear codes, and area codes, that do not change on a frequent basis. Look-up tables and their contents are listed in Appendix 4.

For more information on the design and use of the front-end codebase, see the *Albacore Tuna Catch and Effort Database User’s Guide* (Stiff in prep.).

3.3. Data Entry, Editing, Validation and Quality Control

Data entry, modification, viewing, and verification are performed through forms built into the user-interface. Validation rules are programmed into data entry fields to eliminate gross errors in data entry. Drop-down “combo-box” fields are employed in data selection fields to minimize typing and reduce errors. FAO and DFO-areas are applied automatically based on latitude and longitude data entry. Calculated fields and summary statistics for numbers of fish and catch weight are displayed in the forms where appropriate to eliminate unnecessary data entry and to provide verification of data entered.

Verification and validation procedures involve summary reports of descriptive statistics, distribution plots for size and location data for outlier analyses, and reconciliation procedures between saleslip, triplog, and hail data records.

3.3.1. Triplogs

Triplog data are processed in the *Triplog* data form, which allows data entry and updating of trip information for all three triplog data tables on one screen, including trip meta-data (vessel registration number, trip number, offload date, etc.), daily catch meta-data (date, hours fished), and catch event data (number of fish, average weight, etc.). See Figure 2 and Table 6 for more logbook data entry details.

3.3.2. Saleslips

The majority of saleslips for Canadian tuna fishery operations are submitted to DFO by seafood processors on an annual basis, under agreements between the companies and DFO. These data are keypunched (and verified) by DFO in-season into the *Fishslips* data form, which provides data entry capabilities for slip information for both saleslip meta-data (vessel registration number, trip number, landing date, etc.), and species data (size class, total landed weight, etc.). See Figure 3 and Table 7 for more saleslip data entry details. Further verification and error analysis for saleslip data are performed during the process of data reconciliation with triplog data (see Section 3.3.4).

3.3.3. Hails

Hail data are retrieved from DFO Coast Guard logs which are used to document the involvement of Canadian fishing vessels in the albacore tuna fishery, including the movement between Canadian and U.S. fishing zones. The hail data are retrieved in spreadsheet format, which is imported into the Albacore Catch and Effort Relational Database, where the data are sorted and filtered for unique annual vessel entries.

3.3.4. Data Reconciliation

The post-season reconciliation process serves to match saleslip and triplog records by year, vessel, and trip. This is required because saleslip records may not be identified by trip number. Discrepancies in fishing location and catch weights must also be resolved.

A program utility displays all triplog and saleslip data available for a given vessel. By comparing triplog start and end dates with saleslip record landing dates the user can assign trip number to saleslip records. In the case where saleslip records are missing trip information, but no corresponding trip exists in the triplog data, an arbitrary trip number (e.g., 99) must be assigned to distinguish between the saleslip trip records.

It is important to also examine daily fishing area information from the triplogs to ensure that the matching saleslip record is assigned an appropriate fishing area (generally the fishing area with the majority of dates fished in the triplog data) to ensure that the match-merge process on vessel-trip-area is enabled.

Discrepancies in triplog and saleslip catch weights that are not attributable to errors in triplog daily average fish weight or triplog catch (pcs) (verified against the hardcopy logbook data), may be resolved by omitting the saleslip record from analysis, in cases where the saleslip landed weight is obviously underestimating catch. This may be the case where *dock sales* or *take-home* catches are a significant factor, and becomes apparent when the two weight estimates are divided by the logbook verified catch in pieces. Average saleslip fish weights of less than 10 lbs are a good indicator that the saleslip data are not fully accounting for the catch provided there are no obvious errors in daily triplog catch numbers, and the saleslip record should be omitted from analysis.

4. DATA ANALYSIS

As indicated above, the objective of the DBMS is to provide an annual⁶ estimate of the reported catch and effort by FAO and DFO geographical areas, and to expand upon those estimates where necessary to accommodate missing elements of the Canadian albacore tuna fishery. By way of example, the compilation of catch and effort for 1998 is reported in Table 4 for the *triplog-based* ($T + Y$) analysis, and Table 5 for the *saleslip-based* ($S + X$) analysis.

4.1. Reported Catch and Effort

The general sequence of analysis begins with a synthesis of annual reported triplog and saleslip data to identify unique contributions to catch and effort from those sources (Tables 4-5 A). Where both triplog and saleslip information exist for key variables, precedence is given to the triplog information for preliminary analyses (Table 4) and to the saleslip information for the final analysis (Table 5). The total reported catch, days fished, and unique number of vessels fishing are designated by hemisphere and area (Tables 4-5 B, C). Note that the number of unique vessels by area (in B & C) do not sum up to the total number of unique vessels fishing (A), because a vessel may fish in more than one area within a fishing season. However, both the catch and days fished by area are additive and equal to the total reported catch weight and days fished.

The reported catch may be segregated into geographically-defined areas (D), and undefined areas (E) for which geographical details are not available for the catch and effort information. The undefined area catches, the number of vessels fishing, and the number of days fished are geographically apportioned, assuming the distribution of catch and effort for undefined areas will follow the distributions for defined area catches.

The area-specific fishing days per boat is calculated from the total reported days fishing divided by the number of boats in the area (F). From this, the principle indicator of effort, catch per day (G), is then based on the total area-specific reported catch divided by the area-specific fishing days per boat.

4.2. Expanded Catch and Effort

If unreported fishing effort is known to exist, then the total catch estimates can be extrapolated based on the estimated number of the unreported vessels times the reported days per vessel and catch per day indices. Counts for vessels hailing without submitting triplog or saleslip data (Tables 4-5, H), and “ghost” vessel estimates obtained from interview data (I), are allocated to fishing areas based on the distribution of reporting vessels for defined areas (D). The total “expanded” vessels (J) by area is then calculated as the sum of the reported vessels plus unaccounted “hail” vessels (H) plus “ghost” vessels (I). Multiplying the expanded vessel count (J) by the days per vessel yields expanded days fished (K). This in turn is combined with catch per day to produce expanded catch weight (L).

The expanded values for days fished and catch can be summed across DFO areas to provide total estimates for FAO areas, which, when converted to tonnes, can be added to

⁶ Monthly catch and effort analyses are also available.

trans-shipment totals (M) to estimate total expanded catch by FAO area. Total catch across FAO areas and hemispheres (north and south) yields the total expanded catch in tonnes (Tables 4-5, N). Some rounding error may be evident in the grand total.

Annual expanded catch estimate summary reports (1995-2005) can be found in Appendix 2. Estimates of expanded albacore tuna catch by Canadian fishers (Figure 8) show an increasing trend for the years 1995-2004, followed by a drop in catch in response to the introduction of the limited vessel entry program in 2005.

4.3. Reporting Details

The analytical flow that links tables and queries for the estimation of *annual reported and expanded catch and effort estimates* (Tables 4-5) can be compartmentalized as follows:

- a) Calculation of total reported catch (pcs, lbs) and effort (jigs, hours, days fished) by vessel-area from triplog data (circle *T*). Albacore catch weight is calculated by vessel, trip, and date by multiplying the daily total number of fish by the average weight of fish for that vessel, trip, and date. If the average weight is missing, then the weighted mean weight of albacore for that trip is applied, or (if absent), the weighted mean weight of albacore caught in that area is applied. Although for the purposes of this analysis, fishing effort is based on vessel-days, other effort indices such as hours fished, number of jigs, and jig-hours, are also calculated. Total catch and effort indices are compiled by vessel and area in an intermediate table for input to other components of the analysis. See Figures 9a-b for a flowchart documenting triplog data compilation and query execution.
- b) Calculation of total reported catch (lbs) and effort (days fished) by vessel-area from saleslip data (circle *S*) (Figures 10a-b). Saleslip data are filtered for albacore species, and the catch weight is summed by trip across size class and fish condition codes. For effort calculations, the number of days fished is used, or (if absent), an estimate of days fished based on length of trip and the ratio of trip length to days fished for the vessel, or (if absent) the fleet, is used. Triplog data, if available for this vessel and area, are merged in for calculation of other catch effort estimates, such as catch per jig hour – however these CPUE indices are currently not employed in any further calculations. Total catch and effort indices are compiled by vessel and area in an intermediate table for input to other components of the analysis.
- c) Calculation of unique vessels fishing by area based on triplog and saleslip sources (Figures 11a-b). Compare and contrast unique vessel list from triplog and saleslip sources, and compile total number of unique vessels by hemisphere for input to other components of the analysis.
- d) Calculation of unique unreported vessels based on intersection of hail dataset in circle *H*, triplog dataset circle *T*, and saleslip dataset circle *S* (Figures 12a-b). Get list of vessels that have hailed but not submitted catch data (segment *Q*). Store count of *Q* vessels in an intermediate table for input to other components of the analysis.
- e) Calculation of expanded catch (t) and effort (days fished) based on *saleslip-based* (*S + X*) analysis (circle *S*, segment *X*, and segment *Q*, by area and hemisphere

- (Figures 13a-b)). Combine unique year-vessel-trip catch and effort summaries from triplog and saleslip sources (a and b, above), such that total reported catch by vessel for defined areas is based on saleslip catch (circle *S*) plus unmatched triplog catch from segment *X*. Calculate proportion of unique vessels and reported catch by DFO area for defined areas. Use proportions to distribute undefined area catch and days fished data. Calculate area-specific days-per-vessel and catch-per-day statistics based on reported data. Use vessel distribution for defined areas to distribute unreported vessels (hail-ins and “ghost” vessels) to DFO area. Sum reported and unreported vessels to get “expanded” total vessels by area. Use expanded vessels with reported days-per-vessel and catch-per-day to generate expanded catch by DFO area. Summarize by FAO area, convert to tonnes, and incorporate supplemental and trans-shipment tonnage (if any) for estimate of total annual catch (expanded).
- f) Calculation of expanded catch (*t*) and effort (days fished) based on *triplog-based* (*T* + *Y*) analysis (circle *T*, segment *Y*, and segment *Q*, by area and hemisphere (Figures 14a-b)). Combine unique vessel-trip-area catch and effort summaries from triplog and saleslip sources (a and b, above), such that total reported catch by vessel for defined areas is based on triplog catch (circle *T*) plus saleslip catch from segment *Y*. Calculate proportion of unique vessels and reported catch by DFO area for defined areas. Use proportions to distribute undefined area catch and days fished data. Calculate area-specific days-per-vessel and catch-per-day statistics based on reported data. Use vessel distribution for defined areas to distribute unreported vessels (hail-ins and “ghost” vessels) to DFO area. Sum reported and unreported vessels to get “expanded” total vessels by area. Use expanded vessels with reported days-per-vessel and catch-per-day to generate expanded catch by DFO area. Summarize by FAO area, convert to tonnes, and incorporate supplemental and trans-shipment tonnage (if any) for estimate of total annual catch (expanded).
- g) The *triplog-based* (*T* + *Y*) analysis (f, above) applies also to monthly summaries. See Table 8 for a monthly summary report.

4.4. Catch and Effort Spatial Analysis for Export to GIS

Albacore catch and effort data from triplog sources can be summarized for export to Geographic Information System (GIS) software for further analysis and plotting. Triplog catch is summarized by year, vessel, trip, and date at latitude/longitude coordinates adjusted to identify lat-long cell midpoints. The adjustment involves converting the standard lat-longs to decimal lat-longs by removing minutes and seconds from the coordinates (integerizing) and adding 0.5. Decimal longitudes greater than 180 are adjusted down to 179.5. The individual year-vessel-trip-date catch and effort records are merged with summaries of annual total catch, effort (vessels), and catch per vessel estimates. Grouping and label fields necessary are set up for GIS program purposes. The resulting query can be exported to Excel for upload into the GIS program, or summarized at the month and lat-long level for export to U.S. tuna fisheries authorities. See Figures 15a-b for a flowchart documenting lat-long data compilation and query execution.

5. DISCUSSION

5.1. Post-season and In-season Analyses

Although this database management system was originally developed as a post-season summary tool for the albacore fishery, it may also be used in-season based on its ability to forecast fleet catch based on (a) *reported catch* for a subset of the vessels, and (b) *expansion factors* associated with known fleet size and reported catch per unit effort. Assuming representative reporting of *days fished per vessel trip* and *catch per day* in the early months of the season, the program can forecast reasonable total catch estimates (expanded) based on as little as 30-40% of the vessels reporting (Figure 16). Accuracy improves as the proportion of *vessels reporting* asymptotically approaches 100%.

However, since saleslip data are not available until post-season, the in-season estimates must rely almost entirely on triplog-based information ($T+Y$ method), which raises the question of the reliability of triplog catch data vs. saleslip records. Effort estimates (*unique vessels fishing*, and *days fished* by area) are the same for each estimation method, however catch weight estimates may vary between the two approaches. To examine this issue, regression analyses were performed treating saleslip and triplog catch weight data matched by year, vessel, and trip as independent estimates of catch weight.

5.2. Comparison of Estimates

Catch estimates by year, vessel, and trip are highly correlated between saleslip and triplogs (Table 9, Figure 17). Correlations (r^2) range from 0.99 in recent years (1998-2005) to a low of 0.95 in 1995-1996.

Triplog data *overestimated* saleslip catch weights by 1% on average. Significant overestimates occurred in 1997 (15%), 1999-2000 (5%), and 2001-2003 (1-2%). Triplog data significantly *underestimated* saleslip catch weights in 1995 (11%) and 2005 (6%).

It is no surprise that post-season (final) estimates from triplog-based ($T+Y$) and saleslip-based ($S+X$) are highly correlated (Figure 18); the data are not independent at that stage in the season. However the relationship does suggest that triplog data overestimate saleslip catch weights by a mere 1% on average, which indicates that triplog data are a reasonable substitute for saleslip-based estimates of albacore landing weights.

5.3. Addressing Potential Sources of Error

Some of the potential sources of error mentioned above may be remedied simply via raised awareness in vessel operators concerning data quality, or via management directives aimed at securing more detailed information from fish processing facilities.

For example, saleslip records are considered the most accurate estimate of landed fish weight for a trip. However, these pooled trip totals cannot be readily assigned to geographical area or date. Fishers could be encouraged to include saleslip data when submitting trip logbook information to DFO. Saleslip data could be annotated with trip information (trip number) to facilitate post-season data reconciliation processes.

Limitations in the triplog dataset pertain to logbook availability. Prior to 2003, triplog data submission to DFO was voluntary, ranging from 13% of vessels fishing in 1994, to over 90% in recent years (Table 3). Since 2003, albacore tuna fishers have been required

to submit a copy of their triplogs to DFO at the end of the fishing season (November 30th) as part of the fishing license agreement. In recent years, compliance has been above 95%.

A source of bias in saleslip weight estimates involves *public sales* or *take-home* totals, which are neither captured on saleslips nor uniquely identified in triplog data. This source of bias is potentially significant because the bias always occurs in one direction, resulting in an underestimate of total landings when catch estimates are based principally on saleslip data ($S+X$ method). Furthermore, when matched with total pieces caught from triplogs saleslip weights, – biased low – will overestimate average fish size. This data issue could be reconciled by educating fishers on the importance of these data to the management of the albacore tuna fishery.

Errors in allocating saleslip catch to fishing area may also be incurred due to the difficulty of partitioning aggregated saleslip data for catch from multiple geographical areas. For this reason, geographical area of catch is specified from the logbook information for the corresponding trip wherever reconciliation of the two sources of data is possible. In lieu of logbook data, catch area for a saleslip data will be assigned based on relevant information on the saleslip.

6. ACKNOWLEDGEMENTS

Thanks to Lisa Mijacika and Cory Paterson for information on the management of albacore tuna fisheries in recent years, and to Laili Amar, Meghan Bowes, and Sheila Malcolmson for astute data entry and triplog and saleslip data reconciliation efforts.

Lorne Clayton of the *Canadian Highly Migratory Species Foundation* (CHMSF) managed the distribution and retrieval process for tuna logbooks in recent years. Sam Boehner (Archipelago Marine Research, Inc.) assisted with triplog data collection. Thanks also go to the Canadian fishing vessel operators for submitting logbook data on a timely basis; to the Canadian and U.S. fish processors for the timely provision of saleslip records; and to the DFO catch statistics group for assisting in saleslip data recovery and reconciliation.

We are grateful to Jon Schnute and Jake Schweigert from DFO for reviewing the manuscript. Their constructive comments on the analysis and content have greatly aided the comprehensibility of this report.

7. LITERATURE CITED

- Anonymous. 1917-1950. *Annual Reports of the Fisheries Statistics of Canada*. Government of Canada, Dominion Bureau of Statistics.
- Anonymous. 1952-1996. *British Columbia catch statistics: by species, gear, month and area*. Canada Department of Fisheries and Oceans.
- Argue, A.W., and Shepard, M.P. 2005. *Historical commercial catch statistics for Pacific salmon (*Oncorhynchus spp.*) in British Columbia, 1828 to 1950*. Can. Tech. Rep. Fish. Aquat. Sci. 2601: 595 pp.
- Argue, A.W., and Shaw, W. 2000. *An update for Canadian tuna fisheries in the north and south Pacific Ocean*. 17 pp. Document submitted by DFO to the Thirteenth Meeting of the Standing Committee on Tuna and Billfish in Noumea, New Caledonia, July 5-12, 2000.
- Argue, A.W., Shaw, W., and Williscroft, N. 1999. *An update for Canadian albacore fisheries in the north and south Pacific Ocean*. 8 pp. Document submitted by DFO to the Twelfth Meeting of the Standing Committee on Tuna and Billfish in Tahiti, French Polynesia, June 16-23, 1999.
- Bourque, S.C. 1974. *Status of the 1973 British Columbia albacore fishery*. Environment Canada, Fisheries and Marine Service, Southern Operations Branch Pacific Region. Technical Report Series No. PAC/T-74-3. 61 pp.
- Bourque, S.C. 1975. *Status of the 1974 British Columbia albacore fishery*. Environment Canada, Fisheries and Marine Service, Southern Operations Branch Pacific Region. Technical Report Series No. PAC/T-75-10. 75 pp.
- Bourque, S.C. and R.D. Humphreys. 1973. *Status of the 1972 British Columbia albacore fishery*. Environment Canada, Fisheries and Marine Service, Pacific Region. Technical Report 1973-3. 62 pp.
- Clemens, H.B. 1961. *The migration, age, and growth of Pacific albacore (*Thunnus germo*)*, 1951-1958. Fish Bull. Calif. Dept. Fish and Game 115: 128 pp.
- Joseph, J, Klawe, W., and Murphy, P. 1988. *Tuna and billfish – fish without a country*. Inter-American Tropical Tuna Commission, La Jolla CA. 69 pp.
- Lockner, B.J. 1977a. *Status of the 1976 British Columbia albacore fishery*. Fisheries and Environment Canada, Field Services Directorate Pacific Region. Technical Report Series No. PAC/T-77-9. 30 pp.
- Lockner, B.J. 1977b. *Status of the 1975 British Columbia albacore fishery*. Environment Canada, Fisheries and Marine Service, Southern Operations Branch Pacific Region. Technical Report Series No. PAC/T-77-1. 26 pp.
- Shaw, W., and Argue, A.W. 2000. *The 1999 Canadian north Pacific albacore troll fishery*. Document submitted by DFO to the Seventeenth Meeting of the North Pacific Albacore Workshop, Taipei, Taiwan, December 6-13, 2000. NPALB/00/04:17 pp.
- Stiff, H. (in prep). *Albacore Tuna Catch and Effort Database User's Guide*.

- Stocker, M. (Ed). 2005. *Report of the Nineteenth North Pacific Albacore Workshop*. Nanaimo, B.C., Canada, November 25-December 2, 2004. Fisheries and Oceans Canada, Pacific Biological Station. 127 pp.
- Stocker, M., and Shaw, B. 2005. *Canadian albacore tuna fisheries in the north and south Pacific Ocean in 2004*. WCPFC-SC1 FR WP-4, Noumea, New Caledonia. 6 pp.
- Ware, D.M., and Yamanaka, K.L. 1991. *Catch statistics for the Canadian albacore tuna fishery: 1945-1990*. 4 pp. Document submitted by DFO to the Annual Meeting of the International North Pacific Fisheries Commission, Tokyo, Japan, November 1991.

8. FIGURES




Figure 1. Albacore tuna catch and effort relational database diagram.

ALBACORE LOG BOOK

Albacore Tuna Trip Logs

Vessel Name: **Wave-Length Rider** CFV #: **12345** Start Date: **06-Aug-98**

Start Date: **06-Aug-98** Year: **1998** Captain's Name: **L. M. Jiggin** Trip #: **2**

Crew Size: **2** Gear Type: **30** Days Fished: **7** Offloading Date: **14-Aug-98** Fish Slip #: **9876-5432**

Plant: **NORTH SEA PRODUCTS LTD (TOFINO BUYIN)** **TOFINO**

Matching Fish Slip ☒ Data Verified ☒

Date	Time 24hrs	Latitude	Longitude	Fishing Area/FAO	# of Jigs Used	Water Temp. (°F)	# of Fish Caught	Ave. WT (lbs)	Remarks (weather, feed, fleet size, etc.)
06-08-98	8:00	47300	N	8	64.0	2	11		
		125 37.0	W						
		W/A		67					
	20:00	46 12.0	N	8	65.0	22	12		
		125 26.0	W						
		OR		67					
			N	13		0			
			W						

Daily Fish: **24** Daily Weight: **286**

Record: 1 of 7 Record: 1 of 2

Fish Offloaded: **547** lbs Offloaded: **6,649** mt Offloaded: **3.016**

Calculate

Record: 27 of 32

Figure 2. Triplog data entry form.

Fish Slips

Vessel: Wave-Length Rider Date: 14-Aug-98 Slip: 9876-5432

Date: 14-Aug-1998 Slip #: 9876-5432

Fisher's Name: I.M. Jiggin Gear: Troll

Area of Catch: OR FAO: 67

Length of Trip: 10 Fishing Days: 8

Plant: NORTH SEA PRODUCTS LTD (TOFINO BUYING ST) Company: 034

Landing Status: Fish Quality: Port: TOFINO

Matching Trip Log? ☒ Trip #: 2 Data Verified: ☒

Species Code	Species Name	Size Code	DFD Code	Pieces	Weight (lbs)	DFD Form	Price per lb	Value
1051	Tuna	+15	60		580		\$1.40	\$812.00
1052	Tuna	-15	60		5,876		\$1.10	\$6,463.60
376	Albacore							
Total (all species)								\$7,275.60

Record: 18 of 21

Figure 3. Saleslip data entry form.




Figure 4. Food and Agricultural Organization (FAO) fishing areas.




Figure 5. DFO Albacore Tuna Fishing Areas within the FAO fishing areas.




Figure 6. Venn diagram demonstrating the intersection of fisher logbook (triplogs) and landed catch (saleslip) data subsets.




Figure 7. Expanded Venn diagram demonstrating the intersection of fisher logbook (triplogs) and landed catch (saleslip), and hail data subsets.




Figure 8. Albacore tuna catch estimates, 1995-2005.




Figure 9a. Triplog data analytical flow model: pseudo-code.




Figure 9b. Triplog data analytical flow model: database objects.




Figure 10a. Saleslips analytical flow model: pseudo-code.




Figure 10b. Saleslips analytical flow model: database objects.




Figure 11a. Unique vessels analytical flow model: pseudo-code.




Figure 11b. Unique vessels analytical flow model: database objects.




Figure 12a. Hail data analytical flow model: pseudo-code.




Figure 12b. Hail data analytical flow model: database objects.




Figure 13a. Catch and effort analytical flow model (saleslips-based approach): pseudo-code.




Figure 13b. Catch and effort analytical flow model (saleslips-based approach): database objects.




Figure 13a (cont'd.). Catch and effort analytical flow model (saleslips-based approach): pseudo-code.




Figure 13b (cont'd.). Catch and effort analytical flow model (saleslips-based approach): database objects.




Figure 14a. Catch and effort analytical flow model (triplog-based approach): pseudo-code.




Figure 14b. Catch and effort analytical flow model (triplog-based approach): database objects.




Figure 15a. Analytical flow model for catch, effort, and CPUE by year-vessel-trip-date at latitude and longitude coordinates, for export to GIS: pseudo-code.




Figure 15b. Analytical flow model for catch, effort, and CPUE by year-vessel-trip-date at latitude and longitude coordinates, for export to GIS: database objects.




Figure 16. Reported and expanded albacore catch as a function of percent of vessels reporting (2005 simulation). Expanded catch estimates begin to stabilize and approach the final reported catch estimate after approximately 30-40% of the vessels have reported.




Figure 17. Saleslip trip catch weight as a function of triplog trip catch weight, 1995-2005. ($r^2 = 0.987$, $b = 0.991$, $\text{Prob}(b=1) = 0.326$, $N = 2730$)




Figure 18. Comparison of post-season catch estimates based on triplog-based approach ($T+Y$) and saleslip-based approach ($S+X$).

9. TABLES

Table 1. Total reported catch, supplemental catch, and percent supplemental of total catch from the Albacore Tuna Relational Database, 1995-2005 (see Appendix 1 for historical catch, 1939-1994).

Year	Total Catch (lbs)	Supplemental Catch (lbs)	Percent Supplemental
1995	2,690,294	19,618	0.73%
1996	4,093,977	295,465	7.22%
1997	3,418,374	49,186	1.44%
1998	7,818,823	139,073	1.78%
1999	5,703,129	171,059	3.00%
2000	8,691,469	169,019	1.94%
2001	11,322,378	153,267	1.35%
2002	10,019,259	1,037,463	10.35%
2003	14,772,224	N/A	N/A
2004	15,804,135	N/A	N/A
2005	10,560,676	N/A	N/A


Table 2. DFO Albacore Tuna Fishing Areas and associated FAO areas.

Area	Area Name	FAO	FAO Area	Nation
BC	British Columbia	67	Northeast Pacific	CANADA
WA	Washington	67	Northeast Pacific	USA
OR	Oregon	67	Northeast Pacific	USA
CA-N	California	67	Northeast Pacific	USA
CA-S	California	77	Eastern Central Pacific	USA
US	United States	67	Northeast Pacific	USA
MX	Mexico	77	Eastern Central Pacific	OFFSHORE
InConv	InConvention	UNK	Unknown	OFFSHORE
OutConv	OutConvention	UNK	Unknown	OFFSHORE
NWP-IC	NW Pacific - InConv	61	Northwest Pacific	OFFSHORE
NEP-IC	NE Pacific - InConv	67	Northeast Pacific	OFFSHORE
NEP-OC	NE Pacific - OutConv	67	Northeast Pacific	OFFSHORE
WCP-IC	WC Pacific - InConv	71	West Central Pacific	OFFSHORE
ECP-IC	EC Pacific - InConv	77	Eastern Central Pacific	OFFSHORE
ECP-OC	EC Pacific - OutConv	77	Eastern Central Pacific	OFFSHORE
SWP-OC	SW Pacific – OutConv	81	Southwest Pacific	OFFSHORE
SWP-IC	SW Pacific - InConv	81	Southwest Pacific	OFFSHORE
SEP-IC	SE Pacific - InConv	87	Southeast Pacific	OFFSHORE
INT	International	UNK	Unknown	UNKNOWN
UNK	Unknown	UNK	Unknown	UNKNOWN

Table 3. Annual record of unique vessels fishing and percent of vessels submitting triplogs in Canadian albacore tuna fishing fleet. Unique Vessels Fishing represents the number of unique vessels active in the fishery based on triplog data, saleslip data, or other data sources. Percent Triplog Submission represents percent of vessels submitting triplogs as a function of total unique vessels fishing.

Year	Vessels Submitting Triplogs	Vessels Submitting Saleslips	Other Vessels	Unique Vessels Fishing	% Triplog Submission
1990	15	73		76	20%
1991	9	42		45	20%
1992	18	118		120	15%
1993	13	90		90	14%
1994	11	84	1	98	13%
1995	51	177	94	285	18%
1996	72	146	134	295	24%
1997	60	124	56	200	30%
1998	110	167	32	217	51%
1999	170	188	18	238	71%
2000	166	184	40	243	68%
2001	200	201	11	248	81%
2002	170	100	22	232	73%
2003	185	162	1	193	96%
2004	202	196	7	221	91%
2005	198	174	3	212	93%

Table 4. Example catch and effort summary report (T+Y method).

		Fisheries and Oceans	Pêches et Océans	Canada	
1998 Albacore Tuna Catch Summary Report					[Preliminary]
North & South Pacific Source Data					
Source	Vessels¹ Submitting	Unique² Vessels	Reported Catch³ (lbs) (t)	Unique Vessels	Expanded Vessels
Triplogs	110	110	4,984,365 2,261	CAN 93	124
Saleslips	167	75	2,733,747 1,240	USA 90	120
Hails	118	19			
Estimated		13			
		217	7,718,112 3,501		

North Pacific Vessels & Catch															
AREA		REPORTED⁴				Days	Catch	Phone-In⁵	Estimated⁶	EXPANDED⁷				Other⁸	Total
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch	per Vessel	per Day	Vessels	Vess.	Days	Catch (lbs)	(t)	
61	NWP-IC	4	64	180,030	0.6	6.0	11,914	15.23	2,741.4	0.47	5	81	223,070	101	
61			64	180,030		6	11,914					81	223,070	101	0 101
67	BC	93	1,035	565,818	13.9	97.3	37,446	10.60	532.8	10.98	6.35	124	1,316	701,090	318
	WA	60	613	676,872	8.9	57.6	44,795	9.72	1,076.5	7.08	4.10	80	779	838,694	380
	OR	72	1,843	2,389,625	10.7	173.3	158,145	24.37	1,263.7	8.50	4.92	96	2,343	2,960,922	1,343
	CA-N	2	11	14,011	0.3	1.0	927	5.24	1,241.3	0.24	0.14	3	14	17,361	8
	US	3	26	114,460	0.4	2.4	7,575	8.18	4,330.3	0.35	0.20	4	33	141,824	64
	NEP-IC	21	209	861,358	3.1	19.7	57,004	9.49	4,011.1	2.48	1.43	28	266	1,067,286	484
	NEP-OC	32	809	2,063,899	4.8	76.0	136,588	24.06	2,487.0	3.78	2.19	43	1,028	2,557,323	1,160
67			4,545	6,686,043		427	442,481					5,779	8,284,501	3,758	63 3,821
77	CA-S	4	52	17,962	0.6	4.9	1,189	12.38	336.6	0.47	0.27	5	66	22,256	10
	ECP-IC	8	93	209,296	1.2	8.7	13,851	11.04	2,198.9	0.94	0.55	11	118	259,333	118
	ECP-OC	6	32	30,279	0.9	3.0	2,004	5.08	922.1	0.71	0.41	8	41	37,518	17
77			177	257,537		17	17,044					225	319,107	145	102 247
Subtotal			4,786	7,123,610		450	471,439					6,085	8,826,679	4,004	165 4,169


South Pacific Vessels & Catch															
AREA		REPORTED⁴				Days	Catch	Phone-In⁵	Estimated⁶	EXPANDED⁷				Other⁸	Total
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch	per Vessel	per Day	Vessels	Vess.	Days	Catch (lbs)	(t)	
81	SWP-IC	1	37	123,063	0.0	0.0	0	37.00	3,326.0	0.00	2.00	3	111	369,189	167
81			37	123,063		0	0					111	369,189	167	0 167
Subtotal			37	123,063		0	0					111	369,189	167	0 167

Total North & South Pacific Vessels and Catch															
AREA		REPORTED				Total		Catch	Days	Catch	EXPANDED				Other
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch	per Vessel	per Vessel	per Day	Vess.	Catch (lbs)	(t)	(t)	Total
Total		161	4,823	7,246,673	24	450	471,439	185	5,273	7,718,112	41,720	29	1,464	217	9,195,868 4,171 165 4,336

NOTES:

- 1 Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- 2 The no. of unique vessels fishing. Since Triplog catch is used instead of saleslip catch when both exist for a given vessel and fishing area, Triplog vessels may include vessels from other sources. However, Saleslip vessels represent vessels that only submitted saleslips and Phone-Ins represent vessels that only phoned in. Total unique saleslip vessels (75) is partitioned into defined area vessels (52) and undefined area vessels (23).
- 3 Triplog catch is based on no. fish x daily mean weight, or (if absent) weighted annual area mean weight. Triplog and Saleslip weights include Albacore only.
- 4 Reported Vessels corresponds to the number of unique vessels that fished in each area. Note that some vessels may fish in more than one area. Days=Days Fished. Saleslip vessels & catch from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- 5 Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- 6 Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- 7 Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel.
Expanded Catch = Expanded Days x Catch per Day.
- 8 OTHER includes Supplemental catch tomes (63) and Trans-shipment tomes (102), which were not recorded in either saleslips or triplog catch data.

Table 5. Example catch and effort summary report (S+X method).



Fisheries and Oceans

Pêches et Océans

Canada

1998 Albacore Tuna Catch Summary Report

[Final]

North & South Pacific Source Data					
Source	Vessels ¹ Submitting	Unique Vessels ²	Reported Catch ³ (lbs) (t)		
Triplogs	110	110	5,085,076	2,307	
Saleslips	167	75	2,733,747	1,240	
Hails	118	19			
Estimated		13			
	217		7,818,823	3,547	

	Unique Vessels	Expanded Vessels
CAN	93	124
USA	90	120

North Pacific Vessels & Catch																			
AREA		REPORTED ⁴						Days per Vessel		Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED ⁷			Other ⁸ (t)	Total		
FAO	DFO	Defined Areas		Undefined Areas		Vess.	Days	Catch	Vess.	Days	Catch	Vessel	Day	Vessels	Vessels	Vess.	Days	Catch (lbs)	(t)
61	NWP-IC	4	64	169,682	0.6	6.0	11,073	15.23	2,581.6	0.47	0.27	5	81	210,067	95				
61			64	169,682		6.0	11,073									81	210,067	95	0
67	BC	93	1,035	556,165	13.9	97.3	36,294	10.60	523.2	10.98	6.35	124	1,316	688,533	312				
	WA	60	613	666,521	8.9	57.6	43,495	9.72	1,059.1	7.08	4.10	80	779	825,154	374				
	OR	72	1,843	2,373,115	10.7	173.3	154,863	24.37	1,253.9	8.50	4.92	96	2,343	2,937,921	1,333				
	CA-N	2	11	16,715	0.3	1.0	1,091	5.24	1,479.6	0.24	0.14	3	14	20,693	9				
	US	3	26	114,460	0.4	2.4	7,469	8.18	4,326.6	0.35	0.20	4	33	141,702	64				
	NEP-IC	21	209	982,219	3.1	19.7	64,097	9.49	4,570.0	2.48	1.43	28	266	1,215,989	552				
	NEP-OC	32	809	2,090,670	4.8	76.0	136,431	24.06	2,517.1	3.78	2.19	43	1,028	2,588,253	1,174				
67			4,545	6,799,865		427	443,740						5,779	8,418,244	3,818			63	3,882
77	CA-S	4	52	16,194	0.6	4.9	1,057	12.38	303.2	0.47	0.27	5	66	20,048	9				
	ECP-IC	8	93	209,773	1.2	8.7	13,689	11.04	2,202.0	0.94	0.55	11	118	259,699	118				
	ECP-OC	6	32	28,807	0.9	3.0	1,880	5.08	876.6	0.71	0.41	8	41	35,663	16				
77			177	254,774		17	16,626						225	315,410	143			102	245
Subtotal			4,786	7,224,321		450	471,439						6,085	8,943,721	4,057			165	4,222

South Pacific Vessels & Catch																		
AREA		REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	Vess.	EXPANDED ⁷				Other ⁸ (t)	Total
		Defined Areas		Undefined Areas									Days Catch (lbs) (t)					
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch						Days	Catch (lbs)	(t)			
81	SWP-IC	1	37	123,063	0.0	0.0	0	37.00	3,326.0	0.00	2.00	3	111	369,189	167			
81			37	123,063		0	0						111	369,189	167	0	167	
Subtotal			37	123,063		0	0						111	369,189	167	0	167	

Total North & South Pacific Vessels and Catch																	
	REPORTED																
	Defined Areas			Undefined Areas			Total		Catch per Vessel	Days per Vessel	Catch per Day	EXPANDED			Other (t)	Total	
	Vess.	Days	Catch	Vess.	Days	Catch	Vess.	Days				Catch	Vess.	Catch (lbs)			(t)
Total	161	4,823	7,347,384	24	450	471,439	185	5,273	7,818,823	42,264	29	1,483	217	9,312,910	4,224	165	4,389

NOTES:

- Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (75) is partitioned into defined area vessels (52) and undefined area vessels (23).
- Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catchweight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- OTHER includes Supplemental catch tonnes (63) and Trans-shipment tonnage (102), which were not recorded in either saleslips or triplog catch data.

Table 6. Logbook data form used on albacore tuna fishery vessels, 2006.

2006 CANADIAN PACIFIC ALBACORE TUNA LOG BOOK										SUBMISSION DEADLINE: OCT 31, 2006	
Vessel Name: _____		Gear Type: _____		FIN		Captain's Name: _____		Hall #: _____		Buyer: _____	
Sales Slip # _____		Total # of Fish Offloaded: _____		Offload Date: _____		Total Weight Offloaded: _____		Offloading Port: _____		Page _____ of _____ for this trip	
Date (mm/dd)	Total Hours Fished	Position Latitude/Longitude	# of Jigs Used	Water Temp (F)	Species	# of Fish	Ave. Wt. Per Fish	Remarks (Weather, Fleet Size, etc.)			
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						
			S								
			E								
			W								
			N		ALB						

Table 7. Saleslip data form used in the albacore tuna fishery.

BOOK/CASH FISH (SMALL FORM)
COMPANY NAME & ADDRESS ▶

97

FISHERMAN'S NAME _____

ADDRESS _____

BOAT NAME _____

CFV # _____

NAME OF PLANT, PACKER OR COLLECTOR _____

PACKER C.F.V. # _____

TRIP LENGTH IN DAYS _____

FISH QUALITY

SLUSH	
ICE	

HAIL IN NO. _____

113601

AREA OF CATCH _____

DAYS FISHING _____

TOTAL _____

GEAR:

OILNET ☐

SEINE ☐

TROLL ☐

FREEZER ☐

OTHER _____

SKIPPER'S NAME _____ S.I. # _____ %

CREW NAME _____ S.I. # _____ %

CREW NAME _____ S.I. # _____ %

CREW NAME _____ S.I. # _____ %

PCS	WEIGHT	CODE	SPECIES	PRICE	VALUE
			SOCKEYE RD.		
			COHO RD.		
			"		
			PINKS RD.		
			"		
			CHUMS RD.		
			RED SPRING RD.		
			"		
			WHITE SPRING RD.		
			"		
			STEELHEADS RD.		
			JACKS		
			RED SPRING DR.		
			RED SPRING DR.		
			WHITE SPRING DR.		
			WHITE SPRING DR.		
			SOCKEYE DR.		
			COHO DR.		
			COHO DR.		
			PINKS DR.		
			CHUMS DR.		
			STEELHEADS DR.		
			LINGCOD		

CASH ☐ **BOOK** ☐

I certify that the above information is complete and correct.

TOTAL CATCH VALUE _____

M.C.L. _____

U.I.C. DEDUCTIONS _____


CASH PAYMENT _____

CREDIT TO ACCOUNT _____

FISHERMAN'S SIGNATURE _____ FALYMAN'S SIGNATURE _____


NOT NEGOTIABLE **FIRM'S COPY**

Table 8. Example catch and effort summary report for the month of July (*T+Y* method).



Fisheries and Oceans

Pêches et Océans



Jul 2005 Albacore Tuna Catch Summary Report

[Preliminary]

North & South Pacific Source Data

Source	Vessels ¹ Submitting	Unique ² Vessels	Reported Catch ³ (lbs) (t)		Unique Vessels	Expanded Vessels
Triplogs	120	120	2,316,153	1,051	CAN 21	36
Saleslips	53	2	48,117	22	USA 117	199
Hails	207	85				
Estimated		0				
		207	2,364,270	1,072		

North Pacific Vessels & Catch

A R E A		R E P O R T E D ⁴						D a y s		C a t c h	Phone-In ⁵	Estimated ⁶	E X P A N D E D ⁷			Other ⁸	Total	
FAO	DFO	Defined Areas		Undefined Areas		Total		per Vessel	per Day	per Vessel	Vessels	Vessels	Vess.	Days	Catch (lbs)	(t)	(t)	Total
67	BC	21	38	20,182	0.0	0.0	0	1.81	531.1	14.63	0.00	36	64	34,243	16			
	WA	106	508	373,955	0.0	0.0	0	4.79	736.1	73.85	0.00	180	862	634,497	288			
	OR	114	1,438	1,968,314	0.0	0.0	0	12.62	1,368.5	79.43	0.00	193	2,440	3,339,681	1,515			
	CA-N	1	1	405	0.0	0.0	0	1.00	405.0	0.70	0.00	2	2	687	0			
	NEP-OC	1	1	1,414	0.0	0.0	0	1.00	1,414.0	0.70	0.00	2	2	2,399	1			
67		1,986	2,364,270		0	0							3,370	4,011,507	1,820	0	1,820	
Subtotal		1,986	2,364,270		0	0							3,370	4,011,507	1,820	0	1,820	

South Pacific Vessels & Catch

Total North & South Pacific Vessels and Catch

		R E P O R T E D						Total		Catch per Vessel	Days per Vessel	Catch per Day	E X P A N D E D			Other (t)	Total
		Defined Areas		Undefined Areas		Total							Vess.	Catch (lbs)	(t)		
Total		122	1,986	2,364,270	0	0	122	1,986	2,364,270	19,379	16	1,190	207	4,011,507	1,820	0	1,820

NOTES:

- 1 Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- 2 The no. of unique vessels fishing. Since Triplog catch is used instead of saleslip catch when both exist for a given vessel and fishing area, Triplog vessels may include vessels from other sources. However, Saleslip vessels represent vessels that only submitted saleslips and Phone-Ins represent vessels that only phoned in. Total unique saleslip vessels (2) is partitioned into defined area vessels (2) and undefined area vessels (0).
- 3 Triplog catch is based on no. fish x daily mean weight, or (if absent) weighted annual area mean weight. Triplog and Saleslip weights include Albacore only.
- 4 Reported Vessels corresponds to the number of unique vessels that fished in each area. Note that some vessels may fish in more than one area. Days=Days Fished. Saleslip vessels & catch from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- 5 Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- 6 Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- 7 Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- 8 OTHER includes Supplemental catch tonnes (0) and Trans-shipment tonnes (0), which were not recorded in either saleslips or triplog catch data.

Table 9. Correlation and regression analysis of saleslip landings as a function of triplog landings, by year, vessel, and trip.

Year	N	r²	b	Prob b=0	Prob b=1
1995	37	0.95	1.110	0.0001	0.0086
1996	89	0.95	0.970	0.0001	0.1928
1997	44	0.97	0.854	0.0001	0.0001
1998	155	0.99	1.007	0.0001	0.1874
1999	216	0.99	0.953	0.0001	0.0001
2000	317	0.99	0.944	0.0001	0.0001
2001	357	0.99	0.982	0.0001	0.0001
2002	115	0.99	0.991	0.0001	0.0342
2003	425	0.99	0.982	0.0001	0.0019
2004	590	0.99	0.996	0.0001	0.4135
2005	385	0.99	1.064	0.0001	0.0001

Left Blank on Purpose

10. APPENDICES


Appendix 1. Historical Canadian albacore tuna jig-troll catch (tonnes), 1939-1994.¹

Year	Catch (t)	Year	Catch (t)
1939	129	1967	161
1940	2	1968	1,028
1941	35	1969	1,365
1942	-	1970	390
1943	13	1971	1,746
1944	210	1972	3,921
1945	648	1973	1,400
1946	196	1974	1,331
1947	36	1975	111
1948	984	1976	278
1949	1,012	1977	53
1950	961	1978	23
1951	86	1979	521
1952	71	1980	212
1953	5	1981	200
1954	-	1982	104
1955	-	1983	225
1956	17	1984	50
1957	8	1985	56
1958	74	1986	30
1959	212	1987	104
1960	5	1988	155
1961	4	1989	140
1962	1	1990	302
1963	5	1991	139
1964	3	1992	363
1965	15	1993	494
1966	44	1994	1,998

¹ 1939-1946 Anon. (1917-1950); 1947-1969, 1977-1990 Ware and Yamanaka (1991); 1970-1976, 1991-1994 Stocker (2005).


Appendix 2. Canadian albacore tuna jig-troll catch estimates from the Albacore Tuna Relational Database, 1995-2005.

Appendix 2.1. Canadian albacore tuna jig-troll catch (tonnes), 1995.



Fisheries and Oceans

Pêches et Océans



1995 Albacore Tuna Catch Summary Report

[Final]

North & South Pacific Source Data

Source	Vessels Submitting ¹	Unique Vessels ²	Reported Catch (lbs) ³		Unique Vessels	Expanded Vessels	
Triplogs	51	51	958,044	435	CAN	175	284
Saleslips	177	142	1,732,250	786	USA	9	15
Hails	176	81					
Estimated		13					
		287	2,690,294	1,220			

North Pacific Vessels & Catch

AREA	REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In Vessels ⁵	Estimated Vessels ⁶	EXPANDED ⁷				Other (t) ⁸	Total
	Defined Areas		Undefined Areas		Vess.	Days					Catch	Vess.	Days	Catch (lbs)		
FAO DFO	Vess.	Days	Catch	Vess.			Days	Catch								
67 BC	175	3,504	2,179,264	18.0	350.4	180,923	19.97	612.3	73.45	11.00	277	5,541	3,392,869	1,539		
WA	9	119	60,760	0.9	11.9	5,044	13.19	502.7	3.78	0.57	14	188	94,596	43		
OR	2	2	982	0.2	0.2	82	1.00	483.4	0.84	0.13	3	3	1,529	1		
NEP-IC	2	11	32,480	0.2	1.1	2,697	5.49	2,907.1	0.84	0.13	3	17	50,568	23		
NEP-OC	6	114	210,580	0.6	11.4	17,482	18.95	1,818.7	2.52	0.38	10	180	327,850	149		
67	3,750	2,484,066		375	206,228						5,930	3,867,411	1,754	9	1,763	
Subtotal	3,750	2,484,066		375	206,228						5,930	3,867,411	1,754	9	1,763	

South Pacific Vessels & Catch

Total North & South Pacific Vessels and Catch

	REPORTED						Total	Catch per Vessel	Days per Vessel	Catch per Day	EXPANDED				Other (t)	Total	
	Defined Areas		Undefined Areas		Vess.	Days					Catch	Vess.	Days	Catch (lbs)			(t)
	Vess.	Days	Catch	Vess.			Days	Catch									
Total	175	3,750	2,484,066	18	375	206,228	193	4,125	2,690,294	13,939	21	652	287	3,867,411	1,754	9	1,763

NOTES:

1

Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).

2

The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (142) is partitioned into defined area vessels (124) and undefined area vessels (18).

3

Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catchweight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.

4

Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.

5

Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.

6

Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.



7

Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.

8

OTHER includes Supplemental catch + lost logbook tonnage estimates (9) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.



Appendix 2.2. Canadian albacore tuna jig-troll catch (tonnes), 1996.

 Fisheries and Oceans / Pêches et Océans															
1996 Albacore Tuna Catch Summary Report															
[Final]															
North & South Pacific Source Data															
Source	Vessels ¹ Submitting		Unique Vessels ²		Reported Catch ³ (lbs) (t)				Unique Vessels		Expanded Vessels				
Triplogs	72		72		2,382,910	1,081			CAN	90	177				
Saleslips	146		89		1,711,067	776			USA	83	163				
Hails	255		121												
Estimated			13												
			295		4,093,977	1,857									
North Pacific Vessels & Catch															
AREA	REPORTED ⁴		DEFINED AREAS		UNDEFINED AREAS		Days per Vessel	Catch per Day	Phone-In Vessels ⁵	Estimated Vessels ⁶	EXPANDED ⁷		Other ⁸	Total	
FAO DFO	Vess.	Days	Catch	Vess.	Days	Catch					Vess.	Days	Catch (lbs) (t)		
61 NWP-IC	2	21	10,936	0.3	2.5	1,058	10.28	511.0	1.50	0.16	4	41	20,708	9	
61		21	10,936		2	1,058						41	20,708	9	
67 BC	90	1,455	625,585	12.8	171.2	60,520	15.83	421.8	67.64	7.02	177	2,808	1,184,570	537	
WA	34	105	96,754	4.8	12.4	9,360	3.02	904.2	25.55	2.65	67	203	183,207	83	
OR	80	1,924	1,533,019	11.3	226.3	148,305	23.54	782.1	60.12	6.24	158	3,712	2,902,831	1,317	
CA-N	5	8	4,631	0.7	0.9	448	1.57	568.1	3.76	0.39	10	15	8,770	4	
US	4	125	53,868	0.6	14.7	5,211	30.59	422.9	3.01	0.31	8	241	102,001	46	
NEP-IC	13	214	907,338	1.8	25.2	87,777	16.11	4,160.5	9.77	1.01	26	413	1,718,081	779	
NEP-OC	18	258	364,389	2.6	30.4	35,251	14.03	1,385.9	13.53	1.40	35	498	689,985	313	
67		4,089	3,585,584		481	346,872						7,890	6,789,446	3,080	
77 CA-S	1	23	6,985	0.1	2.7	676	22.51	298.0	0.75	0.08	2	44	13,226	6	
ECP-IC	3	35	24,836	0.4	4.1	2,403	11.42	696.3	2.25	0.23	6	68	47,027	21	
ECP-OC	6	56	49,456	0.9	6.6	4,784	9.14	866.6	4.51	0.47	12	108	93,647	42	
77		114	81,276		13	7,863						220	153,900	70	
Subtotal		4,224	3,677,797		497	355,793						8,151	6,964,054	3,159	
South Pacific Vessels & Catch															
AREA	REPORTED ⁴		DEFINED AREAS		UNDEFINED AREAS		Days per Vessel	Catch per Day	Phone-In Vessels ⁵	Estimated Vessels ⁶	EXPANDED ⁷		Other ⁸	Total	
FAO DFO	Vess.	Days	Catch	Vess.	Days	Catch					Vess.	Days	Catch (lbs) (t)		
81 SWP-IC	1	56	60,387	0.0	0.0	0	56.00	1,078.3	0.00	2.00	3	168	181,161	82	
81		56	60,387		0	0						168	181,161	82	
Subtotal		56	60,387		0	0						168	181,161	82	
Total North & South Pacific Vessels and Catch															
AREA	REPORTED ⁴		DEFINED AREAS		UNDEFINED AREAS		Total		Catch per Vessel	Days per Vessel	Catch per Day	EXPANDED ⁷		Other ⁸	Total
FAO DFO	Vess.	Days	Catch	Vess.	Days	Catch	Vess.	Days				Vess.	Catch (lbs) (t)		
Total	141	4,280	3,738,184	20	497	355,793	161	4,777	4,093,977	25,428	30	857	295	7,145,215	3,241
														157	3,398

NOTES:

- Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (89) is partitioned into defined area vessels (69) and undefined area vessels (20).
- Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch weight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days = Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- OTHER includes Supplemental catch + lost logbook tonnage estimates (134) and Trans-shipment tonnage (23) not recorded in either saleslips or triplog catch data.



Appendix 2.3. Canadian albacore tuna jig-troll catch (tonnes), 1997.

		Fisheries and Oceans / Pêches et Océans																					
1997 Albacore Tuna Catch Summary Report														[Final]									
North & South Pacific Source Data																							
Source		Vessels Submitting ¹		Unique Vessels ²		Reported Catch (lbs) (t) ³								Unique Vessels		Expanded Vessels							
Triplogs		60		60		2,123,862 963								CAN 67		119							
Saleslips		124		84		1,294,512 587								USA 59		104							
Hails		130		43																			
Estimated				13																			
				200		3,418,374 1,551																	
North Pacific Vessels & Catch																							
AREA		REPORTED ⁴						Days per Vessel		Catch per Day		Phone-In ⁵ Vessels		Estimated ⁶ Vessels		EXPANDED ⁷			Other ⁸ (t)		Total		
FAO DFO		Vess. Days		Catch		Vess. Days		Catch								Vess. Days		Catch (lbs) (t)					
61 NWP-IC		7 59		149,338		2.2 13.5		31,178		7.91 2,489.2		2.09		0.70		12 95		235,482		107			
61		59		149,338		14		31,178								95		235,482		107		0 107	
67 BC		67 631		217,014		20.7 144.5		45,306		8.84 338.5		20.01		6.70		114 1,011		342,196		155			
WA		44 520		380,136		13.6 119.2		79,362		11.10 718.9		13.14		4.40		75 834		599,413		272			
OR		35 608		467,329		10.8 139.4		97,565		16.32 755.6		10.45		3.50		60 975		736,900		334			
CA-N		2 7		2,043		0.6 1.6		426		3.29 286.9		0.60		0.20		3 11		3,221		1			
US		2 33		3,498		0.6 7.6		730		15.49 104.2		0.60		0.20		3 53		5,516		3			
NEP-IC		20 496		1,164,474		6.2 113.7		243,109		23.29 2,308.8		5.97		2.00		34 795		1,836,185		833			
NEP-OC		17 138		159,338		5.3 31.6		33,265		7.62 1,135.5		5.08		1.70		29 221		251,251		114			
67		2,433		2,393,833		558		499,764								3,901		3,774,681 1,712		22		1,734	
77 CA-S		4 63		22,114		1.2 14.4		4,617		14.79 345.2		1.19		0.40		7 101		34,870		16			
ECP-IC		8 79		95,626		2.5 18.1		19,964		9.27 1,190.4		2.39		0.80		14 127		150,787		68			
ECP-OC		7 63		76,245		2.2 14.4		15,918		8.45 1,190.2		2.09		0.70		12 101		120,227		55			
77		205		193,986		47		40,499								329		305,883 139		188		327	
Subtotal		2,697		2,737,157		618		571,440								4,324		4,316,047 1,958		210		2,168	
South Pacific Vessels & Catch																							
AREA		REPORTED ⁴						Days per Vessel		Catch per Day		Phone-In ⁵ Vessels		Estimated ⁶ Vessels		EXPANDED ⁷			Other ⁸ (t)		Total		
FAO DFO		Vess. Days		Catch		Vess. Days		Catch								Vess. Days		Catch (lbs) (t)					
81 SWP-IC		1 57		109,777		0.0 0.0		0		57.00 1,925.9		0.00		2.00		3 171		329,331		149			
81		57		109,777		0		0								171		329,331 149		0		149	
Subtotal		57		109,777		0		0								171		329,331 149		0		149	
Total North & South Pacific Vessels and Catch																							
		REPORTED ⁴						Total		Catch per Vessel		Days per Vessel		Catch per Day		EXPANDED ⁷			Other ⁸ (t)		Total		
		Defined Areas		Undefined Areas												Vess. Catch (lbs) (t)							
		Vess. Days		Catch		Vess. Days		Catch		Vess. Days		Catch											
Total		110 2,754		2,846,934		34 618		571,440		144 3,372		3,418,374		23,739		23 1,014		200 4,645,378		2,107		210 2,317	

NOTES:

- Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (84) is partitioned into defined area vessels (50) and undefined area vessels (34).
- Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch weight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- OTHER includes Supplemental catch + lost logbook tonnage estimates (22) and Trans-shipment tonnage (188) not recorded in either saleslips or triplog catch data.

Appendix 2.4. Canadian albacore tuna jig-troll catch (tonnes), 1998.

 Fisheries and Oceans / Pêches et Océans																					
1998 Albacore Tuna Catch Summary Report												[Final]									
North & South Pacific Source Data																					
Source		Vessels Submitting ¹		Unique Vessels ²		Reported Catch ³ (lbs) (t)				Unique Vessels		Expanded Vessels									
Triplogs		110		110		5,085,076 2,307				CAN 92		119									
Saleslips		167		75		2,733,747 1,240				USA 91		118									
Hails		118		19																	
Estimated				13																	
				217		7,818,823 3,547															
North Pacific Vessels & Catch																					
AREA		REPORTED ⁴						Days per Vessel		Catch per Day		Phone-In ⁵ Vessels		Estimated ⁶ Vessels		EXPANDED ⁷		Other ⁸ (t)		Total	
FAO DFO		Vess. Days		Catch		Vess. Days		Catch		Vess. Days		Catch (lbs) (t)		Vess. Days		Catch (lbs) (t)					
61 NWP-IC		4 64		169,682		0.6 5.6		10,638		15.23 2,591.5		0.41		0.27		5 80		207,259 94			
61				64 169,682		6 10,638										80 207,259 94		0		94	
67 BC		92 1,020		539,568		13.1 89.0		33,826		10.56 517.1		9.45		6.25		121 1,275		659,055 299			
WA		60 613		666,521		8.5 53.4		41,785		9.72 1,063.2		6.16		4.07		79 766		814,123 369			
OR		72 1,843		2,373,115		10.2 160.7		148,773		24.37 1,258.7		7.39		4.89		95 2,303		2,898,642 1,315			
CA-N		2 11		16,715		0.3 1.0		1,048		5.24 1,485.3		0.21		0.14		3 14		20,416 9			
US		5 71		148,503		0.7 6.2		9,310		13.47 2,051.4		0.51		0.34		7 88		181,389 82			
NEP-IC		21 209		982,219		3.0 18.3		61,576		9.49 4,587.5		2.16		1.43		28 262		1,199,732 544			
NEP-OC		32 809		2,090,670		4.5 70.5		131,066		24.06 2,526.7		3.29		2.17		42 1,011		2,553,649 1,158			
67		4,575		6,817,311		399 427,383										5,718 8,327,006 3,777		63		3,840	
77 CA-S		4 52		16,194		0.6 4.5		1,015		12.38 304.4		0.41		0.27		5 65		19,780 9			
ECP-IC		8 93		209,773		1.1 8.1		13,151		11.04 2,210.5		0.82		0.54		11 116		256,227 116			
ECP-OC		6 32		28,807		0.9 2.8		1,806		5.08 879.9		0.62		0.41		8 40		35,187 16			
77		177		254,774		15 15,972										221 311,193 141		102		243	
Subtotal		4,816		7,241,767		420 453,993										6,018 8,845,458 4,012		165		4,177	
South Pacific Vessels & Catch																					
AREA		REPORTED ⁴						Days per Vessel		Catch per Day		Phone-In ⁵ Vessels		Estimated ⁶ Vessels		EXPANDED ⁷		Other ⁸ (t)		Total	
FAO DFO		Vess. Days		Catch		Vess. Days		Catch		Vess. Days		Catch (lbs) (t)		Vess. Days		Catch (lbs) (t)					
81 SWP-IC		1 37		123,063		0.0 0.0		0		37.00 3,326.0		0.00		2.00		3 111		369,189 167			
81		37		123,063		0 0		0								111 369,189 167		0		167	
Subtotal		37		123,063		0 0		0								111 369,189 167		0		167	
Total North & South Pacific Vessels and Catch																					
AREA		REPORTED ⁴						Total		Catch per Vessel		Days per Vessel		Catch per Day		EXPANDED ⁷		Other (t)		Total	
Vess. Days		Catch		Vess. Days		Catch		Vess. Days		Catch		Vess. Days		Catch (lbs) (t)		Vess. Days		(t)		Total	
Total		162 4,853 7,364,830		23 420 453,993		185 5,273 7,818,823		42,264 29 1,483		217 9,214,647 4,180		165 4,345									

NOTES:

- 1 Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- 2 The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (75) is partitioned into defined area vessels (52) and undefined area vessels (23).
- 3 Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch weight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- 4 Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- 5 Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- 6 Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- 7 Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- 8 OTHER includes Supplemental catch + lost logbook tonnage estimates (63) and Trans-shipment tonnage (102) not recorded in either saleslips or triplog catch data.

Appendix 2.5. Canadian albacore tuna jig-troll catch (tonnes), 1999.

1999 Albacore Tuna Catch Summary Report

[Final]

North & South Pacific Source Data

Source	Vessels ¹ Submitting	Unique Vessels ²	Reported Catch ³ (lbs) (t)		Unique Vessels	Expanded Vessels
Triplogs	170	170	4,722,815	2,142	CAN 162	173
Saleslips	188	50	980,314	445	USA 176	188
Hails	161	9				
Estimated		9				
		238	5,703,129	2,587		

North Pacific Vessels & Catch

REPORTED ⁴																	EXPANDED ⁷							Other ⁸	
AREA		Defined Areas				Undefined Areas				Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	Vess.	Days	Catch (lbs)	(t)	(t)	Total						
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch																		
61	NWP-IC	5	85	96,846	0.0	0.0	0	17.00	1,139.4	0.21	0.16	5	91	103,922	47										
61		85	96,846		0	0							91	103,922	47		0	47							
67	BC	162	1,574	900,084	0.0	0.0	0	9.72	571.8	6.66	5.18	174	1,689	965,843	438										
	WA	164	1,812	1,604,915	0.0	0.0	0	11.05	885.6	6.74	5.24	176	1,945	1,722,169	781										
	OR	107	1,796	1,547,806	0.0	0.0	0	16.78	861.9	4.40	3.42	115	1,927	1,660,887	753										
	CAN	17	77	53,190	0.0	0.0	0	4.50	695.3	0.70	0.54	18	82	57,076	26										
	NEP-IC	13	80	79,005	0.0	0.0	0	6.15	987.6	0.53	0.42	14	86	84,777	38										
	NEP-OC	29	514	483,715	0.0	0.0	0	17.72	941.1	1.19	0.93	31	552	519,055	235										
67		5,853	4,668,714		0	0						6,280	5,009,807	2,272		78	2,350								
77	CAS	18	337	377,291	0.0	0.0	0	18.72	1,119.6	0.74	0.58	19	362	404,856	184										
	ECP-IC	7	140	135,046	0.0	0.0	0	20.00	964.6	0.29	0.22	8	150	144,913	66										
	ECP-OC	10	80	89,202	0.0	0.0	0	8.00	1,115.0	0.41	0.32	11	86	95,719	43										
77		557	601,540		0	0						598	645,488	293		44	337								
Subtotal		6,495	5,367,100		0	0						6,969	5,759,217	2,612		122	2,734								

South Pacific Vessels & Catch

REPORTED ⁴																	EXPANDED ⁷					Other ⁸	
AREA		Defined Areas				Undefined Areas				Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	Vess.	Days	Catch (lbs)	(t)	(t)	Total				
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch																
81	SWP-IC	3	118	336,029	0.0	0.0	0	39.40	2,842.8	0.00	2.00	5	197	560,048	254								
81		118	336,029		0	0							197	560,048	254	0	254						
Subtotal		118	336,029		0	0							197	560,048	254	0	254						



Total North & South Pacific Vessels and Catch

Total North Atlantic Vessels and Catch																	
	REPORTED										EXPANDED						
	Defined Areas		Undefined Areas		Total		Catch per Vessel	Days per Vessel	Catch per Day				Other (t)	Total			
	Vess.	Days	Catch	Vess.	Days	Catch				Vess.	Days	Catch			Vess.	Catch (lbs)	(t)
Total	220	6,613	5,703,129	0	0	0	220	6,613	5,703,129	25,923	30	862	238	6,319,265	2,866	122	2,988

NOTES:

- Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (50) is partitioned into defined area vessels (50) and undefined area vessels (0).
- Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catchweight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- OTHER includes Supplemental catch + lost logbook tommage estimates (78) and Trans-shipment tommage (44) not recorded in either saleslips or triplog catch data.



Appendix 2.6. Canadian albacore tuna jig-troll catch (tonnes), 2000.

		Fisheries and Oceans	Pêches et Océans		
2000 Albacore Tuna Catch Summary Report					[Final]
North & South Pacific Source Data					
Source	Vessels¹ Submitting	Unique Vessels²	Reported Catch³ (lbs)	Reported Catch³ (t)	
Triplogs	166	166	7,361,023	3,339	
Saleslips	184	37	1,330,446	603	
Hails	205	38			
Estimated		2			
		243	8,691,469	3,942	
					Unique Vessels
					Expanded Vessels
					CAN 131 155
					USA 184 218
North Pacific Vessels & Catch					
AREA	REPORTED⁴				
	Defined Areas		Undefined Areas		
FAO DFO	Vess.	Days	Catch	Vess.	Days
				Catch	
				Days	Catch
				per Vessel	per Day
				Phone-In⁵ Vessels	Estimated⁶ Vessels
					EXPANDED⁷
					Vess.
					Days
					Catch (lbs)
					(t)
					Other⁸ (t)
					Total
61 NWP-IC	10	280	380,030	0.0	1.0
				1,109	27.96
				1,356.6	1.87
				0.00	12
				333	452,134
				205	205
61		280	380,030	1	1,109
					333
					452,134
					205
					0
					205
67 BC	131	999	788,453	0.6	3.4
				2,301	7.61
				788.9	24.52
				0.00	156
				1,189	938,048
				425	425
WA	147	1,214	880,774	0.7	4.1
				2,570	8.25
				725.2	27.52
				0.00	175
				1,445	1,047,884
				475	475
OR	161	4,325	5,078,586	0.8	14.7
				14,820	26.82
				1,173.7	30.14
				0.00	192
				5,148	6,042,155
				2,741	2,741
CA-N	6	28	2,362	0.0	0.1
				7	4.67
				84.0	1.12
				0.00	7
				33	2,810
				1	1
US	2	29	14,929	0.0	0.1
				44	14.48
				514.5	0.37
				0.00	2
				35	17,762
				8	8
NEP-IC	8	178	493,979	0.0	0.6
				1,442	22.24
				2,771.2	1.50
				0.00	10
				212	587,703
				267	267
NEP-OC	18	192	358,068	0.1	0.7
				1,045	10.65
				1,864.1	3.37
				0.00	21
				229	426,004
				193	193
67		6,965	7,617,150	24	22,229
					8,291
					9,062,366
					4,111
					77
					4,188
77 CA-S	6	65	45,044	0.0	0.2
				131	10.82
				692.7	1.12
				0.00	7
				77	53,591
				24	24
ECP-IC	7	53	210,572	0.0	0.2
				614	7.57
				3,964.9	1.31
				0.00	8
				63	250,525
				114	114
ECP-OC	3	4	528	0.0	0.0
				2	1.33
				131.9	0.56
				0.00	4
				5	628
				0	0
77		122	256,145	0	747
					145
					304,743
					138
					0
					138
Subtotal		7,367	8,253,325	25	24,085
					8,769
					9,819,243
					4,454
					77
					4,531
South Pacific Vessels & Catch					
AREA	REPORTED⁴				
	Defined Areas		Undefined Areas		
FAO DFO	Vess.	Days	Catch	Vess.	Days
				Catch	
				Days	Catch
				per Vessel	per Day
				Phone-In⁵ Vessels	Estimated⁶ Vessels
					EXPANDED⁷
					Vess.
					Days
					Catch (lbs)
					(t)
					Other⁸ (t)
					Total
81 SWP-IC	3	209	414,059	0.0	0.0
				0	69.67
				1,981.1	0.00
				2.00	5
				348	690,098
				313	313
81		209	414,059	0	0
					348
					690,098
					313
					0
					313
Subtotal		209	414,059	0	0
					348
					690,098
					313
					0
					313
Total North & South Pacific Vessels and Catch					
AREA	REPORTED⁴				
	Defined Areas		Undefined Areas		
FAO DFO	Vess.	Days	Catch	Vess.	Days
				Catch	
				Days	Catch
				per Vessel	per Day
				Phone-In⁵ Vessels	Estimated⁶ Vessels
					EXPANDED⁷
					Vess.
					Days
					Catch (lbs)
					(t)
					Other⁸ (t)
					Total
Total	202	7,576	8,667,384	1	25
				24,085	203
				7,601	8,691,469
				42,815	37
				1,143	243
				10,509,341	4,767
				77	4,844

NOTES:

- 1 Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- 2 The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (37) is partitioned into defined area vessels (36) and undefined area vessels (1).
- 3 Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catchweight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- 4 Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- 5 Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- 6 Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- 7 Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- 8 OTHER includes Supplemental catch + lost logbook tonnage estimates (77) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.

Appendix 2.7. Canadian albacore tuna jig-troll catch (tonnes), 2001.

		Fisheries and Oceans / Pêches et Océans																							
2001 Albacore Tuna Catch Summary Report														[Final]											
North & South Pacific Source Data																									
Source		Vessels ¹ Submitting		Unique Vessels ²		Reported Catch ³ (lbs)		Reported Catch ³ (t)						Unique Vessels		Expanded Vessels									
Triplogs		200		200		9,541,246		4,328						CAN 176		185									
Saleslips		201		36		1,781,132		808						USA 207		217									
Hails		224		12																					
Estimated				0																					
				248		11,322,378		5,136																	
North Pacific Vessels & Catch																									
AREA		REPORTED ⁴						Days per Vessel		Catch per Day		Phone-In ⁵ Vessels		Estimated Vessels ⁶		EXPANDED ⁷				Other ⁸ (t)		Total			
FAO DFO		Vess.		Days		Catch		Vess.		Days		Catch		Vess.		Days		Catch (lbs)		(t)					
61 NWPC-IC		7 169		698,083		0.0 0.0		0		24.14 4,130.7		0.36		0.00		7 178		733,579		333					
61		169		698,083		0 0		0								178		733,579		333		0		333	
67 BC		176 1,669		1,459,549		0.0 0.0		0		9.48 874.3		8.95		0.00		185 1,754		1,533,763		696					
WA		194 3,303		3,280,824		0.0 0.0		0		17.03 993.2		9.86		0.00		204 3,471		3,447,646		1,564					
OR		169 3,616		4,198,628		0.0 0.0		0		21.40 1,161.1		8.59		0.00		178 3,800		4,412,118		2,001					
CA-N		18 110		127,767		0.0 0.0		0		6.11 1,161.5		0.92		0.00		19 116		134,264		61					
US		3 60		47,743		0.0 0.0		0		20.00 795.7		0.15		0.00		3 63		50,171		23					
NEP-IC		7 96		428,133		0.0 0.0		0		13.72 4,458.3		0.36		0.00		7 101		449,902		204					
NEP-OC		16 151		316,604		0.0 0.0		0		9.44 2,096.7		0.81		0.00		17 159		332,702		151					
67		9,006		9,859,248		0 0		0								9,464		10,360,566		4,700		70		4,770	
77 CA-S		16 235		230,073		0.0 0.0		0		14.69 979.0		0.81		0.00		17 247		241,772		110					
ECP-IC		5 99		66,948		0.0 0.0		0		19.80 676.2		0.25		0.00		5 104		70,352		32					
ECP-OC		4 27		8,516		0.0 0.0		0		6.75 315.4		0.20		0.00		4 28		8,949		4					
77		361		305,538		0 0		0								379		321,074		146		0		146	
Subtotal		9,536		10,862,869		0 0		0								10,021		11,415,218		5,178		70		5,248	
South Pacific Vessels & Catch																									
AREA		REPORTED ⁴						Days per Vessel		Catch per Day		Phone-In ⁵ Vessels		Estimated Vessels ⁶		EXPANDED ⁷				Other ⁸ (t)		Total			
FAO DFO		Vess.		Days		Catch		Vess.		Days		Catch		Vess.		Days		Catch (lbs)		(t)					
81 SWP-IC		4 168		459,509		0.0 0.0		0		42.09 2,729.6		0.00		0.00		4 168		459,509		208					
81		168		459,509		0 0		0								168		459,509		208		0		208	
Subtotal		168		459,509		0 0		0								168		459,509		208		0		208	
Total North & South Pacific Vessels and Catch																									
AREA		REPORTED ⁴						Days per Vessel		Catch per Day		Phone-In ⁵ Vessels		Estimated Vessels ⁶		EXPANDED ⁷				Other ⁸ (t)		Total			
FAO DFO		Vess.		Days		Catch		Vess.		Days		Catch		Vess.		Days		Catch (lbs)		(t)					
Total		236 9,704		11,322,378		0 0		0		236 9,704		11,322,378		47,976 41		1,167		248 11,874,728		5,386		70		5,456	

NOTES:

1. Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
2. The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (36) is partitioned into defined area vessels (36) and undefined area vessels (0).
3. Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch weight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
4. Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
5. Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
6. Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
7. Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
8. OTHER includes Supplemental catch + lost logbook tonnage estimates (70) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.

[Final]

Source	Vessels ¹ Submitting	Unique Vessels ²	Reported Catch (lbs)	Reported Catch (t) ³	Unique Vessels	Expanded Vessels	
Triplogs	171	171	9,073,075	4,116	CAN	124	135
Saleslips	100	40	1,012,754	459	USA	200	218
Hails	217	19					
Estimated		2					
		232	10,085,829	4,575			


REPORTED ⁴																			
AREA		Defined Areas						Undefined Areas		Days per Vessel	Catch per Day	Phone-In Vessels	Estimated Vessels	EXPANDED ⁷				Other (t)	Total
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch	Vessels	(t)					Vess.	Days	Catch (lbs)	(t)		
61	NWP-IC	6	97	307,196	0.0	0.0	0	16.17	3,167.0	0.54	0.00	7	106	334,858	152				
61			97	307,196		0	0						106	334,858	152	0	152		
67	BC	124	629	770,301	0.0	0.0	0	5.07	1,224.6	11.17	0.00	135	686	839,664	381				
	WA	191	4,443	5,325,052	0.0	0.0	0	23.26	1,198.6	17.20	0.00	208	4,843	5,804,560	2,633				
	OR	133	1,979	2,745,451	0.0	0.0	0	14.86	1,387.3	11.98	0.00	145	2,157	2,992,672	1,357				
	CA-N	15	128	376,855	0.0	0.0	0	8.53	2,944.2	1.35	0.00	16	140	410,790	186				
	US	1	3	3,178	0.0	0.0	0	3.00	1,059.3	0.09	0.00	1	3	3,464	2				
	NEP-IC	3	19	14,925	0.0	0.0	0	6.33	785.5	0.27	0.00	3	21	16,269	7				
	NEP-OC	11	137	104,961	0.0	0.0	0	12.45	766.1	0.99	0.00	12	149	114,412	52				
67			7,336	9,340,723		0	0					7,998	10,181,831	4,618	471	5,089			
77	CA-S	6	59	69,927	0.0	0.0	0	9.83	1,185.2	0.54	0.00	7	64	76,224	35				
	ECP-IC	4	113	202,447	0.0	0.0	0	28.25	1,791.6	0.36	0.00	4	123	220,677	100				
	ECP-OC	8	29	6,882	0.0	0.0	0	3.63	230.4	0.72	0.00	9	32	7,284	3				
77			201	279,056		0	0					219	304,184	138	0	138			
Subtotal			7,636	9,926,975		0	0					8,323	10,820,873	4,908	471	5,379			

AREA		REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED ⁷				Other ⁸		Total
		Defined Areas		Undefined Areas								Vess.	Days	Catch	Vess.	Days	Catch (lbs)	
FAO	DFO	Vess.	Days	Catch	Vess.	Days	Catch					Vess.	Days	Catch (lbs)	(t)	(t)		
81	SWP-IC	2	79	158,854	0.0	0.0	0	39.50	2,010.8	0.00	2.00	4	158	317,708	144			
81		79	158,854		0	0						158	317,708	144	0	144		
Subtotal		79	158,854		0	0						158	317,708	144	0	144		

	REPORTED										EXPANDED				Other (t)	Total	
	Defined Areas		Undefined Areas		Total		Catch per Vessel	Days per Vessel	Catch per Day			(t)					
	Vess.	Days	Catch	Vess.	Days	Catch				Vess.	Catch (lbs)						
Total	211	7,715	10,085,829	0	0	0	211	7,715	10,085,829	47,800	37	1,307	232	11,138,581	5,052	471	5,523

- 1 Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- 2 The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (40) is partitioned into defined area vessels (40) and undefined area vessels (0).
- 3 Reported Catch represents allocated catch attributed to Unique Vessels only. Catch is based on saleslip catch weights where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- 4 Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- 5 Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- 6 Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- 7 Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- 8 OTHER includes Supplemental catch + lost logbook tonnage estimates (471) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.

Appendix 2.9. Canadian albacore tuna jig-troll catch (tonnes), 2003.



Fisheries and Oceans
Pêches et Océans

Canada

[Final]

2003 Albacore Tuna Catch Summary Report

North & South Pacific Source Data

Source	Vessels Submitting ¹	Unique Vessels ²	Reported Catch ³ (lbs) (t)		Unique Vessels	Expanded Vessels
Triplogs	185	185	14,678,566	6,658	CAN 119	125
Saleslips	162	7	380,443	173	USA 177	185
Hails	173	1				
Estimated		0				
		193	15,059,009	6,831		

North Pacific Vessels & Catch

AREA	DFO	REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED ⁷			Other ⁸ (t)	Total
		Defined Areas		Undefined Areas		Vess.	Days					Catch	Vess.	Days		
61	NWP-IC	8	291	737,539	0.3			6.6	9,935	35.66	2,511.3				0.04	0.00
61			291	737,539		7	9,935					299	751,205	341	0	341
67	BC	119	868	1,192,555	5.2	19.8	16,065	7.15	1,362.0	0.62	0.00	125	892	1,214,652	551	
	WA	172	2,718	4,390,846	7.5	62.1	59,148	15.49	1,600.7	0.90	0.00	180	2,794	4,472,206	2,029	
	OR	157	4,126	8,135,226	6.8	94.2	109,587	25.76	1,953.6	0.82	0.00	165	4,241	8,285,966	3,758	
	CA-N	7	51	137,488	0.3	1.2	1,852	7.14	2,671.2	0.04	0.00	7	52	140,036	64	
	NEP-IC	7	42	52,695	0.3	1.0	710	5.88	1,243.2	0.04	0.00	7	43	53,672	24	
	NEP-OC	6	24	6,226	0.3	0.5	84	3.92	257.0	0.03	0.00	6	25	6,341	3	
67			7,829	13,915,036		179	187,445					8,047	14,172,872	6,429	0	6,429
77	CA-S	2	23	7,069	0.1	0.5	95	11.27	304.5	0.01	0.00	2	24	7,200	3	
	ECP-IC	8	46	189,904	0.3	1.1	2,558	5.64	4,090.5	0.04	0.00	8	47	193,423	88	
	ECP-OC	4	11	232	0.2	0.3	3	2.70	20.9	0.02	0.00	4	11	237	0	
77			80	197,205		2	2,656					82	200,859	91	0	91
Subtotal			8,200	14,849,781		187	200,037					8,429	15,124,936	6,861	0	6,861

South Pacific Vessels & Catch

AREA	DFO	REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED ⁷			Other ⁸ (t)	Total
		Defined Areas		Undefined Areas		Vess.	Days					Catch	Vess.	Days		
71	WP-IC	1	4	0	0.0			0.0	0	4.00	0.0				0.00	0.00
71			4	0	0	0	0					4	0	0	0	0
Subtotal			4	0	0	0	0					4	0	0	0	0

Total North & South Pacific Vessels and Catch

Total	REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED ⁷			Other ⁸ (t)	Total		
	Defined Areas		Undefined Areas		Total	Vess.					Days	Catch	Vess.			Days	Catch (lbs)
	Vess.	Days	Catch	Vess.			Days	Catch	Vess.	Days				Catch	Vess.		
Total	184	8,204	14,849,781	8	187	200,037	192	8,391	15,049,818	78,384	44	1,794	193	15,124,936	6,861	0	6,861

NOTES:

1

Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).

2

The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (7) is partitioned into defined area vessels (-1) and undefined area vessels (8).

3

Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.

4

Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.

5

Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.

6

Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.

7

Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.


8

OTHER includes Supplemental catch & lost logbook tommare estimates (0) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.

NOTES:

- 1 Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- 2 The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (7) is partitioned into defined area vessels (-1) and undefined area vessels (8).
- 3 Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch weight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- 4 Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- 5 Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- 6 Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- 7 Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- 8 OTHER includes Supplemental catch + lost logbook tonnage estimates (0) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.

Appendix 2.10. Canadian albacore tuna jig-troll catch (tonnes), 2004.



Fisheries
and Oceans

Pêches
et Océans

Canada

[Final]

2004 Albacore Tuna Catch Summary Report

North & South Pacific Source Data

Source	Vessels ¹ Submitting	Unique Vessels ²	Reported Catch ³		Unique Vessels	Expanded Vessels
			(lbs)	(t)		
Triplogs	203	203	16,514,523	7,491	CAN	172
Saleslips	196	11	395,045	179	USA	202
Hails	217	7				
Estimated		0				
		221	16,909,568	7,670		

North Pacific Vessels & Catch

AREA	DFO	REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED ⁷			Other ⁸ (t)	Total
		Vess.	Days	Catch	Vess.	Days	Catch					Vess.	Days	Catch (lbs)		
61	NWP-IC	3	60	94,172	0.0	0.0	0	20.00	1,569.5	0.10	0.00	3	62	97,253	44	
61			60	94,172	0	0						62	97,253	44	0	44
67	BC	172	2,058	2,826,734	0.0	0.0	0	11.97	1,373.5	5.63	0.00	178	2,125	2,919,197	1,324	
	WA	190	2,274	3,001,274	0.0	0.0	0	11.97	1,319.8	6.21	0.00	196	2,348	3,099,447	1,406	
	OR	184	5,003	10,475,088	0.0	0.0	0	27.19	2,093.8	6.02	0.00	190	5,167	10,817,731	4,907	
	CA-N	11	34	61,182	0.0	0.0	0	3.09	1,799.5	0.36	0.00	11	35	63,184	29	
	NEP-IC	2	14	1,398	0.0	0.0	0	7.00	99.9	0.07	0.00	2	14	1,444	1	
	NEP-OC	8	72	93,416	0.0	0.0	0	9.00	1,297.4	0.26	0.00	8	74	96,471	44	
67			9,455	16,459,092	0	0						9,764	16,997,474	7,710	0	7,710
77	CA-S	1	1	100	0.0	0.0	0	1.00	99.7	0.03	0.00	1	1	103	0	
	ECP-IC	3	80	165,979	0.0	0.0	0	26.67	2,074.7	0.10	0.00	3	83	171,408	78	
	ECP-OC	5	32	51,211	0.0	0.0	0	6.40	1,600.3	0.16	0.00	5	33	52,886	24	
77			113	217,289	0	0						117	224,397	102	0	102
Subtotal			9,628	16,770,554	0	0						9,943	17,319,123	7,856	0	7,856

South Pacific Vessels & Catch

AREA	DFO	REPORTED ⁴						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED ⁷			Other ⁸ (t)	Total
		Vess.	Days	Catch	Vess.	Days	Catch					Vess.	Days	Catch (lbs)		
81	SWP-IC	1	67	139,014	0.0	0.0	0	67.00	2,074.8	0.00	0.00	1	67	139,014	63	
81			67	139,014	0	0						67	139,014	63	0	63
Subtotal			67	139,014	0	0						67	139,014	63	0	63

Total North & South Pacific Vessels and Catch

	DFO	REPORTED						Days per Vessel	Catch per Day	Phone-In ⁵ Vessels	Estimated ⁶ Vessels	EXPANDED			Other (t)	Total		
		Vess.	Days	Catch	Vess.	Days	Catch					Vess.	Days	Catch (lbs)			(t)	
Total		214	9,695	16,909,568	0	0	0	214	9,695	16,909,568	79,017	45	1,744	221	17,458,137	7,919	0	7,919

NOTES:

1

Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).

2

The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (11) is partitioned into defined area vessels (11) and undefined area vessels (0).

3

Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch weight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.

4

Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days = Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.

5

Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.

6

Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.


7

Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.

8


OTHER includes Supplemental catch + lost logbook tonnage estimates (0) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.

Appendix 2.11. Canadian albacore tuna jig-troll catch (tonnes), 2005.



Fisheries and Oceans

Pêches et Océans



2005 Albacore Tuna Catch Summary Report

[Final]

North & South Pacific Source Data

Source	Vessels ¹ Submitting	Unique Vessels ²	Reported Catch ³ (lbs)	(t)	Unique Vessels	Expanded Vessels
Triplogs	201	201	10,403,502	4,719	CAN	201
Saleslips	174	9	251,562	114	USA	158
Hails	207	3				
Estimated		0				
		213	10,655,064	4,833		

North Pacific Vessels & Catch

AREA	DFO	REPORTED ⁴			Days per Vessel	Catch per Day	Phone-In Vessels ⁵	Estimated Vessels ⁶	EXPANDED ⁷			Other (t) ⁸	Total		
		Vess.	Days	Catch					Vess.	Days	Catch (lbs)			(t)	
61	NWP-IC	1	55	24,530	0.0	0.1	24	54.57	445.7	0.01	0.00	1	56	24,902	11
61			55	24,530		0	24					56	24,902	11	11
67	BC	196	2,894	3,473,292	1.9	5.2	3,444	14.65	1,199.2	2.80	0.00	201	2,940	3,525,931	1,599
	WA	144	1,174	887,233	1.4	2.1	880	8.09	755.1	2.06	0.00	147	1,193	900,679	409
	OR	144	4,004	5,606,222	1.4	7.2	5,559	27.59	1,399.0	2.06	0.00	147	4,068	5,691,186	2,582
	CA-N	19	47	65,251	0.2	0.1	65	2.45	1,387.2	0.27	0.00	19	48	66,240	30
	NEP-OC	21	252	430,085	0.2	0.5	426	11.91	1,705.3	0.30	0.00	22	256	436,603	198
67			8,371	10,462,082		15	10,375					8,505	10,620,638	4,817	0
77	ECP-IC	1	1	30	0.0	0.0	0	0.99	30.0	0.01	0.00	1	1	30	0
	ECP-OC	2	2	68	0.0	0.0	0	0.99	34.0	0.03	0.00	2	2	69	0
77			3	98		0	0					3	99	0	0
Subtotal			8,429	10,486,710		15	10,399					8,564	10,645,640	4,829	0

South Pacific Vessels & Catch

AREA	DFO	REPORTED ⁴			Days per Vessel	Catch per Day	Phone-In Vessels ⁵	Estimated Vessels ⁶	EXPANDED ⁷			Other (t) ⁸	Total		
		Vess.	Days	Catch					Vess.	Days	Catch (lbs)			(t)	
81	SWP-IC	2	111	157,955	0.0	0.0	0	55.50	1,423.0	0.00	0.00	2	111	157,955	72
81			111	157,955		0	0					111	157,955	72	72
Subtotal			111	157,955		0	0					111	157,955	72	72

Total North & South Pacific Vessels and Catch

	REPORTED ⁴			Total Vess.	Total Days	Total Catch	Catch per Vessel	Days per Vessel	Catch per Day	EXPANDED ⁷		Other (t)	Total		
	Defined Areas	Undefined Areas	Total							Vess.	Catch (lbs)			(t)	
Total	208	8,540	10,644,665	2	15	10,399	210	8,555	10,655,064	50,738	41	1,245	213	10,803,595	4,900

NOTES:

- 1 Total number of vessels submitting source information in calendar year (northern hemisphere) or November 01 to March 31 (southern hemisphere).
- 2 The no. of unique vessels fishing. Triplog vessels include vessels from saleslip sources where both triplog and saleslip records exist for a vessel-trip. Unique Saleslip vessels represent vessels that only submitted saleslips and Unique Hails represent vessels that only phoned in. Total unique saleslip vessels (9) is partitioned into defined area vessels (7) and undefined area vessels (2).
- 3 Reported Catch represents albacore catch attributed to Unique Vessels only. Catch is based on saleslip catch weight where landing records exist. If saleslip data are missing, then triplog catch is based on #fish x daily mean weight, or (if absent), weighted mean annual weight for area of catch.
- 4 Reported Vessels corresponds to the number of unique vessels that fished in each area. Some vessels may fish in more than one area. Days=Days Fished. Catch & effort from Undefined locations are allocated to specific fishing areas based on the distribution of vessels & catch for defined areas.
- 5 Counts for vessels for which only a phone-in record exists are allocated to fishing areas based on the distribution of vessels & catch for defined areas.
- 6 Estimated vessels collected from interview information are allocated to fishing areas based on the distribution of vessels for defined areas.
- 7 Expanded Vessels = Reported Vessels + Phone-in Vessels + Estimated Vessels. Expanded Days = Expanded Vessels x Days per Vessel. Expanded Catch = Expanded Days x Catch per Day.
- 8 OTHER includes Supplemental catch + lost logbook tonnage estimates (0) and Trans-shipment tonnage (0) not recorded in either saleslips or triplog catch data.

Appendix 3. Albacore Relational Database Data Table Field Properties and Data Structure – Field names, structure, data type and description, by data table.

3.1. Logbook Data – Table *Triplog*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	CFV#	Number, Long Integer	4	Canadian Fishing Vessel Number	<input checked="" type="checkbox"/>	Yes (Duplicates OK)
2	TRIPLogID	AutoNumber, Long Integer	4	Trip Log ID Number	<input checked="" type="checkbox"/>	No
3	Trip#	Number, Integer	2	Trip number for a specific year	<input checked="" type="checkbox"/>	No
4	StartDate	Date/Time	8	Start of trip	<input type="checkbox"/>	No
5	Year	Text	4	Fishing Year	<input type="checkbox"/>	No
6	Captain'sName	Text	45	Name of vessel Captain during the trip	<input type="checkbox"/>	No
7	CrewSize	Number, Long Integer	4	Number of crew members	<input type="checkbox"/>	No
8	GearType	Text	4	Type of gear used	<input type="checkbox"/>	No
9	DaysFished	Number, Integer	2	Number of days spent fishing	<input type="checkbox"/>	No
10	AreaFished	Text	10	Using the Latitude and Longitude given, Area Fished can be calculated	<input type="checkbox"/>	No
11	TotalFish	Number, Integer	2	Total fish offloaded according to saleslip	<input type="checkbox"/>	No
12	TotalWeight	Number, Integer	2	Total weight offloaded according to saleslip	<input type="checkbox"/>	No
13	OffloadingDate	Date/Time	8	Date fish were offloaded	<input type="checkbox"/>	No
14	PortCode	Text	30	Port fish were offloaded	<input type="checkbox"/>	No
15	Buyer	Text	80	Buyer	<input type="checkbox"/>	No
16	Hail	Text	15	Hail number	<input type="checkbox"/>	No
17	MatchingFishSlip	Yes/No	1	Has a matching Fish Slip	<input type="checkbox"/>	No
18	Slip#	Text	12	Fish sales slip number	<input type="checkbox"/>	No
19	Plant	Text	10	DFO - Plant Number Code	<input type="checkbox"/>	No
20	DataVerified	Yes/No	1	Has the data been verified?	<input type="checkbox"/>	No
21	ReceiptSent?	Yes/No	1	Has acknowledgement of triplog receipt been sent?	<input type="checkbox"/>	No
22	Verified?	Yes/No	1	Has this trip been checked?	<input type="checkbox"/>	No
23	ImportID	Number, Long Integer	4	Archipelago TripHeaderID	<input type="checkbox"/>	No
24	Comment	Memo	0	Remarks associated with this trip	<input type="checkbox"/>	No

3.2. Logbook Data – Table *CatchData*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	TripLogID	Number, Long Integer	4	Trip Log ID Number	<input checked="" type="checkbox"/>	No
2	CatchDataID	AutoNumber, Long Integer	4	Catch Data ID Number	<input checked="" type="checkbox"/>	Yes (No Duplicates)
3	Date	Date/Time	8	Date of fishing	<input type="checkbox"/>	No
4	HoursFished	Number, Integer	2	Hours fished that day	<input type="checkbox"/>	No
5	ImportID	Number, Long Integer	4	TripHeaderID for import purposes	<input type="checkbox"/>	No

3.3. Logbook Data – Table *CatchDataSets*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	TripLogID	Number, Long Integer	4	Trip Log ID Number	<input checked="" type="checkbox"/>	No
2	CatchDataID	Number, Long Integer	4	Catch Data ID Number	<input checked="" type="checkbox"/>	Yes (Duplicates OK)
3	CatchDataSetID	AutoNumber, Long Integer	4	Catch Data Sets ID Number	<input checked="" type="checkbox"/>	No
4	Time	Date/Time	8	Time of sets	<input type="checkbox"/>	No
5	Latitude	Text	8	Start latitude	<input type="checkbox"/>	No
6	LatHemisphere	Text	1	Enter S or N	<input type="checkbox"/>	No
7	Longitude	Text	9	Start Longitude	<input type="checkbox"/>	No
8	LongHemisphere	Text	1	Enter E or W	<input type="checkbox"/>	No
9	FishingArea	Text	7	Fishing Area determined by Latitude and Longitude	<input type="checkbox"/>	No
10	FAOArea	Text	4	FAO Statistical Areas determined by Latitude and Longitude	<input type="checkbox"/>	No
11	JigsUsed	Number, Integer	2	Number of jigs used	<input type="checkbox"/>	No
12	WaterTemp	Text	5	Water temperature in F	<input type="checkbox"/>	No
13	NumberOfFish	Number, Long Integer	4	Number of fish caught	<input type="checkbox"/>	No
14	AvgWtofFish	Number, Long Integer	4	Average weight of fish caught	<input type="checkbox"/>	No
15	OtherSpeciesName	Text	25	Name of other Species besides Albacore Tuna that was caught	<input type="checkbox"/>	No
16	Other SpeciesNumber	Number, Long Integer	4	Number of other species caught	<input type="checkbox"/>	No
17	Remarks	Memo	0	Remarks on weather, feet, feet size, etc	<input type="checkbox"/>	No
18	Omit?	Yes/No	1	Omit this record from catch analysis?	<input type="checkbox"/>	No
19	ImportID	Number, Long Integer	4	TripHeaderID for import purposes	<input type="checkbox"/>	No

3.4. Saleslip Data – Table *FishSlipHeader*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	CFV#	Number, Long Integer	4	Canadian Fishing Vessel Number	<input checked="" type="checkbox"/>	Yes (Duplicates OK)
2	FishSlipID	AutoNumber, Long Integer	4	Fish Slip ID Number	<input checked="" type="checkbox"/>	Yes (No Duplicates)
3	Year	Text	4	Fishing Year	<input type="checkbox"/>	No
4	Slip#	Text	12	Slip number	<input type="checkbox"/>	No
5	TripNumber	Number, Integer	2	Trip number for a specific year	<input type="checkbox"/>	No
6	Date	Date/Time	8	Date of Fish Slip	<input type="checkbox"/>	No
7	Fisherman'sName	Text	30	Fisherman Name	<input type="checkbox"/>	No
8	Buyer'sName	Text	80	Name of Buyer	<input type="checkbox"/>	No
9	AreaOfCatch	Text	7	Area of catch	<input type="checkbox"/>	No
10	CatchAreaFAO	Text	4	FAO Statistical Areas determined by Latitude and Longitude.	<input type="checkbox"/>	No
11	DaysFished	Number, Integer	2	Number of days fished	<input type="checkbox"/>	No
12	LengthOfTrip	Number, Integer	2	Length of trip	<input type="checkbox"/>	No
13	Gear	Text	3	DFO - Gear Code	<input type="checkbox"/>	No
14	Plant	Text	10	DFO - Plant Number Code	<input type="checkbox"/>	No
15	Company	Text	4	DFO - Company Code	<input type="checkbox"/>	No
16	LandingStatus	Text	3	DFO - Landing Status Code	<input type="checkbox"/>	No
17	TotalWeight	Number, Long Integer	4	Total weight (lbs) of Tuna	<input type="checkbox"/>	No
18	Total Value	Currency	8	Total amount paid for Tuna	<input type="checkbox"/>	No
19	Quality	Text	10	Quality of fish	<input type="checkbox"/>	No
20	MatchingTripLog	Yes/No	1	Fish Slip has a matching Trip Log	<input type="checkbox"/>	No
21	DataVerified	Yes/No	1	Has data entered been verified?	<input type="checkbox"/>	No
22	ImportID	Number, Long Integer	4	Import tag for identification purposes	<input type="checkbox"/>	Yes (Duplicates OK)

3.5. Saleslip Data – Table *FishSlipData*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	FishSlipID	Number, Long Integer	4	Fish Slip ID Number	<input checked="" type="checkbox"/>	Yes (Duplicates OK)
2	FishSlipDataID	AutoNumber, Long Integer	4	Fish Slip Data ID Number	<input checked="" type="checkbox"/>	No
3	SpeciesCode	Number, Long Integer	4	Fish Species	<input type="checkbox"/>	No
4	SpeciesName	Text	25	Name of Species	<input type="checkbox"/>	No
5	SizeCode	Text	10	Code for fish size	<input type="checkbox"/>	No
6	DFOCode	Number, Long Integer	4	DFO code	<input type="checkbox"/>	No
7	Pieces	Number, Long Integer	4	Number of Fish	<input type="checkbox"/>	No
8	Weight	Number, Long Integer	4	Weight (lbs) of fish	<input type="checkbox"/>	No
9	DFOForm	Number, Long Integer	4	DFO Form Code	<input type="checkbox"/>	No
10	PricePerPound	Currency	8	Ex-vessel price per pound	<input type="checkbox"/>	No
11	Value	Currency	8	Ex-vessel dollar value	<input type="checkbox"/>	No
12	ImportID	Number, Long Integer	4	Import tag for identification purposes	<input type="checkbox"/>	Yes (Duplicates OK)

3.6. Hail Data – Table *VesselFishingYears*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	VesselID	Number, Long Integer	4	Vessel ID Number	<input checked="" type="checkbox"/>	No
2	FishingYears	Text	4	The year a vessel fished tuna based on information received by F&O (e.g. trip log, sales slip, fishticket, record from other agency, phone-in to F&O at start of season, etc).	<input type="checkbox"/>	No
3	CalledIn	Yes/No	1	This means that the vessel owner/operator called in to say that they would be tuna fishing, or for 1998, they completed a questionnaire stating that they fished albacore in 1998.	<input type="checkbox"/>	No
4	DateCalledIn	Date/Time	8	Date vessel called in	<input type="checkbox"/>	No
5	TripLog	Yes/No	1	A check means that F&O has a minimum of one Trip Log on file for that vessel.	<input type="checkbox"/>	No
6	NumberOfTripLogs	Number, Long Integer	4	The number of Trip Logs F&O has for that vessel for that year.	<input type="checkbox"/>	Yes (Duplicates OK)
7	FishSlip	Yes/No	1	A check means that F&O has a minimum of one Sales Slip (Fish Ticket) on file for that vessel.	<input type="checkbox"/>	No
8	NumberOfFishSlips	Number, Long Integer	4	Lists how many Fish Slips F&O has for that vessel for that year.	<input type="checkbox"/>	Yes (Duplicates OK)
9	FishingYearID	AutoNumber, Long Integer	4	Fishing Year ID Number	<input checked="" type="checkbox"/>	No

3.7. Supplemental Data – Table *SupplementalData*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	ID	AutoNumber, Long Integer	4	Record ID	<input type="checkbox"/>	Yes (Duplicates OK)
2	Vessel	Text	255	Vessel name	<input type="checkbox"/>	No
3	VRN	Number, Long Integer	4	Vessel registration number	<input checked="" type="checkbox"/>	No
4	Year	Number, Integer	2	Year of catch	<input checked="" type="checkbox"/>	No
5	AddToTotalCatch	Number, Single	4	Supplemental catch to add to total landings	<input type="checkbox"/>	No
6	AddToUSCatch	Number, Single	4	Supplemental catch to add to U.S. landings only	<input type="checkbox"/>	No
7	NewTotalCatch	Number, Single	4	Total landings including supplemental catch	<input type="checkbox"/>	No
8	NewTotalUSCatch	Number, Single	4	Total U.S. landings including supplemental catch	<input type="checkbox"/>	No
9	Comment	Text	100	Comments associated with this record	<input type="checkbox"/>	No
10	Updated	Date/Time	8	Date of update	<input type="checkbox"/>	No

3.8. Trans-shipment Data – Table *Transshipments*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	Year	Text	4	Catch year	<input checked="" type="checkbox"/>	No
2	FAOArea	Text	7	FAO area where transshipments were obtained	<input checked="" type="checkbox"/>	No
3	Tonnage	Number, Long Integer	4	Transshipments from this FAO area in metric tonnes	<input type="checkbox"/>	No
4	Update	Date/Time	8	Date of update	<input type="checkbox"/>	No
5	Comment	Text	255	Remarks	<input type="checkbox"/>	No

3.9. Estimated (unreported) Vessels Data – Table *UnreportedVesselCount*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	Year	Text	4	Fishing year	<input checked="" type="checkbox"/>	No
2	Hemisphere	Text	5	Fishing location	<input checked="" type="checkbox"/>	No
3	UnreportedVessels	Number, Long Integer	4	Number of vessels	<input type="checkbox"/>	No
4	Update	Date/Time	8	Date of update	<input type="checkbox"/>	No
5	Comment	Text	255	Comment associated with this record	<input type="checkbox"/>	No

Appendix 4. Albacore Relational Database Lookup Table Field Properties and Data Structure – Table field names, data field properties, structure, data type and description, by lookup data table.

4.1. Area Codes – DFO Fishing Areas – Table *CatchAreaOrder*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	CatchOrderID	AutoNumber, Long Integer	4	Record ID	<input checked="" type="checkbox"/>	Yes (No Duplicates)
2	CatchAreaOrder	Number, Long Integer	4	DFO Catch Area sort sequence	<input type="checkbox"/>	No
3	CatchAreaShort	Text	7	DFO Catch Area abbreviated (ex. BC)	<input type="checkbox"/>	No
4	CatchAreaLong	Text	25	DFO Catch Area (ex. British Columbia waters)	<input type="checkbox"/>	No
5	FAOAreaShort	Text	4	FAO Catch Area abbreviated (ex. 67)	<input type="checkbox"/>	No
6	FAOAreaLong	Text	25	FAO Catch Area (ex. Northeast Pacific)	<input type="checkbox"/>	No
7	FAOAreaOrder	Number, Long Integer	4	FAO Catch Area sort sequence	<input type="checkbox"/>	No

4.2. Area Codes – FAO Fishing Areas – Table *FAOAreaOrder*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	FAOCatchOrderID	AutoNumber, Long Integer	4	Record ID	<input checked="" type="checkbox"/>	Yes (No Duplicates)
2	FAOCatchAreaOrder	Number, Long Integer	4	FAO Catch Area sort sequence	<input type="checkbox"/>	No
3	FAOCatchAreaShort	Text	4	FAO Catch Area abbreviated (ex. 67)	<input type="checkbox"/>	No
4	FAOCatchAreaLong	Text	25	FAO Catch Area (ex. Northeast Pacific)	<input type="checkbox"/>	No

4.3. Buyer Codes – Table *BuyerInformation*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	BuyerID	AutoNumber, Long Integer	4	Buyer Information ID Number	<input checked="" type="checkbox"/>	Yes (Duplicates OK)
2	Buyer	Text	80	Name of Buyer (Company)	<input type="checkbox"/>	No
3	Contact Person	Text	30	Person to contact at Company	<input type="checkbox"/>	No
4	Address	Text	45	Address of Company	<input type="checkbox"/>	No
5	City	Text	30	City of Company	<input type="checkbox"/>	No
6	Province/State	Text	4	Province or State of Company	<input type="checkbox"/>	No
7	PostalCode	Text	8	Postal Code of Company	<input type="checkbox"/>	Yes (Duplicates OK)
8	ZipCode	Text	15	US Postal Code	<input type="checkbox"/>	Yes (Duplicates OK)
9	Telephone	Text	14	Telephone number of Company	<input type="checkbox"/>	No
10	Fax	Text	14	Fax number of Company	<input type="checkbox"/>	No
11	Notes	Memo	0	Comments relevant to Buyer (alternate contact, etc.)	<input type="checkbox"/>	No
12	LastUpdated	Date/Time	8	When the Buyer was last updated	<input type="checkbox"/>	No

4.4. Port Codes – Table *OffloadPorts*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	PortID	AutoNumber, Long Integer	4	Offloading Port ID Number	<input checked="" type="checkbox"/>	Yes (Duplicates OK)
2	PortCode	Text	5	Code to describe offloading ports	<input type="checkbox"/>	Yes (No Duplicates)
3	OffloadingPort	Text	30	Name of offloading port	<input type="checkbox"/>	No

4.5. Vessel Codes – Table *VesselInformation*

#	Field Name	Data Type	Size	Description	Primary Key	Indexed
1	VesselID	AutoNumber, Long Integer	4	Vessel ID Number	<input checked="" type="checkbox"/>	No
2	CFV#	Number, Long Integer	4	The unique Canadian Fishing Vessel number assigned by F&O.	<input checked="" type="checkbox"/>	Yes (No Duplicates)
3	VesselName	Text	30	The name of the vessel as provided by the owner/operator to F&O through the Vessel Licensing Unit.	<input type="checkbox"/>	No
4	HomePort	Text	25	Vessel Home Port	<input type="checkbox"/>	No
5	RadioSign	Text	10	The radio call sign provided by the vessel owner/operator.	<input type="checkbox"/>	No
6	Registration#	Text	15	The unique Canadian Vessel registration number assigned by the Canada Department of Transport.	<input type="checkbox"/>	No
7	Licence	Text	10	Type of Fishing Licence	<input type="checkbox"/>	No
8	GrossV/d	Number, Double	8	Gross Volume	<input type="checkbox"/>	No
9	GRT	Number, Double	8	Vessel Gross Registered Tonnage (GRT) in metric tonnes. Estimate of the Gross Metric Tonnage of Vessel from Conversion Table.	<input type="checkbox"/>	No
10	OALen	Number, Double	8	Vessel overall length in metres.	<input type="checkbox"/>	No
11	Breadth	Text	10	Also called Beam. Vessel width in metres.	<input type="checkbox"/>	No
12	Depth	Text	10	Vessel draft in metres	<input type="checkbox"/>	No
13	Hull	Text	10	Vessel hull material: steel (STL), fibreglass (FBGS), wood (WOOD)	<input type="checkbox"/>	No
14	EngineHP	Number, Double	8	Horsepower of the vessel's main engine.	<input type="checkbox"/>	No
15	Prop	Text	10	Propulsion	<input type="checkbox"/>	No
16	GearCode	Number, Double	8	Type of Fishing Method or Methods	<input type="checkbox"/>	Yes (Duplicates OK)
17	CompanyName	Text	40	Company Name	<input type="checkbox"/>	No
18	Salutation	Text	2	Salutation	<input type="checkbox"/>	No
19	OwnerLast	Text	30	Owner Name	<input type="checkbox"/>	No
20	OwnerFirst	Text	30	Owner Name	<input type="checkbox"/>	No
21	Address	Text	50	Address	<input type="checkbox"/>	No
22	City	Text	30	City	<input type="checkbox"/>	No
23	Province	Text	2	Province	<input type="checkbox"/>	No
24	PostalCode	Text	7	Postal Code	<input type="checkbox"/>	Yes (Duplicates OK)
25	Telephone	Text	15	Telephone Number	<input type="checkbox"/>	No
26	Fax	Text	15	Fax Number	<input type="checkbox"/>	No
27	Email	Text	50	E-mail Address	<input type="checkbox"/>	No
28	Comments	Memo	0	Comments associated with this vessel	<input type="checkbox"/>	No
29	CurrentlyLicensedUS	Yes/No	1	Yes if the vessel is currently licensed & has fished in the U.S. (with two exceptions (Freeland and Melville))	<input type="checkbox"/>	No
30	TriplogReceiptSent	Yes/No	1	Owner has been sent an acknowledgement of receipt of triplogs? (cleared annually)	<input type="checkbox"/>	No