

October 7 to October 13, 2012 (Week 41)


Overall Influenza Summary

- Influenza activity in Canada remains low but has increased slightly from last week as more regions have reported localized and sporadic influenza activity
- In week 41, a total of 17 laboratory detections of influenza were reported, the majority (94%) of which were for influenza A viruses [8 A(H3), 1 A(H1N1)pdm09, 7 A(un-subtyped) and 1 B]
- Four new influenza outbreaks were reported in week 41: 3 in LTCFs and 1 other
- One influenza A-associated hospitalization (in a person ≥ 65 years of age) was reported in week 41
- The ILI consultation rate declined in week 41 to 10.2 per 1,000 patient visits and is below the expected level for this time of year

Influenza Activity (geographic spread) and Outbreaks


In week 41, 3 regions (within AB & ON) reported localized activity, 6 regions (within BC, AB, ON & QC) reported sporadic activity and the rest reported no activity (see Figure 1). Note that data was not received from SK for week 41. Four new influenza outbreaks were reported in week 41: 3 in long-term care facilities (LTCF) and 1 other type of facility (Figure 3).

Figure 1. Map of overall Influenza activity level by province and territory, Canada, Week 41




Note: Influenza activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates (see graphs and tables) and reported outbreaks. Please refer to detailed definitions on the last page. For areas where no data is reported, late reports from these provinces and territories will appear on the FluWatch website.

Figure 2. Number of influenza surveillance regions† reporting widespread or localized influenza activity, Canada, by report week, 2012-2013 (Total number of influenza surveillance regions in Canada=58)



† sub-regions within the province or territory as defined by the provincial/territorial epidemiologist. Graph may change as late returns come in.

Figure 3. Overall number of influenza outbreaks, Canada, by report week, 2012-2013



Influenza and Other Respiratory Virus Detections

The percentage of positive influenza tests increased slightly from the previous week from 0.4% in week 40 to 0.9% in week 41 (Figures 4 and 5). Of the influenza viruses detected this week (n=17), 94% were positive for influenza A viruses (of which 50% were A(H3), 6.2% were A(H1N1)pdm09 and 43.8% were A untyped) and 6% were influenza B viruses (Table 1).

Cumulative influenza virus detections by type/subtype to date are as follows: 90.7% influenza A (51.0% - A(H3), 12.2% - A(H1N1)pdm09 and 36.7% - untyped) and 9.3% influenza B (Table 1).

To date this season, individuals aged 65 years and older make up the largest proportion of all influenza cases. Additional information on age and type/subtype of the positive influenza cases will be presented in Table 2 once more information becomes available.

The percentage positive for rhinovirus detections declined from the previous week (20.6% in week 41) but remains the highest compared to the other respiratory viruses. The percentage positive for the other respiratory viruses in week 41 remained low: RSV-2.0%; parainfluenza-2.4%; adenovirus-1.2%; hMPV-0.2%; and coronavirus-0.7% (Figure 5). For more details, see the weekly [Respiratory Virus Detections in Canada Report](#).

Table 1. Weekly & Cumulative numbers of positive influenza specimens by Provincial Laboratories, Canada, 2012-2013

Reporting provinces	October 7 to October 13, 2012						Cumulative (August 26, 2012 to October 13, 2012)					
	Influenza A					B	Influenza A					B
	A Total	A(H1)	A(H3)	Pand H1N1	A (UnS)*	Total	A Total	A(H1)	A(H3)	Pand H1N1	A (UnS)*	Total
BC	1	0	0	0	1	0	5	0	3	0	2	0
AB	6	0	2	1	3	0	14	0	4	6	4	1
SK	0	0	0	0	0	0	1	0	0	0	1	1
MB	0	0	0	0	0	0	3	0	3	0	0	0
ON	7	0	6	0	1	0	21	0	14	0	7	1
QC	2	0	0	0	2	1	4	0	0	0	4	2
NB	0	0	0	0	0	0	0	0	0	0	0	0
NS	0	0	0	0	0	0	0	0	0	0	0	0
PE	0	0	0	0	0	0	0	0	0	0	0	0
NL	0	0	0	0	0	0	1	0	1	0	0	0
Canada	16	0	8	1	7	1	49	0	25	6	18	5

*Unsubtyped: The specimen was typed as influenza A, but no test for subtyping was performed. Specimens from NT, YT, and NU are sent to reference laboratories in other provinces. Note: Weekly data is based on week of positive lab detection. Cumulative data includes updates to previous weeks; due to reporting delays, the sum of weekly report totals do not add up to cumulative totals.

Table 2. Weekly & Cumulative numbers of positive influenza specimens by age groups reported through case-based laboratory reporting, Canada, 2012-2013*

Age groups	Weekly (October 7 to October 13, 2012)					Cumulative (Aug. 26, 2012 to October 13, 2012)				
	Influenza A				B	Influenza A				B
	A Total	Pandemic H1N1	A/H3N2	A unsubtyped	Total	A Total	Pandemic H1N1	A/H3N2	A unsubtyped	Total
<5	-	-	-	-	-	-	-	-	-	-
5-19	-	-	-	-	-	-	-	-	-	-
20-44	-	-	-	-	-	-	-	-	-	-
45-64	-	-	-	-	-	-	-	-	-	-
65+	-	-	-	-	-	-	-	-	-	-
Unknown	-	-	-	-	-	-	-	-	-	-
Total	-	-	-	-	-	-	-	-	-	-

*Please note that the table will be populated once more data becomes available.

Figure 4. Influenza tests reported and percentage of tests positive, Canada, by report week, 2012-2013






Figure 5. Percent positive influenza tests, compared to other respiratory viruses, Canada, by reporting week, 2012-2013



Influenza Strain Characterizations

Since the start of the 2012-2013 influenza season, the National Microbiology Laboratory (NML) has antigenically characterized 1 influenza B virus; the virus was antigenically similar to the vaccine strain B/Wisconsin/01/2010 (Yamagata lineage).

Figure 6. Influenza strain characterizations, Canada, 2012-2013, N = 1



Note: The recommended components for the 2012-2013 Northern Hemisphere influenza vaccine include: an A/Victoria/361/2011 (H3N2)-like virus; an A/California/7/2009 (H1N1)pdm09-like virus; and a B/Wisconsin/1/2010-like virus.

Antiviral Resistance

Since the beginning of the season, NML has tested 1 influenza B virus for resistance to oseltamivir and 1 for zanamivir and it was found that the virus tested was sensitive to oseltamivir and zanamivir. A total of 3 influenza A(H3N2) viruses were tested for amantadine resistance and all 3 viruses tested were resistant (Table 3).

Table 3. Antiviral resistance by influenza virus type and subtype, Canada, 2012-2013


Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	0	0	0	0	3	3 (100%)
A (H1N1)	0	0	0	0	0	0
B	1	0	1	0	NA*	NA*
TOTAL	1	0	1	0	3	3 (100%)

* NA – not applicable

Influenza-like Illness Consultation Rate

The national influenza-like-illness (ILI) consultation rate in week 41 declined slightly from the previous week to 10.2 ILI consultations per 1,000 patient visits. However this rate is below the expected levels for this time of year (Figure 7). The highest consultation rates were observed in those between 5 and 19 years of age (16.6/1,000) and those less than 5 years of age (14.2/1,000).

Figure 7. Influenza-like illness (ILI) consultation rates, Canada, by report week, 2012-2013 compared to 1996/97 through to 2011/12 seasons (with pandemic data suppressed)



Note: No data available for mean rate in previous years for weeks 19 to 39 (1996-1997 through 2002-2003 seasons). Delays in the reporting of data may cause data to change retrospectively.

Severe Respiratory Illness Surveillance

Paediatric Influenza Hospitalizations and Deaths (IMPACT)

No laboratory-confirmed influenza-associated paediatric (16 years of age and under) hospitalizations have been reported by the Immunization Monitoring Program Active (IMPACT) network since the start of the 2012-2013 influenza season.

Influenza Hospitalizations and Deaths (Aggregate Surveillance System)

In week 41, one laboratory-confirmed influenza A-associated hospitalization was reported (from ON). The case was ≥65 years of age.

To date this season, 5 influenza A-associated hospitalizations have been reported from 3 provinces (AB, ON & NL); all were ≥ 65 years of age. Of the influenza A hospitalizations where subtype was available (4), 50% were due to influenza A(H1N1)pdm09 and 50% were due to A(H3N2). There has been one hospitalisation for which admission to ICU was required (from NL). To date this season, no influenza-associated deaths have been reported.

Note: Influenza-associated hospitalizations are not reported to PHAC by the following Provinces: BC, NU, QC, NS, & NB. Only hospitalizations that require intensive medical care are reported by SK. ICU admissions are not reported in ON.

International Influenza Updates

WHO: No new updates have been reported by the WHO since October 12, 2012.

[World Health Organization influenza update](#)

United States: The proportion of tests positive for influenza viruses decreased slightly in week 40 (2.6%) compared to the previous week. Of the positive influenza detections reported during week 40, 52% were positive for influenza A viruses. Of the 30 influenza A viruses for which subtype information was available, 90% were A(H3) and 10% were A(H1N1)pdm09. All other indicators of influenza activity remained low. [Centers for Disease Control and Prevention seasonal influenza report](#)

Europe: In week 41, surveillance data showed no evidence of increasing influenza activity in Europe and influenza activity remained low. A total of 21 samples tested positive for influenza in week 41 (0.5%): 62% were influenza A (38% were influenza A(H3N2) and 62% were A un-subtyped) and 38% were influenza B. ILI and ARI consultation rates are at usual levels for this time of year.

[EuroFlu weekly electronic bulletin](#)

Human Avian and Swine Influenza Updates

Human Avian Influenza

No cases of human avian influenza A/H5N1 infection have been reported by the WHO since August 10, 2012. A summary of the latest WHO [Human-Animal Interface Risk Assessment](#) (as of October 1, 2012) is available on the WHO webpage.

[WHO Avian influenza situation updates](#)

Human Swine Influenza

No new human cases of infection with swine influenza viruses or variants were reported this week.

FluWatch reports include data and information from the following sources: laboratory reports of positive influenza tests in Canada (National Microbiology Laboratory), sentinel physician reporting of influenza-like illness (ILI), provincial/territorial assessment of influenza activity based on various indicators, including laboratory surveillance, ILI reporting, and outbreaks, influenza-associated paediatric and adult hospitalizations, antiviral sales in Canada, and WHO and other international reports of influenza activity.

Abbreviations: Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

ILI definition for the 2012-2013 season

ILI in the general population: Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

Definitions of ILI/Influenza outbreaks for the 2012-2013 season

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

Hospitals and residential institutions: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Institutional outbreaks should be reported within 24 hours of identification. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Workplace: Greater than 10% absenteeism on any day which is most likely due to ILI.

Other settings: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

Influenza Activity Levels Definition for the 2012-2013 season

Influenza Regional Activity levels are defined as:

- 1 = No activity: no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported
- 2 = Sporadic: sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†
- 3 = Localized: (1) evidence of increased ILI* and
(2) lab confirmed influenza detection(s) together with
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**
- 4 = Widespread: (1) evidence of increased ILI* and
(2) lab confirmed influenza detection(s) together with
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **greater than or equal to 50% of the influenza surveillance region†**

Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

* More than just sporadic as determined by the provincial/territorial epidemiologist.

† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.

We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.

This report is available on the Public Health Agency website at the following address: <http://www.phac-aspc.gc.ca/fluwatch/index.html>. Ce rapport est disponible dans les deux langues officielles.