

Canadian Data Report of  
Fisheries and Aquatic Sciences 1166

2005

DATA REPORT ON A BENTHIC SURVEY CONDUCTED NEAR  
KENT ISLAND, BRITISH COLUMBIA

by

T.F. Sutherland<sup>1</sup>, C.D. Levings<sup>1</sup>, S.A. Petersen<sup>1</sup>, D. Sinclair<sup>2</sup>, S. Jepps<sup>2</sup>,  
B. Gillard<sup>3</sup>, J. Knight<sup>3</sup>, R. McPhie<sup>1</sup>, P. Walton<sup>1</sup>, J. Lessard<sup>4</sup>, E. McGreer<sup>5</sup>,  
and B. Taekema<sup>5</sup>

<sup>1</sup>Fisheries and Oceans Canada  
Science Branch – Pacific Region  
DFO-UBC Centre for Aquaculture and Environmental Research  
4160 Marine Drive, West Vancouver, B.C., V7V 1N6, Canada

<sup>2</sup>Fisheries and Oceans Canada  
Oceans, Habitat and Enhancement Branch  
Campbell River Office, #315 – 940 Alder Street  
Campbell River, B.C., V9W 2P8, Canada

<sup>3</sup>Fisheries and Oceans Canada  
Fisheries Management Branch  
P.O. Box 10, Port Hardy, B.C., V0N 2P0, Canada

<sup>4</sup>Fisheries and Oceans Canada  
Science Branch – Pacific Region  
Pacific Biological Station  
3190 Hammond Bay Rd., Nanaimo, B.C., V9T 6N7, Canada

<sup>5</sup>B.C. Ministry of Water, Land, and Air Protection  
2080-A Labieux Rd., Nanaimo, B.C., V9T 6J9, Canada

© Her Majesty the Queen in Right of Canada, 2005  
Cat. No. Fs 97-13/1166E ISSN 0706-6465

Correct citation for this publication:

Sutherland, T.F., Levings, C.D., Petersen, S.A., Sinclair, D., Jepps, S., Gillard, B.,  
Knight, J., McPhie, R., Walton, P., Lessard, J., McGreer, E., and Taekema, B.  
2005. Data report on a benthic survey conducted near Kent Island, British  
Columbia. Can. Data. Rep. Fish. Aquat. Sci. 1166: v + 27 p.

## LIST OF TABLES

<u>Table</u>	<u>Page</u>
1 Invertebrate species observed during a SCUBA dive at a fish farm at Kent Island, British Columbia on May 7, 2002.....	8
2 Flora and fish species observed during a SCUBA dive at a fish farm at Kent Island, British Columbia on May 7, 2002.....	9
3 Abalone observed during a SCUBA dive at a fish farm at Kent Island, British Columbia.....	10
4 GPS coordinates of the corners of the netpen structure at Kent Island, British Columbia.....	11
5 Invertebrate, fish and algae species observed during ROV survey at Kent Island, British Columbia.....	13
6 Description of grab sampling information at both farm (S, KI) and reference (RF) stations at Kent Island, British Columbia.....	14
7 Sediment grain size fractions (Ocean Dumping and Ocean Dumping Extra Points Protocols) observed at grab stations at Kent Island, British Columbia.....	15
8 Sediment total carbon and total nitrogen contents observed at farm and reference stations at Kent Island, British Columbia.....	19
9 Sediment trace-metal concentrations observed at farm and reference stations at Kent Island, British Columbia.....	20
10 Detection limits for sediment trace-metal analysis conducted on an Optima 4300 inductively coupled plasma emission spectrometer (PESC. SEDMET Method V 6.0).....	21
11 Benthic debris descriptions observed at farm and reference stations at Kent Island, British Columbia.....	24
12 Macrofaunal abundance observed at farm and reference stations at Kent Island, British Columbia.....	25

## LIST OF FIGURES

<u>Figure</u>	<u>Page</u>
1    Location of Kent Island, British Columbia.....	6
2    Location of shoreline reconnaissance dives and abalone transect dives at Kent Island, British Columbia.....	7
3    Location of ROV survey transects at Kent Island, British Columbia....	12
4    Location of farm and reference stations near Kent Island, British Columbia.....	14
5    Approximate locations of grab sampling stations relative to the netpen layout at Kent Island, British Columbia.....	16
6    Sediment grain size fractions observed at farm sampling stations at Kent Island, British Columbia.....	17
7    Sediment grain size fractions observed at reference stations at Kent Island, British Columbia.....	18
8    Sediment minor metals observed at farm sampling stations at Kent Island, British Columbia.....	22
9    Sediment minor metals observed at reference stations at Kent Island, British Columbia.....	23
10   Macrofaunal abundance (annelids excluded) found at farm sampling stations at Kent Island, British Columbia.....	26
11   Macrofaunal abundance (annelids excluded) found at reference stations at Kent Island, British Columbia.....	27



## ABSTRACT

Sutherland, T.F., Levings, C.D., Petersen, S.A., Sinclair, D., Jepps, S., Gillard, B., Knight, J., McPhie, R., Walton, P., Lessard, J., McGreer, E., and Taekema, B. 2005. Data report on a benthic survey conducted near Kent Island, British Columbia. Can. Data Rep. Fish Aquat. Sci. 1166: v + 27 p.

A grab sampling survey was carried out at a finfish site situated on the southwest side of Kent Island located in Queen Charlotte Strait, British Columbia. Replicate grab samples were also collected at a nearby reference site. The sediment samples were analyzed for sediment grain size, total nitrogen and carbon content, trace-metal concentration, and macrofaunal abundance. A SCUBA survey was also carried out approximately 6 months prior to the grab sampling to assess the fish, flora, and epifaunal communities present in the nearshore environment. The data resulting from these surveys are presented in this report.

## RESUMÉ

Sutherland, T.F., Levings, C.D., Petersen, S.A., Sinclair, D., Jepps, S., Gillard, B., and Knight, J., McPhie, R., Walton, P., Lessard, J., McGreer, E., and Taekema, B. 2005. Data report on a benthic survey conducted near Kent Island, British Columbia. Can. Data Rep. Fish Aquat. Sci. 1166: v + 27 p.

Une campagne d'échantillonnages ponctuels ponctuels a été effectuée sur un site piscicole se trouvant au sud-ouest de l'île Kent, dans le détroit de la Reine-Charlotte (Colombie-Britannique). Une campagne de contrôle parallèle par échantillonnages ponctuels a été effectuée. Des échantillons de référence ont également été recueillis de la même façon sur un site de référence se trouvant à proximité. Les sédiments recueillis échantillons ont été soumis à des analyses granulométriques, chimiques ( et à des analyses visant à établir la teneurs en azote, et en carbone et, la concentration de en métaux en traces) et biologiques (abondance de la macrofaune). l'abondance macrofaunique du sédiment. Environ six mois avant la campagne d'échantillonnage ponctuel, des relevés un programme d'étude en plongée autonome ont avait été effectués exécuté pour caractériser recenser les communautés piscicoles, floristiques et épifauniques présentes à proximité du rivage. Ce rapport présente orte sur les résultats données obtenues.

## INTRODUCTION

Environmental impacts may be associated with fish farm activities through changes in sedimentation rates of organic particulate matter, sediment oxygen demand and sulphide reduction, and benthic community structure (Fisheries and Oceans Canada 2004). The relationship between organic enrichment and the behavioural responses of benthic macrofauna have been reviewed by Diaz and Rosenberg (1995), while the recovery of benthic macrofaunal populations influenced by organic enrichment associated with fish farm activities has been documented by Lu and Wu (1998). This report provides a data summary of a suite of sediment environmental parameters as well as benthic macrofauna taxa observed at a salmon netpen location southwest of Kent Island, British Columbia. This farm had been in operation for a short period (March 13 to August 31, 2002) and consisted of twelve 24 x 24 m netpens in two rows of six (held 1.008 M Atlantic salmon smolts). The documentation of this data set may be useful in providing information for potential changes in macrofaunal communities associated with this level of production under current conditions in the British Columbia environment.

## SITE CHARACTERISTICS

Kent Island is part of the Walker group of islands located in Queen Charlotte Strait, British Columbia (Figure 1). The farm site is situated on the southwest side of Kent Island next to Bolivar Passage in a steep sided bay which drops to 100 m in depth within 600 m of the shore. The bay is approximately 735 m wide and 400 m long. Reference stations were located on the southeast side of Kent Island where Ripple Passage and Shelter Passage meet. The depth beneath the netpen system ranged from approximately 38 m to 90 m while the depths of the reference stations ranged from 60 m to 70 m (Table 6). A report by Brooks (2002) describes the benthic substrate consisting of bedrock, boulders, large cobble, and patches of sand near the proposed tenure site with predominantly sandy substrates observed 1600 m from the site. In addition, rich macroalgal beds and invertebrate communities, including Northern abalone (*Haliotis kamtschatkana*), were observed in the highly diverse habitat in the nearshore environment.

## BENTHIC SURVEY DESIGN

The staff of the Habitat Enhancement Branch (HEB) and Conservation and Protection Division (C&P) made plans to conduct a benthic grab survey at the fish farm site at Kent Island. Advice was sought from the Science Branch (C.D. Levings) regarding the sampling design of the grab survey. Sediment samples were collected along the perimeter of the netpen system as well as in a central location between the two netpen arrays. Three reference stations were located on the southwest side of Kent Island where Ripple Passage and Shelter Passage meet.

## MATERIALS AND METHODS

### SCUBA SURVEY

Shelley Jepps, Bryce Gillard, and Joe Knight carried out a reconnaissance SCUBA survey along the nearshore (896 metres) on May 7, 2002. Two transects made up of 3 dives were surveyed along the shore in a zig zag pattern between highwater and 20 feet of depth as measured by the diver's depth gauge. The primary diver recorded observed fauna and flora on an underwater slate and made general habitat observations. The secondary diver used a Sony Digital Handycam (model DCR-TRV17 NTSC) with an Amphibico Dive Buddy housing to video the habitat and species present. A species list of fish, invertebrates, and algae was generated from observations made along this dive.

Due to the noted presence of abalone during the reconnaissance SCUBA survey, a second set of SCUBA surveys were conducted between May 28<sup>th</sup> and 31<sup>st</sup>, June 10<sup>th</sup> and 12<sup>th</sup> of 2002 with the specific intent of estimating abalone population at the site. This survey was conducted using methodology outlined in Lessard et al. (2002). Transect locations for both dives are shown in Figure 2.

### ROV SURVEY

An ROV survey of the site was conducted with a Deep Ocean Phantom HD2 between June 4 and 6, 2002. The GPS locations of the four corners of the netpen structure were recorded using a Garmin II Plus (model GR-145-00214) from fisheries support vessels on June 4 (Port Hardy 1) and June 5 (Chimaera IV). A series of nine transects was then filmed with the ROV around the perimeter of the netpen structure as well as northward and southward along the predominant current in the area. The locations of these transects relative to the netpen structure are shown in Figure 3. Observations of habitat and fauna within the tenure were made from these transects by Shelley Jepps.

### GRAB SURVEY

On September 12<sup>th</sup> and 13<sup>th</sup>, 2002, three Ponar grab (0.04 m<sup>2</sup>) samples were collected by HEB and C&P staff at the Kent Island aquaculture facility. Although the individual nets had been removed from the two netpen arrays, walkway structures and 2 large predator nets remained in place. Only one Ponar grab (Station S-17) was successfully deployed from the netpen walkways using a portable capstan winch on September 12, 2002, due to net entanglements during deployments. As a result, perimeter sampling was resumed on September 13, 2002, with the use of a capstan gear puller and a 733-Hurricane Zodiac rigid hull inflatable alongside the netpen structure. Two more grab samples were successfully obtained using this technique (Stations S-9 and S-10).

When the Ponar grab was retrieved, the contents were carefully placed into a bucket and three 150 mL subsamples were removed from the intact surface layer using a plastic scoop. Each of these samples were placed in 180 mL cylindrical amber glass sample jars with plastic lids and stored in a cooler for geotechnical and chemical analyses. The remainder of the grab sample was then washed through a 1.0 mm sieve using screened surface seawater for macrofaunal analysis. The sieved samples were then placed in 1 L plastic sample jars and preserved with a solution of 10% buffered formalin.

The preserved samples were later transferred into isopropyl alcohol prior to laboratory analysis.

Further grab sampling was carried out using a Van Veen grab (0.1 m<sup>2</sup>) aboard the B.C. provincial vessel, Grizzly Coast, along with the support of the Ministry of Water, Land and Air Protection (MWLAP) on October 1, 2002. Once the grab samples were retrieved, four subsamples were removed from the surface layer and homogenized in a plastic container before being split into three 180 mL amber sample bottles according to MWLAP monitoring protocols. Ten sediment grab samples were collected along the netpen perimeter (Figure 5), while 3 reference samples were collected on the opposite side (southeast) of Kent Island. The macrofauna samples were sampled in a similar manner to those described above.

## **LABORATORY ANALYSIS**

The surficial sediment subsamples were delivered to Environment Canada's Pacific Environmental Science Centre (PESC), North Vancouver, British Columbia, for trace-metal, total nitrogen, total carbon and particle size analysis. The preserved macrofauna samples were sent to Biologica Environmental Services Ltd., Victoria, British Columbia, for invertebrate enumeration and general sample description.

### **Trace Metal Analysis**

A subsample of sediment from one of the three sample jars collected at each station was removed and placed in a 16 dram plastic vial with a snap closure, ¼ to ½ full. These subsamples were then analyzed according to the PESC SEDMET Method V 6.0, using an Optima 4300 Inductively Coupled Plasma Emission Spectrometer (PerkinElmer Life and Analytical Science, Woodbridge, ON). This procedure tests for content of: aluminum, antimony, arsenic, barium, beryllium, boron, cadmium, calcium, chromium, cobalt, copper, iron, lead, magnesium, manganese, molybdenum, nickel, phosphorous, potassium, selenium, silicon, silver, sodium, strontium, sulphur, tin, titanium, vanadium, and zinc.

### **Total Carbon and Nitrogen Analysis**

The total nitrogen concentration of the sediment samples was determined using the PESC TN Method V 2.0 automated, colorimetric, persulphate digest cadmium/copper reduction. Percent total carbon determinations were made by Pacific Soil Analysis Inc., Richmond, British Columbia. Determinations were made using a 12% CaCO<sub>3</sub> standard and performed in a 2500° F oven.

### **Particle Size Analysis**

The particle size analysis on the benthic sediment samples was carried out according to the Ocean Dumping Extra Points format (Soilcon, 2003). Each sample was first put through a 2.0 mm dry sieve. The portion <2.0 mm was then wet sieved through 1.0 mm, 0.5 mm, 0.25 mm and 0.125 mm sieves and finally the remaining solution was stirred and 20 mL subsamples removed with a volumetric pipette at predetermined time intervals to determine the soil fractions less than 0.063 mm, 0.004 mm and 0.002 mm. The particle size data was also merged and presented in the standard Ocean Dumping

format which consists of four broader categories: <0.004 mm, 0.004 - 0.63 mm, 0.63 - 2.00 mm, >2.00 mm.

### **Macrofauna Analysis**

For the macrofauna analysis, a Rose Bengal stain was added to the samples to stain organisms pink and help facilitate the sorting process. Five mL fractions of each sample were sorted in a gridded dish using a dissection microscope (10 to 30 x magnification) removing all organisms with forceps. Each dish was sorted at least twice. For sorting quality assurance, the debris was retained and a portion from each jar was spread in a 22.5 x 32.5 cm white pan and examined under a dissection microscope at 12 x magnification. If greater than 2 organisms were recovered the entire sample was resorted and subjected to another spot check. A sorting efficiency was calculated based on the results of the resorts. The entire sample was sorted in this manner. In terms of taxonomic verification, sorting identifications of the observed organisms were verified by a taxonomist. When incomplete specimens were present only anterior portions were counted. Colonial organisms such as bryozoa, hydrozoa and cnidarians were counted as colony fragments, not as specimens or individuals. Identified organisms were stored in 70% ethanol in labelled vials.

### **Debris Description**

Estimates of the retained debris portions of the preserved samples were provided by Biologica Environmental Services Ltd. Wood, mineral material and shell fragments were measured using 0.5 cm increments, while mineral material was measured using 0.2 cm. If the mineral portion of the debris was <0.2 cm it was identified as sand, silt or clay depending on a subjective judgement by the observer. Materials > 0.2 and < 0.5 cm were identified as gravel (<0.5 cm) and gravel larger than 0.5 cm was measured to the nearest 0.5 cm.

## **ACKNOWLEDGEMENTS**

We would like to thank the Pacific Environmental Science Centre (PESC), Pacific Soil Analysis Inc, Soilcon Laboratories Ltd., and Biologica Environmental Services Ltd. for analysis of various chemical, geotechnical, and biological sediment components, respectively. We would also like to thank the crew of the Grizzly Coast for their assistance during the field survey. Thanks to Scott Morrison for providing the maps and Beth Piercey for her assistance in editing this report. Funding for this project was provided by the HEB, C&P, and Science branches.

## REFERENCES

- Brooks, K.M. 2002. Predicted environmental response to the Kent Island aquaculture facility with emphasis on pinto abalone (*Haliotis kamtschatkana*). Report submitted to Omega Salmon Group Ltd., October 18, 2002, 22 p.
- Diaz, R.J., and Rosenberg, R. 1995. Marine benthic hypoxia: a review of its ecological effects and the behavioural responses of the benthic macrofauna. *Oceanogr. Mar. Biol. Annu. Rev.* 33: 245–303.
- Fisheries and Oceans Canada. 2004. A scientific review of the potential environmental effects of aquaculture in aquatic ecosystems. Volume III. Near-field organic enrichment from marine finfish aquaculture (D.J. Wildish, M. Dowd, T.F. Sutherland, and C.D. Levings); Environmental fate and effect of chemicals associated with Canadian freshwater aquaculture (R.J. Scott). *Can. Tech. Rep. Fish. Aquat. Sci.* 2450: ix + 117 pp.
- Lessard, J., Campbell, A. and Hajas, W. 2002. Survey protocol for the removal of allowable numbers of northern abalone, *Haliotis kamtschatkana*, for use as broodstock in aquaculture in British Columbia. *Can. Sci. Advisory Secretariat Res. Doc.* 2002/126.
- Lu, L. and Wu, R.S.S. 1998. Recolonization and succession of marine macrobenthos in organic-enriched sediment deposited from fish farms. *Environmental Pollution*, 101: 241 – 251.
- Pacific Environmental Science Centre. SEDMET Method V 6.0.
- Pacific Environmental Science Centre. TN Method V 2.0.
- Soilcon, 2003. Lab Methods, Soil Physical Properties, Particle Size Analysis. <http://www.soilconlabs.com> (accessed Jan 15, 2003).






Figure 1. Location of Kent Island, British Columbia.




Figure 2. Location of shoreline reconnaissance dives and abalone transect dives at Kent Island, British Columbia.



Table 1. Invertebrate species observed during a SCUBA survey at a fish farm at Kent Island, British Columbia on May 7, 2002.

Common Name	Latin Name
California Sea Cucumber	<i>Parastichopus californicus</i>
Orange Sea Cucumber	<i>Cucumaria</i> sp.
California Mussel	<i>Mytilus californianus</i>
Green False-jingle	<i>Pododesmus macrochisma</i>
Rock Scallop	<i>Crassadoma gigantea</i>
Northern Abalone	<i>Haliotis kamtschatkana</i>
Rough Keyhole Limpet	<i>Diodora aspera</i>
Brown Turban Snail	<i>Tegula pulligo</i>
Topsnail	<i>Calliostoma</i> sp.
Whitecap Limpet	<i>Acmaea mitra</i>
Limpets	Various unidentified
Leafy Hornmouth	<i>Ceratastoma foliatum</i>
Oregon Triton	<i>Fusitriton oregonensis</i>
Giant Pacific Octopus	<i>Octopus dofleini</i>
Gumboot Chiton	<i>Cryptochiton stelleri</i>
Black Katy Chiton	<i>Katharina tunicata</i>
Lined Chiton	<i>Tonicella</i> sp.
Chitons	Various unidentified
Sunflower Star	<i>Pycnopodia helianthoides</i>
Painted Star	<i>Orthasterias koehleri</i>
Morning Star	<i>Solaster stimpsoni</i>
Leather Star	<i>Dermasterias imbricata</i>
Mottled Star	<i>Evasterias troscheli</i>
Blood Star	<i>Henricia leviuscula</i>
Ochre Star	<i>Pisaster ochraceus</i>
Brittle Star	<i>Ophiopholis aculeata</i>
Puget Sound King Crab	<i>Lopholithodes mandtii</i>
Decorator Crab	Unidentified
Kelp Crab	<i>Pugettia producta</i>
Hermit Crabs	Unidentified
Dock (Coonstripe) Shrimp	<i>Pandalus danae</i>
Shrimp	Unidentified
Red Sea Urchin	<i>Strongylocentrotus franciscanus</i>
Purple Sea Urchin	<i>Strongylocentrotus purpuratus</i>
Green Sea Urchin	<i>Strongylocentrotus droebachiensis</i>
Slime Tube Worm	<i>Myxicolla infundibulum</i>
Calcareous Tube Worms	<i>Serpula vermicularis</i>
Coralline Fringed Tube Worm	<i>Dodecaceria concharum</i>
Northern Feather Duster Worms	<i>Eudistylia vancouveri</i>
Double Crowned Feather Dusters	Unidentified
Brooding Anemone	<i>Epiactis prolifera</i>
Buried Anemone	<i>Urticina coriacea</i>
Giant Plumose Anemone	<i>Metridium giganteum</i>
Giant Green Anemone	<i>Anthopleura xanthogrammica</i>
Painted Anemone	<i>Urticina crassicornis</i>
White-spotted Anemone	<i>Urticina lofotensis</i>
Fish-Eating Anemone	<i>Urticina piscivora</i>
Tube-dwelling Anemone	<i>Pachycerianthus fimbriatus</i>
Anemones	Unidentified anemones
Pink Mouth Hydroid	<i>Ectopleura</i> sp. (probably other sp. as well)
Glassy Plume Hydroid	<i>Plumeria</i> sp.
Orange Cup Coral	<i>Balanophyllia elegans</i>
Tritonia	<i>Tritonia festiva</i>
Odhner's Dorid	<i>Archidoris odhneri</i>
Tochni	<i>Tochuina tetraquetra</i>
Opalescent Nudibranch	<i>Hermisenda crassicornis</i>
Unidentified Nudibranchs	Various sp.
Staghorn Bryozoan	<i>Heteropora magna</i>
Goose Barnacle	<i>Pollicipes polymerus</i>
Giant Barnacle	<i>Balanus nubilus</i>
Stalked Sea Squirt	<i>Styela montereyensis</i>
Glassy Sea Squirt	<i>Ascidia paratropa</i>
Colonial Purple Tunicate	Unidentified
Sea Squirts	Various unidentified

Table 2. Flora and fish species observed during a SCUBA survey at a fish farm at Kent Island, British Columbia on May 7, 2002.

### Algae and Marine Plants

Common Name	Latin Name
Surf Grass	<i>Phyllospadix scouleri</i>
Fringed Sieve Kelp	<i>Agarum</i> sp.
Unidentified Red Filamentous Algae	Various sp.
Unidentified Brown Filamentous Algae	Various sp.
Iridescent Red	Various sp.
Encrusting Coralline Algae	Sp.?
Coralline Algae	Sp.?
Giant Kelp	<i>Macrocystis integrifolia</i>
Bull Kelp	<i>Nereocystis leutkeana</i>
Kelp	Laminarians
Split Kelp	<i>Laminaria groenlandica</i>
Sugar Kelp	<i>Laminaria saccharina</i>
Woody-stemmed Kelp	<i>Pterygophora californica</i>
Double Pom-pom Kelp	<i>Eisenia arborea</i>
Ribbon Kelp	<i>Alaria</i> sp.
Seersucker	<i>Costaria costata</i>
Acid Kelp	<i>Desmarestia</i> sp.
Three-ribbed Kelp	<i>Cymathere triplicata</i>

### Fish

Common Name	Latin Name
Buffalo Sculpin	<i>Enophrys bison</i>
Kelp Greenling	<i>Hexagrammos decagrammus</i>
Painted Greenling	<i>Oxylebius pictus</i>
Yellowtail Rockfish	<i>Sebastes flavidus</i>
Black Rockfish	<i>Sebastes melanops</i>
Quillback Rockfish	<i>Sebastes maliger</i>
Copper Rockfish	<i>Sebastes caurinus</i>
Tube Snout	<i>Aulorhynchus flavidus</i>
Unidentified Sculpins	Various sp.
Black-eyed Goby	<i>Coryphopterus nicholsi</i>
Goby	Unidentified

Table 3. Abalone observed during a SCUBA dive at a fish farm at Kent Island, British Columbia.

Transect	Transect Length (m)	Number of Quadrats	Number of Abalone
1	18	9	2
2	28	14	2
3	12	6	0
4	14	7	4
5	18	9	2
6	12	6	0
7	12	6	0
8	22	11	0
9	18	9	0
10	18	9	3
11	16	8	2
12	28	14	0
13	18	9	0
14	30	15	0
15	30	15	2
16	18	9	3
17	16	8	1
18	18	9	1
19	30	15	1
20	24	12	1
21	34	17	6
22	16	8	1
23	14	7	2
24	26	13	0
25	40	20	1

Table 4. GPS coordinates of the corners of the netpen structure at Kent Island, British Columbia (NP = netpen).

Corner	GPS coordinates	Date	Vessel
NP-1	N 50° 53.925' W 127° 32.438'	June 4, 2002	Port Hardy 1
NP-2	N 50° 53.942' W 127° 32.298'	June 4, 2002	Port Hardy 1
NP-3	N 50° 53.977' W 127° 32.310'	June 5, 2002	Chimaera IV
NP-4	N 50° 53.960' W 127° 32.454'	June 5, 2002	Chimaera IV




Figure 3. Location of ROV survey transects at Kent Island, British Columbia.

Table 5. Invertebrate, fish and algae species observed during ROV survey at Kent Island, British Columbia.

Common Name	Latin Name
<b><u>Invertebrates</u></b>	
California Sea Cucumber	<i>Parastichopus californicus</i>
Unidentified White Cucumber on sand	Possibly <i>Pentamara</i> sp.
Pedal Cucumber	<i>Psolus chitonoides</i>
Swimming Scallop	<i>Chlamys</i> sp.
Oregon Triton	<i>Fusitriton oregonensis</i>
Red Flabellina Nudibranch	<i>Flabellina triophina</i>
Red Octopus	<i>Octopus rubescens</i>
Pink Pisaster	<i>Pisaster brevispinus</i>
Painted Star	<i>Orthasterias koehleri</i>
Striped Sunstar	<i>Solaster stimpsoni</i>
Morning Sunstar	<i>Solaster dawsoni</i>
Blood Star	<i>Henricia leviuscula</i>
Vermillion Star	<i>Mediaster aequalis</i>
Rose Star	<i>Crossaster papposus</i>
Wrinkled Star	<i>Pteraster militaris</i>
Brittle Star	Unidentified
Sun Star	<i>Solaster</i> sp.
Red Sea Urchin	<i>Strongylocentrotus franciscanus</i>
Calcareous Tube Worms	<i>Serpula</i> sp.
Unidentified Sediment Tube Worms	
Snakelock Anemone	<i>Cribrinopsis fernaldi</i>
Giant Plumose Anemone	<i>Metridium giganteum</i>
Sea Pen	<i>Ptilosarcus gurneyi</i>
Orange Cup Coral	<i>Balanophyllia elegans</i>
Staghorn Bryozoan	<i>Heteropora magna</i>
Unidentified Hydrozoans/Bryozoans	
Sea Peach	<i>Halocynthia aurantium</i>
Sea Squirts	Various unidentified
Hexactinellid sponge	
Yellow Boring Sponge	<i>Cliona celata</i>
Rough Scallop Sponge	<i>Myxilla incrustans</i>
Unidentified White Finger-like Sponge	
Unidentified Stalked Trumpet-like Sponge	Possibly <i>Stylissa</i>
Sea Spider	<i>Pycnogonida</i>
<b><u>Fish</u></b>	
Kelp Greenling	<i>Hexagrammos decagrammus</i>
Cod	<i>Gadidae</i>
Yellowtail Rockfish	<i>Sebastes flavidus</i>
Quillback Rockfish	<i>Sebastes maliger</i>
Tiger Rockfish	<i>Sebastes negrocinctus</i>
Rat Fish	<i>Hydrolagus collieri</i>
Rock Sole	<i>Lepidopsetta bilineata</i>
Unidentified Rockfish	Most likely Sharpchin
<b><u>Algae</u></b>	
Encrusting Coralline Algae	Unidentified

Table 6. Description of grab sampling information at both farm (S, KI) and reference (RF) stations, Kent Island, British Columbia.

Station	Date	GPS Location	Grab Type	Depth (m)	Time (PDT)	General Observations
S-9	Sept 13, 2002	N 50° 53.933' W 127° 32.447'	Ponar	71	13:51	Light odour, small sample, some dark surface area, red worms
S-10	Sept 13, 2002	N 50° 53.950' W 127° 32.451'	Ponar	84	12:06	Moderate odour, black surface layer, red worms, small sample
S-17	Sept 12, 2002	N 50° 53.953' W 127° 32.376'	Ponar	84	9:21	Strong odour, black surface layer, red worms
KI-1	Oct 1, 2002	N 50° 53.975' W 127° 32.317'	Van Veen	65	8:49	Shell hash, moderate odour, brown colour, gas bubbles
KI-3	Oct 1, 2002	N 50° 53.943' W 127° 32.306'	Van Veen	39	13:10	No odour, worm tubes
KI-5	Oct 1, 2002	N 50° 53.935' W 127° 32.381'	Van Veen	71	12:26	Strong odour, black colour, gritty texture, dead clams
KI-6	Oct 1, 2002	N 50° 53.931' W 127° 32.405'	Van Veen	77	12:05	Strong odour, black surface, gritty texture, dead clams
KI-7	Oct 1, 2002	N 50° 53.928' W 127° 32.433'	Van Veen	75	11:49	Moderate odour
KI-8	Oct 1, 2002	N 50° 53.936' W 127° 32.443'	Van Veen	85	11:05	N/A
KI-9	Oct 1, 2002	N 50° 53.959' W 127° 32.442'	Van Veen	87	10:41	N/A
KI-10	Oct 1, 2002	N 50° 53.963' W 127° 32.414'	Van Veen	79	10:12	N/A
KI-11	Oct 1, 2002	N 50° 53.968' W 127° 32.376'	Van Veen	72	9:42	Rocks, little shell hash, small crab
KI-12	Oct 1, 2002	N 50° 53.971' W 127° 32.344'	Van Veen	67	9:15	Brown colour, moderate odour, shell hash, worm tubes
RF-1	Oct 1, 2002	N 50° 54.123' W 127° 30.973'	Van Veen	68	14:43	Brown colour, no odour
RF-2	Oct 1, 2002	N 50° 54.144' W 127° 30.914'	Van Veen	69	15:12	Brown colour, no odour
RF-3	Oct 1, 2002	N 50° 54.092' W 127° 30.958'	Van Veen	60	15:26	Brown colour, no odour

Table 7. Sediment grain size fractions (Ocean Dumping and Ocean Dumping Extra Points Protocols) of grab samples collected at Kent Island, British Columbia.

Sampling Station															
Ocean Dumping Protocol	S-9	S-10	S-17	KI-1	KI-3	KI-6	KI-7	KI-8	KI-9	KI-10	KI-11	KI-12	RF-1	RF-2	RF-3
Gravel, >2 mm	0.82	2.12	3.34	17.77	2.91	9.00	45.68	0.70	0.35	0.94	12.04	18.28	9.27	0.37	0.27
Sand, <2.00 mm >0.063 mm	87.92	86.93	79.26	48.00	76.12	76.06	48.47	90.05	91.01	89.92	70.91	47.59	61.93	93.92	93.32
Silt, <0.063 mm >0.004 mm	5.49	4.78	7.39	13.43	7.22	6.01	2.41	3.68	3.42	3.75	6.48	13.20	10.53	1.87	2.28
Clay, <0.004 mm	5.77	6.17	10.01	20.80	13.75	8.93	3.44	5.57	5.23	5.50	10.56	20.94	18.28	3.85	4.13

Sampling Station															
Ocean Dumping Extra Points Protocol	S-9	S-10	S-17	KI-1	KI-3	KI-6	KI-7	KI-8	KI-9	KI-10	KI-11	KI-12	RF-1	RF-2	RF-3
Gravel, >2 mm	0.82	2.12	3.34	17.77	2.91	9.00	45.68	0.70	0.35	0.94	12.04	18.28	9.27	0.37	0.27
<2.00 mm	0.10	0.06	0.41	5.58	1.41	1.18	4.23	0.08	0.10	0.05	0.52	2.85	1.63	0.04	0.01
<1.00 mm	0.23	0.11	0.68	3.75	2.15	1.89	0.26	0.21	0.15	0.15	0.68	2.44	1.49	0.06	0.07
<0.500 mm,	2.42	1.36	3.50	4.31	8.12	6.79	3.87	2.79	1.74	1.84	3.06	3.56	2.72	0.69	1.03
<0.250 mm	48.13	43.42	40.51	12.37	38.07	38.71	27.30	51.08	48.49	48.90	34.28	15.71	18.50	48.06	44.71
<0.125 mm	37.04	41.98	34.16	21.99	26.37	27.49	17.05	35.89	40.52	38.87	32.37	23.03	37.59	45.07	47.49
<0.063 mm	5.49	4.78	7.39	13.43	7.22	6.01	2.40	3.68	3.42	3.75	6.49	13.19	10.52	1.86	2.29
<0.004 mm	0.97	1.05	1.93	3.89	2.78	2.04	0.71	0.91	0.80	0.87	2.12	3.85	3.09	0.45	0.55
<0.002 mm	4.80	5.12	8.08	16.91	10.97	6.89	2.73	4.66	4.43	4.63	8.44	17.09	15.19	3.40	3.58






Figure 5. Approximate locations of grab sampling stations relative to the netpen layout at Kent Island, British Columbia.




Figure 6. Sediment grain size fractions observed at farm sampling stations at Kent Island, British Columbia.




Figure 7. Sediment grain size fractions observed at reference stations at Kent Island, British Columbia.

Table 8. Sediment total carbon and total nitrogen contents observed at farm and reference stations at Kent Island, British Columbia.

Sampling Station	Percent Total Carbon	Total Nitrogen
	(%)	(µg g <sup>-1</sup> )
S-9	2.04	761
S-10	1.90	637
S-17	4.09	1273
KI-1	7.93	2360
KI-3	4.26	1772
KI-5	5.12	1552
KI-6	4.19	1895
KI-7	3.25	776
KI-8	1.92	689
KI-9	2.43	758
KI-10	1.83	876
KI-11	4.26	1222
KI-12	7.96	2254
RF-1	3.05	1636
RF-2	0.68	641
RF-3	0.55	590

Table 9. Sediment trace-metal concentrations observed at farm and reference stations at Kent Island, British Columbia.

Element	Stations Sampled															
	S-9	S-10	S-17	KI-1	KI-3	KI-5	KI-6	KI-7	KI-8	KI-9	KI-10	KI-11	KI-12	RF-1	RF-2	RF-3
	(μg g <sup>-1</sup> )															
Aluminum	16667	16425	18626	20593	16442	18208	14826	15780	14573	14600	12470	16711	17852	13757	11062	12619
Antimony	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Arsenic	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Barium	26.7	29.0	30.4	61.6	35.7	37.9	25.9	22.6	22.2	21.3	18.5	32.6	48.9	33.6	17.8	20.6
Beryllium	0.4	0.5	0.4	0.3	0.3	0.3	0.3	0.4	0.5	0.5	0.5	0.5	0.4	0.5	0.5	0.5
Boron	0	0	0	10	2	0	0	0	0	0	0	0	8	7	0	0
Cadmium	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Calcium	35262	25271	41782	80506	52301	49968	44621	30059	27811	27176	24641	33600	68427	35008	13918	15245
Chromium	19.0	18.9	22.0	24.0	18.8	21.0	18.3	19.2	18.3	18.7	15.9	19.7	22.5	16.8	14.0	14.6
Cobalt	8.4	8.5	9.1	7.9	7.5	7.9	7.3	8.5	8.4	8.3	7.8	8.4	7.8	6.9	6.8	7.3
Copper	11.3	13.7	14.4	13.8	18.5	12.0	11.7	11.2	10.6	10.4	10.7	12.9	15.9	10.6	8.2	8.3
Iron	19687	19551	23247	21076	18149	20736	18811	20789	19059	19127	16608	19665	20111	15586	13770	14988.0
Lead	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Magnesium	6237	5928	7292	9666	7555	7479	6808	6261	5978	5872	5467	6515	8068	6028	4541	4855
Manganese	274.6	279.1	300.6	257.7	221.2	271.8	218.5	260.1	245.5	246.6	215.5	264.0	244.2	211.2	191.8	218.2
Molybdenum	0	0	3	0	2	3	3	0	0	0	0	0	0	0	0	0
Nickel	13	12	13	18	13	13	12	12	12	13	12	14	16	12	10	10
Phosphorus	720	701	1848	1106	1365	1518	2913	1052	746	1461	834	874	959	640	576	590
Potassium	1489	1419	1793	3455	2134	2246	1703	1329	1299	1264	1086	1769	2656	1869	1053	1176
Selenium	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Silicon	609	656	478	661	469	541	633	672	702	865	631	897	620	771	1760	756
Silver	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Sodium	5038	3937	6483	17569	10246	8768	8465	4933	5432	5122	4497	5502	8956	7811	4734	4606
Strontium	236.5	164.4	317.1	852.8	469.0	441.7	343.0	204.5	178.3	169.6	148.6	241.5	609.9	285.2	86.3	97.6
Sulfur	1306	1203	2196	3677	2429	2706	2765	1560	1332	1410	1303	1628	2992	1904	1080	1112
Tin	0	0	0	0	0	0	0	0	0	0	0	8	0	0	0	0
Titanium	2293.6	2313.7	2439.4	1891.3	1775.5	1993.8	1758.8	2286.3	2224.7	2197.9	1970.5	2173.2	1898.8	1764.6	1803.5	1999.0
Vanadium	77	77	92	75	65	76	69	82	76	77	66	79	75	59	54	60
Zinc	35.4	34.0	50.0	58.4	49.2	52.5	53.2	37.1	35.2	36.2	34.4	41.3	52.9	37.7	25.7	27.5

Table 10. Detection limits for sediment trace-metal analysis conducted on an Optima 4300 inductively coupled plasma emission spectrometer. (PESC. SEDMET Method V 6.0).

Element	Detection Limit ( $\mu\text{g g}^{-1}$ )
<b>(ICP Total)</b>	
Aluminum	8
Antimony	8
Arsenic	8
Barium	0.2
Beryllium	0.2
Boron	2
Cadmium	0.8
Calcium	20
Chromium	0.8
Cobalt	0.8
Copper	0.8
Iron	0.8
Lead	8
Magnesium	20
Manganese	0.2
Molybdenum	2
Nickel	3
Phosphorus	20
Potassium	20
Selenium	8
Silicon	8
Silver	2
Sodium	20
Strontium	0.2
Sulfur	8
Tin	8
Titanium	0.3
Vanadium	2
Zinc	0.3




Figure 8. Sediment minor metals observed at farm sampling stations at Kent Island, British Columbia.




Figure 9. Sediment minor metals observed at reference stations at Kent Island, British Columbia.



Table 11. Benthic debris descriptions observed at farm and reference sampling stations at Kent Island, British Columbia.

Station	Benthic Debris Description
S-9	Worm tube fragments, woody debris (to 9.0 cm), shell fragments (to 4.5 cm), organic debris, urchin spine fragments, foraminiferan tests, and gravel (to 2.5 cm). 150 mL
S-10	Worm tube fragments, woody debris (to 1.5 cm; <5% charcoal), organic debris, shell fragments (to 3.0 cm), foraminiferan tests, and algae. 100 mL
S-17	Worm tube fragments, shell fragments (to 3.0 cm), organic debris, woody debris (to 3.5 cm; <5% charcoal), and urchin spine fragments. 400 mL
KI-1	Shell fragments (to 4.5 cm), organic debris, wood fragments (to 3.0 cm), urchin spine fragments, sand, foraminiferan tests, and gravel (to 2.0 cm). 1450 mL
KI-3	Shell fragments (to 6.5 cm), worm tube fragments (including <i>Phyllochaetopterus</i> tubes), sand, foraminiferan tests, bryozoan tests, urchin spine fragments, organic debris, woody debris (to 4.0 cm), and algae. 1450 mL
KI-5	Shell fragments (to 2.5 cm), worm tube fragments, foraminiferan tests, sand, organic debris, woody debris (to 2.0 cm), urchin spine fragments, bryozoan tests, algae, and gravel (to 1.0 cm). 1300 mL
KI-6	Shell fragments (to 3.0 cm), worm tube fragments, woody debris (to 3.0 cm; <25% charcoal), organic debris, foraminiferan tests, gravel (to 2.0 cm), sand, and algae. 900 mL
KI-7	Shell fragments (to 2.5 cm), organic debris, worm tube fragments, woody debris (to 2.5 cm; 10% charcoal), foraminiferan tests, sand, and algae. 550 mL
KI-8	Organic debris, woody debris (to 1.5 cm; <5% charcoal), worm tube fragments, and shell fragments (to 3.5 cm). 350 mL
KI-9	Organic debris, worm tube fragments, woody debris (to 2.5 cm; <5% charcoal), and shell debris (to 4.0 cm). 400 mL
KI-10	Worm tube fragments, shell fragments (to 4.0 cm), wood fragments (to 2.0 cm; <5% charcoal), organic debris, foraminiferan tests, and urchin spine fragments. 400 mL
KI-11	Shell fragments (to 2.5 cm), worm tube fragments, woody debris (to 5.5 cm; <5% charcoal), organic debris, urchin spine fragments, gravel (to 2.5 cm), and algae. 550 mL
KI-12	Shell fragments (to 4.0 cm), organic debris, woody debris (to 2.5 cm; 5% charcoal), foraminiferan tests, urchin spine fragments, worm tube fragments, bryozoan tests, sand, and gravel (to 2.0 cm). 1650 mL
RF-1	Shell fragments (to 4.5 cm), worm tube fragments, organic debris, woody debris (to 1.5 cm), bryozoan tests, urchin spine fragments, sand, foraminiferan tests, and gravel (to 1.5 cm). 750 mL
RF-2	Worm tube fragments, shell fragments (to 5.0 cm), woody debris (to 2.0 cm; <5% charcoal), organic debris, and foraminiferan tests. 100 mL
RF-3	Worm tube fragments, shell fragments (to 4.5 cm), woody fragments (to 2.0 cm; <5% charcoal), and ophiuroid fragments. 100 mL

Table 12. Macrofaunal abundance observed at farm and reference stations at Kent Island, British Columbia.

Taxon	Sampling Stations															
	S-9	S-10	S-17	KI-1	KI-3	KI-5	KI-6	KI-7	KI-8	KI-9	KI-10	KI-11	KI-12	RF-1	RF-2	RF-3
	(Organisms m <sup>-2</sup> )															
PORIFERA	25	25	275	1100	190	0	0	10	50	0	0	10	0	10	30	60
CNIDARIA																
Hydrozoa	25	300	75	380	40	80	80	150	0	10	20	110	40	10	190	450
PLATYHELMINTHES	0	0	0	0	0	0	0	20	0	0	0	0	0	0	0	0
NEMERTEA	0	75	0	570	30	0	0	0	50	20	10	100	80	30	30	80
NEMATODA	25	300	175	3540	540	70	10	250	40	50	50	170	30	1250	310	170
ANNELIDA																
Polychaeta	3725	8900	5875	9100	3590	9390	16000	12310	6270	5650	3560	3790	2890	2150	3330	3510
Oligochaeta	25	0	75	50	10	10	0	20	0	0	0	0	30	0	0	0
SIPUNCULA	0	0	0	0	0	0	0	0	0	0	0	0	0	10	0	10
MOLLUSCA																
Polyplacophora	0	0	0	100	0	0	0	0	0	0	0	10	50	40	10	0
Gastropoda	25	0	75	300	30	100	20	30	40	10	0	50	100	190	310	200
Bivalvia	75	100	0	60	80	0	0	30	80	90	30	100	20	220	430	380
Scaphopoda	0	0	0	0	0	0	0	0	0	0	0	0	0	0	30	20
ARTHROPODA																
Pycnogonida	0	0	0	0	0	0	0	0	0	0	0	0	0	0	10	20
Arachnida (Acarida)	0	0	0	60	0	0	0	0	0	0	0	0	0	0	0	0
Insecta	0	0	0	10	0	0	0	0	0	0	0	0	0	0	0	0
Crustacea	50	150	125	2650	200	360	30	60	110	30	30	60	40	380	530	410
ENTOPROCTA	0	0	0	40	0	0	0	10	0	0	0	0	0	0	0	10
BRYOZOA	0	25	1250	350	0	20	0	130	20	30	20	100	40	110	120	10
ECHINODERMATA																
Ophiuroidea	0	0	0	30	0	0	0	10	10	0	10	20	0	50	210	60
Holothuroidea	0	0	0	20	0	0	0	0	0	0	0	0	0	90	50	60
UROCHORDATA																
Ascidacea	0	0	0	0	0	0	0	0	0	0	0	20	0	0	0	0
Unidentified eggs and larvae	0	50	0	560	10	0	0	0	430	0	0	0	0	0	10	100
Total organisms m <sup>-2</sup>	3975	9875	7925	18360	4710	10030	16060	13030	6670	5890	3730	4540	3320	4540	5590	5450
Total number of taxa	8	8	8	16	9	7	4	12	9	8	8	12	10	13	14	15




Figure 10. Macrofaunal abundance (annelids excluded) found at farm sampling stations at Kent Island, British Columbia. \*Note: stations S-17 and KI-1 standardized to 4000 to accomodate larger range.




Figure 11. Macrofaunal abundance (annelids excluded) found at reference stations at Kent Island, British Columbia.