

Adult Chinook Escapement Assessment Conducted on the Cowichan River During 2001

N.K. Hop-Wo, D.A. Nagtegaal, E.W. Carter, and K. Jones

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, British Columbia
V9T 6N7

2003

**Canadian Manuscript Report of
Fisheries and Aquatic Sciences 2645**



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Canada

Canadian Manuscript Report of Fisheries and Aquatic Sciences

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 1426 - 1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport manuscrit canadien des sciences halieutiques et aquatiques

Les rapports manuscrits contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui traitent des problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Manuscript Report of
Fisheries and Aquatic Sciences 2645

2003

ADULT CHINOOK ESCAPEMENT ASSESSMENT CONDUCTED ON THE
COWICHAN RIVER DURING 2001

by

N.K. Hop-Wo, D. A. Nagtegaal, E. W. Carter, and K. Jones

Fisheries and Oceans Canada

Sciences Branch, Pacific Region

Pacific Biological Station

Nanaimo, British Columbia

V9T 6N7

©Her Majesty the Queen in Right of Canada, 2003,
as represented by the Minister of Fisheries and Oceans
Cat. No. Fs 97-4/2645E ISSN 0706-6473

Correct citation for this publication:

Hop-Wo, N.K., D.A. Nagtegaal, E.W. Carter, and K. Jones. 2003. Adult chinook escapement assessment conducted on the Cowichan River during 2001. Can. Manuscr. Rep. Fish. Aquat. Sci. 2645: 45 p.

List of Tables

Table 1.	Daily counts at the Cowichan River fence site, 2001 ..	16
Table 2.	Counts, by time interval, at the Cowichan River fence, 2001	17
Table 3.	Visual survey data collected for the Cowichan River for the years 1981 to 2001	18
Table 4.	Estimated catch of chinook adults and jacks in the Cowichan River Native food fishery for the years 1981 to 2001	22
Table 5.	Summary, by day and location, of chinook broodstock collected by Cowichan River hatchery staff during 2001	23
Table 6.	Annual hatchery broodstock collection of adult and jack Cowichan River chinook for the years 1981 to 2001	24
Table 7.	Summary of age, by sex, for Cowichan River chinook collected for hatchery broodstock during 2001	25
Table 8.	Length-frequency of chinook sampled on the Cowichan River spawning grounds, 2001	25
Table 9.	Summary of age, by sex, for Cowichan River chinook collected from the spawning grounds during 2001	27
Table 10.	Length-frequency of chinook hatchery broodstock collected during 2001	27
Table 11.	Release and recovery data for coded wire tags recovered from Cowichan River chinook sampled on the spawning grounds during 2001	28
Table 12.	Cowichan River hatchery chinook release data for the years 1979 to 2001 ..	29
Table 13.	Cowichan River daily discharge (c ³ /s) during 2001 ..	33
Table 14a.	Daily summary of carcasses examined, tags applied and tagged recoveries, by sex, for chinook in the upper Cowichan River, 2001	34
Table 14b.	Tags applied carcasses examined and marks recovered, by sex, for chinook in the upper Cowichan River, 2001	34
Table 15.	Petersen mark recapture estimates, stratified by sex, for Cowichan River chinook, 2001	35
Table 16.	Incidence of tagged adult chinook carcasses recovered on the spawning grounds by recovery period and sex, on the Cowichan River, 2001	35
Table 17.	Proportion of the tag application sample recovered on the spawning grounds, by application period and sex, in the Cowichan River, 2001	36
Table 18.	Sex composition of chinook in the tag application and recovery samples on the upper Cowichan River, 2001	36

Table 19. Sex composition of chinook in the tag application and recovery samples on the upper Cowichan River, 2001 (jacks included).....	36
Table 20. Total adult chinook returns to the Cowichan River for the years 1975 to 2001.....	37

List of Figures

Figure 1. Cowichan River study area.....	39
Figure 2. River management zones for the Native food fishery	41
Figure 3a. Daily fence counts of adult and jack chinook and water temperature at the fence site	42
Figure 3b. Daily fence counts of adult and jack chinook and water depth at the fence site	42
Figure 4. Adult chinook catch in the Cowichan First Nations fishery for the years 1971 to 2001	43
Figure 5. Monthly Cowichan River discharge (m^3/s) in 2001 along with the historical minimum, maximum and mean values	43
Figure 6. Adult and jack chinook length frequencies collected from the Cowichan River spawning grounds, 2001.....	44
Figure 7. Annual adult chinook escapement estimates for the Cowichan River for the years 1953 to 2001	44
Figure 8. Annual releases of hatchery chinook into the Cowichan River as fry (3 g) and as pre-smolts (6 g)	45
Figure 9. Natural and enhanced contribution to the annual Cowichan River chinook escapement for the years 1982 to 2001	45

ABSTRACT

Hop-Wo, N.K., D.A. Nagtegaal, E.W. Carter, and K. Jones. 2003. Adult chinook escapement assessment conducted on the Cowichan River during 2001. Can. Manuscr. Rep. Fish. Aquat. Sci. 2645: 45 p.

In 2001, the Biological Sciences Branch, Pacific Biological Station, continued a study of chinook salmon (*Oncorhynchus tshawytscha*) productivity in the Cowichan River. This in-depth escapement assessment project has been in place since 1988. Major components of this study included: i) enumerating spawners and total return, ii) estimating Native food fish catch, iii) recording hatchery broodstock removals, and iv) collecting biological, environmental and coded-wire tag data. Population estimates for adult and jack chinook were determined based on the fence count data since this was considered to be the most accurate enumeration method. A carcass mark-recapture study was conducted on the spawning grounds to augment the collection of biological data and to supplement the fence count population estimate. The total return of adult chinook to the Cowichan River was estimated to be 5,998 fish of which 3,282 spawned naturally in the river. A total of 1,732 adult chinook were collected for hatchery broodstock and 918 adults were reported as captured in the Native food fishery.

RÉSUMÉ

Hop-Wo, N.K., D.A. Nagtegaal, E.W. Carter, and K. Jones. 2003. Adult chinook escapement assessment conducted on the Cowichan River during 2001. Can. Manuscr. Rep. Fish. Aquat. Sci. 2645: 45 p.

En 2001, la Direction des sciences biologiques, à la Station biologique du Pacifique, a poursuivi son étude sur la productivité du saumon quinnat (*Oncorhynchus tshawytscha*) dans la rivière Cowichan. Ce projet d'évaluation approfondie des échappées est en cours depuis 1988. Ses principaux volets sont les suivants: i) dénombrement des géniteurs et des remontes totales, ii) estimation des captures à des fins alimentaires par les Autochtones, iii) consignation du nombre de géniteurs prélevés pour les écloséries et iv) collecte de données sur la biologie, l'environnement et les micromarques codées. On a estimé la population d'adultes et de saumons d'un an en mer à partir des données sur le dénombrement à la barrière, cette méthode étant considérée comme la plus exacte. On a procédé à la récupération des micromarques sur les carcasses dans la fraye afin d'enrichir les données biologiques recueillies et de compléter l'estimation de la population dénombrée à la barrière. La remonte totale de quinnats adultes dans la rivière Cowichan a été estimée à 5 998 unités, dont 3 232 qui s'y sont reproduites naturellement. Au total, 1 732 adultes ont été recueillis pour servir de géniteurs dans des écloséries, tandis que 918 adultes auraient été capturés dans le cadre de la pêche à des fins alimentaires par les Autochtones.

INTRODUCTION

Chinook stocks are invaluable to both commercial and recreational fisheries of the Pacific Northwest (Collicut and Shardlow 1995). In spite of protective measures, chinook salmon abundance has continued to decline. This trend has resulted in the recent addition of chinook to the list of threatened and endangered species in the United States (Waples 1991). The problem of declining stocks is similarly serious on the West Coast of Canada, and has potential ramifications regarding the sustainability of British Columbia's fishing industry (Argue *et al* 1983). Over the past several years, considerable interest has been focused on the chinook stocks of the southern portion of the Strait of Georgia due to the decline in these stocks and their importance to local fisheries (Farlinger *et al.* 1990). The Stock Assessment Division, Pacific Biological Station, initiated a study of chinook productivity to assess rebuilding strategies and to evaluate the effects of harvest management policies for these stocks. In the fall of 1988, a study was implemented on Cowichan River chinook with additional information collected from the Squamish and Nanaimo River stocks. These three stocks were identified as exploitation and escapement indicators and deemed to represent the status of all lower Georgia Strait chinook stocks. Since then, due to logistical reasons the Squamish River system was dropped as an indicator.

Major hatchery production of chinook on the Cowichan River began in 1980 (Cross *et al.* 1991). Chinook fry releases have increased from 64,681 in 1980 to 2.41 million in 2001. Coded wire tag releases also began in 1980 and by 2000 approximately 9 % of the chinook released carried coded wire tags.

This report presents the results of the study completed during 2001. The objectives included:

1. enumerating chinook, coho and chum salmon migrating past the counting fence,
2. estimating the Native food fishery catch,
3. recording hatchery broodstock removals,
4. collecting biological data and sampling coded-wire tag (CWT) recoveries, and
5. implementing a carcass mark-recapture study for both adult and jack chinook.

METHODOLOGY

A detailed description of the methodology is presented in Nagtegaal *et al.* (1994b). A summary of the methods is presented below along with any changes that were incorporated during 2001.

STUDY AREA

The Cowichan River watershed is located on the Southeast coast of Vancouver Island and drains an area totalling 826 km². The Cowichan River system includes Cowichan, Bear, Mesachie, Somenos, and Quamichan lakes. Cowichan Lake (62 km²), the largest of the five lakes, is situated approximately 50 km west of the Cowichan Bay estuary. Discharge from a flow

control dam situated at the outlet of Cowichan Lake ranges from 7 to 326 m³·s⁻¹, and averages 44.9 m³·s⁻¹ (Feilden and Holtby 1987). A total of 26 tributaries drain into the Cowichan River. The largest of these is the Koksilah River, which intersects the mainstem of the Cowichan River approximately 2.5 km upstream of the estuary. The Cowichan River watershed system is a typical Vancouver Island and coastal British Columbia stream in which maximum flows occur during winter months due to heavy rainfall (McDougall 1985).

The Cowichan River supports many salmonid species including chinook (*Oncorhynchus tshawytscha*), coho (*Oncorhynchus kisutch*), chum (*Oncorhynchus keta*), sockeye (*Oncorhynchus nerka*), and pink (*Oncorhynchus gorbuscha*) salmon; as well as cutthroat trout (*Salmo clarkii*), steelhead trout (*Salmo gairdnerii*), kokanee salmon (*Oncorhynchus nerka*), and dolly varden char (*Salvelinus malma*). Attempts have been made to introduce several other species including: atlantic salmon (*Salmo salar*), brown trout (*Salmo trutta*), and speckled char (*Salvelinus fontinalis*) (Perrin *et al.* 1988). The salmonids of the Cowichan River support several vital fisheries, which include a Native food fishery, tidal sport fishery, and a large commercial ocean fishery.

ENUMERATION FENCE

The counting fence was placed upstream of the city of Duncan, in the same location as in previous years (Figure 1). The design incorporated a resistance board weir with a counting raceway (adjustable flashboard) and trap box adjacent to a counting tower equipped with floodlights. Counts were continuously recorded by 15-minute interval for adult and jack chinook, adult and jack coho, and chum salmon. If identification was in doubt fish were recorded as unknown. Water depth, temperature, and clarity, and weather condition were recorded three times per day. The fence was checked daily for any breeches and cleaned of leaves and other debris. Any removals of broodstock at the fence site by hatchery staff were recorded.

SWIM SURVEYS

In an effort to maintain consistency with historical data sets, swim surveys were conducted, in conjunction with Cowichan Tribes Aboriginal Fisheries Management (CTAF), to estimate the spawning population of chinook. Swims were generally made in the upper section of the river (Figure 1) with counts extrapolated to the total system. Each survey was conducted by three experienced swimmers while one person in a canoe recorded the data. Each swimmer (one in the middle and one on each side of the river) counted the fish seen within their range of visibility. The three swimmers attempted to keep abreast as they approached each pool while the person in the canoe lagged behind within hailing distance. Counts were recorded by pool/riffle and then compiled by river section. To maintain consistency in counting procedures the same swim team was used for each survey whenever possible. Based on the historical distribution of spawners, swim counts were expanded by a factor of 3.4 to derive an estimate of total escapement (Nagtegaal *et al.* 1994a). This expansion factor was consistently applied to swim counts with no adjustments made for run timing or changes to spawner distribution. A final escapement estimate was then

determined based on consultation with Fishery Officers using the swim counts in conjunction with other anecdotal information.

It was intended that the swim survey estimates remain independent of the fence count. However, even though fence counts were not passed on to the swim teams during the season, general trends in escapement numbers were known.

NATIVE FOOD FISHERY

In 1990, a systematic approach was developed by the Cowichan Tribes Aboriginal Fisheries Management (CTAF) program to monitor the fishery more closely and to better estimate the Native food fish catch (Paige 1992, 1997). This approach involved recording catch and effort by management zone within the Native fishing boundaries (Figure 2). A crew of four observers patrolled the fishery on a daily basis and interviewed fishermen for numbers caught by area and total time spent fishing. In this way, weekly estimates of catch per unit effort (CPUE) were obtained. CPUE was adjusted for daily changes in fishing effort and differences in effort among fishing zones. These data were then extrapolated over time and area to estimate total catch by week and summed over all weeks to estimate the total catch for 2001.

$$CATCH \equiv \sum_n^{w=1} CPUE_w \times EFFORT_d$$

where w refers to the time interval for catch (week), and d refers to the time interval for effort (day). No confidence limits were calculated (Paige 1997).

For some years since 1988, an observer was employed to independently collect catch and biological data from the in-river Native fishery. Due to budget constraints no observers were employed from 1999 - 2001 and CTAF catch estimates could not be independently verified.

BIOLOGICAL DATA

Biological data for chinook were collected from hatchery broodstock samples and from carcasses recovered on the spawning grounds. Data collected included sex, post-orbital hypural (POH) length, and adipose fin status (present/absent). Scales were also taken for age analysis and the heads were removed from all adipose fin clipped fish for recovery and decoding of coded-wire tags (CWT). Hatchery staff randomly collected biological data from approximately 25% of the chinook broodstock and from all chinook identified as carrying a CWT. All chinook recovered on the spawning grounds were sampled and spawning condition was noted.

MARK-RECAPTURE

A mark-recapture program involving the tagging and subsequent recovery of chinook jack and adult carcasses was conducted on the spawning grounds. Crews consisting of two or three individuals surveyed the upper section of the river each day using an inflatable boat (Figure 1). All

chinook carcasses encountered were individually tagged with a Ketchum¹ aluminum sheep ear tag on the left operculum and immediately released in the same area as captured. Location of capture and release, tag number, spawning condition, POH length, sex, and adipose fin status (present or absent) were recorded for each carcass. Tag number and recovery location were recorded for all previously marked carcasses, which were then returned to the river at the capture site.

The section of the river selected for the mark recapture study is located above Skutz Falls and represents the area where the majority of chinook spawning has typically occurred. A 4.2 m pole with a gaff hook attached to the end was used to recover carcasses. Carcasses that ended up in deep pools were occasionally unable to be retrieved.

POPULATION ESTIMATE

Adult chinook salmon escapement estimates were generated from the carcass mark-recapture data using the Petersen model (Chapman modification) stratified by sex (Ricker 1975). Data was stratified to minimize the effects of differential tagging and tag recovery between sexes. This study follows the estimation procedure as outlined in previous reports (Nagtegaal *et al.* 1994a, 1994b, 1994c).

To determine the validity of the mark recapture estimate, potential biases were assessed to test for violations of the assumptions inherent in the Petersen mark recapture method (Ricker 1975; Seber 1982; Krebs 1989). Sex related bias in the application sample was examined by comparing the sex ratio of the marked versus the unmarked recoveries. Sex related bias in the recovery sample was assessed by comparing the sex ratio of recovered samples with unrecovered samples. Size bias in the application sample was examined by comparing the lengths of marked versus unmarked recoveries. Size bias in the recovery sample was examined by comparing the lengths of the recovered and unrecovered tag samples. Statistical comparisons of the length data were carried out using pooled variance Student's t-test. Temporal bias in the application samples was assessed by stratifying sampling dates into approximately equal periods and comparing mark incidence among strata. Similarly, temporal bias in recovery samples was stratified by dates and the ratio of tagged recoveries to tags applied was compared. Statistical assessments of sex and period biases were carried out using Chi-square tests.

RESULTS

ENUMERATION FENCE

The enumeration fence was installed upstream from the City of Duncan at the traditional site (Figure 1) and was operational from September 4th to November 1st. Water conditions were clear for most of the study with only two days noted as cloudy and nine days as partially cloudy (Table 1). As a result, conditions were ideal for viewing fish migration past the counting tower

¹Ketchum Manufacturing Ltd., Ottawa, Canada.

and all counts are deemed to be reliable. A total of 3,361 chinook adults, 1,454 chinook jacks, 15,436 coho adults and 1,296 coho jacks were counted migrating past the enumeration fence. In addition, 7,859 chum and 243 unidentified fish were enumerated at the fence site (Table 1). Since there were no breeches of the fence during the project it was assumed that all fish migrating past the fence during the study period were enumerated.

In past years, chinook jacks have been prominent during the early segment of the return migration often matching adult chinook counts. In 2001, the proportion of jack chinook mirrored the overall temporal pattern of adult chinook migration past the fence (Table 1, Figures 3a, 3b).

The pattern of daily migration past the fence was examined by summarizing hourly counts throughout the program (Table 2). A major peak in migration for both adult and jack chinook occurred between 0700 and 1000 hrs. During this period, 32.9% of adult chinook and 42.0% of jack chinook moved past the fence site (Table 2).

In past years, tower counts and species identification were verified on several occasions throughout the run by capturing all fish migrating past the fence during one counting segment in the upstream trap box. For logistical reasons, this practise was not carried out in 2001.

SWIM SURVEYS

A summary of all visual surveys conducted by DFO Fishery Officers and by the Cowichan Tribes Aboriginal Fisheries Management group from 1981 to 2001 is presented in Table 3. Total escapement estimates for each year include adult chinook only. In 2001, two swim surveys were conducted in the upper section of the river (Figure 1) and one survey conducted in the lower section of the river (Figure 2).

NATIVE FOOD FISHERY

Historical estimates of the number of chinook captured in the in-river Native food fishery are presented in Table 4 and Figure 4. The 2001 catch estimate of 918 adults and 120 jacks was determined by the Cowichan Tribes Aboriginal Fisheries Management group (Figure 2). While it is difficult to assess the quality of the data collected, the catch estimate is considered to be a minimum. Also, if non-catch mortality (fish that escaped capture but died before spawning due to fishery induced injuries) were included, the impact of the fishery would likely be much more significant.

HATCHERY COMPONENT

Between September 18 and October 3, Cowichan River hatchery staff collected 680 male and 818 female chinook from the river downstream of the fence. In addition, 108 male and 126 female chinook were collected upstream of the fence (Table 5). The total number of chinook removed from the river for hatchery broodstock was within the range required for hatchery

production (Table 6). Age analysis of scale samples revealed that the majority of adult fish (95.6%) used in broodstock were either three or four years old (Table 7).

BIOLOGICAL DATA

A total of 229 male, 463 female and 182 jack chinook carcasses were recovered on the spawning grounds and measured for post orbital-hypural length (Table 8). Conditions for the recovery of carcasses were ideal with discharge levels below the long-term average during October and November (Figure 5). The length of adult male chinook carcasses ranged from 40.9 to 82.5 cm and averaged 63.0 cm, while female carcasses ranged from 40.9 to 82.5 cm and averaged 65.2 cm. Jack chinook carcasses ranged in length from 28.6 to 55.5 cm and averaged 38.6 cm. A total of 11 male, 15 female and three jack chinook carcasses recovered from the spawning grounds were missing an adipose fin indicating the presence of a coded-wire tag (Table 8). These values represent 4.8%, 3.2% and 1.6% of the total male, female and jack carcasses recovered. The majority of adult chinook were three or four years old (54.1% and 41.4%, respectively) while most fish identified as jacks (98.9%) were two years old (Table 9).

A total of 178 male, 217 female and seven jack chinook were randomly collected for broodstock sampling (Table 10). Post orbital-hypural length measurements of male chinook ranged from 48.0 to 78.5 cm and averaged 59.3 cm, female chinook ranged from 49.5 to 83.3 cm and averaged 64.6 cm, and jack chinook ranged in length from 38.0 to 51.0 cm and averaged 44.4 cm. A total of zero male, 16 female (representing 7.4%) and zero jack chinook were missing an adipose fin indicating the presence of a coded wire tag (Table 10).

There was a statistically significant difference between the length of both male and jack chinook carcasses recovered on the spawning grounds and those sampled for hatchery broodstock (Student's t-test: $t = 4.948$; $p < 0.01$ and $t = 3.832$; $p < 0.01$, for males and jacks respectively). In both cases, the broodstock sample was significantly larger than the samples collected on the spawning ground. However, there was no significant difference between the lengths of female chinook sampled on the spawning grounds or sampled at the hatchery (Student's t-test: $t = 1.235$; $p < 0.01$).

A significant difference is evident when comparing the rate of adipose fin clips of chinook sampled from the spawning grounds and chinook sample from hatchery broodstock (Chi-square 8.788; $p = 0.05$ for males and Chi-square = 5.802; $p = 0.05$, for females). Adult male chinook have a higher rate of clipped adipose fins while female chinook have a lower rate of clipped adipose fins when sampled on the spawning grounds compared to broodstock sampling (Table 8, 10). The broodstock sampling size of jack chinook was too small for statistical comparison.

Coded wire tags were recovered from 26 chinook carcasses sampled on the spawning grounds. The majority of these fish (84.6%) were adults from the 1998 brood, of which 59.1% were released as part of the upper Cowichan River (late) release strategy (Table 11). Three CWT recoveries (11.5%) were jack chinook from the 1999 brood year. A summary of all chinook releases from the Cowichan hatchery since 1979 is presented in Table 12.

ENVIRONMENTAL INFORMATION

Environmental data collected at the fence site included water temperature, visibility and river depth (Table 1). Water temperature was generally higher at the start of the study and decreased discontinuously to the end of October. Water temperature over this period ranged from 9.0 to 18.9 °C and averaged 15.3 °C. Water depth at the fence site was consistently low until the last week of September where water levels rose and fell throughout the first three weeks of October. During the last week of October the water level continued to rise until fence removal (Figure 3b). Water depth over the study period ranged from 42.7 to 89.0 cm and averaged 55.0 cm. Water clarity was recorded in the form of a visibility code. For the majority of the study, visibility at the fence site was clear. Only nine days were recorded as moderately cloudy (code 1-2) and two days as cloudy (Table 1).

River discharge was recorded at the Water Survey Canada station below the Island Highway Bridge in the City of Duncan (Table 13, Figure 1). River discharge during the course of the 2001 study was lower than the 30-year average during the September and October but above average during November and December (Figure 5).

CARCASS MARK-RECAPTURE

Between November 5 and December 7 a total of 227 male, 465 female and 181 jack chinook carcasses were tagged and released in the upper Cowichan River (Table 14a, 14b). Of the 163 carcasses recovered with tags, 41 (25.2%) were male, 109 (66.9%) were female, and 13 (8.0%) were jack. Using the Petersen estimator, the adult chinook spawning ground population size was determined to be 3,869 fish (95% CI: 3,311 to 4,427), while the jack population was estimated to be 2,535 fish (95% CI: 1,299 to 3,771) (Table 15).

Potential Biases

The assessment of sampling selectivity had several potential biases in the carcass mark recapture study.

1. Temporal bias: Temporal bias in the tagging sample was examined by stratifying the mark incident rate into four recovery periods (Table 16). There was a significant temporal bias in the application sample for both male and female adult chinook when the data was stratified into four equal recovery periods (Chi-square = 14.86; $p=0.05$ and Chi-square = 21.13; $p=0.05$, for males and females, respectively). Mark incidence increased during each recovery period.

Temporal bias in the recovery sample was analysed by stratifying the recovery rates into four application periods (Table 17). A statistical difference in the recovery sample for both male and female adult chinook was observed (Chi-square = 24.19; $p=0.05$ and Chi-square = 15.77; $p=0.05$, males and females, respectively). The recovery rate increased during each successive application period.

2. Fish Sex: Sex related bias was examined by comparing the sex ratio of the marked and unmarked spawning ground recoveries by application sample and by recovery sample. No sex related bias was evident when comparing male and female chinook populations (Chi-square = 2.60; $p=0.05$ and Chi-square = 1.85; $p=0.05$, application sample and recovery sample, respectively) (Table 18). However, when jack chinook were included in the assessment both the application and recovery samples showed significant bias (Chi-square = 22.76; $p=0.05$, and Chi-square = 16.61; $p=0.05$, application sample and recovery sample, respectively) (Table 19).

3. Size bias: Size related bias was examined by comparing the POH mean lengths of unrecovered marked chinook and recaptured chinook by sex. No size bias was evident in the recovery sample for adult males and jack chinook (Student's t-test: $t=1.481$; $p<0.01$ and $t=1.771$; $p<0.01$ for males and jacks respectively). However, there was a significant size difference in the recovery sample of adult females (Student's t-test: $t=2.658$; $p<0.01$).

POPULATION ESTIMATE

Escapement and total return estimates for 2001 were determined using fence count data since this is considered to be the most accurate enumeration method. However, after reviewing both spawning ground carcass recovery and hatchery broodstock collection data, it became evident that the chinook fence count did not accurately reflect the true jack to adult ratio. Analyzing the lengths of jack and adult chinook with the traditionally accepted jack designated length of 450mm revealed an overlapping of age groups (Figure 6). As a result, the spawning ground carcass recovery data was utilized to apportion the total chinook fence count with a more reflective ratio of jack and adult chinook populations. This was accomplished by comparing age data with length data to calculate the proportion of 2 years old jack chinook with lengths greater than 450mm (9.2%) and the proportion of 3 years old or greater adult chinook with lengths of 450mm or less (0.6%). The fence count data was then adjusted by these proportions yielding total chinook fence counts of 3,057 adults and 1,758 jacks. Since previous studies have indicated that approximately 15% of the chinook run arrives after late October (Nagtegaal and Carter 1998), the total fence count was expanded by this value and the final estimate of adult chinook migration past the fence site was 3,516 fish.

The number of natural spawning adult chinook in the Cowichan River during 2001 was determined to be the fence count, plus an additional 15% estimate for chinook arrival after fence removal (Nagtegaal and Carter 1998), minus any broodstock removals from areas above the fence. Following this methodology, the total number of adult chinook spawning in the Cowichan River was estimated to be 3,282 fish (Table 20).

The total return of adult chinook to the Cowichan River was determined to be the sum of the total fence count with the addition of broodstock removals and the Native fishery catch. A further 300 fish were added to this total to account for chinook lost to seal predation in the Cowichan estuary (see discussion for details). Following this methodology, the total return of adult chinook to the Cowichan River was estimated to be 5,197 fish (Table 20).

DISCUSSION

ENUMERATION FENCE

The floating fence design functioned well in the flow conditions that existed in the lower Cowichan River during the fall of 2001. While the fence was intended to be self-cleaning, field staff were required to regularly remove leafy debris from the fence panels during periods of heavy rain and related high flows. During such periods, fence panels became over burdened and could not remain afloat without the removal of debris. While this difficulty has been observed in other fence studies (Cousens *et al.* 1982, Johnston *et al.* 1986) it is likely that the intensity of the problem varies by year and location.

During many of the past monitoring studies, high flows have caused fence damage or forced early removal resulting in incomplete fence counts. During the current study, the enumeration fence was operational throughout the monitoring period and a complete count of all fish passing the site was attained. However, since the fence was removed on November 1 there was likely a portion of the chinook run that arrived after fence removal. Past studies have indicated that approximately 15% of the chinook run arrives after October 30 (Nagtegaal and Carter 1998). Based on this observation, the adult chinook fence count was expanded by 15% to account for unassessed spawners.

Several of the past fence enumeration studies conducted on the Cowichan River have noted the mis-identification of jack versus adult chinook by observers monitoring the migration of chinook past the fence (Nagtegaal and Carter 1998, 2000). While trap box verifications of observer counts were not carried out in 2001, it became clear from both spawning ground and hatchery broodstock data that the adult to jack ratio recorded at the fence was not reflective of the true population parameters. Since there were a large number of jack chinook at the upper end of the size distribution, it appears as if many of these fish were counted as adults. This seems likely as it would be very difficult to determine the size of a moving fish to within a few centimetres when viewing them from a counting tower well above the river. As a result, the total chinook fence count was adjusted by the ratio of adult and jack chinook sampled during the spawning ground carcass recovery.

SWIM SURVEYS

Among the biases typically associated with swim surveys, the extrapolation of actual swim counts to total estimates warrants some consideration (Burns unpubl.). Assumptions concerning the distribution of chinook in the river at the time of the survey are the basis for expanding these counts to estimate total escapement (T. Fields, 230 Underwood St., Duncan, B.C. V9L-3X3; pers. comm.). In 1991, it became apparent that during high water flow conditions in early fall, expansions based on the swim survey results overestimated total escapement (Nagtegaal *et al.* 1994b, 1994c). The results of the 1992 swim surveys support the hypothesis that during low water flow conditions in late fall, expansions based on swim survey results underestimate the number of spawners. Low flow conditions lead to an underestimation of spawners because the distribution of fish in the river

is affected by water flow. Generally, in low water years, not as many fish make it to the traditional spawning areas above Skutz Falls. The expansion of swim surveys conducted in the upper area alone tends to underestimate the number of fish. Conversely, during high water years most of the fish make it above Skutz Falls so the expansion factor tends to overestimate the number of fish.

NATIVE FOOD FISHERY

Catch estimation procedures developed by the Cowichan Tribes Aboriginal Fisheries Management unit have not been assessed by stock assessment staff. As a result, no comments can be made regarding the methodologies used. The 2001 estimate of 918 adult chinook was a considerable increase over 1999, and 2000 years (Table 4). In past years, independent observer estimates of adult chinook catch have been 2 to 3.5 times higher than the CTAF estimate. Since no observers were employed during 2001, CTAF catch estimates could not be independently verified.

BIOLOGICAL DATA

Male and female chinook collected by the hatchery for broodstock were significantly larger than the carcasses recovered on the spawning ground. This suggests that either one or both sampling methods were size selective or that different size components of the returning chinook population were available for sampling at each site. Since deadpitch carcass recovery is generally selective for larger fish (Tscharplinski and Hyatt 1991) and carcass sampling occurred throughout the spawning period, it seems unlikely that selectivity in this sampling method could explain the result. Also, since broodstock collection occurred over an extended period and beach seines were employed in the capture of fish (a relatively non-selective gear), selectivity seems unlikely. However, since not all chinook captured in each set were taken for broodstock, it may be that the smaller fish were released back into the river. This would result in larger than average fish being included in the broodstock sample while smaller than average fish were returned to the river to contribute to the spawning ground carcass recovery sample. The selective retention of larger broodstock could explain the observed result.

MARK RECAPTURE

The carcass mark-recapture portion of the Cowichan River chinook stock assessment project was implemented many years ago for several reasons. First, since handling fish at the fence site caused significant migration delays this practice was abandoned to ensure chinook reached the spawning grounds without undue stress. As a result, no representative biological samples were available for assessment. While length and age data were collected by hatchery staff during broodstock capture, this data was not consistently representative of the spawning population. The sampling of carcasses from the spawning ground provided an additional source of biological data, which when pooled with the hatchery sample was more representative of the true population. Second, the recovery of coded-wire tags from hatchery broodstock did not provide an adequate sample size for a rigorous assessment. Spawning ground carcass recoveries supplemented the hatchery sample and strengthened the coded-wire tag analysis. Finally, since

high flows in past years have caused fence damage that resulted in incomplete enumeration of spawners, an additional method of providing a population estimate was required.

Population estimates for adult and jack chinook were determined using the pooled Petersen estimator. Since the true population size was not known, a direct measure of the accuracy of the estimates was not possible. However, an assessment of the underlying assumptions of equal probability of capture, simple random recovery sampling and complete mixing can usually be made by testing recovery and application samples for temporal, spatial, sex and size related biases (Schubert 2000). To carry out most of the bias assessments, different gear types must be utilized for capturing the tag application and the recovery samples. In the current study, spawning ground carcass recovery was used to attain both samples thus limiting the ability to assess sample biases.

The assessment of sampling selectivity revealed several biases in the carcass mark-recapture study. First, there was a significant temporal bias in the application sample for both male and female chinook. The assessment revealed that tag incidence was very low during the early period and extremely high during the last period. This is likely due to the nature of the carcass recovery study, since tagging and recovery were concurrent activities. As a result, there were very few tagged carcasses available for recovery in the early period and as the number of tags in the population accumulated towards the end of the study, the tag incidence in the later periods was much higher. There was also a temporal bias in the recovery sample for males and females. Once again this bias was primarily due to the nature of the study since tags applied in later periods were less likely to be recovered as they were only available for recovery for a short period.

No sex related bias was evident in the application or recovery samples for adult chinook; however, when jacks were included in the assessment both samples showed significant bias. This is likely due to the size difference between adult and jack chinook, which effects the way carcasses behave in the river. Since jack carcasses are much smaller, they are more easily flushed out of the recovery area by small increases in river flow. As a result, they are less likely to be recovered than adult chinook carcasses.

Size bias testing did not provide an assessment of the size selectivity of the sampling method since both application and recovery samples were attained using the same method. Rather, the size bias assessment provided an evaluation of the recoverability, based on size, of tagged carcasses that were redistributed back into the river after tagging. Testing revealed that there was no size bias for male or jack chinook, yet statistically longer female chinook were recaptured after being tagged. However, length distributions of tagged female chinook and recaptured female chinook are similar and bias to the mark-recapture estimate would be minimal.

The detection of sampling biases usually results in the use of a stratified estimator; however, Schubert (2000) compared the performance of several mark-recapture population estimators for a sockeye salmon population of known abundance and concluded that the pooled Petersen estimator was less biased and preferred over stratified estimators. In that study, the Schaeffer estimator would not improve accuracy and it was recommended that the method be abandoned for use in population estimation. Also, it was determined that while the maximum

likelihood Darroch estimator could potentially improve accuracy there was no obvious way of selecting between accurate and highly biased estimates. Parken and Atagi (2000) found that pooled and stratified estimators of Nass River summer steelhead produced similar escapement estimates but that the pooled estimator was more precise and had less statistical bias than the stratified estimator. These findings indicate the robust nature of the pooled Petersen estimator and suggest that its use to determine population abundance from mark-recapture data is generally appropriate under a wide range of circumstances.

SEAL PREDATION

Although seal predation was not directly assessed in this study, it is worthwhile to examine the impact seals have on chinook in Cowichan Bay. In 1988, the number of seals gradually increased from a low of 30 in April to a peak of about 100 in December. According to Olesiuk *et al.* (1990) harbour seals consume an estimated 9 tonnes of salmon annually in Cowichan Bay. An estimated 23% (Sept.) to 48% (Nov.) of the harbour seals' diet in Cowichan Bay was comprised of salmon (Bigg *et al.* 1990). Based on these data, consumption of chinook salmon could potentially range from 100 to 500 adults. These data were collected in 1988 when low flows in the Cowichan River persisted until the end of October. Predation likely increases the longer chinook salmon remain in the estuary waiting for high water to allow upstream migration. While low flow conditions occurred during September and October, 2001, Fishery Officers observations indicated that seal and sea lion populations have remained constant for the last couple of years in Cowichan Bay. As a result, predation on chinook was estimated to be approximately 300 adults.

POPULATION ESTIMATE

The 2001 Cowichan River chinook population estimate was based on the enumeration fence count. Moderate river flows allowed the fence to remain in place without significant breeches for the duration of the study and it was felt that a near complete census was attained. The estimate of the number of natural spawning adult chinook migrating past the fence site of 3,282 was just below the 95% confidence limits of the 3,311-4427 adult chinook mark-recapture estimate. This over-estimation by the mark-recapture estimate may be due to higher than average water flow during the mark-recapture sampling period (Figure 5). Higher than normal water flow would decrease the number of carcasses recaptured resulting in an over-estimation of the actual chinook population.

The estimated number of adult spawners was the third lowest since 1975, below the period average of 5,651 and well below the escapement goal (Table 20, Figure 7). This could partially be due to low hatchery contribution in 1997, which would have decreased the return of four year old hatchery chinook (Figure 8). Natural and enhanced contributions to Cowichan River chinook escapement from 1982 to the present can be found in Table 9.

ACKNOWLEDGEMENTS

We would like to thank Ron Zenner, James Patterson, Michael Folkes, Candace Boyle, Phillip Joe, Dave Key, Dave Burton, Kevin Timothy, Ashley Pifko, Frank Sutherland, Phil Burrash, Chad Stewart, and Stephan DeLeeuw for their involvement in the implementation of the enumeration fence, collection of migration and biological data. We thank W. Jansen and E. Teskey, Fisheries and Oceans Canada Fishery Officers stationed in Duncan, for their helpful assistance. We thank the Cowichan Tribes Fisheries Management Unit, in particular Wayne Paige, Jed and Doug August, and Wayne Paige Jr. for their cooperation and assistance in acquiring Native food fishery data and swim survey results. We thank Doug Millerd, manager of the Cowichan hatchery, for providing broodstock capture and biosampling data. We thank the conscientious sport fishers who returned the Ketchum tags from the chinook they caught in the Cowichan River. We thank the City of Duncan for allowing us to use their access road to the counting fence site and granting us storage space, power and water from their pumphouse facility. We thank C. Harlow, from Fisheries and Oceans Canada, B. Hollingshead, the Regional Water Manager for British Columbia, and A. Jezierski from Norske Canada for coordinating the water release management plan for the Cowichan River.

REFERENCES

- Argue, A.L., R. Hilborn, R.M. Peterman, M. J. Staley, and C.J. Walters. 1983. Strait of Georgia chinook and coho fishery. *Can. Bull. Fish. Aquat. Sci.* 211: 91 p.
- Bigg, M.A., G.M. Ellis, P. Cottrell, and L. Milette. 1990. Predation by Harbour seals and Sea lions on adult salmon in Comox Harbour and Cowichan Bay, British Columbia. *Can. Tech. Rep. Fish. Aquat. Sci.* 1769: 31 p.
- Burns, T. An assessment of Chinook salmon enumeration methods in the Cowichan River. Unpublished manuscript.
- Collicut, L.D. and T.F. Shardlow. 1995. Strait of Georgia sport fishery creel survey statistics for salmon and groundfish, 1991. *Can. Tech. Rep. Fish. Aquatic Sci.* 2137: 75 p.
- Cousens, N.B.F., G.A. Thomas, C.G. Swann, and M.C. Healey. 1982. A review of salmon escapement estimation techniques. *Can. Tech. Rep. Fish. Aquat. Sci.* 1108.
- Cross, C.L., L. Lapi, E. A. Perry. 1991. Production of chinook and coho salmon from British Columbia hatcheries, 1971 through 1989. *Can. Tech. Rep. Fish. Aquat. Sci.* 1816: 48 p.
- Farlinger, S., N. Bourne, B. Riddell, D. Chalmers, and A. Tyler (Eds). 1990. Pacific stock assessment review committee (PSARC) annual report for 1989. *Can. Manuscr. Rep. Fish. Aquat. Sci.* 2064: 236 p.

- Fielden, R.J. and L.B. Holtby. 1987. Standing crop and habitat characteristics of juvenile salmonids at sites in the Cowichan River system. Can. Manuscr. Rep. Fish. Aquat. Sci. 1950: 65 p.
- Johnston, N.T., J.R. Irvine, and C.J. Perrin. 1986. A comparative evaluation of fence count, mark-recapture and Bendix sonar estimates of salmon escapements in the Keogh River, a variable -flow coastal B.C. stream. Can. Tech. Rep. Fish. Aquat. Sci. 2111: 44 p.
- Krebs, C.J. 1989. Ecological methodology. Harper Collins Publishers, New York. N.Y.
- McDougall, R.D. 1985. A habitat management area plan for salmon stocks in the Cowichan watershed. Prepared for Field Services Branch, Dept. Fish. Ocean, Pacific Region.
- Nagtegaal, D.A., P. J. Starr, and B. Riddell. 1994a. A preliminary report on the chinook productivity study conducted on the Cowichan River, 1988 and 1989. Can. Manuscr. Rep. Fish. Aquat. Sci. 2233: 53 p.
- Nagtegaal, D.A., J. Candy, and B. Riddell. 1994b. A preliminary report on the chinook productivity study conducted on the Cowichan River during 1990 and 1991. Can. Manuscr. Rep. Fish. Aquat. Sci. 2265: 71 p.
- Nagtegaal, D.A., J. Candy, and B. Riddell. 1994c. A preliminary report on the chinook productivity study conducted on the Cowichan River during 1992. Can. Manuscr. Rep. Fish. Aquat. Sci. 2268: 73 p.
- Nagtegaal, D.A., and E.W. Carter. 1998. Adult chinook escapement assessment conducted on the Cowichan River during 1996. Can. Manuscr. Rep. Fish. Aquat. Sci. 2449: 65 p.
- Nagtegaal, D.A., and E.W. Carter. 2000. Adult chinook escapement assessment conducted on the Cowichan River during 1999. Can. Manuscr. Rep. Fish. Aquat. Sci. 2544: 59 p.
- Olesiuk, P.F., M.A. Bigg, G.M. Ellis, S.J. Crockford, and R.J. Wigen. 1990. An assessment of the feeding habits of Harbour seals (*Phoca vitulina*) in the Strait of Georgia, British Columbia, based on scat analysis. Can. Tech. Rep. Fish. Aquat. Sci. 1730: 135 p.
- Paige, W. 1992. Cowichan River Management Unit: Salmon Catch Statistical Program for 1991/1992. Cowichan Indian Band. Unpublished manuscript.
- Paige, W. 1997. Cowichan Tribes Aboriginal Fisheries: Salmon Catch Statistical Program for 1994/1996. Cowichan Indian Band. Unpublished manuscript.
- Parken, C.K., and D.Y. Atagi.. 2000. Preliminary estimate of the escapement of summer steelhead to the Nass River, 1998. British Columbia Ministry of Environment, Lands and Parks, Smithers, BC, Skeena Fisheries Report SK-124.

- Perrin, C.J., N.T. Johnston, and S.C. Samis. 1988. Effects of treated sewage effluent on periphyton and zoobenthos in the Cowichan River, British Columbia. Can. Tech. Rep. Fish. Aquat. Sci. 1591: 64 p.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Board Can. 191: 382 p.
- Schubert, N.D. 2000. The 1994 Stellako River sockeye (*Oncorhynchus nerka*) escapement: evaluation of pooled Petersen and stratified mark-recapture estimates of a known population. Can. Tech. Rep. Fish. Aquat. Sci. 2303: 56 p.
- Seber, G.A.F. 1982. The estimation of animal abundance and related parameters, Second edition. Griffin, London.
- Tschaplinski, P.J., and K.D. Hyatt. 1991. A comparison of population assessment methods employed to estimate the abundance of sockeye salmon (*Oncorhynchus nerka*) returning to Henderson Lake, Vancouver Island, during 1989. Can. Tech. Rep. Fish. Aquat. Sci. 1978: 101 p.
- Waples, R.S. 1991. Genetic interactions between hatchery and wild salmonids: lessons from the Pacific Northwest. Can. J. Fish. Aquat. Sci. 48 (Suppl. 1): 124-133.

Table 1. Daily counts at the Cowichan River fence site, 2001.

Date	Visibility	Temp (°C)	Depth (cm)	Chinook		Coho		Chum	Unknown
				Adult	Jack	Adult	Jack		
4-Sep	1	18.0	60.0	0	0	0	0	0	0
5-Sep	1	16.9	53.3	0	0	0	0	0	0
6-Sep	1	17.0	43.7	1	0	0	0	0	0
7-Sep	1	16.1	43.7	4	0	0	0	0	0
8-Sep	1	16.7	43.7	1	0	0	0	0	0
9-Sep	1	16.8	43.6	9	3	0	0	0	0
10-Sep	1	17.2	43.3	3	7	0	0	0	0
11-Sep	1	17.0	43.0	1	0	0	0	0	0
12-Sep	1	17.9	43.8	3	2	0	0	0	0
13-Sep	1	18.3	43.7	4	5	0	0	0	0
14-Sep	1	18.4	43.7	21	1	0	0	0	0
15-Sep	1	18.4	43.3	3	1	0	0	0	0
16-Sep	1	18.9	43.0	8	5	0	0	0	0
17-Sep	1	18.3	43.0	25	13	0	0	0	0
18-Sep	1	17.6	42.7	5	5	0	0	0	0
19-Sep	1	17.0	42.7	13	5	0	0	0	0
20-Sep	1	17.0	42.7	9	6	0	0	0	0
21-Sep	1	17.2	43.8	41	28	1	0	0	0
22-Sep	1	17.2	45.0	44	41	0	1	0	0
23-Sep	1	17.8	45.0	41	12	0	1	0	0
24-Sep	1	18.7	45.0	113	46	0	0	0	0
25-Sep	1	18.3	45.1	32	30	1	0	0	0
26-Sep	1	18.1	46.0	38	40	0	2	0	0
27-Sep	1-2	15.3	47.3	47	25	2	1	0	1
28-Sep	1-2	16.0	62.4	612	418	24	17	0	7
29-Sep	1	16.0	62.7	139	96	1	0	0	6
30-Sep	1	16.1	63.0	35	15	1	0	0	1
1-Oct	1	15.9	61.8	168	31	14	2	0	0
2-Oct	1	15.4	59.4	28	3	10	0	0	0
3-Oct	1	15.7	54.1	15	2	3	0	0	1
4-Oct	1	15.5	50.8	38	6	24	3	0	0
5-Oct	1	13.5	49.6	13	1	13	5	0	1
6-Oct	1	14.5	49.8	3	0	8	2	0	0
7-Oct	1	14.1	49.7	13	1	27	7	1	0
8-Oct	1	14.3	49.6	16	1	37	6	0	0
9-Oct	1	13.7	49.3	96	5	115	17	0	3
10-Oct	1	14.0	49.4	124	8	28	16	1	1
11-Oct	1	13.7	50.0	79	6	119	13	1	0
12-Oct	1-2	14.0	51.3	149	17	78	13	1	2
13-Oct	1	14.0	62.8	660	151	240	83	3	24
14-Oct	1	14.0	63.6	104	26	347	92	10	6
15-Oct	1	13.7	63.0	56	8	294	30	4	3
16-Oct	1	13.3	62.1	9	13	78	13	4	1
17-Oct	1	12.9	59.1	8	1	54	5	1	0
18-Oct	1	13.0	56.2	17	10	146	6	3	1
19-Oct	1	13.8	52.0	6	10	13	4	0	0
20-Oct	1	12.6	50.8	6	2	71	15	6	0

Table 1. (continued)

Date	Visibility	Temp ($^{\circ}\text{C}$)	Depth (cm)	Chinook		Coho		Chum	Unknown
				Adult	Jack	Adult	Jack		
21-Oct	1	12.3	50.6	12	3	66	12	10	0
22-Oct	1	12.0	52.1	9	4	215	16	25	0
23-Oct	1	11.6	59.7	56	11	2661	190	451	6
24-Oct	1-2	10.4	65.3	69	36	643	150	362	12
25-Oct	1-2	10.1	69.4	122	77	677	72	298	9
26-Oct	2	9.0	75.0	77	57	1367	95	399	13
27-Oct	1-2	9.0	79.4	37	42	1507	84	708	2
28-Oct	1	na	83.1	15	13	912	51	975	3
29-Oct	1-2	na	84.0	17	10	426	49	519	2
30-Oct	1-2	na	87.3	39	37	848	62	1225	8
31-Oct	1-2	na	88.5	48	56	4190	155	2735	130
1-Nov	2	na	89.0	0	2	175	6	117	0
Totals				3361	1454	15436	1296	7859	243

Visibility Code: 1 = clear; 2 = cloudy.

Table 2. Counts, by time interval, at the Cowichan River fence, 2001.

Time Period	Chinook				Coho				Chum	
	Adult		Jack		Adult		Jack		Count	%
	Count	%	Count	%	Count	%	Count	%		
0000 - 0100	125	3.7	43	3.0	270	1.7	49	3.8	169	2.2
0100 - 0200	142	4.2	48	3.3	324	2.1	29	2.2	137	1.7
0200 - 0300	140	4.2	48	3.3	341	2.2	28	2.2	179	2.3
0300 - 0400	205	6.1	44	3.0	322	2.1	23	1.8	115	1.5
0400 - 0500	114	3.4	44	3.0	243	1.6	20	1.5	122	1.6
0500 - 0600	93	2.8	33	2.3	318	2.1	23	1.8	92	1.2
0600 - 0700	108	3.2	45	3.1	230	1.5	22	1.7	47	0.6
0700 - 0800	325	9.7	159	10.9	645	4.2	83	6.4	242	3.1
0800 - 0900	422	12.6	235	16.2	844	5.5	79	6.1	771	9.8
0900 - 1000	360	10.7	217	14.9	1002	6.5	63	4.9	801	10.2
1000 - 1100	216	6.4	71	4.9	925	6.0	108	8.3	515	6.6
1100 - 1200	92	2.7	55	3.8	1351	8.8	57	4.4	675	8.6
1200 - 1300	96	2.9	27	1.9	1116	7.2	53	4.1	602	7.7
1300 - 1400	63	1.9	24	1.7	1569	10.2	60	4.6	684	8.7
1400 - 1500	59	1.8	23	1.6	1257	8.1	99	7.6	634	8.1
1500 - 1600	73	2.2	27	1.9	1133	7.3	127	9.8	624	7.9
1600 - 1700	96	2.9	50	3.4	1169	7.6	120	9.3	581	7.4
1700 - 1800	158	4.7	73	5.0	692	4.5	89	6.9	289	3.7
1800 - 1900	80	2.4	38	2.6	336	2.2	47	3.6	158	2.0
1900 - 2000	32	1.0	19	1.3	179	1.2	13	1.0	65	0.8

Table 2. (continued)

Time Period	Chinook				Coho				Chum	
	Adult		Jack		Adult		Jack		Chum	
	Count	%	Count	%	Count	%	Count	%	Count	%
2000 - 2100	46	1.4	13	0.9	272	1.8	21	1.6	56	0.7
2100 - 2200	61	1.8	35	2.4	309	2.0	21	1.6	69	0.9
2200 - 2300	99	2.9	38	2.6	292	1.9	28	2.2	114	1.5
2300 - 2400	156	4.6	45	3.1	297	1.9	34	2.6	118	1.5
Total	3361	100	1454	100	15436	100	1296	100	7859	100

Table 3. Visual survey data collected for the Cowichan River for the years 1981 to 2001

Chinook							
Method ¹	Date	Jacks		Adults		River Segment ²	
		Count	Estimate	Count	Estimate		
1981	S	Sept.	12	175	208	1000	2-4
	S	Oct.	2	103	93	1500	2-4
	S		14	364	1160	4000	2-4
	H		22		2000		1-7
	S		23		3200	5000	2-4
Estimate for Season ³						5500	
1982	S	Sept.	14	199	131	600	2-4
	S	Oct.	13		153		2-4
	H		19	saw few fish on spawning grounds			1-13
	F	Nov.	8			4000	
Estimate for Season						4500	
1983	S	Sept.	8	38	61	254	2-6
	S		15	62	121	504	2-6
	S		28	190	470	1838	1-2
	S	Oct.	7	207	425	1804	2-6
	S		14	802	997	2836	2-7
	S		25	901	1113	4500	1-6
Estimate for Season						4500	
1984	S	Aug.	28	80	84	400	2-5
	S	Sept.	6	25	72		
	S		13	79	80		3-11
	S		19	35	71		2-6
	S		26	291	434		2-6
	S	Oct.	3	205	283		3-7
	S	"		206	282	2200	8-11
	S		23	525	1300	5000	1-6
	S	Nov.	1	350	1276		1-6
Estimate for Season						5000	

Table 3. (continued)

Chinook								
Method ¹	Date	Jacks		Adults		River Segment ²		
		Count	Estimate	Count	Estimate			
1985	S	Sept. 12	39		46	220		2-6
	S		17	42		10		12-13
	S		18	210		33		2-6
	S		27	245		104	456	2-6
	S	Oct. 3	244			99	360	2-6
	S		10	285		219		2-6
	S		16	293		347		2-6
	S		31	229		934	3500	1-6
Estimate for Season							3500	
1986	S	Sept. 9	295		85	300		2-6
	S		18	46	29	300		3-6
	S		24	161	56	350		12-13
	S	Oct. 7	1310		223	1000		2-6
	S		29	613	473	1200		1-6
	S	Nov. 6	1178		491	1200		
	H		8		515			1-13
Estimate for Season							1200	
1987	S	Sept. 9	30	300	10	50		3-8
	S		17	111	16	75		2-6
	S		25	112	16	75		3-6, 11-12
	S	Oct. 6	196	800	115	400		2-6
	S		15	196	96			1-6
	H		16		saw very few spawners			1-13
	S		28	417	468			1-6
	S	Nov. 6	329		649			1-6
Estimate for Season							1200	
1988	S	Aug. 25	100		50			2-6
	S	Sept. 1	271		149	700		2-6
	S		23	1464	271	1000		2-6
	S	Oct. 3	821	1600	1094	3500		2-6
	S		14	2008	2076	4000		1-6
Estimate for Season							5500	
1989	S	Sept. 11	151		58	300		2-6
	S		21	95	39	350		3-6
	S	Oct. 5	95		48	700		2-3
	S		18	719	350	1200		2-6
	S	Nov. 1	1537		2267			2-6
Estimate for Season							5000	
1990	S	Aug. 29	254		54	250		2-6
	S	Sept. 14	385		89	1000		3-6
	S		27	3169	477	2200		2-3
	S	Oct. 19	4297		2382	5000		2-6
Estimate for Season							5300	

Table 3. (continued)

	Method ¹	Date	Chinook				River Segment ²
			Jacks		Adults		
			Count	Estimate	Count	Estimate	
1991	S	Sept.	19		1882	6000	2-6
	S	Oct.	2		2873	7500	2-6
	S		17		2924	8700	2-6
	S		31		3502 ⁴	9000	2-6
Estimate for Season						10000	
1992	S	Sept.	16	5	8		2-5
	S	Oct.	2	124	46	200	2-6
	S		15	359	291	700	2-6
	S		15	113	162		2-6
	S		27	514	797	2000	1-6
	S		28	591	767		1-6
	S	Nov.	13	506	467		1-6
	S		13	450	640 ⁵		1-6
Estimate for Season						7500	
1993	S	Sept.	23	23	14	47	2-6
	S		30	81	62	210	2-6
	S	Oct.	14	207	199	676	2-6
	S		28	127	327	1111	2-6
	S	Nov.	4	480	987	3355	
Estimate for Season ⁶						5200	
1994	S	Aug.	24	39	3		2-6
	S	Sept.	14	67	46	156	2-6
	S		28	421	323	1098	2-6
	S	Oct.	13	1253	1146	3896	2-6
	S		26	442	1450	4930	2-6
Estimate for Season ⁶						5500	
1995	S	Sept.	28	294	267	1170	2-6
	S	Oct.	25	490	1798	6653	2-6
Estimate for Season ⁶						15500	
1996	S	Sept.	13	45	46	147	2-6
	S		26	166	150	510	2-6
	S	Oct.	2	254	534	1815	2-6
	S		9	579	1157	3933	2-6
	S		15	195	707	2403	2-6
	S		22	557	1699	5776	2-6
Estimate for Season ⁶						6500	
1997	S	Sept.	23	165	358	1217	2-6
	S	Sept.	25	87	404	1373	2-6
	S	Sept.	30	54	509	1730	2-6
	S	Oct.	16	84	289	980	2-6
	S	Oct.	23	1036	1831	6225	2-6
Estimate for Season ⁵						6500	

Table 3. (continued)

Chinook							
Method ¹	Date	Jacks		Adults		River Segment ²	
		Count	Estimate	Count	Estimate		
1998	S	Sept.	25	72	37		2-6
	S	Oct.	13	54	53		2-6
	S	Oct.	20	130	857	2913	2-6
	S	Oct.	26	317	1260	4284	2-6
Estimate for Season						4284	
1999	S	Sept.	10	88	46	221	2-6
	S	Oct.	13	321	342	1641	2-6
Estimate for Season						4500	
2000	S	Sept.	13	33	25	N/A ⁷	2-6
Estimate for Season						3069	
	S	Sept.	26	80	230	782	Lower ⁸
	S	Oct.	19	439	940	3196	1-6
2001	S	Oct.	23	135	255	867	2-6
Estimate for Season						3282	

¹S - Swim survey, H - Helicopter survey, F - boat survey

²Refer to Figure 1.

³Total escapement estimate for adult chinook.

⁴516 chinook carcasses were counted in this total.

⁵28 chinook carcasses were counted in this total.

⁶Swim surveys conducted by Cowichan Tribes River Management Unit, total escapement determined by Fishery officers.

⁷Adult population estimate based on swim survey records was not feasible due to extremely low water conditions during swim survey.

⁸Swim Survey took place in the lower Cowichan River from Black Bridge to Green Point (Figure 2).

Table 4. Estimated catch of chinook adults and jacks in the Cowichan River native food fishery¹ for the years 1981 to 2001.

Year ²	Chinook Catch	
	Adult	Jack ³
1971	725	
1972	700	
1973	900	
1974	1000	
1975	900	
1976	1000	
1977	1000	
1978	500	
1979	500	
1980	1500	
1981	1500	1500
1982	1000	1000
1983	250	1000
1984	355	700
1985	1000	1000
1986	800	800
1987	800	800
1988	681	450
1989	1055	250
1990	820	150
1991	450	70
1992	900	12
1993	650	22
1994	700	227
1995	533	120
1996	810	150
1997	191	N/A
1998	1073	N/A
1999	233	89
2000	89	N/A
2001	918	120

¹ Includes chinook caught in both the spear fishery and the in-river gillnet fishery.

² Since 1988, data has been collected by the Cowichan Tribes River Management unit. Prior to 1988, data were collected by local Fishery Officers.

³ Estimates for jack chinook were not provided in 1997, 1998 and 2000.

Table 5. Summary, by day and location, of chinook broodstock collected by the Cowichan River hatchery during 2001.

Date	Below Fence			Above Fence		
	Male	Jack	Female	Male	Jack	Female
9/18/01	31	0	54	2	0	9
9/19/01	45	0	69	5	0	10
9/20/01	33	0	60	16	0	27
9/21/01	26	0	37	6	0	5
9/24/01	40	0	32	0	0	0
9/25/01	89	0	116	13	0	23
9/26/01	143	0	195	1	0	1
9/27/01	86	0	90	0	0	5
9/28/01	97	0	99	2	0	0
10/1/01	42	0	42	24	0	33
10/2/01	17	0	24	6	0	9
10/3/01	31	0	0	33	0	4
Total	680	0	818	108	0	126

Table 6. Annual hatchery broodstock collection of adult and jack Cowichan River chinook for the years 1981 to 2001.

Year	Chinook Broodstock Collection	
	Adult	Jack ¹
1981	282	
1982	534	
1983	242	
1984	278	
1985	175	
1986	315	
1987	582	
1988	678	30
1989	535	96
1990	327	1
1991 ²	1755	347
1992	1850	77
1993	2200	228
1994	1357	145
1995	2149	512
1996	1615	258
1997	125	79
1998	1485	201
1999	1659	1
2000	1529	14
2001	1732	0

¹ Barry Cordecedo (Salmonid Enhancement Program) provided numbers on broodstock collection from 1981 to 1987. The broodstock numbers provided included jacks, but no reliable records were kept. It was estimated that for most years about 10 to 15 jacks were collected. These estimates were subtracted from the broodstock numbers resulting in an estimate of the number of adult chinook removed from the system.

² In addition, 284 males were removed for broodstock but later returned to the river.

Table 7. Summary of age, by sex, for Cowichan River chinook collected for hatchery broodstock during 2001.

Age	Male		Female		Total Adult		Jack	
	#	%	#	%	#	%	#	%
2	2	1.3%	0	0.0%	2	0.6%	6	85.7%
3	128	80.0%	103	51.5%	231	64.2%	1	14.3%
4	29	18.1%	84	42.0%	113	31.4%	0	0.0%
5	1	0.6%	13	6.5%	14	3.9%	0	0.0%
Total	160	100%	200	100%	360	100%	7	100%

Table 8. Length-frequency of chinook sampled on the Cowichan River spawning grounds, 2001.

Length (cm)	Males	Jacks	Females
29	0	2	0
30	0	4	0
31	0	7	0
32	0	3	0
33	0	4	0
34	0	4	0
35	0	14	0
36	0	10	0
37	0	19	0
38	0	20	0
39	0	12	0
40	0	24	0
41	1	19	1
42	0	17	0
43	0	11	0
44	0	6	0
45	1	4	1
46	3	1	0
47	4	0	0
48	6	0	0
49	4	0	0
50	5	0	2
51	5	0	9
52	2	0	1
53	5	0	4
54	0	0	6
55	4	1	8
56	6	0	13
57	5	0	11
58	8	0	20
59	14	0	17

Table 8. (continued)

Length (cm)	Males	Jacks	Females
60	15	0	32
61	15	0	27
62	10	0	16
63	13	0	21
64	11	0	26
65	10	0	23
66	10	0	13
67	5	0	27
68	5	0	22
69	8	0	24
70	9	0	32
71	9	0	22
72	3	0	22
73	7	0	16
74	3	0	18
75	4	0	7
76	2	0	8
77	5	0	7
78	2	0	2
79	3	0	0
80	2	0	4
81	3	0	0
82	2	0	1
83	0	0	0
Total	229	182	463
Mean Length (cm)	63.0	38.6	65.2
Adipose Fin Clips	11	3	15
Fin Clip Rate	4.8%	1.6%	3.2%

Table 9. Summary of age, by sex, for Cowichan River chinook collected from the spawning grounds during 2001.

Age	Male		Female		Total Adult		Jack	
	#	%	#	%	#	%	#	%
2	7	6.7%	2	1.0%	9	2.9%	89	98.9%
3	65	62.5%	105	50.0%	170	54.1%	1	1.1%
4	32	30.8%	98	46.7%	130	41.4%	0	
5			5	2.4%	5	1.6%	0	
Total	104	100%	210	100%	314	100%	90	100%

Table 10. Length-frequency of chinook hatchery broodstock collected during 2001.

Length (cm)	Males	Jacks	Females
38	0	1	0
39	0	0	0
40	0	0	0
41	0	1	0
42	0	0	0
43	0	2	0
44	0	0	0
45	0	1	0
46	0	0	0
47	0	0	0
48	1	0	0
49	1	1	1
50	2	0	2
51	6	1	0
52	8	0	1
53	4	0	0
54	12	0	1
55	14	0	8
56	19	0	7
57	18	0	7
58	10	0	11
59	15	0	9
60	11	0	17
61	10	0	14
62	6	0	15
63	4	0	14
64	2	0	6
65	7	0	12
66	7	0	16
67	4	0	7
68	3	0	9
69	3	0	16

Table 10. (continued)

Length (cm)	Males	Jacks	Females
74	0	0	10
75	0	0	1
76	1	0	2
77	1	0	0
78	1	0	0
79	0	0	1
80	0	0	2
81	0	0	0
82	0	0	0
83	0	0	1
Total	178	7	217
Mean Length (cm)	59.3	44.4	64.6
Adipose Fin Clips	0	0	16
Fin Clip Rate	0.0%	0.0%	7.4%

Table 11. Release and recovery data for coded wire tags recovered from Cowichan River chinook sampled on the spawning grounds during 2001.

Recovery Data				Release Data			
Date (dd/mm/yy)	Location	POH Length (mm)	Sex	Brood Year	Tag Code	Location ¹	Date (dd/mm/yy)
05/11/01	10	652	F	98	183727	Cowichan Lake Pen	07/05/99
06/11/01	16	560	F	98	183730	Upper Cowichan R. (late)	10/05/99
06/11/01	16	664	M	98	183726	Cowichan Lake Pen	07/05/99
06/11/01	14	650	M	98	182804	Hatchery (late)	21/05/98
06/11/01	14	665	F		No Pin		
07/11/01	18	655	M	98	183733	Upper Cowichan R. (late)	10/05/99
07/11/01	84	607	F		No Data		
08/11/01	10	648	F	98	183730	Upper Cowichan R. (late)	10/05/99
08/11/01	12	650	M	98	183729	Upper Cowichan R. (early)	31/03/99
09/11/01	52	620	M	98	183733	Upper Cowichan R. (late)	10/05/99
09/11/01	17	652	F	98	183731	Upper Cowichan R. (late)	10/05/99
09/11/01	14	640	M	98	183730	Upper Cowichan R. (late)	10/05/99
09/11/01	53	702	M	98	182804	Hatchery (late)	21/05/98
13/11/01	36	600	F	98	183728	Upper Cowichan R. (early)	31/03/99
13/11/01	61	586	F	98	183726	Cowichan Lake Pen	07/05/99
13/11/01	26	426	J	99	183126	Hatchery (late)	05/05/00
13/11/01	26	474	M	99	183123	Upper Cowichan R. (late)	28/04/00
13/11/01	26	382	J	99	183119	Upper Cowichan R. (early)	07/03/00
14/11/01	10	600	F	98	183731	Upper Cowichan R. (late)	10/05/99
14/11/01	11	640	F	98	183730	Upper Cowichan R. (late)	10/05/99
14/11/01	11	496	M	98	183727	Cowichan Lake Pen	07/05/99
14/11/01	53	428	J	99	183124	Upper Cowichan R. (late)	28/04/00

Table 11. (continued)

Recovery Data				Release Data			
Date (dd/mm/yy)	Location	POH Length (mm)	Sex	Brood Year	Tag Code	Location ¹	Date (dd/mm/yy)
21/11/01	11	570	F	98	183731	Upper Cowichan R. (late)	10/05/99
21/11/01	14	642	F	98	183731	Upper Cowichan R. (late)	10/05/99
23/11/01	14	605	F	98	183733	Upper Cowichan R. (late)	10/05/99
23/11/01	13	550	F	98	183734	Seapen	17/05/99
26/11/01	36	605	M	98	183730	Upper Cowichan R. (late)	10/05/99
26/11/01	8	509	F		No Pin		
04/12/01	26	691	F	98	183730	Upper Cowichan R. (late)	10/05/99
04/12/01	14	648	M		No Pin		

¹ Cowichan Hatchery release strategies for chinook:

Upper Cowichan River (late): raised to pre-smolt size (5-6 g) prior to release approximately 3 km below the weir in May.

Upper Cowichan River (early): raised to fry (3 g) prior to release approximately 3 km below the weir in early April.

Cowichan Lake Pen: raised to pre-smolt size (5-6 g) prior to release just above the weir in May.

Hatchery (late): raised to pre-smolt size (5-6 g) prior to release at the hatchery in May.

Seapen: raised to smolt size (6+ g) prior to release from the netpens in Cowichan Bay in early June.

Table 12. Cowichan River hatchery chinook release¹ data for the years 1979 to 2001.

Tagcode	Brood Year	Number Tagged	Number Released	CWT % Marked	Weight (g)	Start Release Date	End Release Date	Release Site
021846	1979	31628	32134	98.4	2.8		1980/05/07	COWICHAN R
022060	1979	32034	32547	98.4	2.8		1980/05/07	SKUTZ FALLS
022158	1980	52519	65000	80.8	2.3		1981/06/09	COWICHAN R
022307	1981	30179	30373	99.4	3.1		1982/05/12	COWICHAN R
022339	1982	49135	224944	21.8	2.9		1983/05/14	SKUTZ FALLS
022831	1983	50613	101000	50.1	4.27		1984/05/25	KOKSILAH R
NOCN8311	1983		200000	0.0	4.27		1984/05/31	COWICHAN R
NOCN8411	1984		187823	0.0	4.8	1985/05/13	1985/05/14	COWICHAN R
023803	1985	25365	25804	98.3	4.26	1986/05/23	1986/05/24	COWICHAN R
023804	1985	25455	25895	98.3	4.26	1986/05/23	1986/05/24	COWICHAN R
023911	1985	11980	12187	98.3	4.26	1986/05/23	1986/05/24	COWICHAN R
NOCN8619	1986		321172	0.0	4	1987/05/13	1987/05/22	COWICHAN R
NOCN8620	1986		54608	0.0	3.48		1987/05/21	KOKSILAH R
024334	1987	14298	14334	99.7	3.41		1988/04/18	COWICHAN R
024729	1987	25360	25424	99.7	3.4		1988/04/18	COWICHAN R
024730	1987	25869	25934	99.7	3.4		1988/04/18	COWICHAN R
024731	1987	27428	27497	99.7	7.1	1988/04/18	1988/05/18	COWICHAN LK
024732	1987	27271	27339	99.8	7.1		1988/05/18	COWICHAN LK
024733	1987	26911	26978	99.8	7.1		1988/05/18	COWICHAN LK

Table 12. (continued)

Tagcode	Brood Year	Number Tagged	Number Released	CWT % Marked	Weight (g)	Start Release Date	End Release Date	Release Site
024734	1987	23521	23580	99.7	7.1		1988/05/18	COWICHAN LK
024735	1987	26719	26786	99.7	3.4	1988/04/18	1988/05/18	COWICHAN R
024945	1987	26461	123361	21.5	7.49	1988/05/25	1988/05/26	COWICHAN R UP
024946	1987	26658	123560	21.6	7.49	1988/05/25	1988/05/26	COWICHAN R UP
024947	1987	26761	123663	21.6	7.49	1988/05/25	1988/05/26	COWICHAN R UP
025008	1987	26817	123720	21.7	7.49	1988/05/25	1988/05/26	COWICHAN R UP
024860	1988	25117	25243	99.5	3.66		1989/04/28	COWICHAN R
025012	1988	26595	54768	48.6	6.49		1989/05/21	COWICHAN R
025013	1988	25982	54154	48.0	6.49		1989/05/21	COWICHAN R
025015	1988	23058	24894	92.6	3.66		1989/04/28	COWICHAN R
025016	1988	26821	26821	100.0	3.66		1989/04/28	COWICHAN R
025017	1988	27611	28175	98.0	3.66		1989/04/28	COWICHAN R
025523	1988	27531	56123	49.1	6.49		1989/05/21	COWICHAN R
025524	1988	27205	55378	49.1	6.49		1989/05/21	COWICHAN R
025749	1988	26922	133331	20.2	6.06		1989/05/15	COWICHAN LK
025750	1988	27036	133446	20.3	6.06		1989/05/15	COWICHAN LK
025751	1988	23106	130107	17.8	6.06		1989/05/15	COWICHAN LK
025752	1988	26169	132842	19.7	6.06		1989/05/15	COWICHAN LK
020352	1989	28287	28573	99.0	3.4	1990/04/12	1990/04/12	COWICHAN R
020522	1989	27072	36800	73.6	6.53	1990/05/22	1990/05/23	COWICHAN R
020622	1989	27787	37242	74.6	6.53	1990/05/22	1990/05/23	COWICHAN R
020623	1989	28164	37619	74.9	6.53	1990/05/22	1990/05/23	COWICHAN R
020624	1989	28331	37786	75.0	6.53	1990/05/22	1990/05/23	COWICHAN R
020938	1989	28312	28312	100.0	3.4	1990/04/12	1990/04/12	COWICHAN R
020939	1989	26218	26218	100.0	3.4	1990/04/12	1990/04/12	COWICHAN R
026103	1989	27145	27145	100.0	3.4	1990/04/12	1990/04/12	COWICHAN R
026255	1989	26400	119674	22.1	7.19		1990/05/14	COWICHAN LK
026256	1989	25693	119497	21.5	7.19		1990/05/14	COWICHAN LK
026257	1989	25790	119325	21.6	7.19		1990/05/14	COWICHAN LK
026258	1989	25219	118748	21.2	7.19		1990/05/14	COWICHAN LK
020333	1990	25687	94172	27.3	8.43	1991/05/15	1991/05/15	COWICHAN LK
020334	1990	25898	94384	27.4	8.43	1991/05/15	1991/05/15	COWICHAN LK
020335	1990	25739	94224	27.3	8.43	1991/05/15	1991/05/15	COWICHAN LK
020336	1990	27135	27135	100.0	3.31	1991/04/17	1991/04/17	COWICHAN R
020337	1990	26631	26631	100.0	3.31	1991/04/17	1991/04/17	COWICHAN R
020338	1990	27046	27046	100.0	3.31	1991/04/17	1991/04/17	COWICHAN R
020339	1990	26721	34318	77.9	6.4	1991/05/21	1991/05/22	COWICHAN R
020340	1990	26993	34592	78.0	6.4	1991/05/21	1991/05/22	COWICHAN R
020341	1990	26533	33995	78.0	6.4	1991/05/21	1991/05/22	COWICHAN R
020342	1990	25437	92182	27.6	4.75	1991/06/17	1991/06/18	COWICHAN R
020343	1990	25391	92136	27.6	4.75	1991/06/17	1991/06/18	COWICHAN R
NOCN9044	1990		5086	0.0	5.41	1991/06/26	1991/06/26	COWICHAN ESTUARY
180513	1991	26972	336330	8.0	5.04	1992/05/17	1992/05/17	COWICHAN LK
180514	1991	25964	335584	7.7	5.04	1992/05/17	1992/05/17	COWICHAN LK
180515	1991	27694	254287	10.9	4.01	1992/04/21	1992/04/22	COWICHAN R LOW
180516	1991	27148	254015	10.7	4.01	1992/04/21	1992/04/22	COWICHAN R LOW
180517	1991	27471	505110	5.4	5.47	1992/05/19	1992/05/21	COWICHAN R UP
180518	1991	27277	504916	5.4	5.47	1992/05/19	1992/05/21	COWICHAN R UP

Table 12. (continued)

Tagcode	Brood Year	Number Tagged	Number Released	CWT % Marked	Weight (g)	Start Release Date	End Release Date	Release Site
180519	1991	27432	160695	17.1	3.75	1992/04/21	1992/04/22	COWICHAN R LOW
180520	1991	27001	160262	16.8	3.75	1992/04/21	1992/04/22	COWICHAN R LOW
180521	1991	26871	27444	97.9	6.29	1992/05/29	1992/05/29	COWICHAN ESTUARY
180522	1991	26852	27424	97.9	6.29	1992/05/29	1992/05/29	COWICHAN ESTUARY
NOCN9145	1991		513053	0.0	5.69	1992/05/19	1992/05/20	COWICHAN R UP
180209	1992	24770	98974	25.0	6.3	1993/05/25	1993/05/25	COWICHAN ESTUARY
180210	1992	26383	327416	8.1	5.86	1993/05/17	1993/05/19	COWICHAN R UP
180550	1992	25311	326344	7.8	5.86	1993/05/17	1993/05/19	COWICHAN R UP
181042	1992	53620	412953	13.0	6.52	1993/05/25	1993/05/25	COWICHAN R
181043	1992	54235	901937	6.0	5.64	1993/05/10	1993/05/10	COWICHAN LK
181044	1992	55027	907719	6.1	3.56	1993/04/07	1993/04/07	COWICHAN R UP
021211	1993	24875	103900	23.9	6.17	1994/05/25	1994/05/25	COWICHAN BAY
181319	1993	49966	1001002	5.0	6.29	1994/05/05	1994/05/05	COWICHAN LK
181320	1993	50420	684279	7.4	3.79	1994/04/18	1994/04/18	COWICHAN R UP
181321	1993	50045	652354	7.7	6.11	1994/05/18	1994/05/18	COWICHAN R UP
181322	1993	50285	490079	10.3	6.06	1994/05/24	1994/05/24	COWICHAN R
181329	1994	25023	103815	24.1	6.08	1995/05/31	1995/05/31	COWICHAN BAY
181436	1994	50133	100252	50.0	5.44	1995/05/30	1995/05/30	COWICHAN R
181437	1994	49962	418750	11.9	4	1995/05/02	1995/05/02	COWICHAN R UP
181438	1994	49610	939287	5.3	6.32	1995/05/15	1995/05/17	COWICHAN R UP
181439	1994	49846	101763	49.0	6.48	1995/05/25	1995/05/25	COWICHAN LK
182023	1995	25114	109088	23.0	6.76	1996/05/10	1996/05/10	COWICHAN BAY
182024	1995	25653	297360	8.6	6.56	1996/05/06	1996/05/06	COWICHAN LK
182025	1995	24488	283856	8.6	6.56	1996/05/06	1996/05/06	COWICHAN LK
182026	1995	25183	355089	7.1	6.26	1996/05/07	1996/05/07	COWICHAN R UP
182027	1995	25218	355583	7.1	6.26	1996/05/07	1996/05/07	COWICHAN R UP
182028	1995	25052	344597	7.3	3.47	1996/04/02	1996/04/02	COWICHAN R UP
182029	1995	25129	345657	7.3	3.47	1996/04/02	1996/04/02	COWICHAN R UP
182030	1995	25196	245910	10.2	6.37	1996/05/09	1996/05/09	COWICHAN R
182031	1995	25020	244193	10.2	6.37	1996/05/09	1996/05/09	COWICHAN R
182737	1996	25235	100196	25.2	6.79	1997/05/07	1997/05/07	COWICHAN BAY
182738	1996	25108	318583	7.9	5.44	1997/04/30	1997/04/30	COWICHAN LK
182739	1996	25205	319814	7.9	5.44	1997/04/30	1997/04/30	COWICHAN LK
182740	1996	25218	448340	5.6	6.29	1997/04/28	1997/04/29	COWICHAN R UP
182741	1996	25649	456002	5.6	6.29	1997/04/28	1997/04/29	COWICHAN R UP
182742	1996	25457	401644	6.3	3.34	1997/04/01	1997/04/01	COWICHAN R UP
182743	1996	25019	394733	6.3	3.34	1997/04/01	1997/04/01	COWICHAN R UP
182744	1996	25154	219780	11.4	5.89	1997/05/05	1997/05/05	COWICHAN R
182745	1996	25082	219151	11.4	5.89	1997/05/05	1997/05/05	COWICHAN R
182761	1997	25213	25213	100.0	3.68	1998/04/09	1998/04/09	COWICHAN R UP
182762	1997	25206	25206	100.0	3.68	1998/04/09	1998/04/09	COWICHAN R UP
182763	1997	25698	25698	100.0	3.68	1998/04/09	1998/04/09	COWICHAN R UP
182801	1997	24817	28209	88.0	6.47	1998/05/13	1998/05/13	COWICHAN R UP
182802	1997	24890	28282	88.0	6.47	1998/05/13	1998/05/13	COWICHAN R UP
182803	1997	24923	28316	88.0	6.47	1998/05/13	1998/05/13	COWICHAN R UP
182804	1997	24971	24971	100.0	6.46	1998/05/21	1998/05/21	COWICHAN R
182805	1997	25026	25026	100.0	6.46	1998/05/21	1998/05/21	COWICHAN R
183213	1997	24915	51754	48.1	6.27	1998/05/25	1998/05/25	COWICHAN BAY

Table 12. (continued)

Tagcode	Brood Year	Number Tagged	Number Released	CWT % Marked	Weight (g)	Start Release Date	End Release Date	Release Site
183107	1998	25163	224868	11.2	3.07	1999/03/31	1999/03/31	COWICHAN R UP
183108	1998	25201	225208	11.2	3.07	1999/03/31	1999/03/31	COWICHAN R UP
183109	1998	24803	132012	18.8	6.56	1999/05/10	1999/05/10	COWICHAN R UP
183110	1998	24927	132676	18.8	6.56	1999/05/10	1999/05/10	COWICHAN R UP
183111	1998	25163	75629	33.3	6.31	1999/05/10	1999/05/10	COWICHAN R
183112	1998	24875	74763	7.0	6.31	1999/05/10	1999/05/10	COWICHAN R
183726	1998	25135	356567		5.93	1999/05/07	1999/05/07	COWICHAN LK
183727	1998	25136	356568	7.0	5.93	1999/05/07	1999/05/07	COWICHAN LK
183728	1998	25234	225504	11.2	3.07	1999/03/31	1999/03/31	COWICHAN R UP
183729	1998	25087	224189	11.2	3.07	1999/03/31	1999/03/31	COWICHAN R UP
183730	1998	24867	132354	18.8	6.56	1999/05/10	1999/05/10	COWICHAN R UP
183731	1998	24921	132644	18.8	6.56	1999/05/10	1999/05/10	COWICHAN R UP
183732	1998	24959	75015	33.3	6.31	1999/05/10	1999/05/10	COWICHAN R
183733	1998	25024	75211	33.3	6.31	1999/05/10	1999/05/10	COWICHAN R
183734	1998	25127	99928	25.1	5.1	1999/05/17	1999/05/17	COWICHAN BAY
183119	1999	24855	270757	9.2	3.19	2000/03/07	2000/03/07	COWICHAN R UP
183120	1999	24917	271436	9.2	3.19	2000/03/07	2000/03/07	COWICHAN R UP
183121	1999	24933	271609	9.2	3.19	2000/03/07	2000/03/07	COWICHAN R UP
183122	1999	25024	272601	9.2	3.19	2000/03/07	2000/03/07	COWICHAN R UP
183123	1999	24776	481197	5.1	6.58	2000/04/27	2000/04/28	COWICHAN R UP
183124	1999	24839	482428	5.1	6.58	2000/04/27	2000/04/28	COWICHAN R UP
183125	1999	25118	215385	11.7	6.99	2000/05/05	2000/05/05	COWICHAN R
183126	1999	25039	215306	11.6	6.99	2000/05/05	2000/05/05	COWICHAN R
183127	1999	25078	99936	25.1	8.66	2000/05/17	2000/05/17	COWICHAN BAY
182811	2000	25175	99829	25.2	7.98	2001/05/23	2001/05/23	COWICHAN BAY
183216	2000	25152	504558	5.0	6.55	2001/05/01	2001/05/01	COWICHAN R UP
183217	2000	24833	503194	4.9	6.55	2001/05/01	2001/05/01	COWICHAN R UP
184539	2000	50166	338640	14.8	6.21	2001/05/03	2001/05/03	COWICHAN R
184546	2000	49972	481337	10.4	3.19	2001/03/19	2001/03/20	COWICHAN R UP
184547	2000	50054	482162	10.4	3.19	2001/03/19	2001/03/20	COWICHAN R UP

Table 13. Cowichan River daily discharge¹ measured in cubic meters per second, for 2001.

Day	Month											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
1	61.4	52.8	30.3	42.3	62.7	24.2	14.4	4.24	7	15.4	46.1	152
2	59.4	59.1	32	49.5	69.8	23.7	14.1	4.57	7.08	13.9	46.9	151
3	58	65.6	30.8	48.2	67.3	23.2	13.7	5.53	7.24	11.2	46.9	127
4	65.1	66.1	29.1	46.2	65.3	22.3	13.3	4.73	7	9.18	47.2	116
5	139	67.8	26.7	44.5	64.3	21.5	13.1	4.67	6.4	9.1	47.8	107
6	108	66.1	22.1	43.5	61.1	21	13.4	5.13	6.06	9.22	47.7	105
7	94.4	63.6	18.9	41	57.6	20.2	13.5	4.93	5.89	9.09	47.4	99.5
8	87	61.8	19.1	39.2	54.8	19.6	12.4	4.73	5.82	9.42	46.9	113
9	83.8	60.1	19.1	37.1	45.4	19.5	12.4	4.5	5.7	9.19	45.8	119
10	83.4	58	16.9	35.3	39.1	19.2	11.2	4.44	5.7	9.53	44.9	110
11	81	55.7	16.4	33.2	34.1	19	10.8	4.42	5.63	9.82	44	102
12	78.3	53.9	15.8	31.4	33	18.9	10.3	4.33	5.65	12.5	44.5	98.8
13	76	52	16.9	30.5	31.8	18.5	9.96	4.3	5.6	18	48.2	114
14	72.1	50.2	17	28.7	32.7	17.9	9.59	4.28	5.55	18	67.5	131
15	69	48	17.4	26.8	34.5	17.5	9.32	4.23	5.4	18	108	122
16	65.4	46	17.5	25.5	38.7	17	9.19	4.33	5.38	17.6	114	250
17	62.2	44.4	16.9	26.6	38.8	16.8	8.81	4.33	5.33	15.8	105	255
18	60.3	43.8	19.7	31.6	37.6	16.4	6.19	4.21	5.39	13.8	95.2	217
19	60.6	42.9	31.2	34	37	15.9	5.54	4.24	5.48	11	109	208
20	60.6	41.3	29.2	34.6	35.8	15.5	6.33	4.23	5.57	10.9	154	188
21	71	39.6	27	33.9	33.9	15.2	6.18	4.7	5.7	11.2	160	171
22	71.4	38.8	22.4	33.3	32.9	15	6.02	5.81	5.61	12.1	162	155
23	67.2	37.4	20.7	35.4	32.2	14.8	5.73	5.7	5.56	17.8	150	142
24	63.6	36.1	19.3	37.4	31.8	14.6	5.71	6.33	5.35	21.6	138	133
25	61.3	34.4	21	43.1	29.9	14.3	5.73	6.18	5.61	24.9	130	122
26	58.3	32.9	26.8	45	28.8	13.9	5.72	5.04	6.1	29.8	120	113
27	56.1	31.8	28.9	45.4	27.4	14.1	5.06	5.47	9.07	35.2	111	104
28	54.4	30.5	35.9	45.2	27	14.9	4.47	5.94	15.9	37.9	107	96.6
29	55.5		37.7	46.3	27.1	14.8	4.48	6.46	16.2	39.1	110	89.1
30	54.2		39.3	50.9	25.7	14.7	4.74	6.48	16.3	41.3	111	85.7
31	53.5		40.8		24.5		4.43	6.54		45.3		82.3
Total	2192	1381	762.8	1146	1263	534.1	275.8	155	210.3	566.9	2656	4179
Mean	70.7	49.3	24.6	38.2	40.7	17.8	8.9	5.0	7.0	18.3	88.5	134.8
Max	139	67.8	40.8	50.9	69.8	24.2	14.4	6.54	16.3	45.3	162	255
Min	53.5	30.5	15.8	25.5	24.5	13.9	4.43	4.21	5.33	9.09	44	82.3

¹ Water Survey of Canada data recorded at the Island Highway bridge in Duncan, BC.

Table 14a. Daily summary of carcasses examined, tags applied and tagged recoveries, by sex, for chinook in the upper Cowichan River, 2001.

Date	Carcasses Examined				Tags Applied				Tagged Carcasses Recovered			
	Unkn.	Male	Female	Jack	Unkn.	Male	Female	Jack	Unkn.	Male	Female	Jack
5-Nov	0	10	32	6	0	10	32	6	0	0	0	0
6-Nov	0	22	31	6	0	22	31	6	0	0	0	0
7-Nov	0	28	29	20	0	25	26	20	0	3	3	0
8-Nov	0	15	56	5	0	14	48	3	0	0	8	1
9-Nov	0	36	38	25	0	28	26	25	0	7	12	0
13-Nov	0	17	19	39	0	16	17	37	0	1	2	2
14-Nov	0	21	65	22	0	20	59	21	0	1	6	1
16-Nov	0	9	30	3	0	8	25	3	0	1	5	0
19-Nov	0	6	30	9	0	3	23	8	0	2	6	1
20-Nov	0	11	7	6	0	10	6	6	0	1	1	0
21-Nov	0	4	29	5	0	4	24	5	0	0	5	0
22-Nov	0	13	31	6	0	9	22	3	0	4	9	3
23-Nov	0	20	44	10	0	15	30	9	0	5	14	1
26-Nov	0	11	22	9	0	9	17	6	0	2	5	3
27-Nov	0	14	38	9	0	11	28	9	0	3	9	0
29-Nov	0	4	20	2	0	3	11	2	0	1	8	0
30-Nov	0	3	9	0	0	3	6	0	0	0	3	0
3-Dec	0	12	20	6	0	10	17	5	0	2	3	1
4-Dec	0	4	7	4	0	2	6	4	0	1	1	0
5-Dec	0	3	4	1	0	3	2	1	0	0	2	0
6-Dec	2	7	13	1	0	2	7	1	0	5	6	0
7-Dec	0	2	3	1	0	0	2	1	0	2	1	0
Total	2	272	577	195	0	227	465	181	0	41	109	13

Table 14b. Tags applied, carcasses examined and marks recovered, by sex, for chinook in the upper Cowichan River, 2001.

Sex	Tags Applied	Carcasses Examined	Marks Recovered	Percent Recovered
Male	227	272	41	18.1%
Female	465	577	109	23.4%
Jack	181	195	13	7.2%

Table 15. Petersen mark recaptures estimates, stratified by sex, for Cowichan River chinook, 2001.

Sex	Population Estimate	95% Confidence Limits	
		Lower	Upper
Male	1460	1059	1862
Female	2436	2028	2844
Total Adult	3869	3311	4427
Jack	2535	1299	3771
Total Population	5526	4752	6301

Table 16. Incidence of tagged adult chinook carcasses recovered on the spawning grounds by recovery period and sex, on the upper Cowichan River, 2001.

Recovery Period	Days of Recovery	Tagged Recoveries			Total Recoveries			Tag Incidence (%)		
		Male	Female	Total	Male	Female	Total	Male	Female	Total
Nov 5 - Nov 12	8	10	23	33	111	186	297	9.01	12.37	11.11
Nov 13 - Nov 20	8	6	20	26	64	151	215	9.38	13.25	12.09
Nov 21 - Nov 28	8	14	42	56	62	164	226	22.58	25.61	24.78
Nov 29 - Dec 7	9	11	24	35	35	76	111	31.43	31.58	31.53
Total	33	41	109	150	272	577	849	15.07	18.89	17.67
Chi-Square Test Result:								14.86	21.13	
Critical Chi-Square (df = 3; alpha = 0.05)								7.82	7.82	

Table 17. Proportion of the tag application sample recovered on the spawning grounds, by application period and sex, in the Cowichan River, 2001.

Application Period	Days of Application	Tags Applied			Tagged Recoveries			Percent Recovered		
		Male	Female	Total	Male	Female	Total	Male	Female	Total
Nov 5 - Nov 12	8	99	163	262	10	23	33	10.10	14.11	12.60
Nov 13 - Nov 20	8	57	130	187	6	20	26	10.53	15.38	13.90
Nov 21 - Nov 28	8	48	121	169	14	42	56	29.17	34.71	33.14
Nov 29 - Dec 7	9	23	51	74	11	24	35	47.83	47.06	47.30
Total		227	465	692	41	109	150	18.06	23.44	21.68
Chi-Square test result:								24.19	15.77	
Critical Chi-Square (df = 3; alpha = 0.05)								7.82	7.82	

Table 18. Sex composition of chinook in the tag application and recovery samples on the upper Cowichan River, 2001.

Sex	Application sample by recovery status				Recovery sample by mark status			
	Sample Size	Recovered	Not Recovered	Total	Sample Size	Marked	Not Marked	Total
Male	227	27.3%	34.3%	32.8%	272	27.3%	32.8%	32.0%
Female	465	72.7%	65.7%	67.2%	577	72.7%	67.2%	68.0%
Chi-Square test result:				2.60	1.85			
Critical Chi-Square (df = 1; alpha = 0.05)				3.84	3.84			

Table 19. Sex composition of chinook in the tag application and recovery samples on the upper Cowichan River, 2001 (jacks included).

Sex	Application sample by recovery status				Recovery sample by mark status			
	Sample Size	Recovered	Not Recovered	Total	Sample Size	Marked	Not Marked	Total
Male	227	25.2%	26.2%	26.0%	272	25.2%	26.0%	26.1%
Female	465	66.9%	50.1%	53.3%	577	66.9%	53.3%	55.3%
Jack	181	8.0%	23.7%	20.7%	195	8.0%	20.7%	18.7%
Chi-Square test result:				22.76	16.61			
Critical Chi-Square (df = 2; alpha = 0.05)				5.99	5.99			

Table 20. Total adult chinook returns to the Cowichan River for the years 1975 to 2001.

Year	Natural Spawners	Broodstock Removal	Native Catch	Total Return
1975	6500		900	7400
1976	3460		1000	4460
1977	4150		1000	5150
1978	4370		500	4870
1979	8750	195	500	9445
1980	5950	337	1500	7787
1981	5500	282	1500	7282
1982	4500	534	1000	6034
1983	4500	242	250	4992
1984	5000	278	355	5633
1985	3500	175	1000	4675
1986	1832	315	800	2947
1987	1937	582	800	3319
1988	6200	678	681	7559
1989	5000 ¹	535 ²	1055	6590
1990	5300	326	820	6446
1991	6000 ³	1755	250	8005
1992	8500	1850	260	10610
1993	5058	1970	295	7323
1994	5050	1357	345	6752
1995	14300	2149	533	16982
1996	12980	1615	810	15405
1997	9845	125	191	10161
1998	4371	1485	1073	6929
1999	4500	1659	233	6692 ⁴
2000	5109 ⁵	1529	89	7024 ⁴
2001	3282 ⁵	1732	918	5998 ⁴

¹ For the years 1989 to the present, the number of natural spawners was calculated as the number of adults recorded at the fence minus the adults removed for broodstock above the fence. In years when fence counts were incomplete, the cumulative run timing curve was used to expand the count.

² This number is the total broodstock removed and may include some jacks.

³ Due to early flooding, estimate is based on expansion of swim surveys and weir counts.

⁴ Includes an estimated 300 chinook lost to seal predation in the Cowichan estuary.

⁵ For the years 2000 to the present, the adult fence count totals used in calculating natural spawners was adjusted using jack/adult ratios obtained from the spawning grounds.

Fig. 1 Cowichan River Survey Areas:

Swim survey locations were:

- 1 - Bird House Pool
- 2 - Road Pool
- 3 - Train Trestle (mile 70.2)
- 4 - Old Pick-up Site
- 5 - Maple Tree
- 6 - Three Firs Pool
- 7 - Skutz Falls
- 8 - Marie Canyon
- 9 - Bible Camp
- 10 - Cowichan Side channel
- 11 - Sandy Pool
- 12 - Sewer
- 13 - JC Pool

Swim survey areas:


Bird House (1) to Three Firs pool (6) represents the Upper survey section.

Marie Canyon (8) to enumeration fence (A) represents the Middle survey section.

A - refers to the adult enumeration fence


Tag recovery locations:

Locations numbered 1 to 45 are in the upper river section, those numbered 46 to 83 are in the middle river section.



**Fig. 2 River Management Zones for
Native Food Fishery**

- A-Cliffs to Silver bridge
- B-Silver bridge to JC's place
- C-Quamichan to Black creek
- D-Powerline to Elliot's barn
- E-Elliot's barn to Brian's pool
- F-Brian's pool to Clem Clem and
part of Koksilah
- G-Clem Clem to mouth
- H-North side to Four plex
- I-Four plex to Meriner's
slough
- J-Meriner's slough to mouth






Figure 3a. Daily fence counts of adult and jack chinook and water temperature at the fence site.




Figure 3b. Daily fences count of adult and jack chinook and water depth at the fence site.




Figure 4. Adult chinook catch in the Cowichan First Nations fishery for the years 1971 to 2001.




Figure 5. Monthly Cowichan River discharge (m^3/s) in 2001 along with historical values.




Figure 6. Adult and jack chinook length frequencies collected from the Cowichan River spawning grounds, 2001.




Figure 7. Annual adult chinook escapement estimates for the Cowichan River for the years 1953 to 2001.




Figure 8. Annual releases of hatchery chinook into the Cowichan River as fry (3 g) and as pre-smolts (6 g).




Figure 9. Natural and enhanced contribution to the Cowichan River chinook escapement for the years 1982 to 2001.