

Adult and Juvenile Coho Salmon Enumeration and Coded-wire Tag Recovery Analysis for Zolzap Creek, BC, 2002

B.E. Baxter

**Fisheries and Oceans Canada
Room 202, 417-2nd Ave.
Prince Rupert, BC
V8J 1G8**

2003

**Canadian Manuscript Report of
Fisheries and Aquatic Sciences No. 2646**



**Fisheries
and Oceans**

**Pêches
et Océans**

Canada

Canadian Manuscript Report of Fisheries and Aquatic Sciences

Manuscript reports contain scientific and technical information that contributes to existing knowledge but which deals with national or regional problems. Distribution is restricted to institutions or individuals located in particular regions of Canada. However, no restriction is placed on subject matter, and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Manuscript reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 1426 - 1550 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Manuscript Reports. The current series name was changed with report number 1551.

Manuscript reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

Rapport manuscrit canadien des sciences halieutiques et aquatiques

Les rapports manuscrits contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui traitent de problèmes nationaux ou régionaux. La distribution en est limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports manuscrits peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports manuscrits sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1 à 900 de cette série ont été publiés à titre de manuscrits (série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés comme manuscrits (série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros 901 à 1425 ont été publiés à titre de rapports manuscrits de l'Office des recherches sur les pêcheries du Canada. Les numéros 1426 à 1550 sont parus à titre de rapports manuscrits du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 1551.

Les rapports manuscrits sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Manuscript Report of
Fisheries and Aquatic Sciences 2646

2003

ADULT AND JUVENILE COHO SALMON ENUMERATION AND CODED-WIRE TAG
RECOVERY ANALYSIS FOR ZOLZAP CREEK, BC, 2002

prepared by

Bruce E. Baxter

LGL Limited
environmental research associates¹

for the

Nisga'a Lisims Government²

¹9768 Second St., Sidney, BC V8L 3Y8

²PO Box 228, New Aiyansh, BC V0J 1A0

© Minister of Supply and Services Canada 2003
Cat. No. Fs 97-4/2646E ISSN 0706-6473

Correct citation for this publication:

Baxter, B.E. 2003. Adult and juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 2002. Can. Manusc. Rep. Fish. Aquat. Sci. 2646: vii + 44 p.

TABLE OF CONTENTS

LIST OF TABLES	iv
LIST OF FIGURES	v
LIST OF APPENDICES	v
ABSTRACT	vi
RÉSUMÉ	vii
INTRODUCTION	1
STUDY STREAM	1
JUVENILE COHO STUDIES	2
METHODS	2
Trapping Operations	2
Physical Observations	2
Fish Enumerations	2
Biosampling	3
Coded-wire Tagging	3
RESULTS	4
Physical Observations	4
Fish Enumerations	4
Coho Smolts	4
Non-coho Species	4
Biosampling: Length, Weight, and Age	4
Coded-wire Tagging	5
ADULT COHO STUDIES	5
METHODS	5
Population Estimates	5
Biosampling	5
Coded-wire Tag Recoveries	6
Escapement	6
Commercial and Sport Harvests	6
Geographic Distribution of Harvest	6
RESULTS	7
Physical Observations	7
Adult enumerations	7
Mark-recapture Estimates	7
Biosampling - Age and Length	8
Coded-wire Tag Recoveries	8
Escapement	8
Commercial and Sport Harvests	8
DISCUSSION	9
ACKNOWLEDGMENTS	11
REFERENCES	12
TABLES	14
FIGURES	28
APPENDICES	38

LIST OF TABLES

Table 1.	Age-length distribution of Zolzap Creek coho smolts, 2002.....	15
Table 2.	Coho smolt catch at Zolzap Creek enumeration fence, by week, in 2002.....	16
Table 3.	Non-coho catch at the spring juvenile and fall adult fences at Zolzap Creek, 1992-2002.....	17
Table 4.	Coded-wire tag retention rates for Zolzap Creek coho smolts, 2002.....	18
Table 5.	Coded-wire tagged coho smolt releases from Zolzap Creek, 2002.....	19
Table 6.	Fence enumerations, carcass recoveries, and Peterson population estimates for adult coho escapement at Zolzap Creek, 2002.....	20
Table 7.	Freshwater age distribution of adult coho at Zolzap Creek, 2002.....	21
Table 8.	Estimates of total escapement of adipose clipped coho and contribution to escapement at Zolzap Creek, 1993-2002.....	22
Table 9.	Estimated Canadian and American commercial and sport harvest of Zolzap Creek CWT coho in 2002 using tag recovery data (Mark Recovery Program, Fisheries and Oceans, Canada and ADF&G mark tag and age lab, online searchable database).....	23
Table 10.	Expanded Canadian and American commercial and sport harvest of Zolzap Creek coho and estimated total return in 2002 from 2001 smolt year.....	24
Table 11.	Estimated commercial harvest distribution of Zolzap Creek CWT coho by area and gear type, 2002.....	25
Table 12.	Adult and juvenile coho abundance and smolt-adult survival, by smolt year at Zolzap Creek, 1992-2002.....	26
Table 13.	Adult and juvenile coho production and age composition by brood year, Zolzap Creek, 1990-1999 ^a	27

LIST OF FIGURES

Figure 1.	The Nass River watershed, British Columbia.....	29
Figure 2.	Zolzap Creek and location of enumeration fence.....	30
Figure 3.	Water level and temperature at Zolzap Creek, 2002.....	31
Figure 4.	Daily migration of coho smolts at Zolzap Creek, 6 May - 21 June, 2002.....	32
Figure 5.	Length-frequency and calculated age distribution of Zolzap Creek coho smolts, 2002.....	33
Figure 6.	Daily counts of adult coho at the Zolzap Creek enumeration fence, 31 Aug - 5 Oct, 2001.....	34
Figure 7.	Length-frequency distribution of coho, by sex, Zolzap Creek, 2002.....	35
Figure 8.	Fisheries Statistical Areas for the north coast of British Columbia and southeast Alaska, and commercial harvest distribution of Zolzap Creek CWT coho, 2002.....	36
Figure 9.	Exploitation rates for three wild coho indicator stocks.....	37
Figure 10.	Total percent survivals for three wild coho indicator stocks.....	37
Figure 11.	Canadian and Alaskan exploitation rates on Zolzap Creek coho, 1993-2002.....	37

LIST OF APPENDICES

Table A-1.	Juvenile coho catch at Zolzap Creek enumeration fence, 2002.....	39
Table B-1.	Non-coho catch at Zolzap Creek enumeration fence, 2002.....	41
Table C-1.	Coded-wire tagging data for coho smolts at Zolzap Creek, 2002.....	43
Table D-1.	Daily counts of adult coho at Zolzap Creek enumeration fence, 2002.....	44

ABSTRACT

Baxter, B.E. 2003. Adult and juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 2002. *Can. Manusc. Rep. Fish. Aquat. Sci.* 2646: vii + 44 p.

Adult and juvenile coho migrations were monitored at Zolzap Creek, British Columbia, as part of the 2002-2003 Nisga'a Fisheries Program. The 2002 season is the eleventh year of continuous operation of the Zolzap Creek fences since 1992. This report includes eleven year summaries of the most pertinent data. Smolt trapping was conducted from 29 April to 26 June 2002 using an in-stream wire-mesh fence. A total of 15,001 coho smolts were captured during the trapping period, and an unknown number migrated out during periods when the fence was not operational. Of those captured, 12,318 were released with coded-wire tags. Migration timing, mean length and weight at age, and age composition are presented.

Adult coho escapement was monitored using an in-stream fence and carcass surveys. The counting fence was operational between 21 August and 5 October. A total of 1,918 adult coho were counted at the fence with an estimated escapement of 3,233 using the adjusted Peterson model. Adipose clip rate was 46.4% for adult coho. Age and length characteristics of adult males and females are presented.

Canadian and US commercial harvests were examined using coded-wire tag recovery data obtained from the Mark-Recovery Program and the Alaska Department of Fish and Game (ADF&G) mark tag and age lab online searchable database. Total exploitation rate on Zolzap Creek coho in 2002 was 19.9% (3.3% Canadian, 16.6% US). Of the total commercial catch of Zolzap Creek coho, Canadian catch accounted for 16.5% and the US catch accounted for an estimated 83.5%. Harvests occurred over a wide area ranging from SE. Alaska to the US Northern Outside Statistical Area in Alaska (northwest of Juneau, AK). Limited Canadian commercial harvests occurred in Areas 1-5 for Zolzap Creek in 2002. US harvests of Zolzap coho in Alaska were largest in the Southern Inside Statistical Area for the net fishery and the Central Outside Statistical Area for the troll fishery. Total survival was 8.4% and smolt-to-spawner survival was 6.7%.

RÉSUMÉ

Baxter, B.E. 2003. Adult and juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 2002. Can. Manusc. Rep. Fish. Aquat. Sci. 2646: vii + 44 p.

Les migrations de saumons coho, adultes et jeunes, ont été mesurées au ruisseau Zolzap en Colombie-Britannique, dans le cadre de la Stratégie des pêcheries autochtones des Nisga'a en 2002-2003. L'année 2002 marque la 11^{ème} saison d'opération continue des barrières en fil métallique du ruisseau Zolzap depuis 1992. Ce rapport contient 11 ans de sommaires des données les plus intéressantes. Le piégeage des saumoneaux prit place entre le 29 avril et le 26 juin 2002 à l'aide d'une barrière en fil métallique installée dans le ruisseau. En tout, 15,001 saumoneaux coho ont été capturés pendant la période de piégeage tandis qu'un nombre inconnu a migré quand la barrière n'était pas opérationnelle. Sur l'ensemble des saumoneaux capturés, 12,318 ont été remis à l'eau avec une marque magnétique codée. La période de migration, la longueur moyenne, le poids et la composition selon l'âge sont présentées.

L'échappée de saumons coho adultes a été surveillée grâce à une barrière installée dans le ruisseau et à l'observation des carcasses. La barrière de comptage fut opérationnelle entre le 21 août et le 5 Octobre. Un total de 1,918 saumons coho adultes ont été dénombrés à la barrière avec une échappée estimée à 3,233 utilisant le model Peterson ajusté. Le taux d'ablation de la nageoire adipeuse était de 46.4% pour les saumons coho adultes. Nous présentons les caractéristiques d'âge et de longueur pour les mâles et les femelles adultes.

Les récoltes commerciales canadiennes et américaines ont été examinées grâce aux données de récupération des marques magnétiques codées provenant du Programme de marquage-récupération et en direct de la base de données du Département de Pêche et Chasse de l'Alaska. En 2002 le taux total d'exploitation commerciale de saumon coho au ruisseau Zolzap fut évalué à 19.9% (3.3% pour le Canada, 16.6% pour les États-Unis). Sur le total de prises commerciales de saumon coho au ruisseau Zolzap, le Canada en comptait 16.5% et les États-Unis, une estimation de 83.5%. Les récoltes couvraient un vaste secteur, s'étendant à partir du sud-est de l'Alaska jusqu'à la zone statistique nord extérieure de l'Alaska aux États-Unis (au nord-ouest de Juneau, AK). Des récoltes commerciales Canadiennes limitées ont prit place dans le secteur 1-5 pour le ruisseau Zolzap in 2002. Les saumons coho du Zolzap récoltés par les États-Unis en Alaska furent plus nombreux dans la zone statistique sud intérieure pour la pêche au filet, et, dans la zone statistique centrale extérieure pour la pêche à la traîne. Le taux total de survie fut 8.4% tandis que pour les saumoneaux/géniteurs le taux de survie fut 6.7%.

INTRODUCTION

As part of the Aboriginal Fisheries Strategy (AFS), a program was established for fisheries research in the Nisga'a Traditional Territory, British Columbia. One component of this large research initiative focused on the assessment of juvenile and adult coho populations in tributaries to the Nass River. Juvenile and adult coho enumeration studies have been conducted on Zolzap Creek since 1992 (Nass 1996a; Nass 1996b; Nass 1996c; Nass and English 1994; Nass 1997a; Nass 1997b; Nass 2001; Nass and Frith 2001; Baxter et al. 2001; Baxter and Stephens 2002, Baxter and Stephens 2002a, Baxter and Stephens 2002b). This report presents results for studies conducted at Zolzap Creek in 2002.

The objectives of the research were to:

1. Enumerate migrating juvenile coho and estimate escapement;
2. Document the timing, size, and age distribution of migrating coho;
3. Mark coho smolts with coded-wire tags (CWT) to enable the determination of oceanic harvest rates;
4. Monitor the escapement for marked CWT adult coho, and determine ocean exploitation and survival rates; and
5. Collect water temperature and level data for future examination of the relationships between physical environmental factors and coho smolt migration timing, and between adult escapement and smolt production.

Achievement of these objectives involved the construction and operation of in-stream, semi-permanent, panel fences located approximately 0.5 km upstream of the mouth of Zolzap Creek.

STUDY STREAM

Zolzap Creek is a tributary to the Nass River, located in northwestern British Columbia (Figures 1 and 2). Zolzap Creek flows for 6 km in a northwesterly direction between Nisga'a Lava Bed Memorial Park and the Kitimat Mountain Range to its confluence with the Nass River, 5 km downstream of Gitwinksihlkw. The main channel of the creek is regularly interrupted by beaver dams and log jams. The substrate is highly variable and ranges between silty particulate, to granite cobble, and coarse pumice. Major flow contributions come from Lava Creek (3 km in length) which flows from the lava beds and numerous small creeks that flow from the steep alpine. Intermittent flows of water from the Nass River and Vedder Creek are possible during flooding periods. The mouth of Zolzap Creek enters a side channel to the Nass River known as Zolzap Slough. The lower 0.5 km of Zolzap Creek regularly becomes inundated when water levels on the Nass River are high. Zolzap Creek supports many species of salmonids including coho (*Oncorhynchus kisutch*), pink (*O. gorbuscha*), chum (*O. keta*), sockeye (*O. nerka*), rainbow

(*O. mykiss*), cutthroat (*O. clarki*), and Dolly Varden (*Salvelinus malma*). Non-coho species include lampreys (family Petromyzontidae), sticklebacks (family Gasterosteidae), and sculpins (family Cottidae). Coho escapement was estimated to be 1,561 in 1992 (Nass 1996b), 1,048 in 1993 (Nass 1996c), 2,536 in 1994 (Nass 1997a), 908 in 1995 (Nass 1997b), 1,039 in 1996 (Nass 2001), 470 in 1997 (Nass and Frith 2001), 967 in 1998 (Baxter et al. 2001), 1,393 in 1999 (Baxter and Stephens 2002), 456 in 2000 (Baxter and Stephens 2002a), and 1,897 in 2001 (Baxter and Stephens 2002b).

JUVENILE COHO STUDIES

METHODS

Trapping Operations

An in-stream, semi-permanent enumeration fence was located 0.5 km upstream of the creek mouth for the capture of downstream migrating coho smolts. Fence design was based on Conlin and Tutty (1979) and minor modifications were required due to site characteristics and available materials. The fence was built in a W-pattern and spanned the entire creek bed. Three by eight foot panels constructed of 2x4's and covered with 1/4" wire-mesh were laid on their long side in the creek bed to form the fence. Rebar of 3/8" and 1/2" diameter were used to anchor the panels to the stream bed. A second layer of panels were installed on top of the first row of panels to create a fence with a total height of six feet. Burlap sandbags and heavy duty plastic garden sheeting were used to seal the base of the panels. Two hinged panels were installed in each of the fence wings for release of excess water in the event of flooding. Plywood trap boxes with Vexar-screened windows (to allow water exchange) were anchored at each down-stream apex and were connected to the fence with 8" Big-O tubing. Additional boxes were made for holding fish after processing and were designed with a small door for releasing fish. Provisions for upstream migrating adults were made by constructing a simple trap consisting of a wire-mesh panel extending out from the stream bank to one wing of the fence. Plywood was used to cover the adult trap area.

Physical Observations

Crews monitored water temperatures, water levels, and weather daily. Crews recorded temperature to the nearest degree (1 °C) using a maximum-minimum thermometer and water level using staff gauges calibrated to the nearest centimeter (0.01 m). A total of three staff gauges were used; two were located within 50 m of the trapping site (one upstream, one downstream of the fence) and one approximately 1 km upstream of the fence. Precipitation was recorded on a scale of zero to five with zero representing no precipitation and five being heavy precipitation.

Fish Enumerations

Daily numbers of coho smolts captured at the fence were obtained from automatic counters on coded-wire tagging machines or by manual counts. The number of fence mortalities

was added to the total count. Coho juveniles with standard lengths greater than or equal to 70 mm were identified as smolts. Coho smaller than 70 mm tended to be dark with distinct parr marks and lacked the silver colouration typical of smolts. Therefore, this group consisted of pre-smolts and fry. All coho pre-smolts and fry, and non-coho species were counted and released downstream of the fence during sorting. Upstream migrating juveniles caught in the adult traps were counted and released upstream.

Biosampling

A random sample of up to 25 smolts (i.e., coho greater than or equal to 70 mm) were obtained from each day's catch. These smolts were anaesthetized and measured for fork length and weighed using an electronic scale (0.1 g). Scale sampling followed the stratified method of Ketchen, described by Ricker (1975); age sample data (column X on Table 1) included non-random samples, and length sample data (column Y on Table 1) and the calculated age representation was based on random sampling. Crews attempted to collect at least 10 scale samples from each 5 mm size class of coho for the study period. Smolts from under-represented size classes were selected to supplement random samples. Mean length and weight data was determined by multiplying the mean length and weight data for each 5 mm bin class by the total number of length and weight samples in that bin class (factor) to come up with a weighted mean length and weight for that bin class. The average length and weight for all sampled fish was determined by summing all the weighted length and weight measurements and dividing by the overall sum of the factors. Scale samples were interpreted by the Fisheries and Oceans Canada Scale Lab, Nanaimo, BC. Secondary quality control checks were performed to ensure a reliable age designation. Scale ages are reported in Gilbert-Rich notation where freshwater age-2 coho (i.e., having survived two winters from egg deposition) have a single freshwater annulus.

Coded-wire Tagging

Coded-wire tagging at Zolzap Creek was performed using a Mark IV tagging machine (Northwest Marine Technology Ltd. Shaw Island, WA). Smolts were anaesthetized in a MS222 bath prior to tagging. All tagged fish were adipose fin-clipped (AFC). The numbers of coho smolts tagged with each tag code and the number of smolts untagged were recorded. All tagged smolts were placed in a holding box in the stream and allowed to recover from the tagging operation before release.

Tag retention tests were conducted for each tag code. A sample of tagged coho smolts (minimum of 200 smolts) were retained in a holding box from 24 h to 72 h. Following the holding period, smolts were lightly anaesthetized and checked for the presence of a coded-wire tag using the quality control device (QCD) from the coded-wire tagging machine. Coho smolts not possessing a tag were checked a second time. The total number of tags detected for each tag group and the total number of fish tested was recorded.

RESULTS

Physical Observations

Water temperatures during the smolt migration period at Zolzap Creek ranged from a minimum of 3 °C in early May to a maximum of 9 °C in mid-June (Figure 3A). Water level at gauge 2 (50 m upstream of the fence) remained steady at a gauge height of approximately 0.3 m from the beginning of monitoring on 29 April until 19 May. Water level rose steadily to a level of 0.8 m on 23 May and then rose rapidly to a level of 1.4 m on the 29 May. The fence was flooded out for four days (15-18 June) with water levels declining afterwards until the end of the spring monitoring period. High water levels in Zolzap Creek occur when the Nass mainstem flow rises causing water to back-up into the creek. Water flow in Zolzap Creek declines to very low velocities during these flooding events.

Fish Enumerations

The Zolzap Creek juvenile counting fence was operated from 29 April to 26 June 2002. The fence was topped for a four day period from 15-18 June and then operated without further interruptions until the end of the spring monitoring period. Approximately 25 to 30 baited gee traps were used to supplement catches at the fence during periods of high water and low smolt movement.

Coho Smolts: A total of 15,001 coho smolts were counted at the fence and included gee trap catches (Table 2). The maximum daily number of smolts captured at the fence was 1,833 and occurred on 21 May (Table A-1, Figure 4). There were a total of 2,891 fry and pre-smolt coho counted and released during trapping operations and 25 mortalities (Table A-1).

Non-coho Species: Lampreys (larvae and young adults) were caught in the largest numbers, followed by juvenile Dolly Varden, juvenile cutthroat, and juvenile steelhead (Table 3, Table B-1).

Biosampling: Length, Weight, and Age

The mean fork length of age-2 smolts was 98.3 mm and the mean weight was 9.7 g (Table 1). Age-3 smolts averaged 109.7 mm and 13.1 g. The length-frequency distribution showed substantial overlap between age-2 and age-3 coho (Figure 5). Age-2 smolts were most numerous in the 95 - 100 mm length class and age-3 smolts were most numerous in the 100 - 110 mm length class. Age-3 coho smolts were significantly larger than age-2 smolts (t-test, $p < 0.05$). Overall, coho smolts averaged 101.5 mm in length. The calculated freshwater age structure of coho smolts was 69.6% age-2 and 30.4% age-3 (Table 1).

Coded-wire Tagging

Mean tag retention was 98.7% for tag code 28-01-10, and 99.0% for tag code 28-01-11 (Table 4). Crews conducted 10 tests for tag code 28-01-10 for a total of 1,972 samples with 25 tag losses, and 3 tests for tag code 28-01-11 for a total of 600 samples with 6 tag losses.

Releases of adipose fin-clipped coho totalled 12,412 (Table 5; Table C-1). Crews recorded 80 mortalities associated with the tagging process. The total number of coho smolts released with coded-wire tags was 12,318. Approximately 4% (468) of the captured coho smolts were released untagged during the study period and thus the mark rate of coho smolts released was 1.05 (Table 5). The total number of smolts released was 12,880.

ADULT COHO STUDIES

METHODS

Population Estimates

An aluminum conduit fence anchored to a crib-type sill was constructed at Zolzap Creek. All salmonids caught at the fence were counted and classified by sex. Sex was distinguished on the basis of length and body morphology. Previous studies at Zolzap Creek (Nass 1996b, 1996c, 1997a, 1997b, Nass 2001, Nass and Frith 2001, Baxter et al. 2001, Baxter and Stephens 2002, Baxter and Stephens 2002a, Baxter and Stephens 2002b) have shown an absence of jacks in the escapement, and therefore all males were classified as adults. "Jack panels" consisting of 1" wire mesh were used to prevent the passage of small coho through the fence and were used whenever water levels and debris permitted. Each coho was tagged on the operculum with a uniquely numbered Ketchum kurl-lock tag and measured for length. During handling, fish were examined for fin clips or tags that would be associated with coded-wire tagging or mark-recapture studies taking place on the Nass River. All captured fish were released upstream of the fence.

Adult coho abundance downstream of the fence was assessed later in the migration period due to the lack of fish movement past the fence. Delayed migration was the result of persistent low water conditions in Zolzap Creek in the later fall period. During these periods of delayed fish movement, angling was conducted approximately 1 km downstream of the fence in Zolzap Slough to determine relative coho abundance. Live coho were recaptured in upstream surveys and checked for operculum tags. Carcasses were recovered on the fence and during upstream surveys. In 2002, carcasses were recovered primarily in the lower 5 km of the creek.

Biosampling

All live coho captured at the fence were measured for postorbital-hypural length and examined for fin clips and sex. Data recorded from coho captured at the fence were used to calculate sex ratios and mean-length by sex. Crews attempted to sample at least 25 coho a day for scales (5 scales per fish). Scale samples were sent to the Fisheries and Oceans Canada scale

lab, Nanaimo, BC for age determination. Secondary quality control checks were performed at the scale lab to ensure reliability of the age designations. Scale ages are reported in Gilbert-Rich notation where freshwater age-2 coho (i.e., having survived two winters from egg deposition) have a single freshwater annulus.

Adult returns (calculated by escapement method) and smolt production, by CWT and total populations, were calculated for each brood year where data was available. Smolt output and adult escapement were apportioned between brood years (back-calculated) using the age structure observed in the respective yearly migrations. The sum of freshwater age-2, age-3, and age-4 individuals equals total production for a given brood year. Age composition for smolts and adults by brood year were calculated based on the estimated production. Total survival by brood year was calculated as the age specific adult return divided by the respective smolt production. Smolt-to-spawner recruitment for each brood year was calculated as the number of smolts produced divided by the number of adults in the escapement, by brood year. Similarly, the recruit-to-spawner ratio for each brood year was calculated as the number of adults produced divided by the number of adults in the escapement, by brood year.

Coded-wire Tag Recoveries

Coded-wire-tagged smolts were adipose fin-clipped prior to release. Coho smolts at Zolzap Creek were coded-wire tagged in the spring of 2001 (Baxter and Stephens 2002b) during out-migration.

Escapement: Crews examined all coho captured at the fence for the presence or absence of the adipose fin. The contribution and survival of AFC coho to the escapement was determined using methods presented in Bocking et al. (1992) and modified in Nass (1997a). Coded-wire tagged heads were collected from fish captured at the Nass River fishwheels, fish recovered in the native angling fishery below and above the fence, and from carcass recoveries.

Commercial and Sport Harvests: Commercial and sport catches of CWT fish are monitored by the Fisheries and Oceans Canada and various US agencies and compiled in the Mark Recovery Program (MRP) and in the ADF&G mark tag and age lab online searchable database. Data on CWT releases and recaptures are used to estimate the number of fish from a particular stock that have been harvested in the commercial and sport fishery, as well as determining the spatial and temporal distribution of harvests (Kuhn et al. 1988, Nass 1997a). The estimates include catch (observed catch corrected for sampling effort), expanded catch (estimated catch corrected for unmarked fish), exploitation rate (proportion of CWT coho caught in the fishery), and total return (expanded catch plus escapement).

Geographic Distribution of Harvest: Coded-wire tagged fish in the commercial catch are recorded by Canadian and US fishery Statistical Areas. To estimate number of recoveries for each Canadian area, the observed CWT catch was expanded by the mean catch-sampling ratio observed in the Catch Region (e.g., Northern Troll = Stat. Areas 1, 3, 4, and 6). Similarly, US troll catch was expanded using the catch sampling ratio by quadrant (e.g., northwest) and the net catch sampling ratios, by district.

RESULTS

Physical Observations

During the period that the adult fence was operational, water temperatures ranged from a maximum of 10 °C in early September to a minimum of 8 °C in early-October (Figure 3B). Water level ranged from 0.3 m during base flows to 1.2 m during freshets (Figure 3B).

Adult enumerations

The fence was operated continuously from 21 August to 4 October. A rain on snow event that occurred on the 5 October resulted in extreme flow conditions and caused a complete failure of the adult fence. A total of 1,918 adult coho salmon were counted at the fence including 6 coho released untagged (Table 6). Of these, 1,818 adults (adjusted for tag loss) were operculum tagged and released upstream. Maximum daily migration past the fence was 399 adults on 19 September (Table D-1) (Figure 6).

Coho capture surveys were conducted below the fence on 2, 3, 5, 6 and 7 September. A total of 39 coho were caught during these surveys. All coho caught were sexed, opercular tagged, checked for AFC's and then transported by boat above the fence.

For non-coho species captured at the fence, pink salmon had the greatest abundance (17), followed by cutthroat (12), chum salmon (5), and steelhead (4). Dolly Varden (3) and sockeye (1) were also captured at the fence (Table 3). Chum, pink, and sockeye were caught in their greatest numbers in early-mid September. Cutthroat and Dolly Varden were mainly caught in mid-late September. The number of chum, sockeye and pink caught in 2002 were much lower than in 2001 and well below the 1992-2001 averages (44, 13 and 87 respectively). No population estimates were derived for non-coho species.

Mark-recapture Estimates

Crews examined a total of 653 adult coho carcasses collected on the fence, and in twenty-three upstream surveys. Surveys were conducted upstream of the fence from 11 October to 3 December at three access locations along the creek. Upstream surveys were conducted on 15, 24, and 28 October, 6, 12, 19 and 26 November and 3 December at Goat Creek (a tributary); 11, 16, 21, 29 October, 13, 18 and 25 November and 2 December at upper Zolzap Creek. Of the 653 adult coho examined, 367 were tagged, 269 were untagged and 17 had lost their tags, which resulted in a Peterson population estimate of 3,233 adults (2,919 to 3,580; 95% CI) escaping to Zolzap Creek in 2002 (Table 6). An undetermined number of coho were observed spawning below the fence and in the Zolzap Slough area, so our estimate of 3,233 adult coho is likely underestimated.

Biosampling - Age and Length

A total of 216 coho were sampled for scales, of which 183 were successfully aged (Table 7). Unaged samples included marine regenerates. Adult males and females had different age compositions which averaged 71.1% and 78.0% freshwater age-2, and 28.9% and 22.0% freshwater age-3 respectively. The overall age composition was 74.9% age-2, and 25.1% age-3. All aged scales were recorded as marine age-1 (i.e., having 1 marine annulus).

Mean lengths of adult males and females were 46.9 cm (n=753, SD=7.9) and 49.3 cm (n=1,111, SD=6.2), respectively. Adult male coho were widely distributed over the range of 30 to 71 cm with a mode of 52 cm (Figure 7). Female coho also had a mode of 52 cm with a range of 26 to 65 cm. For coho sexed during processing, adult females captured at the fence (n=1,111) were more abundant than males (n=753).

Coded-wire Tag Recoveries

Escapement: Crews examined 1,905 adult coho at the fence for fin clips of which 883 were AFC (46.4%; Table 8). An estimated 1,499 adipose clipped adult coho returned to Zolzap Creek in 2002. An undetermined number of AFC coho spawned below the fence and in the Zolzap slough area. Smolt to spawner survival (i.e., includes natural and harvest mortality) for adult coho was estimated at 6.7%.

Eleven (11) CWT heads were collected at Zolzap Creek. Of these recoveries, 4 were from the native angling fishery below the fence and 7 were from carcass recoveries. In addition, 6 coho with adipose clips were recovered at the Nass River fishwheels. All of the CWT recoveries from Zolzap Creek were from the 2001 release at Zolzap Creek (codes 28-01-06 and 28-01-07). Of the 6 fishwheel CWT recoveries, 2 were from the 2001 release at Zolzap Creek (codes 28-01-06, 28-01-07), 1 was from the 2000 release at Zolzap Creek (code 28-01-05), and 3 were No-Pin.

Commercial and Sport Harvests: Total observed Zolzap Creek coho CWT recoveries were 7 and 102 for Canadian and US (Alaska) fisheries, respectively (Table 9). Observed sport recoveries totalled 0 for the Canadian fisheries and 4 for the Alaskan fishery. All CWT recoveries were from the 2001 release year. US troll and net catch to sample ratios were 2.7 and 4.9, respectively (Table 9). Estimated Zolzap Creek CWT coho catches were 57 (15%) and 312 (85%) for Canadian and US fisheries, respectively, and the Nisga'a food fishery at Zolzap Creek harvested an observed 4 CWT coho (Table 9).

Expanded Canadian and US catches were 78 and 391, respectively, for a total of 469 using the CWT mark ratio at release (i.e., MRP method) (Table 10). Expanded Canadian and US catches were 124 and 673, respectively, for a total of 797 using the adipose clip ratio at recovery (i.e., escapement method). Estimated total adult return for Zolzap Creek coho was 2,264 and 4,030 using the MRP and escapement methods, respectively (Table 10).

Of the total commercial catch of Zolzap Creek coho, Canadian fisheries accounted for 16.1% and the US accounted for 83.9% of the total commercial catch of Zolzap Creek coho (Table 11). US troll and net fisheries accounted for 76.2% and 23.8% of the total US catch, respectively. Canadian troll and net fisheries accounted for 15.6% and 0.6% of the total Canadian catch. Commercial harvest of Zolzap Creek coho occurred over a wide area ranging from Canadian Statistical Area 5 to the US Northern Outside Statistical Area in Alaska (Figure 8). US harvests were largest in the Southern Inside Statistical Area for the net fishery (15.0%) and the Central Outside Statistical Area for the troll fishery (29.6%; Table 11).

Total exploitation rate (Canadian and US combined) on Zolzap Creek coho in 2002 was 19.9% (Table 12). Total Canadian exploitation rate was 3.3% (3.0% troll, 0.1% net, and 0.2% terminal harvest by Nisga'a) and total US exploitation rate was 16.6% (11.5% troll, 4.4% net, and 0.7% sport). Total survival based on CWT returns was 8.4% (Table 12).

DISCUSSION

Over the past ten years of monitoring, the average number of smolts estimated leaving Zolzap Creek was 28,318 (Table 12). For the same time period, the average age composition of the smolt population was 65.7% age-2, 33.6% age-3, and 0.7% age-4 (Table 13).

Adult coho enumerated at the fence in 2002 (1,918) accounted for 59.3% of the Peterson population estimate (3,233). Therefore, approximately 1,315 adults entered Zolzap Creek during the period in which the fence was not operational. An undetermined number of coho were observed spawning below the fence and in Zolzap Slough. The observed native fishery harvested 7 coho below the fence of which 4 were CWT. The observed sport fishery harvested 3 coho below the fence of which 0 were CWT. Average escapement estimates for 1992 - 2001 was 1,395 (Table 12).

Data from 1992 to 2001 have indicated that there are no jacks in the Zolzap Creek escapement (Nass 1996b, 1996c, 1997a, 1997b, 2001, Nass and Frith 2001, Baxter et al. 2001, Baxter and Stephens 2002, Baxter and Stephens 2002a, Baxter and Stephens 2002b). In 2002, CWT and scale ageing data have confirmed again the absence of jacks in the population. There were 11 heads taken at Zolzap Creek for CWT sampling from coho measured between 42 and 58 cm (post-orbital-hypural) and all were found to be from 2001 releases. A post-orbital-hypural length of 35 cm has been used in previous studies at other BC streams to designate jacks in the escapement and is based on CWT analysis. Both the CWT analysis and scale ageing show that coho less than 42 cm from Zolzap Creek in 2002 were marine age-1.

The Department of Fisheries and Oceans, Canada, operates a juvenile and adult fence site at Lachmach River, BC which is used as a Northern BC wild coho indicator stock. Exploitation rates for Lachmach coho have ranged from 21.8% to 70% for the 1994-2001 period (Holtby et al 1999, Barry Finnegan, PBS, Nanaimo, pers. comm.). These exploitation rates are very similar to Zolzap exploitation rates for the 1994-1999 time period (Figure 9). Total survival for Lachmach coho has ranged from 5.5% (1997) to 17.4% (1994) and has been consistently higher than Zolzap Creek survivals (Figure 10).

In Alaska, comprehensive information exists for several southeast stocks, including Hugh Smith Lake (Southern Inside Statistical Area, see Figure 8), which has been monitored since 1982. Preliminary data for the 2002 return suggests exploitation rates of 0% Canadian and 37.4% US (37.4% total; Leon Shaul, Alaska Dept. of Fish and Game, Douglas, AK, pers. comm.). Southeast Alaska and Canadian fisheries accounted for approximately 100% and 0% of the commercial catch of Hugh Smith coho, respectively. The total exploitation rate on Hugh Smith coho (37.4%) was almost double as compared to exploitation rates on Zolzap Creek coho (19.4%) in 2002. Preliminary CWT data for the 2002 return of Hugh Smith coho suggest a survival rate of 14.4% which is higher than for Zolzap Creek coho at 8.4%. Hugh Smith coho have had substantially higher survivals (range 6.6% to 19.4%) compared to Zolzap coho (range 2.4% to 8.9%) in the past nine years.

Zolzap Creek CWT coho have been subjected to total exploitation rates between 19.4% and 72.3% and have had smolt-adult survival rates between 2.1% and 8.9% over the period 1992 to 2002 (Table 12, Figs. 9, 10). 2002 survivals were the second highest recorded since 1992. Canadian fisheries have had exploitation rates between 0% and 21.4% on Zolzap CWT coho, while US fisheries ranged between 17.0% and 54.8% (Figure 11). Of the total catch of Zolzap Creek coho, Canadian fisheries have averaged 17.7% and the US has averaged 82.3%, over ten years (Table 12).

Total smolt production by brood year averaged 28,066 (1990 - 1998) and was composed primarily of freshwater age-2 fish (66.7%; Table 13). Adult production by brood year averaged 3,082 (1990 - 1997) and was 56.7% age-2 fish. Age composition at return was substantially different from that observed in the respective smolt populations and varied widely. Freshwater age-4 fish were absent from all adult escapements with the exception of the 1995 brood year. Total smolt-adult return by brood year of all Zolzap coho (unmarked + CWT) averaged 11.6% (1990-1997; Table 13). Total smolt-adult return of Zolzap CWT coho was substantially lower at 4.8%. Higher survival for all coho compared to CWT coho is likely due to a significant number of unmarked smolts leaving Zolzap during non-operational periods (Nass 1996c). The effects of these conditions are evident from the historical data which shows the AFC at release has been roughly three times that of the AFC rate at return for the period 1993-1999 at Zolzap Creek (Table 8). Therefore, by using only CWT fish, the uncertainty around the number of fish released is eliminated and produces a more accurate estimate of survival for Zolzap coho smolts.

Estimates of total survival and exploitation are based on the assumption that all CWT coho are recovered in fisheries or on the spawning grounds. At Zolzap Creek, it is possible that the escapement of AFC coho is underestimated due to straying. Coho are known to spawn downstream of Zolzap Creek in Zolzap Slough (a side channel to the Nass River) where some CWT coho may return. In addition, a total of six adipose clipped coho were recovered in the fishwheels above Zolzap Creek in 2002 (2 were from the 2001 release at Zolzap Creek, 1 was from the 2000 release at Zolzap Creek and 3 were No-Pin) which tends to confirm our theory of straying. Straying would affect Zolzap Creek survival and exploitation estimates by underestimating survival and overestimating exploitation rates.

Zolzap Creek coho survivals may also be lower than Lachmach and Hugh Smith coho due to predator/prey interactions, with Zolzap Creek coho being more vulnerable to predation during their outmigration. Hugh Smith and Lachmach are both coastal systems and empty directly into marine waters, whereas Zolzap Creek empties into the Nass River. Zolzap Creek smolts must migrate approximately 33 km through Riverine habitat until they reach the ocean and are therefore more susceptible to predation along the way.

Persistent low water conditions at Zolzap Creek in the fall result in coho holding below the fence in Zolzap Slough until water levels rise. During certain low water years, this may result in coho spawning in the Slough area or pulsing through after the fence is demobilized. During these times, the run timing of the returning adult coho may be more an artifact of water levels rather than natural run timing. It is recommended that during subsequent years, the fence be run longer into the fall and increased effort be apportioned to upstream surveys to ensure a complete census of returning coho.

The number of smolts per spawner was 42.7 for the 1998 brood year. This value is conservative as the number of smolts released was likely underestimated. The number of recruits per spawner was 4.6 for the 1998 brood year (Table 13).

ACKNOWLEDGMENTS

The cooperation of many people was essential in meeting the objectives of this study. Special thanks go to Leonard Squires for monitoring all aspects of operations as crew supervisor. Simon Haldane, Vincent Johnson, Dan Gonu, Leonard Guno, Kirby Guno, Pamala Wright, and Lawrence Stephens assisted in constructing and operating the fence. Bob Bocking provided technical support and reviewed the manuscript. Doug Herriot and Brenda Adkins of the Department of Fisheries and Oceans provided the CWT catch data from the Mark Recovery Program. Barry Finnegan (PBS, Nanaimo) and Joel Sawada (DFO, Prince Rupert) provided the Lachmach data. Thanks also to Leon Shaul (ADF&G) for providing Hugh Smith data. Robin Tamasi provided mapping support. Funding for this project was provided by the Canadian Government and Nisga'a Lisims Government as part of the Nisga'a Final Agreement.

REFERENCES

- Baxter, B.E, and C. Y. Stephens. 2002. Adult and Juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 2000. Can. Manusc. Rep. Fish. Aquat. Sci. 2598: viii + 44 p.
- Baxter, B.E, and C. Y. Stephens. 2002a. Adult and Juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 1999. Can. Manusc. Rep. Fish. Aquat. Sci. 2597: viii + 44 p.
- Baxter, B.E, and C. Y. Stephens. 2002b. Adult and Juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 2001. Can. Manusc. Rep. Fish. Aquat. Sci. 2601: viii + 44 p.
- Baxter, B.E, C. Y. Stephens and B. L. Nass.. 2001. Adult and Juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 1998. Can. Manusc. Rep. Fish. Aquat. Sci. 2566: viii + 44 p.
- Bocking, R.C., R.E. Bailey, and J.R. Irvine. 1992. Coho salmon (*Oncorhynchus kisutch*) escapement studies in Black Creek, French Creek, and Trent River, Vancouver Island, 1989. Can. Man. Rep. Fish. Aquat. Sci. 2160: 77p.
- Conlin, K. and B.D. Tutty. 1979. Juvenile salmonid field trapping manual. Fish. Mar. Serv. Manusc. Rep. 1530: 136 p.
- Holtby, L. B., B. O. Finnegan, D. Chen, and D. Peacock. 1999. Biological assessment of Skeena River coho salmon. PSARC Working Paper S99-12:113p.
- Kuhn, B.R., L. Lapi, and J.M. Hamer. 1988. An introduction to the Canadian database on marked Pacific salmonids. Can. Tech. Rep. Fish. Aquat. Sci. 1649: viii + 56p.
- Nass, B.L. 2001. Adult and Juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 1996. Can. Manusc. Rep. Fish. Aquat. Sci. 2564: viii + 44 p.
- Nass, B.L. and H.R. Frith. 2001. Adult and juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 1997. Can. Manusc. Rep. Fish. Aquat. Sci. 2565: viii + 41 p.
- Nass, B.L. 1997a. Adult and Juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 1994. Can. Manusc. Rep. Fish. Aquat. Sci. 2420: viii + 54 p.

- Nass, B.L. 1997b. Adult and Juvenile coho salmon enumeration and coded-wire tag recovery analysis for Zolzap Creek, BC, 1995. Can. Manuscr. Rep. Fish. Aquat. Sci. 2423: viii + 54 p.
- Nass, B.L. 1996a. Enumeration and coded-wire tagging of coho salmon smolts at Zolzap Creek, and enumeration of coho salmon smolts at Seaskinnish and Ginlulak Creeks, 1992. Can. Manuscr. Rep. Fish. Aquat. Sci. 2376: viii + 44 p.
- Nass, B.L. 1996b. Escapement enumeration studies of adult coho salmon at Zolzap Creek, BC, 1992. Can. Manuscr. Rep. Fish. Aquat. Sci. 2374: viii + 30 p.
- Nass, B.L. 1996c. Escapement enumeration studies of adult coho salmon at Zolzap Creek, BC, 1993. Can. Manuscr. Rep. Fish. Aquat. Sci. 2373: viii. + 35 p.
- Nass, B.L. and K.K. English. 1994. Enumeration and coded-wire tagging of coho salmon smolts at Zolzap Creek, 1993. Report NF 93-01 prepared by LGL Limited, Sidney, BC for Nisga'a Tribal Council, New Aiyansh, BC.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Bd. Can. 191. 382 p.

TABLES

Table 1. Age-length distribution of Zolzap Creek coho smolts, 2002.

Size-Class (mm)	Age Sample (X)	Age-groups in X		Length Sample (Y)	Calculated Age Representation in Y	
		2	3		2	3
70	2	2	0	6	6.0	0.0
75	2	2	0	8	8.0	0.0
80	12	12	0	26	26.0	0.0
85	35	33	2	75	70.7	4.3
90	75	71	4	146	138.2	7.8
95	76	68	8	179	160.2	18.8
100	82	51	31	176	109.5	66.5
105	58	28	30	108	52.1	55.9
110	46	22	24	77	36.8	40.2
115	19	7	12	45	16.6	28.4
120	14	2	12	35	5.0	30.0
125	9	3	6	17	5.7	11.3
130	2	1	1	15	7.5	7.5
135	3	1	2	5	1.7	3.3
140	0	0	0	2	0.0	2.0
145	1	0	1	1	0.0	1.0
150	0	0	0	1	0.0	1.0
155	0	0	0	0	0.0	0.0
160	0	0	0	0	0.0	0.0
165	0	0	0	2	0.0	2.0
170	1	0	1	1	0.0	1.0
175	0	0	0	0	0.0	0.0
180	0	0	0	0	0.0	0.0
Mean length				101.5	98.3	109.7
SD				12.4	9.6	7.1
Mean weight (g)				10.7	9.7	13.1
SD				4.4	3.1	2.4
Total samples	437	303	134	925	644	281
% contribution		69.3	30.7		69.6	30.4

Table 2. Coho smolt catch at Zolzap Creek enumeration fence, by week, in 2002.

Week ending	Catch
11-May	493
18-May	2,247
25-May	5,430
1-Jun	2,068
8-Jun	3,095
15-Jun	1,644
22-Jun	24
Total	15,001

Table 3. Non-coho catch at the spring juvenile and fall adult fences at Zolzap Creek, 1992-2002^a.

Species	Time/lifestage	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	92-01 Avg.
Pink	Fall Adult	115	149	251	52	72	44	23	71	33	60	17	87
	Spring Juvenile						344	549	79	5	7	12	197
Chum	Fall Adult	30	111	68	8	19	42	24	32	24	80	5	44
	Spring Juvenile	4	244	328	189	119	0	798	231	98	8	4	202
Sockeye	Fall Adult	4	11	28	7	0	39	10	11	5	15	1	13
	Spring Juvenile	12	69	36	67	121	42	268	141	259	138	118	115
Cutthroat	Spring Adult	308	278	224	43	55	2	117	30	13	45	135	112
	Fall Adult	17	27	14	28	18	12	14	28	18	12	12	19
Dolly Varden	Spring Juvenile	682	309	339	518	711	337	732	647	1095	740	483	611
	Spring Adult	644	728	1,529	28	44	7	25	5	11	10	89	303
	Fall Adult	9	21	10	81	39	21	30	174	27	136	3	55
Steelhead	Spring Juvenile	11	15	36	12	30	4	82	33	41	39	23	30
	Spring Adult	33	0	5	0	0	0	0	0	0	1	0	4
	Fall Adult	5	0	2	0	0	0	4	1	4	4	4	2
Lamprey ^b	Spring Juvenile	749	906	1,277	2,314	1,333	1,794	2,264	1,806	539	1,027	711	1,401
	Spring Adult	-	-	-	-	28	97	144	199	177	295	193	157
	Fall Adult	-	-	-	2	16	4	1	0	0	0	0	3

^a Trapping effort not equal between years.

^b Adults and juveniles not distinguished for period 1992 - 1995.

Table 4. Coded-wire tag retention rates for Zolzap Creek coho smolts, 2002.

Sampling Date	Tagging Date	Tag code	Hours held	Sample size	No. fish no tag	Percent retention
14-May	11-May	28-01-10	72	200	18	91.0
16-May	14-May	28-01-10	48	200	1	99.5
17-May	16-May	28-01-10	24	200	0	100
18-May	17-May	28-01-10	24	200	0	100
19-May	18-May	28-01-10	24	200	3	98.5
20-May	19-May	28-01-10	24	200	1	99.5
21-May	20-May	28-01-10	24	172	0	100
24-May	21-May	28-01-10	72	200	0	100
26-May	24-May	28-01-10	48	200	2	99
06-Jun	04-Jun	28-01-10	48	200	0	100
Subtotal				1,972	25	98.7
09-Jun	08-Jun	28-01-11	48	200	5	97.5
11-Jun	09-Jun	28-01-11	48	200	1	99.5
14-Jun	11-Jun	28-01-11	72	200	0	100
Subtotal				600	6	99.0
Grand Total				2,572	31	98.8

Table 5. Coded-wire tagged coho smolt releases from Zolzap Creek, 2002.

Tag code	Tagging dates	No. AFC	Tag morts	No. released AFC	No. tagged ^a	No. AFC only ^b	No. released untagged ^c	Total release ^d	CWT mark rate ^e
280110	6 May - 7 June	11,551	61	10,174	10,097	77	379	10,553	1.05
280111	8 June - 21 June	2,257	19	2,238	2,221	17	89	2,327	1.05
Total		13,808	80	12,412	12,318	94	468	12,880	1.05

^a No. tagged (corrected for tag loss) = No. released AFC - (No. released AFC * No. lost tags / No. sampled); see Table 4.

^b No. AFC only = No. released AFC - No. tagged

^c No. released untagged = the number of unmarked fish released which belong to the same group as the tagged and AFC only fish.

^d Total release = No. tagged + AFC only + untagged

^e CWT mark rate = Total release / No. tagged

Table 6. Fence enumerations, carcass recoveries, and Petersen population estimates for adult coho escapement at Zolzap Creek, 2002.

Item	Adults	Total
Number live coho captured at fence	1,918 ^a	1,918
Number of live coho released untagged	6	6
Number live coho operculum tagged	1,818 ^b	1,818
Number coho carcasses recovered	653	653
Number of coho carcasses recovered untagged	269	269
Number of coho carcasses recovered tagged	367	367
Petersen estimate	3,233	3,233
Upper 95% CL	3,580	3,580
Lower 95% CL	2,919	2,919

^a Includes 1,879 adult coho examined at fence and 39 adult coho angled and tagged below fence and subsequently released above fence.

^b Adjusted for tag loss.

Table 7. Freshwater age distribution of adult coho at Zolzap Creek, 2002.

Sex	Age-2		Age-3		Total aged	Total unaged	Total sampled
	No.	%	No.	%			
Adult males	59	71.1	24	28.9	83	16	99
Adult females	78	78.0	22	22.0	100	17	117
Total adults	137	74.9	46	25.1	183	33	216

Table 8. Estimates of total escapement of adipose clipped coho and contribution to escapement at Zolzap Creek, 1993-2002.

Escapement Year	No. examined (A)	No. with adipose clips (B)	% AFC (C=B/Ax100)	Population estimate (D)	% sampled (E=A/Dx100)	Estimated adipose clips (F=B/AxD)	No. smolts ^a		Contribution to escap. ^b	Smolt to spawner (%) ^c
							AFC	% AFC		
1993	784	191	24.4	1,048	74.8	255	33,923	83.6	306	0.8
1994	2,416	499	20.7	2,536	95.3	524	22,986	87.3	600	2.3
1995	906	308	34.0	908	99.8	309	29,615	86.0	359	1.0
1996	1,030	218	21.2	1,039	99.1	220	10,166	82.2	268	2.2
1997	462	201	43.5	470	98.3	204	20,625	94.2	206	0.9
1998	963	212	22.0	967	99.6	213	13,566	93.2	228	1.6
1999	1,294	451	34.9	1,393	92.9	486	13,950	88.7	547	3.5
2000	409	260	63.6	456	89.7	290	14,591	98.4	295	2.0
2001	1,893	1,155	61.0	1,897	99.8	1,157	30,304	90.4	1,280	3.8
2002	1,905	883	46.4	3,233	58.9	1,499	22,385	83.5	1,795	6.7
Avg.	1,206	438	37	1,395	91	516	21,211	89	588	2.5

^a smolt releases of the previous migration year; an unknown number of additional unclipped releases were likely.

^b marked contribution to escapement = estimated adipose clips * (clipped + unclipped) / clipped.

^c % survival = estimated AFC + AFC below the fence / AFC smolts* 100.

Table 9. Estimated Canadian and American commercial and sport harvest of Zolzap Creek CWT coho in 2002 using tag recovery data (Mark Recovery Program, Fisheries and Oceans, Canada and ADF&G mark tag and age lab, online searchable database).

Tag code	Observed CWT catch ^a				Catch-sample ratio ^b			Estimated CWT catch ^c			
	N. Troll	N. Net	Sport	Total	N. Troll	N. Net	Sport	N. Troll	N. Net	Sport	Total
<u>Canadian</u>											
28-01-06	2	0	0	2	6.3	0.0	0.0	13	0	0	13
28-01-07	3	2	0	5	14.3	1.0	0.0	43	2	0	45
Total	5	2	0	7	11.1	1.0	0.0	55	2	0	57
<u>American</u>											
28-01-06	35	11	1	47	2.8	4.9	4.1	98	54	4	156
28-01-07	46	6	3	55	2.6	4.8	3.0	118	29	9	156
Total	81	17	4	102	2.7	4.9	3.3	216	83	13	312
<u>Total</u>	86	19	4	109	3.2	4.5	3.3	271	85	13	369
Total commercial											356
Total sport											13
Total native fishery ^d											4
Total escapement ^e											1,159
Total CWT											1,533

^a Observed CWT = CWT's recovered from the commercial and sport catch

^b Cumulative catch-sample ratio = total coho catch / total coho sampled

^c Estimated CWT = observed CWT catch * catch sampling ratio

^d observed harvest

^e Estimated CWT's (adipose clips corrected for tag loss at return) including those below the fence, and at the fishwheels; see Table 8

Table 10. Expanded Canadian and American commercial and sport harvest of Zolzap Creek coho and estimated total return in 2002 from 2001 smolt year.

Tag code	Total release	Smolts tagged ^a	Mark rate ^b	Expanded catch ^c						Contri- bution		Total return ^d
				Canadian			American			Grand		
				Troll	Net	Sport	Total	Troll	Net	Sport	Total	
28-01-06	14,578	13,759	1.06	13	0	0	13	104	57	4	165	179
28-01-07	12,234	8,457	1.45	62	3	0	65	170	42	13	225	290
Total 1 ^e	26,812	22,216	1.21	75	3	0	78	274	99	18	391	469
												1,795
												2,264
Total 2 ^f			2.16	120	4	0	124	466	178	29	673	797
												3,233
												4,030

^a Number smolts released with tags (corrected for tag loss), Nass and Frith 1997.

^b Mark rate at release (= No. released / No. marked) for smolts and Total 1 (MRP method), and mark rate at return for total 2 (Escapement method).

^c Expanded catch = EST * mark rate at release

^d Total return = expanded catch + escapement

^e Total 1 expanded catch is calculated using the total mark rate at release and the total estimated catch for all tag codes (Table 9).

^f Total 2 expanded catch is calculated using the total adipose clip rate at recovery and the total estimated catch for all tag codes (Table 9).

Table 11. Estimated commercial harvest distribution of Zolzap Creek CWT coho by area and gear type, 2002. Percentage is of total commercial harvest (does not include sport recoveries).

Area ^a	Net	%	Troll	%	Total	%
<u>Canada</u>						
Areas 1-5	2	0.6	55	15.6	57	16.1
subtotal	2	0.6	55	15.6	57	16.1
<u>U.S.A. (Alaska)</u>						
Northern Outside	0	0.0	0	0.0	0	0.0
Central Outside	0	0.0	105	29.6	105	29.6
Southern Outside	0	0.0	21	5.9	21	5.9
Southern Inside	54	15.0	42	11.8	95	26.8
Central Inside	15	4.2	0	0.0	15	4.2
Southern Intermediate	14	3.9	37	10.4	51	14.3
Central Intermediate	0	0.0	0	0.0	0	0.0
Unknown	0	0.0	10	2.9	10	2.9
subtotal	83	23.2	216	60.6	298	83.9
TOTAL	85	23.8	271	76.2	356	100.0

^a includes respective sub-areas

Table 12. Adult and juvenile coho abundance and smolt-adult survival, by smolt year, at Zolzap Creek, 1992-2002.

Smolt Year	Smolt Out-migration				Resulting Escapement				Total Return				Catch				Smolt-total return Surv. (%)			
	A		B		Return		Count		Estimate		Year		Return		Year		C		D	
	Count	Estimate	CWT		Year		Count		Estimate		Year		Year		Year		Expanded		Estimated CWT	D/B
1992	40,601	53,000	33,150		1993		794		1,048		1993		1993		1993		2,832		690	2.1
1993	26,334	51,000	22,649		1994		2,438		2,536		1994		1994		1994		9,645		2,025	8.9
1994	34,419	41,000	29,319		1995		908		908		1995		1995		1995		3,057		1,069	3.6
1995	12,369	13,000	10,156		1996		1,039		1,039		1996		1996		1996		3,159		674	6.6
1996	20,745	23,000	20,519		1997		470		470		1997		1997		1997		1,072		486	2.4
1997	15,099	18,000	13,566		1998		967		967		1998		1998		1998		1,986		400	2.9
1998	15,937	19,000	13,900		1999		1,302		1,393		1999		1999		1999		2,808		980	7.1
1999	15,153	16,000	14,572		2000		409		456		2000		2000		2000		955		623	4.3
2000	33,934	34,500	30,132		2001		1,897		1,897		2001		2001		2001		3,765		2,315	7.7
2001	27,948	28,000	22,216		2002		1,918		3,233		2002		2002		2002		4,030		1,874	8.4
2002	15,001	15,000	12,318		2003		-		-		2003		2003		2003		-		-	-
Average	23,413	28,318	20,227				1,214		1,395								3,331		1,114	5.4

Estimate A = Best estimate of total smolt outmigration based on fence counts and migration patterns.

CWT B = Number of smolts that were Coded-wire tagged during their outmigration.

Estimate C = Expanded catch using total adipose clip rate at recovery and the total estimated catch for all tag codes.

CWT D = Estimated catch and escapement of Zolzap Creek CWT coho by tag code, including commercial, sport, native, fishwheel recoveries.

Table 13. Adult and juvenile coho production by freshwater age class and brood year, Zolzap Creek, 1990-1999^a.

All fish

Brood Year	Smolt Production (by freshwater age)			Adult Return (by freshwater age)			Smolts (freshwater age)			Escapement (freshwater age) ^c			Smolt to Adult Return Survival (% by freshwater age)		
	Age 2	Age 3	Age 4	Total	Age 2	Age 3	Age 4	Total	% Age 2	% Age 3	% Age 4	% Age 2	% Age 3	% Age 4	Overall
1990	28,779	16,371	287	45,437	1,651	3,819	0	5,470	63.3	36.0	0.6	30.2	69.8	5.7	23.3
1991	34,629	28,495	910	64,034	5,826	1,794	0	7,620	54.1	44.5	1.4	76.5	23.5	16.8	6.3
1992	12,218	4,927	161	17,306	1,263	1,478	0	2,741	70.6	28.5	0.9	46.1	53.9	10.3	30.0
1993	7,163	6,233	0	13,396	1,681	313	0	1,994	53.5	46.5	0.0	84.3	15.7	23.5	5.0
1994	16,606	6,282	228	23,116	759	1,348	0	2,107	71.8	27.2	1.0	36.0	64.0	4.6	21.5
1995	11,718	7,695	256	19,669	638	868	4	1,509	59.6	39.1	1.3	42.2	57.5	5.4	11.3
1996	11,077	6,624	0	17,701	1,940	481	0	2,422	62.6	37.4	0.0	80.1	19.9	17.5	7.3
1997	9,088	1,553	0	10,641	465	331	0	796	85.4	14.6	0.0	58.4	41.6	5.1	21.3
1998	32,948	8,344	0	41,292	3,434	1,011	-	4,446	79.8	20.2	0.0	77.2	22.8	10.4	12.1
1999 ^a	19,656	4,560	-	24,216	3,018	-	-	3,018	81.2	18.8	-	-	-	-	-
Avg. ^b	18,247	9,614	205	28,066	1,962	1,272	1	3,082	66.7	32.7	0.6	56.7	43.2	11.1	15.8

CWT's

Brood Year	Smolt Production (by freshwater age)			Adult return (by freshwater age)			Smolts (freshwater age)			Escapement (freshwater age) ^c			Smolt to Adult Return Survival (% by freshwater age)		
	Age 2	Age 3	Age 4	Total	Age 2	Age 3	Age 4	Total	% Age 2	% Age 3	% Age 4	% Age 2	% Age 3	% Age 4	Overall
1990	18,000	7,270	205	25,476	402	802	0	1,204	70.7	28.5	0.8	33.4	66.6	2.2	11.0
1991	15,379	20,377	711	36,466	1,223	628	0	1,851	42.2	55.9	1.9	66.1	33.9	8.0	3.1
1992	8,737	3,849	144	12,730	441	315	0	757	68.6	30.2	1.1	58.3	41.7	5.1	8.2
1993	5,596	5,561	0	11,157	359	142	0	500	50.2	49.8	0.0	71.6	28.4	6.4	2.6
1994	14,815	4,735	167	19,716	344	272	0	616	75.1	24.0	0.8	55.9	44.1	2.3	5.7
1995	8,831	5,630	233	14,694	128	303	6	437	60.1	38.3	1.6	29.4	69.3	1.3	5.4
1996	8,104	6,033	0	14,137	677	314	0	991	57.3	42.7	0.0	68.3	31.7	8.4	5.2
1997	8,277	1,356	0	9,633	303	204	0	507	85.9	14.1	0.0	59.8	40.2	3.7	15.0
1998	28,776	6,620	0	35,396	2,112	470	-	2,582	81.3	18.7	0.0	81.8	18.2	7.3	7.1
1999 ^a	15,596	3,745	-	19,340	1,403	-	-	1,403	80.6	19.4	-	-	-	-	-
Avg. ^b	13,211	6,517	162	19,934	739	383	1	858	65.7	33.6	0.7	55.4	44.5	4.7	7.0

Brood Year	All fish			CWT's		
	smolts/spawner	recruits/spawner	smolts/spawner	recruits/spawner	smolts/spawner	recruits/spawner
1990						
1991						
1992	11.1	1.8	8.2	0.5		
1993	12.8	1.9	10.6	0.5		
1994	9.1	0.8	7.8	0.2		
1995	21.7	1.7	16.2	0.5		
1996	17.0	2.3	13.6	1.0		
1997	22.6	1.7	20.5	1.1		
1998	42.7	4.6	36.6	2.7		
Avg.	19.6	2.1	16.2	0.9		

^a (-) Incomplete data for 1999, to be completed with data from subsequent returns.
^b average for "Total" includes years for which complete production data is available.
^c Age composition of adult escapement return for brood year.

FIGURES




Figure 1. The Nass River watershed, British Columbia.




Figure 2. Zolzap Creek and location of enumeration fence.





Figure 3. Water level and temperature at Zolzap Creek, 2002.




Figure 4. Daily migration of coho smolts at Zolzap Creek, 29 April - 26 June, 2002.




Figure 5. Length-frequency and calculated age distribution of Zolzap Creek coho smolts, 2002.




Figure 6. Daily counts of adult coho at the Zolzap Creek enumeration fence, 31 Aug - 5 Oct, 2002.




Figure 7. Length-frequency distribution of coho, by sex, Zolzap Creek, 2002.




Figure 8. Fisheries Statistical Areas for the north coast of British Columbia and southeast Alaska, and commercial harvest distribution of Zolzap Creek CWT coho, 2002.




Figure 9. Exploitation rates for three wild coho indicator stocks.




Figure 10. Total percent survivals for three wild coho indicator stocks.




Figure 11. Canadian and Alaskan exploitation rates on Zolzap Creek coho, 1994-2002.

APPENDICES

Table A-1. Juvenile coho catch at Zolzap Creek enumeration fence, 2002.

Date	fry/presmolts	smolts	morts
30-Apr	27	0	5
01-May	118	0	7
02-May	165	0	0
03-May	118	0	0
04-May	1	0	2
05-May	16	0	0
06-May	0	30	0
07-May	0	38	0
08-May	0	72	0
09-May	0	75	0
10-May	1	155	0
11-May	0	123	0
12-May	0	67	0
13-May	0	210	0
14-May	1	94	0
15-May	0	100	0
16-May	0	545	0
17-May	0	541	0
18-May	2	690	0
19-May	2	806	0
20-May	1	746	0
21-May	1	1,833	0
22-May	2	1,281	0
23-May	0	28	0
24-May	36	187	1
25-May	90	549	0
26-May	76	780	0
27-May	29	94	0
28-May	15	665	0
29-May	6	485	0
30-May	0	0	0
31-May	0	0	0
01-Jun	66	44	1
02-Jun	20	177	0
03-Jun	94	268	0
04-Jun	149	387	0
05-Jun	43	707	0
06-Jun	22	876	0
07-Jun	213	112	0
08-Jun	34	568	0
09-Jun	24	736	0
10-Jun	796	427	0
11-Jun	12	167	0
12-Jun	296	118	5
13-Jun	415	106	3
14-Jun	0	90	0
15-Jun	0	0	0
16-Jun	0	0	0

Table A-1. Juvenile coho catch at Zolzap Creek enumeration fence, 2002.

Date	fry/presmolts	smolts	morts
17-Jun	0	0	0
18-Jun	0	0	0
19-Jun	0	0	0
20-Jun	0	4	1
21-Jun	0	20	0
Total	2,891	15,001	25

Table B-1. Non-coho catch at Zolzap Creek enumeration fence, 2002.

Date	Steelhead		Cutthroat		D. Varden		Sockeye Juvenile	Chum Juvenile	Cottid	Lamprey	Stickleback
	Juvenile	Adult	Juvenile	Adult	Juvenile	Adult					
30-Apr	0	0	0	56	0	19	0	0	8	45	1
01-May	0	0	0	19	0	20	0	0	6	102	0
02-May	6	0	0	16	4	8	0	0	4	97	1
03-May	0	0	0	14	6	18	0	0	14	132	0
04-May	1	0	2	0	27	0	0	1	4	36	0
05-May	0	0	0	0	13	0	0	0	4	15	0
06-May	1	0	0	1	19	18	0	0	1	23	0
07-May	1	0	3	0	23	1	0	0	0	13	1
08-May	2	0	3	3	23	0	0	1	0	19	1
09-May	0	0	1	1	12	1	0	1	0	27	3
10-May	2	0	1	6	10	1	0	1	0	35	1
11-May	0	0	4	4	14	0	0	2	4	85	1
12-May	0	0	0	2	9	0	0	1	0	41	1
13-May	0	0	0	4	12	2	0	3	3	27	0
14-May	1	0	11	4	14	0	0	0	0	26	0
15-May	0	0	1	0	5	0	0	0	0	6	0
16-May	0	0	4	1	18	0	0	0	1	30	0
17-May	1	0	3	1	28	1	0	0	0	26	0
18-May	1	0	6	0	43	0	0	0	0	31	2
19-May	1	0	14	0	51	0	0	1	5	9	0
20-May	2	0	31	1	49	0	0	0	3	2	1
21-May	1	0	23	1	51	0	0	0	0	8	1
22-May	0	0	0	0	2	0	0	0	0	3	1
23-May	0	0	0	0	0	0	0	0	0	0	0
24-May	0	0	0	0	3	0	0	0	0	0	1
25-May	0	0	1	1	10	0	0	0	1	14	1
26-May	0	0	1	0	6	0	0	0	1	3	0
27-May	0	0	0	0	1	0	0	0	0	0	0
28-May	0	0	4	0	1	0	0	0	1	0	0
29-May	0	0	0	0	1	0	0	1	1	0	0
30-May	0	0	0	0	0	0	0	0	0	0	0
31-May	0	0	0	0	0	0	0	0	0	0	0

Table B-1. Non-coho catch at Zolzap Creek enumeration fence, 2002.

Date	Steelhead		Cutthroat		D. Varden		Sockeye Juvenile	Chum Juvenile	Cottid	Lamprey	Stickleback
	Juvenile	Adult	Juvenile	Adult	Juvenile	Adult					
01-Jun	0	0	0	0	0	0	0	0	1	1	0
02-Jun	0	0	1	0	2	0	0	0	1	1	0
03-Jun	0	0	0	0	4	0	0	0	0	0	1
04-Jun	1	0	0	0	6	0	0	0	1	9	0
05-Jun	0	0	0	0	3	0	0	0	1	0	0
06-Jun	0	0	0	0	6	0	0	0	0	5	1
07-Jun	0	0	0	0	1	0	0	0	0	2	0
08-Jun	1	0	2	0	3	0	0	0	0	1	0
09-Jun	1	0	1	0	2	0	0	0	1	2	0
10-Jun	0	0	0	0	0	0	0	0	0	0	0
11-Jun	0	0	0	0	0	0	0	0	0	4	0
12-Jun	0	0	0	0	0	0	0	0	0	2	0
13-Jun	0	0	0	0	0	0	0	0	0	1	0
14-Jun	0	0	0	0	0	0	0	0	0	2	0
15-Jun	0	0	0	0	0	0	0	0	0	0	0
16-Jun	0	0	0	0	0	0	0	0	0	0	0
17-Jun	0	0	0	0	0	0	0	0	0	0	0
18-Jun	0	0	0	0	0	0	0	0	0	0	0
19-Jun	0	0	0	0	0	0	0	0	0	0	0
20-Jun	0	0	0	0	0	0	0	0	0	9	0
21-Jun	0	0	0	0	0	0	4	0	1	11	0
22-Jun	0	0	1	0	0	0	0	0	2	5	0
23-Jun	0	0	0	0	1	0	0	0	4	10	0
24-Jun	0	0	0	0	0	0	0	0	0	4	0
25-Jun	0	0	0	0	0	0	0	0	0	0	0
26-Jun	0	0	0	0	0	0	0	0	0	3	0
Total	23	0	118	135	483	89	4	12	73	927	19

Table C-1. Coded-wire tagging data for coho smolts at Zolzap Creek, 2002.

Date	Total smolts	Fence morts	Tag code	No. AFC	Tag morts	No. rlsd. untagged	No. rlsd. AFC
06-May	30	2	-	0	0	28	0
07-May	38	1	-	0	0	37	0
08-May	72	0	-	0	0	72	0
09-May	75	0	-	0	0	75	0
10-May	155	0	-	0	0	0	0
11-May	123	0	28-01-10	268	5	10	263
12-May	67	2	-	0	0	0	0
13-May	210	1	-	0	0	2	0
14-May	94	0	28-01-10	357	2	9	355
15-May	100	0	-	0	0	1	0
16-May	545	7	28-01-10	621	5	16	616
17-May	541	3	28-01-10	534	3	4	531
18-May	690	0	28-01-10	680	4	10	676
19-May	806	0	28-01-10	796	4	9	792
20-May	746	1	28-01-10	722	5	21	717
21-May	1,833	2	28-01-10	1,819	8	12	1,811
22-May	1,281	157	-	0	0	0	0
23-May	28	0	-	0	0	0	0
24-May	187	0	28-01-10	1,320	4	19	0
25-May	549	2	-	0	0	28	0
26-May	780	4	28-01-10	765	5	9	760
27-May	94	3	-	0	0	0	0
28-May	665	0	-	0	0	1	0
29-May	485	0	28-01-10	1,239	4	0	1,235
30-May	0	0	-	0	0	0	0
31-May	0	0	-	0	0	0	0
01-Jun	44	1	-	0	0	1	0
02-Jun	177	10	-	0	0	1	0
03-Jun	268	0	-	0	0	0	0
04-Jun	387	0	28-01-10	853	6	9	847
05-Jun	707	0	-	0	0	0	0
06-Jun	876	1	28-01-10	1,577	6	5	1,571
07-Jun	112	0	-	0	0	0	0
08-Jun	568	0	28-01-11	672	3	8	669
09-Jun	736	0	28-01-11	715	10	21	705
10-Jun	427	0	-	0	0	0	0
11-Jun	167	0	28-01-11	572	3	21	569
12-Jun	118	0	-	0	0	0	0
13-Jun	106	1	-	0	0	1	0
14-Jun	90	0	28-01-11	298	3	14	295
15-Jun	0	0	-	0	0	0	0
16-Jun	0	0	-	0	0	0	0
17-Jun	0	0	-	0	0	0	0
18-Jun	0	0	-	0	0	0	0
19-Jun	0	0	-	0	0	0	0
20-Jun	4	0	-	0	0	4	0
21-Jun	20	0	-	0	0	20	0
Total	15,001	198	-	13,808	80	468	12,412

Table D-1. Daily counts of adult coho at Zolzap Creek enumeration fence, 2002.

Date	No. examined ^a	No. operculum tagged ^a
31-Aug	1	1
01-Sep	0	0
02-Sep	8	8
03-Sep	8	8
04-Sep	0	0
05-Sep	6	6
06-Sep	3	3
07-Sep	14	14
08-Sep	0	0
09-Sep	0	0
10-Sep	0	0
11-Sep	0	0
12-Sep	0	0
13-Sep	0	0
14-Sep	0	0
15-Sep	0	0
16-Sep	0	0
17-Sep	0	0
18-Sep	0	0
19-Sep	399	392
20-Sep	160	158
21-Sep	14	14
22-Sep	319	317
23-Sep	192	192
24-Sep	106	106
25-Sep	73	73
26-Sep	83	82
27-Sep	29	29
28-Sep	150	150
29-Sep	160	160
30-Sep	42	42
01-Oct	53	53
02-Oct	63	63
03-Oct	25	25
04-Oct	5	5
05-Oct	5	5
Totals	1,918	1,906

^a Includes 1,879 adult coho examined at fence and 39 adult coho angled and tagged below fence and subsequently released above fence.