



Health
Canada Santé
Canada

*Your health and
safety... our priority.*

*Votre santé et votre
sécurité... notre priorité.*

Proposed Re-evaluation Decision

PRVD2015-03

Propamocarb

(publié aussi en français)

11 May 2015

This document is published by the Health Canada Pest Management Regulatory Agency. For further information, please contact:

Publications
Pest Management Regulatory Agency
Health Canada
2720 Riverside Drive
A.L. 6607 D
Ottawa, Ontario K1A 0K9

Internet: pmra.publications@hc-sc.gc.ca
healthcanada.gc.ca/pmra
Facsimile: 613-736-3758
Information Service:
1-800-267-6315 or 613-736-3799
pmra.infoserv@hc-sc.gc.ca

Canada 

ISSN: 1925-0959 (print)
1925-0967 (online)

Catalogue number: H113-27/2015-3E (print)
H113-27/2015-3E-PDF (PDF version)

© Her Majesty the Queen in Right of Canada, represented by the Minister of Health Canada, 2015

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.

Table of Contents

Overview	1
What Is the Proposed Re-evaluation Decision?.....	1
What Does Health Canada Consider When Making a Re-evaluation Decision?	1
What Is Propamocarb?.....	2
Health Considerations.....	2
Environmental Considerations.....	3
Measures to Minimize Risk	3
Next Steps	3
Science Evaluation.....	5
1.0 Introduction.....	5
2.0 Use Description of Propamocarb	5
3.0 The Technical Grade Active Ingredient, Its Properties.....	5
3.1 Identity of the Technical Grade Active Ingredient.....	5
3.2 Physical and Chemical Properties of the Technical Grade Active Ingredient	6
4.0 Human Health	6
4.1 Toxicology Summary	7
4.2 <i>Pest Control Products Act</i> Hazard Considerations	7
4.3 Dermal Absorption.....	8
4.4 Occupational Exposure	8
4.4.1 Mixer/Loader/Applicator Exposure and Risk.....	8
4.4.2 Post-application Exposure and Risk	9
4.5 Non-occupational Exposure	10
4.5.1 Dietary Exposure and Risk	10
4.5.2 Residential Exposure and Risk	11
4.6 Aggregate Exposure and Risk	11
4.7 Cumulative Exposure and Risk	12
5.0 Environment.....	12
5.1 Environmental Fate	12
5.2 Environmental Exposure and Risk Assessment.....	13
5.2.1 Terrestrial Species	13
5.2.2 Aquatic Species	13
5.2.3 Conclusion	13
6.0 Value	14
7.0 Pest Control Product Policy Considerations	14
7.1 Toxic Substances Management Policy Considerations.....	14
7.2 Contaminants and Formulants of Health or Environmental Concern.....	15
8.0 Incident Reports	15
9.0 Organization for Economic Co-operation and Development Status of Propamocarb	15
10.0 Proposed Re-evaluation Decision	16
11.0 Supporting Documentation	16

List of Abbreviations	17
Appendix I Registered Products Containing Propamocarb as of 22 December 2014	19
Appendix II Human Health Toxicity Endpoints for Propamocarb	21
Appendix III Label Amendments for Products Containing Propamocarb	23
References	27

Overview

What Is the Proposed Re-evaluation Decision?

After a re-evaluation of the fungicide propamocarb, Health Canada's Pest Management Regulatory Agency (PMRA), under the authority of the *Pest Control Products Act* and Regulations, is proposing continued registration of products containing propamocarb for sale and use in Canada.

An evaluation of available scientific information found that products containing propamocarb do not present unacceptable risks to human health or the environment when used according to revised label directions.

This proposal affects all end-use products containing propamocarb registered in Canada. Once the final re-evaluation decision is made, the registrant will be instructed on how to address any new requirements.

This Proposed Re-evaluation Decision is a consultation document¹ that summarizes the science evaluation for propamocarb and presents the reasons for the proposed re-evaluation decision.

The information is presented in two parts. The Overview describes the regulatory process and key points of the evaluation, while the Science Evaluation provides detailed technical information on the assessment of propamocarb.

The PMRA will accept written comments on this proposal up to 45 days from the date of publication of this document. Please forward all comments to Publications (please see contact information indicated on the cover page of this document).

What Does Health Canada Consider When Making a Re-evaluation Decision?

The PMRA's pesticide re-evaluation program considers potential risks, as well as value, of pesticide products to ensure they meet modern standards established to protect human health and the environment. Regulatory Directive DIR2012-02, *Re-evaluation Program Cyclical Re-evaluation*, presents the details of the cyclical re-evaluation approach, which is in line with the requirements of the *Pest Control Products Act*.

¹ "Consultation statement" as required by subsection 28(2) of the *Pest Control Products Act*.

What Is Propamocarb?

Propamocarb is a protectant fungicide with systemic activity. It is registered for control of *Pythium* blight and damping off caused by *Pythium spp.* on turfgrass in golf courses and sod farms; downy mildew on cucurbits and lettuce; late blight on potato and tomato; *Pythium* root rot and damping off on greenhouse tomato, cucumber, peppers and eggplant; and *Pythium* and *Phytophthora* root rot on outdoor and greenhouse ornamentals. Propamocarb is formulated as suspension or solution, and is applied using groundboom, aerial, drench, drip irrigation, hand held and/or backpack application equipment.

Health Considerations

Can Approved Uses of Propamocarb Affect Human Health?

Propamocarb is unlikely to affect your health when used according to the label directions.

People could be exposed to propamocarb by consuming food and water, working as a mixer/loader/applicator, by entering treated sites, or by handling treated ornamentals. The PMRA considers two key factors when assessing health risks: the levels at which no health effects occur and the levels to which people may be exposed. The dose levels used to assess risks are established to protect the most sensitive human population (for example, children and nursing mothers). Only uses for which exposure is well below levels that cause no effects in animal testing are considered acceptable for continued registration.

Occupational mixer/loader/applicator exposure is not of concern for workers handling propamocarb according to the current label directions. Further, exposure to workers re-entering treated sites for various activities is not of concern on the day of application. Residential exposure to golfers playing on treated turf and homeowners handling commercially treated ornamentals was not of concern. Dietary exposure to propamocarb through consumption of domestic and imported food commodities and drinking water is not of concern. Currently registered labels include the required mitigation measures, no additional mitigation measures are proposed pertaining to human health exposure to propamocarb.

Maximum Residue Limits

The *Food and Drugs Act* prohibits the sale of adulterated food, in other words, food containing a pesticide residue that exceeds the established maximum residue limit (MRL). Pesticide MRLs are established for *Food and Drugs Act* purposes through the evaluation of scientific data under the *Pest Control Products Act*. Each MRL value defines the maximum concentration in parts per million (ppm) of a pesticide allowed in or on certain foods. Food containing a pesticide residue that is at or below the established MRL does not pose an unacceptable health risk.

MRLs for propamocarb have been established for registered and imported commodities.

Environmental Considerations

What Happens When Propamocarb Is Introduced Into the Environment?

Propamocarb is unlikely to affect non-target organisms when used according to the revised label directions.

Non-target terrestrial and aquatic organisms could be exposed to propamocarb in the environment. Based on the risk assessment conducted by the PMRA, the currently registered uses of propamocarb are not of concern to earthworms and pollinators. However, there is a potential risk identified for birds, small wild mammals, terrestrial plants and amphibians from the use on turf. To minimize the potential risks to these organisms, the registered end-use product label for turf use includes environmental hazard statements and buffer zones.

Measures to Minimize Risk

No additional mitigation measures are proposed pertaining to human health or environmental exposure to propamocarb. However, labels are proposed to be amended as per the current labelling standard.

What Additional Scientific Information Is Required?

No additional data are required.

Next Steps

Before making a final re-evaluation decision on propamocarb, the PMRA will consider all comments received from the public in response to this consultation document. A science-based approach will be applied in making a final decision on propamocarb. The PMRA will then publish a Re-evaluation Decision² that will include the decision, the reasons for it, a summary of comments received on the proposed decision and the PMRA's response to these comments.

² "Decision statement" as required by subsection 28(5) of the *Pest Control Products Act*.

Science Evaluation

1.0 Introduction

Propamocarb is a protectant fungicide with systemic activity.

Following the re-evaluation announcement for propamocarb, the registrant of the technical grade active ingredient in Canada indicated that they intended to provide continued support for all uses included on the label of end-use products in Canada.

Currently registered products containing propamocarb are listed in Appendix I.


2.0 Use Description of Propamocarb

Propamocarb is registered for control of *Pythium* blight and damping off caused by *Pythium spp.* on turfgrass in golf courses and sod farms; downy mildew on cucurbits and lettuce; late blight on potato and tomato; *Pythium* root rot and damping off on greenhouse tomato, cucumber, peppers and eggplant; and *Pythium* and *Phytophthora* root rot on greenhouse and outdoor container ornamentals. Propamocarb can be applied by groundboom, aerial, drench, drip irrigation, handheld and/or backpack application equipment.

3.0 The Technical Grade Active Ingredient, Its Properties

3.1 Identity of the Technical Grade Active Ingredient

Common Name	Propamocarb Hydrochloride
Function	Fungicide
Chemical Family	Carbamate Fungicide
Chemical Name	
1 International Union of Pure and Applied Chemistry (IUPAC)	propyl 3-(dimethylamino)propylcarbamate hydrochloride
2 Chemical Abstracts Service (CAS)	propyl N-[3-(dimethylamino)propyl]carbamate hydrochloride (1:1)
CAS Registry Number	25606-41-1
Molecular Formula	C ₉ H ₂₁ ClN ₂ O ₂

Structural Formula

Molecular Weight 224.73

Purity of the Technical Grade Active Ingredient 72.1%

Registration Number 24543

Based on the manufacturing process used, impurities of human health or environmental concern as identified in the *Canada Gazette*, Part II, Vol. 142, No. 13, SI/2008-67 (2008-06-25), including Toxic Substances Management Policy (TSMP) Track 1 substances, are not expected to be present in the product.

3.2 Physical and Chemical Properties of the Technical Grade Active Ingredient

Property	Result	Interpretation
Vapour pressure at 20°C	3.85×10^{-2} mPa	Non-volatile
Ultraviolet (UV) / visible spectrum	No significant absorption at $\lambda > 300$ nm	—
Solubility in water at 20°C	> 500 g/L (pH 1.6-9.6)	Very soluble in water
n-Octanol/water partition coefficient at 22°C	logP = -1.21 (pH 7) log K_{ow} = -2.6 log K_{ow} = 0.841	Bioaccumulation is unlikely
Dissociation constant (pKa)	9.3 at 20°C	—

4.0 Human Health

Toxicology studies in laboratory animals describe potential health effects resulting from various levels of exposure to a chemical and identify dose levels at which no effects are observed. Unless there is evidence to the contrary, it is assumed that effects observed in animals are relevant to humans and that humans are more sensitive to effects of a chemical than the most sensitive animal species.

Exposure to propamocarb may occur through diet, working as a mixer/loader/applicator, by entering treated sites or by handling treated crops.

4.1 Toxicology Summary

Propamocarb has low acute toxicity by the oral, inhalation and dermal routes. It is minimally irritating to the eyes and skin. Propamocarb is considered to be a potential skin sensitizer in humans. Label statements for skin sensitization potential are proposed for consistency (existing on several labels). No neurotoxic concerns were identified in an acute neurotoxicity study.

In the subchronic and chronic studies, effects on body weight (body-weight gain at the mid and high doses, often correlated with a decrease in food consumption and food efficiency) were observed. The liver, stomach, eyes and brain were identified as target organs.

Propamocarb was not determined to be carcinogenic in rats and mice. No evidence of mutagenic potential was observed in any of the reviewed genotoxicity studies. Propamocarb did not cause adverse reproductive effects nor did it cause malformations.

4.2 *Pest Control Products Act* Hazard Considerations

For assessing risks from potential residues in food or from products used in or around residential areas or schools, the *Pest Control Products Act* requires the application of an additional 10-fold factor to threshold effects. This factor should take into account potential pre- and post-natal toxicity and completeness of the data with respect to the exposure of, and toxicity to, infants and children. A different factor may be determined to be appropriate on the basis of reliable scientific data.

The toxicity database for propamocarb was considered complete and was considered adequate for the assessment of risk for infants and children. Data available include a one-generation reproduction study, multi-generation reproduction studies in rats, and developmental toxicity studies in rats and rabbits.

With respect to potential pre- and post-natal toxicity, there was no indication of increased susceptibility in the offspring compared to parental animals. Offspring toxicity (decreased pup body weights and food consumption) occurred at doses causing parental toxicity (decreased body weights and body-weight gains) in the reproductive studies. No adverse effects were observed on reproduction, and there was no evidence of sensitivity of the young. Signs of fetotoxicity and increased post-implantation loss were observed in the developmental rabbit study at the same doses that caused decreased body-weight gain or body weight in the dams. It is not clear if the increase in abortions and resorptions was maternally or fetally mediated in the rabbits; however, the effect is considered serious (late death in the form of abortion, with an increased incidence of early resorptions at the higher dose). Early resorptions, decreases in mean fetal weight and minor skeletal anomalies also occurred in a rat developmental study, but at the same dose that caused significant maternal toxicity.

In consideration of the information presented above, the *Pest Control Products Act* factor was reduced from 10-fold to 3-fold when using the rabbit developmental toxicity studies for risk assessment for dermal and inhalation exposure for short- to intermediate-term scenarios. The *Pest Control Products Act* factor is reduced to one-fold for other scenarios.

Appendix II provides an overview of toxicology endpoints for propamocarb used in human health risk assessments by the PMRA.

4.3 Dermal Absorption

Based on a chemical-specific in vivo dermal absorption study, a dermal absorption factor of 26.4% was determined for risk assessment purposes for propamocarb.

4.4 Occupational Exposure

Occupational risk is estimated by comparing potential exposures with the most relevant endpoint from toxicology studies being used to calculate a margin of exposure (MOE). This is compared to a target MOE incorporating uncertainty factors protective of the most sensitive subpopulation. If the calculated MOE is less than the target MOE, it does not necessarily mean that exposure will result in adverse effects, but mitigation measures to reduce risk would be required.

Workers can be exposed to propamocarb through mixing, loading and by application using various types of application equipment or when entering a treated site to conduct activities such as scouting and/or handling of treated crops.

The currently registered product labels require that workers must wear long pants, long-sleeved shirts and boots during all activities. In addition, during mixing, loading, cleanup and repair activities, workers must also wear chemical-resistant gloves and safety goggles or a face shield.

4.4.1 Mixer/Loader/Applicator Exposure and Risk

Mixer/loader/applicator exposure is expected to be mainly via dermal and inhalation routes.

Based on propamocarb use pattern, the following scenarios were assessed:

- Short- and intermediate-term exposure from open mixing/loading and groundboom application (open cab) to potato, field tomato, cucurbits and lettuce.
- Short- and intermediate-term exposure from open mixing/loading and aerial application to potato.
- Short- and intermediate-term exposure from open mixing/loading and groundboom (open cab), backpack or turf-gun application to turf (sod farm and golf courses).
- Short- and intermediate-term exposure from open mixing/loading and irrigation, manually pressurized handwand or backpack foliar application to greenhouse cucumber.
- Long-term exposure from open mixing/loading with drench/drip irrigation, backpack sprayer or manually pressurized handwand soil application to greenhouse tomato, pepper, cucumber and eggplant.

- Long-term exposure from open mixing/loading with drench/drip irrigation or manually pressurized handwand soil application to ornamentals (greenhouse and outdoor).

The mixer/loader/applicator exposure assessment was based on workers wearing the personal protection equipment specified on the current registered labels, and applied at the maximum registered application rates for each crop. The potential dermal and inhalation exposures were estimated using data from the Pesticide Handlers Exposure Database (PHED), Version 1.1.

Short- and intermediate-term combined dermal and inhalation exposure is not of concern for the assessed scenarios. For long-term scenarios, the combined dermal and inhalation exposure is also not of concern for the assessed scenarios. Therefore, no further mitigation measures are proposed.

4.4.2 Post-application Exposure and Risk

For workers entering treated fields to conduct post-application activities, dermal exposure is considered to be the primary route of exposure. Propamocarb is relatively non-volatile with a low vapour pressure of 2.9×10^{-7} mmHg at 25°C, meeting the North American Free Trade Agreement (NAFTA) criterion for an inhalation waiver based on low volatility due to a vapour pressure of less than 7.5×10^{-4} mmHg (NAFTA, 1999). Therefore, inhalation exposure is considered to be negligible for outdoor post-application activities compared with dermal exposure.

Post-application occupational risk assessments consider dermal exposure to workers entering treated sites to conduct agronomic activities involving foliar or soil contact. Based on the propamocarb use pattern, there is potential for short- (< 30 days) to long-term (6+ months) exposure.

For foliar and turf contact, potential exposure of post-application workers was estimated using activity-specific transfer coefficients and dislodgeable foliar residue (DFR) or transferable turf residue (TTR) values. The DFR/TTR refers to the amount of residue that can be dislodged or transferred from a surface, such as leaves of a plant. The TC is a measure of the relationship between exposure and DFRs for individuals engaged in a specific activity, and is calculated from data generated in field exposure studies. The transfer coefficients are specific to a given crop and activity combination and reflect standard agricultural work clothing worn by adult workers. Post-application exposure activities include mechanical weeding, mowing, thinning, pruning, scouting, irrigating and harvesting.

For contact with treated soil (soil drench/irrigation), post-application exposure was estimated based on the potential concentration of propamocarb in soil and the expected adherence and absorbance to skin from handling treated soil.

Based on the maximum registered application rates and application intervals, short- and intermediate-term post-application exposure (field vegetables, turf, outdoor ornamentals) is not of concern on the day of application for all assessed uses and handler activities. Long-term post-application exposure (greenhouses) is also not of concern on the day of application for all assessed uses and handler activities.

All required mitigation measures (such as restricted entry intervals) are already on the registered labels. No additional risk mitigation measures are proposed for post-application exposure. Revised wording is required on certain labels for re-entry during the restricted-entry interval period to meet the current label standard.

4.5 Non-occupational Exposure

The non-occupational (residential) risk assessment estimates the risk to the general population, including children/youths, during or after pesticide application.

4.5.1 Dietary Exposure and Risk

In a dietary exposure assessment, the PMRA determines how much of a pesticide residue, including residues in milk and meat, may be ingested with the daily diet (food and drinking water). Exposure to propamocarb from potentially treated imports is also included in the assessment. These dietary assessments are age-specific and incorporate the different eating habits of the population at various stages of life. For example, the assessments take into account differences in children's eating patterns, such as food preferences and the greater consumption of food relative to their body weight when compared to adults.

Plant and animal metabolism of propamocarb is adequately understood. Analytical methods for determination of residues in plants and animals were considered adequate. Storage stability data are available to support the sample storage conditions. Supervised residue trials are available for representative crops to support the registered uses.

The residue definition of propamocarb in plants is the parent, propyl [3-(dimethylamino)propyl]carbamate hydrochloride; in animals, the residue definition is the parent, and the metabolites AE F155306, AE F132675, AE F132679 and propamocarb glucuronide for enforcement and risk assessment purpose.

An acute dietary assessment was not required as an acute reference dose (ARfD) has not been identified for propamocarb.

The chronic dietary risk was calculated by estimating the consumption of different foods and the residue values on those foods. This expected intake of residues was then compared to the acceptable daily intake (ADI). When the expected intake of residues is less than the ADI, then chronic dietary exposure is not of concern. The chronic dietary assessment was refined at the intermediate level using median values from residue field trial data and experimental processing factors where available. Chronic dietary exposure from all supported (import and domestic) propamocarb food uses for the representative population subgroups ranged from

7.1 to 9.7 % of the ADI. Levels in drinking water were estimated based on the field use of propamocarb on turf at a maximum rate of 13.86 kg a.i./ha/year (expected environmental concentration of 58 µg a.i./L, surface water, level I). The estimated drinking water value is conservative to levels of propamocarb detected in the available water monitoring information. Chronic dietary exposure to propamocarb from food and drinking water ranged from 7.9 to 11.2% of the ADI for all subpopulations. The lowest exposed population subgroup was children less than one year old; the highest exposed population subgroup was children 3-5 years old. On this basis, chronic dietary exposure to propamocarb from food and drinking water is not of concern.

A cancer assessment was not required as propamocarb is not carcinogenic.

4.5.2 Residential Exposure and Risk

There are no domestic class products containing propamocarb registered in Canada, therefore, residential mixer/loader/applicator risk assessment was not required.

4.5.2.1 Exposure and Risk for Golfers

There is potential for short- to intermediate-term exposure to golfers exposed to freshly treated turf. Exposure to golfers was estimated using TTR values and transfer coefficients representative of typical golfing activities for adult, youth and child golfers.

The estimated exposure for golfers contacting treated turf was not of concern on the day of application.

4.5.2.2 Exposure and Risk for Handlers of Ornamentals

Propamocarb is registered for greenhouse and outdoor container grown ornamentals. Residential short- and intermediate-term exposure through handling treated ornamentals and soil is possible. Post-application exposure was estimated based on the potential concentration of propamocarb in treated soil and the expected adherence and absorbance to skin.

The estimated exposure from handling treated ornamentals was not of concern.

4.5.2.3 Exposure and Risk for Bystanders

Potential for bystander exposure is considered minimal and is expected to be significantly less than exposures estimated for workers. Based on the worker risk assessment, bystander exposure is not expected to be of concern.

4.6 Aggregate Exposure and Risk

Aggregate exposure is the total exposure to a single pesticide that may occur from food, drinking water, residential and other non-occupational sources as well as from all known or plausible exposure routes (oral, dermal and inhalation).

Aggregate exposure from food and water is considered acceptable and below the level of concern for all population subgroups. The lowest exposed population subgroup was children less than one year old; the highest exposed population subgroup was children 3-5 years old (< 12% of ADI). No acute or cancer aggregate food and water assessment is required for propamocarb.

Based on the exposure estimates for golfers (MOE > 3000, target MOE of 300) and from handling treated ornamentals (MOE > 40,000, target MOE of 300), and considering exposure from food and drinking water (< 12% of the ADI), it is not expected that aggregate exposure (from food, drinking water and residential exposure) would result in risks of concern.

4.7 Cumulative Exposure and Risk

No common mechanism of toxicity has been found with other pest control products. Therefore, no cumulative risk assessment was conducted during the re-evaluation.

5.0 Environment

5.1 Environmental Fate

Propamocarb is stable to hydrolysis and undergoes limited phototransformation (half-life of 35.4 days on soil). Laboratory biotransformation studies have shown that propamocarb is generally non-persistent in aerobic soils, although some studies have shown that this compound may be slightly persistent in some types of soil (overall, the aerobic half-life ranged between 10 and 44 days; corresponding DT₉₀ values ranged from 14 to 63 days). Under field conditions, propamocarb was found to be non-persistent in soil (half-life of 9 days; DT₉₀ of 23 days). Propamocarb will, however, persist in anaerobic soils (half-life of 459 days; corresponding DT₉₀ of > 4 years). No major transformation products are formed in soil.

The mobility of propamocarb was found to be variable. Results from adsorption/desorption studies indicated that propamocarb could have a high mobility in some soils but could be immobile in others. Results from soil column leaching studies were also variable. In some studies, low concentrations of propamocarb were found in leachates (between ≤ 0.07% and < 2%), suggesting a very low potential for leaching. However, in other studies, up to 48% of propamocarb was found in leachates. Drinking water modelling results indicate that propamocarb could reach groundwater. In field studies on turf, however, there was minimal movement of propamocarb to the lower layers of the soil profile. There is limited water monitoring data available. One water monitoring study in Canada that analyzed 53 samples over 3 years indicated no detection of propamocarb in groundwater.

Propamocarb may enter surface water through spray drift and/or runoff. Abiotic processes such as hydrolysis and phototransformation will not affect the degradation of propamocarb in aquatic systems (this compound is stable to hydrolysis and no phototransformation occurred in a laboratory study over a period of 22 days). Propamocarb is slightly persistent in aerobic aquatic systems (in tests performed in aerobic water/sediment systems, the whole system half-life was 23 and 35 days; corresponding DT₉₀ values were 75 and 64 days) but is more persistent under anaerobic aquatic conditions (under these conditions, the biotransformation is biphasic with a

whole system half-life of 50 days for the first phase and 94 days for the second; corresponding DT₉₀ was 260 days). No major transformation products are formed in water. Updated label statements to limit runoff and contamination of aquatic sites are required for all propamocarb products.

Based on the log K_{ow} for propamocarb (-2.6 and 0.841, depending on the study), this compound is unlikely to bioaccumulate. This is consistent with results from bioaccumulation studies, which showed a rapid elimination from fish tissue (depuration half-life of 3 days; DT₉₀ of 14 days).

Based on the vapour pressure and Henry's law constant, propamocarb has a low potential for volatilization.

5.2 Environmental Exposure and Risk Assessment

Based on the propamocarb use-pattern, application to turf was considered to be the worst-case scenario for environmental exposure.

5.2.1 Terrestrial Species

Propamocarb is not acutely toxic to earthworms, honeybees (oral and contact) and other non-target arthropods (spider, hover fly, rove beetle and carabid beetle). Propamocarb has no detrimental effects on seed germination and vegetative vigour in non-target plants; however, negative effects were observed on seedling emergence. Potential risk to terrestrial organisms from uses on field crops, greenhouse vegetables, and greenhouse and outdoor ornamentals were determined to be not of concern. Use on turf could pose a risk to birds and small wild mammals (on an acute dietary and reproductive basis), and non-target plants. Current registered end-use labels for turf use includes environmental hazard statements and buffer zones to minimize the potential risk to non-target organisms.

5.2.2 Aquatic Species

Propamocarb is slightly toxic to aquatic invertebrates on an acute basis. Chronic effects (survival and reproduction) were observed in *Daphnia magna*. Propamocarb is practically non-toxic on an acute basis to freshwater and marine fish. No treatment-related effects were observed in a chronic toxicity test with rainbow trout; however, negative effects on weight and length of fish were noted in an early life-stage toxicity test with fathead minnow. Toxicity to amphibians was extrapolated from the most sensitive fish species. The growth rate, biomass and yield of green algae were adversely affected by propamocarb.

5.2.3 Conclusion

Overall, risks to aquatic species from the use of propamocarb on turf are minimal. However, a potential chronic risk to amphibians from direct overspray was identified. Current labels for turf uses include a statement regarding the potential risk to amphibians (aquatic species) and the requirement for an aquatic buffer zone of one metre around sensitive habitats.

Current labels of registered products have the required risk mitigation measures to protect terrestrial and aquatic organisms and no additional risk mitigation measures are proposed.

Revised environmental hazard label statements to limit runoff and contamination of aquatic sites are proposed for all product labels.

6.0 Value

Propamocarb is a protectant fungicide with systemic activity. It belongs to the carbamate group of fungicides. The proposed biochemical mode of action of propamocarb is disruption of the formation of fungal cell membrane by interfering with synthesis of phospholipids and fatty acids. It reduces mycelial growth and development of sporangia and spores. It is absorbed by roots and leaves, and transported acropetally. Propamocarb is important for the control of water mold fungi (Oomycetes), which may cause diseases such as *Pythium* root rot and downy mildews on greenhouse vegetables. It is also important for the control of *Phytophthora infestans*, the pathogen of late blight on potato and tomatoes. Propamocarb can be rotated with other fungicides from different mode of action groups that control the same pathogen in an Integrated Pest Management program. This will not only prolong the effective life of propamocarb, but also enhance the effective life of other alternative single-site mode of action fungicides.

7.0 Pest Control Product Policy Considerations

7.1 Toxic Substances Management Policy Considerations

The Toxic Substances Management Policy (TSMP) is a federal government policy developed to provide direction on the management of substances of concern that are released into the environment. The TSMP calls for the virtual elimination of Track 1 substances (those that meet all four criteria outlined in the policy: persistent [in air, soil, water and/or sediment], and bioaccumulative, primarily a result of human activity and toxic as defined by the *Canadian Environmental Protection Act*).

Propamocarb was assessed in accordance with the PMRA Regulatory Directive DIR99-03, *The Pest Management Regulatory Agency's Strategy for Implementing the Toxic Substances Management Policy*. It was concluded that propamocarb does not meet TSMP Track-1 criteria.

7.2 Contaminants and Formulants of Health or Environmental Concern

During the re-evaluation of propamocarb, contaminants in the technical grade active ingredient are compared against the *List of Pest Control Product Formulants and Contaminants of Health or Environmental Concern* maintained in the *Canada Gazette*.³ The list is used as described in the PMRA Notice of Intent NOI2005-01 and is based on existing policies and regulations including DIR99-03 and DIR2006-02, and taking into consideration the Ozone-depleting Substance Regulations, 1998, of the *Canadian Environmental Protection Act* (substances designated under the Montreal Protocol). The PMRA has reached the following conclusion:

- Technical grade propamocarb does not contain any contaminants of health or environmental concern identified in the *Canada Gazette*.

8.0 Incident Reports

Starting 26 April 2007, registrants are required by law to report incidents, including adverse effects to health and the environment, to the PMRA within a set time frame.

As of 25 July 2014, the PMRA has received one environmental incident report for this active ingredient. It was classified as minor in severity. The effects observed were considered unrelated to propamocarb exposure.

The United States Environmental Protection Agency (USEPA) Ecological Incident Information System (EIS) was queried for environmental propamocarb incidents. There was one incident report available in the EIS database related to damage to sugar beets.

9.0 Organization for Economic Co-operation and Development Status of Propamocarb

Canada is a member of the Organisation for Economic Co-operation and Development (OECD).

As part of the re-evaluation of an active ingredient, the PMRA takes into consideration recent developments and new information on the status of an active ingredient in other jurisdictions, including OECD member countries.

Propamocarb is currently acceptable for use in other OECD countries, including the United States, Australia and European Union Member States. As of 22 December 2014, no decision by an OECD member country to prohibit all uses of propamocarb for health or environmental reasons has been identified.

³ *Canada Gazette*, Part II, Volume 139, Number 24, pages 2641–2643: *List of Pest Control Product Formulants and Contaminants of Health or Environmental Concern* and in the order amending this list in the *Canada Gazette*, Part II, Volume 142, Number 13, pages 1611–1613. *Part 1 Formulants of Health or Environmental Concern, Part 2 Formulants of Health or Environmental Concern that are Allergens Known to Cause Anaphylactic-Type Reactions and Part 3 Contaminants of Health or Environmental Concern*.

10.0 Proposed Re-evaluation Decision

The PMRA has determined that products containing propamocarb for sale and use in Canada are acceptable for continued registration. Appendix III lists label amendments.

11.0 Supporting Documentation

PMRA documents, such as Regulatory Directive DIR2012-02, *Re-evaluation Program Cyclical Re-evaluation*, and DACO tables (datacode tables) can be found on the Pesticides and Pest Management portion of Health Canada's website. PMRA documents are also available through the Pest Management Information Service. Phone: 1-800-267-6315 within Canada or 1-613-736-3799 outside Canada (long distance charges apply); fax: 613-736-3798; e-mail: pmra.infoserv@hc-sc.gc.ca.

The federal TSMP is available through the Environment Canada website.

List of Abbreviations

ADI	acceptable daily intake
a.i.	active ingredient
ARfD	acute reference dose
bw	body weight
CAS	Chemical Abstracts Service
cm	centimetre(s)
DACO	data code
DFR	dislodgeable foliar residue
EIIS	Ecological Incident Information System
EP	end-use product
g	gram(s)
ha	hectare(s)
IPM	integrated pest management
IUPAC	International Union of Pure and Applied Chemistry
K_{ow}	octanol–water partition coefficient
LC ₅₀	lethal concentration to 50% (a concentration causing 50% mortality in the test population)
LD ₅₀	lethal dose to 50% (a dose causing 50% mortality in the test population)
m	metre(s)
mg	milligram(s)
MOE	margin of exposure
MRL	maximum residue limit
NAFTA	North American Free Trade Agreement
NOAEL	no observed adverse effect level
OECD	Organisation for Economic Co-operation and Development
pH	-log ₁₀ hydrogen ion concentration
PHED	Pesticide Handlers Exposure Database
pK _a	-log ₁₀ acid dissociation constant
PMRA	Pest Management Regulatory Agency
PRVD	Proposed Re-evaluation Decision
ppm	parts per million
REI	restricted entry interval
TC	transfer coefficient
TGAI	technical grade active ingredient
TSMP	Toxic Substances Management Policy
TTR	turf transferable residue
USEPA	United States Environmental Protection Agency

Appendix I Registered Products Containing Propamocarb as of 22 December 2014

Registration Number	Marketing Class	Registrant	Product Name	Formulation Type	Guarantee
24543	TGAI	Bayer CropScience Inc.	Propamocarb HCl Technical	Solution	72.1%
24915	Manufacturing Concentrate	Bayer CropScience Inc.	Tattoo Manufacturing Use Product	Suspension	375 g/L ¹
24544	Commercial	Bayer CropScience Inc.	Tattoo Suspension Concentrate Fungicide	Suspension	375 g/L
26288	Commercial	Bayer CropScience Inc.	Previcur N Aqueous Solution Fungicide	Solution	722 g/L
29156	Commercial	Bayer CropScience Inc.	Banol	Solution	722 g/L
29554	Commercial	Bayer CropScience Inc.	Tattoo Fungicide	Solution	722 g/L

¹ Co-formulated with chlorothalonil with a guarantee of 375 g/L.

Appendix II Human Health Toxicity Endpoints for Propamocarb

Exposure Scenario	Dose ¹ (mg/kg bw/day)	Study	Target MOE ²
Chronic dietary	12	Chronic toxicity and oncogenicity study in mice.	
	ADI = 0.12 mg/kg bw/day, UF ³ = 100× (10 interspecies, 10× intraspecies)		
Acute dietary, general population			
	ARfD : No acute reference dose established.		
Short- and intermediate-term dermal ³	140	Developmental toxicity study in rabbits.	300 (PCPA factor = 3-fold)
Short- and intermediate-term inhalation	140	Developmental toxicity study in rabbits.	300 (PCPA factor = 3-fold)
Long-term dermal ⁴	12	Carcinogenicity study in mice.	100
Long-term inhalation	12	Carcinogenicity study in mice.	100
Cancer	Propamocarb is not carcinogenic.		

¹ NOAEL = no observed adverse effect level; ADI = acceptable daily intake; ARfD = acute reference dose.

² MOE = margin of exposure.

³ UF = uncertainty factor.

⁴ A dermal absorption factor of 26.4% was used in a route-to-route extrapolation.

Appendix III Label Amendments for Products Containing Propamocarb

The label amendments presented below do not include all label requirements for individual end-use products, such as first aid statements, storage, disposal statements, precautionary statements and supplementary protective equipment. Information on labels of currently registered products should not be removed unless it contradicts the above label statements.

A submission to request label revisions will be required within 90 days of finalization of the re-evaluation decision.

1. For end-use products, Reg. No. 26288, 29156, and 29554, on primary panel, add the following statement:

“Potential skin sensitizer”

2. For all end-use products, in the section entitled **PRECAUTION STATEMENTS**, add the following statement:

“Potential skin sensitizer”

3. For end-use product, Reg. No. 29156,

- Change the section title “**ENVIRONMENTAL PRECAUTIONS AND INFORMATION**” to “**ENVIRONMENTAL HAZARDS**”.

- Remove all statements related to greenhouse use:

- Sections on **ENVIRONMENTAL HAZARDS**:

- “DO NOT allow effluent or runoff from greenhouses containing this product to enter lakes, streams, ponds or other waters.”

- Section Re-Entry Restrictions For Turf Applications on Commercial Sod Farms, modify the last sentence as follows:

- “This worker protection restriction provides for the protection of agricultural workers and handlers of agricultural pesticides.”

- In the section entitled **DIRECTIONS FOR USE**, modify the **Buffer Zones** as the following:

“Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Buffer Zones

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer and spot treatment.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of Application	Crop	Buffer Zones (Metres) Required for the Protection of	
		Freshwater Habitat	Terrestrial Habitat
Field Sprayer	Turf	1	1

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.”

- The following statements are required for all products registered for outdoor uses under a section entitled **Environmental Hazards**:

“To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.”

“Avoid application when heavy rain is forecast.”

“Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.”

- The following statements are required for all products under a section entitled **Directions for Use**:

“As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.”

“DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.”

6. For end-use product, Reg. No. 26288, replace the section entitled **Re-Entry Restrictions, For Soil Directed Applications to Ornamentals**:

“**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours. This worker protection restriction provides for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted entry interval.

For Drench Applications: This worker protection restriction allows workers to enter the treated areas following application provided there will be no contact with any treated foliage. Personal protective equipment required for early entry to treated areas that is permitted under this worker protection restriction and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material.
- Shoes plus socks”

With

“**DO NOT** enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.”

References

Studies Considered in the Chemistry Assessment

A. LIST OF STUDIES/INFORMATION SUBMITTED BY REGISTRANT

PMRA Document Number	Reference
1774055	PHY-BCJ-2 - Product Chemistry Information, Formation of Impurities, Beginning Materials and Manufacturing Process, Certification of Ingredient Limits, Verify Certified Limits, Analysis of 5 Product Samples, Colour, Physical State, Odour, Viscosity, Density, DACO 2.99
1774142	1979, PHY-BCJ-2 - Qualitative and Quantitative Analysis of [CBI REMOVED] in Previcur N (Aqueous Solution of Propamocarb-HCl, SN 66 752), DACO: 2.13
1774152	1980, PHY-BCJ-2 - Qualitative and Quantitative Analysis of [CBI REMOVED] in Some Batches of Previcur N (70% Aqueous Solution of Propamocarb-HCl, SN 66 752) and the Precursor Compound [CBI REMOVED], DACO: 2.13
1774173	1980, PHY-BCJ-2 - Synthesis of [CBI REMOVED] Derived from Pro-Pamocarb-HCl 1 (66 752) and Analysis of these Compounds in Previcur N (70 % Aqueous Solution of Propamocarb-HCl), DACO: 2.13
1774182	1986, PHY-BCJ-2 - On the Occurrence of [CBI REMOVED] in Previcur N, DACO: 2.13
1774184	1979, PHY-BCJ-2 - Determination of the Distribution Coefficient of Propamocarb Hydrochloride (Sn 66 752) Between Octanol and Water, DACO: 2.14.11
1774186	1990, PHY-BCJ-2 - The Temperature Dependence of the Vapor Pressure of Propamocarb-HCl (ZK 66752), DACO: 2.14.9
1774189	1990, PHY-BCJ-2 - Propamocarb Hydrochloride - Melting Point, DACO: 2.14.4
1774190	1995, PHY-BCJ-2 - Physical and Chemical Properties of Propamocarb Hydrochloride, DACO: 2.14.1, 2.14.10, 2.14.11, 2.14.13, 2.14.14, 2.14.2, 2.14.3, 2.14.4, 2.14.7, 2.14.8, 2.14.9, 2.16, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9
1782027	1999, Special Data (UV / VIS, IR, MS, ¹ H - NMR, ¹³ C - NMR) And Molar Extinction Coefficient Propamocarb-HCl AE B066752 00 1B97 0001, DACO: 2.14.12 CBI

- 1782028 1998, propamocarb hydrochloride solution 722 g/l Code: AE B066752 00 SL67 A2 (CR 18131/01/940203) Previcur N: Storage Stability up to 48 months, DACO: 2.14.14 CBI
- 1658275 2008, Product Chemistry of Banol Fungicide, DACO: 2.11.1, 2.11.2, 2.11.3, 2.11.4 CBI
- 1366602 2006, Material Accountability of AE B066752 (Propamocarb Hydrochloride) Technical Concentrate. Analytical Profile of Five Production Batches from [CBI REMOVED], DACO: 2.11.4, 2.13.1, 2.13.2, 2.13.3 CBI
- 1366594 2005, Propamocarb-HCl Technical. Material Manufacturing Process. Description of Material Used to Produce the Product and Description of Production Process., DACO: 2.11.1, 2.11.2, 2.11.3 CBI
- 1774064 1996, PHY-BCJ-2 - Hoe 102791, Propamocarb-HCl Conc (CQ 684) Quality Control Data of 10 Typical Consecutively Produced Batches, DACO: 2.13.3

Studies Considered in the Value Assessment

A. ADDITIONAL INFORMATION CONSIDERED

Published Information

PMRA Document Number	Reference
----------------------	-----------

- | | |
|---------|--|
| 2245260 | 2004, BCMAFF. Pesticide Info Sheet – Propamocarb HCL. 2004. British Columbia Ministry of Agriculture, Food and Fisheries. Available from http://www.al.gov.bc.ca/pesticides/infosheets/propamocarb_hcl.pdf (accessed January 4, 2012). |
| 2245263 | FRAC. 2011. FRAC Code List: Fungicides sorted by mode of action (including FRAC Code numbering). Fungicide Resistance Action Committee. Available from http://www.frac.info/frac/publication/anhang/FRAC%20Code%20List%202011-final.pdf (accessed January 4, 2012). |

Studies Considered in the Health Risk Assessment

A. LIST OF STUDIES/INFORMATION SUBMITTED BY REGISTRANT

PMRA Document Number	Reference
1161852	1982, T 53 Propamocarb: Previcur N: Acute Oral Toxicity Study in Rats. (226;27.22;T53;150).(Propamocarb HCl), 226, DACO: 4.2.1
1161855	1982, Propamocarb-HCl: Previcur N: Acute Percutaneous Toxicity Study In Rats. (T56;153;229), 229, DACO: 4.2.2
1161856	1990, Propamocarb HCl: Previcur N (CP604; ZK6752): Acute Inhalation Toxicity Study in The Rat (M+F, 4H Exposure) Revised Final Report. (T7/2;TX76.159;3512;PF1/90). TX76.159, DACO: 4.2.3
1161857	1983, Propamocarb Hcl: Primary Eye Irritation after Single Application with Previcur N in the Rabbit. + Amendment 1. (T47;024333;27.23;603.35)., 024333, DACO: 4.2.4
1403994	1983, Propamocarb Hcl: Primary Eye Irritation after Single Application with Previcur N in the Rabbit. + Amendment 1. (T47;024333;27.23;603.35)., 024333, DACO: 4.2.4
1161859	1983, Propamocarb: Primary Skin Irritation Following A Single 4-Hour Occlusive Application With Previcur N In The Rabbit. (T48;018393;27.18).(Propamocarb HCl)., 018393, DACO: 4.2.5
1403995	1983, Propamocarb: Primary Skin Irritation Following A Single 4-Hour Occlusive Application With Previcur N In The Rabbit. (T48;018393;27.18).(PROPAMOCARB HCL)., 018393, DACO: 4.2.5
1161860	1977, Propamocarb Hcl: Investigation Of A Potential Sensitizing Effect Of CP 604 In Guinea Pigs By Means Of The Optimization Test. (T29/2;266/77;SN66.752;PF)., 266/77, DACO: 4.2.6
625683	1999, USEPA, Guinea-pig Skin Sensitization Study (Magnusson & Kligman method) propamocarb hydrochloride liquid concentrate 71.2% w/w. OPPTS Guideline 870.2600., TOX 98176, DACO: 4.2.6
1161863	1976, TNO, Propamocarb Hcl: Sub-Chronic (90-Day) Feeding Study With Zk17296 in Dogs (Revised Version of Final Report). (T32/2;R5178;B90-8181;TNO/R5178)., R 5178, DACO: 4.3.1

- 1161862 1982, T46 Propamocarb Hcl: Previcur N: Three Month Sub-Chronic Oral Toxicity Study in Rats. (T46;217; 119(015-002))., An-PyQ Centerz & Schenng Report 217, MRID: 43619601, DACO: 4.3.1
- 625717 1998, USEPA, Mouse Dietary 90-Day Toxicity Range Finding Study Propamocarb Hydrochloride Liquid Concentrate. OPPTS Guideline 870.3100. Centre International de Toxicologie (CIT). Study Identification Number TOX 94156. December 17, 1998. 372 pp., TOX 941 56,
- 1403996 1998, Mouse Dietary 90-Day Toxicity Range Finding Study Propamocarb Hydrochloride Liquid Concentrate. OPPTS Guideline 870.3100. Centre International de Toxicologie (CIT). Study Identification Number TOX 94156. December 17,1998. 372 pp., TOX 941 56, 12731
- 1403997 Rat Dietary 90-Day Toxicity Range Finding Study Propamocarb Hydrochloride Liquid Concentrate. OPPTS Guideline 870.3100. Centre International de Toxicologie. Laboratory Project ID Number: . December 17, 1998. 407 pp., TOX 94155,12730 TCR, DACO: 4.3.1
- 1161864 1992, Propamocarb Hcl: Previcur N: Rat 21-Day Dermal Repeat Dose Study. TOX/92/186-16, DACO: 4.3.4
- 1161865 1985, Propamocarb: 24 Month Oral (Feeding) Study with Previcur N in Beagle Dogs, PF-81-812, DACO: 4.4.1
- 1161872 1985, Clinical Biochemistry summary: (Cont'D From Roll#1432) Propamocarb: 24 Month Oral (Feeding) Study With Previcur N In Beagle Dogs, Dated: June 6,1985. + Propamocarb: Dog Two Year Dietary Toxicity (T-66,Addendum 1). Relevance for Humans of Ocular Chan
- 1161873 1981, Propamocarb-HCl: Previcur N (Sn66752) Toxicity and Potential Tumorigenicity in Dietary Administration to Rats for 104 Weeks. (T37/2;SHG/165-G/801056)., 16S-G/801056, DACO: 4.4.1,4.4.2
- 1161890 1987, Propamocarb Hcl: Previcur N (Sn66752) Potential Tumorigenicity to Mice In Dietary Administration For 104 Weeks (Final Report). (T39/2;SHG164/80965/2;603.34)., SHG 164/80965/2, DACO: 4.4.2
- 625698 1998, OECD, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, TOX 941 46, DACO: 4.4.4
- 625720 1999, AgrEvo UK, Propamocarb HCl Liquid Concentrate: Vacuolation of Choroid Plexus Ependymal Cells in Rats From a Chronic Toxicity and Oncogenicity Study., C005985, DACO: 4.4.4
- 625722 1998, AgrEvo UK Limited., Rat Combined Chronic Toxicity and Oncogenicity (dietary). OPPTS Guideline 870.4300, TOX 94145., DACO: 4.4.4

657504	1998, Mouse Dietary Oncogenicity (18 months) Study. 98.0776, DACO: 4.4.4
775525	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
775527	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH. 98.0776, DACO: 4.4.4
775529	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH. 98.0776, DACO: 4.4.4
775531	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
775533	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
775535	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
775537	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
775539	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
775541	1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
775593	1998, Rat Combined Chronic Toxicity and Oncogenicity (dietary). OPPTS Guideline 870.4300., TOX 94145, DACO: 4.4.4
775595	1998, Rat Combined Chronic Toxicity and Oncogenicity (dietary). OPPTS Guideline 870.4300., TOX 94145, DACO: 4.4.4
775597	1998, Rat Combined Chronic Toxicity and Oncogenicity (dietary). OPPTS Guideline 870.4300. AgrEvo UK Limited., TOX 94145, DACO: 4.4.4
775599	1998, Rat Combined Chronic Toxicity and Oncogenicity (dietary). OPPTS Guideline 870.4300. AgrEvo UK Limited., TOX 94145, DACO: 4.4.4
775601	Rat Combined Chronic Toxicity and Oncogenicity (dietary). OPPTS Guideline 870.4300. AgrEvo UK Limited. Study Identification Number TOX 94145. December 17,1998. 1700 pp., TOX98/186-29, DACO: 4.4.4

- 775607 Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH. Report Number 98.0776. November 12, 1998. 4311 pp., DACO: 4.4.4
- 775609 Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH. Report Number 98.0776. November 12, 1998. 4311 pp., 98.0776, DACO: 4.4.4
- 775611 1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
- 775613 1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300. Hoechst Marion Roussel Deutschland GmbH., 98.0776, DACO: 4.4.4
- 775615 1998, Mouse Dietary Oncogenicity (18 months) Study. OPPTS Guideline 870.4300., 98.0776, DACO: 4.4.4
- 1161874 1993, Propamocarb-Hcl: Effect Of Previcur N On Reproductive Function Of Multiple Generations In The Rat. (T40/2;126)., T40/2, DACO: 4.5.1
- 625728 1998, Rat Dietary Reproduction Range-Finding Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800. WIL, TOX 95008, DACO: 4.5.1
- 625729 1998, Rat Dietary Two-Generation Reproductive Toxicity Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800., TOX 96131, DACO: 4.5.1
- 658349 1998, Rat Dietary Two-Generation Reproductive Toxicity Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800., TOX 96131, DACO: 4.5.1
- 775543 1998, Rat Dietary Two-Generation Reproductive Toxicity Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800., TOX 96131, DACO: 4.5.1
- 775545 1998, Rat Dietary Two-Generation Reproductive Toxicity Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800., TOX 96131, DACO: 4.5.1
- 775547 1998, Rat Dietary Two-Generation Reproductive Toxicity Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800., TOX 96131, DACO: 4.5.1
- 775603 1998, Rat Dietary Two-Generation Reproductive Toxicity Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800., TOX 96131, DACO: 4.5.1

-
- 775605 1998, Rat Dietary Two-Generation Reproductive Toxicity Study Propamocarb Hydrochloride Liquid Concentrate 780 g/L. OPPTS Guideline 870.3800., TOX 96131, DACO: 4.5.1
- 1243247 AgrEvo, Table 58 - Rat Dietary Reproduction Range-Finding Study of Propamocarb HCl Individual Gross Description of Organs, DACO: 4.5.1
- 1161875 1990, T41/2 Propamocarb-HCl: Previcur N (CP604) Embryotoxicity including Teratogenicity Study in Rats after Daily Intragastrical Administration from Day 6 to Day 19 of Gestation- Revised Final Report. [T41/2;PF24/90(TX80348);3434;TX80.348].DACO: 4.5.2
- 1161876 1990, T36/2 Propamocarb-HCl: Previcur N (Cp604) Embryotoxicity including Teratogenicity Study in Rabbits after Daily Intragastrical Administration from Day 6 to Day 18 of Gestation- Revised Final Report. [T36/2;PF9/90(TX80122);3433;TX80.122]. DACO: 4.5.2
- 1161877 1987, Technical Propamocarb Hydrochloride: Microbial Metabolic Activation Test To Assess Mutagenic Potential. (T70;TOX/87/186-14;25.04;TOX87250;SMS53/87627). (Propamocarb HCl), TOX/87/186-14, DACO: 4.5.4
- 1161878 1987, T69 -Propamocarb-Hcl: Technical Propamocarb Hydrochloride (Previcur N): Metaphase Chromosome Analysis Of Human Lymphocytes Cultured In Vitro. (T69;TOX/87/186-13;27.17;SMS54/87808;TOX87249)., TOX/87/ 186-13, DACO: 4.5.4
- 1161889 1980, T26 Propamocarb: Previcur N: Mutagenicity Evaluation In Yeast. (T26;27.30;4881;4956F).(Propamocarb HCl), DACO: 4.5.4
- 1162707 1981, Propamocarb-Hcl: Mutagenicity Testing In Bacteria With Previcur N. (T44;142;145).(PROPAMOCARB TECHNICAL), 142, DACO: 4.5.4
- 1162718 1985, Propamocarb-Hcl: Mutagenicity Evaluation of Previcur N in the Gene Conversion And Reverse Mutation Assays With Saccharomyces Cerevisiae Strains. D4,S138 AND S211.(T67;E-9409).(PROPAMOCARB TECHNICAL), E - 9409, DACO: 4.5.4
- 625679 2001, Propamocarb Hydrochloride Liquid Concentrate, 780, Code: AE B066752 00 7K72 A101 in Vitro Mammalian Cell Mutation Test With Mouse Lymphoma Cells. OPPTS Guideline 870.5300., TOX 20042, DACO: 4.5.4
- 1403999 2001, Propamocarb Hydrochloride Liquid Concentrate, 780, Code: AE B066752 00 7K72 A101 in Vitro Mammalian Cell Mutation Test With Mouse Lymphoma Cells. OPPTS Guideline 870.5300., TOX 20042, DACO: 4.5.4
- 625716 1980, Micronucleus Test on CP 604 (SN66752, Previcur N). OPPTS Guideline 870.5395., SHG 191/79737, DACO: 4.5.7
-

-
- 1404000 1980, Micronucleus Test on CP 604 (SN66752, Previcur N). OPPTS Guideline 870.5395., SHG 191/79737, DACO: 4.5.7
- 1161891 1993, Propamocarb: Rat Subchronic (3-Month) Dietary Neurotoxicity Study, TOX/93/186-31, DACO: 4.5.12
- 1161892 1993, Propamocarb: Rat Subchronic (3-Month) Dietary Neurotoxicity Study, TOX/93/186-29;92-4503;TOX/92271, DACO: 4.5.12
- 1161893 1993, Propamocarb: Rat Acute Oral Neurotoxicity Study. + Addendum 1. (T87;TOX/93/186-30;92-4504;TOX/92005;TOX/93/186-31). (PROPAMOCARB HCL), TOX/93/186-30, DACO: 4.5.12
- 1091646 Determination of Transferable Turf Residues on Turf Treated with Propamocarb (Banol). DACO 5.14
- 625685 1998, Propamocarb: Ruminant (Cow) - Metabolism, Distribution and Nature of the Residues in Milk And Edible Tissues. OPPTS Guideline 860.1300. Agrevo USA Company, Analytical Development Corporation and Colorado State University.
- 625692 1994, The Fate Of Propamocarb Hydrochloride In Potato Tubers. OPPTS Guideline 860.1300, DACO: 6.3
- 625709 1998, Statement on the Metabolism of Propamocarb Hydrochloride in Potato Tubers. DACO: 6.3
- 1021057 2000, Propamocarb: Ruminant (Cow) - Metabolism, Distribution and Nature Of The Residues In Milk And (Amended Report Replacing Report Av97E521, Document A91204) Edible Tissues, DACO: 6.2
- 1161879 1991, Propamocarb HCl: The Fate of Propamocarb X HCl In Potato Tubers. DACO: 6.3
- 1161880 1994, Propamocarb HCl: Rotational Plant Uptake in Soybean, Sugarbeet and Oat of [1-14C]-Propyl-Propamocarb Hydrochloride. DACO: 6.3
- 1161895 1994, Propamocarb HCl: Absorption,Distribution And Elimination Studies In The Rat Following Single And Repeated Oral Dosing And Single Intravenous Dosing. DACO: 6.4
- 1161896 1992, Propamocarb HCl: Uptake and Metabolism of Propamocarb Hydrochloride in Spinach After Seed Bed Treatment. DACO: 6.3
- 1162697 1981, Propamocarb-HCl: Formation of Artificial Breakdown Products During The Extraction Process
-

-
- 1162699 1994, Propamocarb-HCl: Metabolism Of Propamocarb HCl In Cucumber Grown In Soil And Hydroculture.(M20;Pf-S91032).(Propamocarb Technical), DACO: 6.3
- 1162700 1991, Propamocarb-HCl: The Fate of Propamocarb X HCl in Potato Tubers. 19Th March 1991. + 1St Addendum To Report No.Upsr14/91 Dated: April 1994. DACO: 6.3
- 1162747 1984, Summaries: Propamocarb-HCl: Propamocarb Hydrochloride: Summary Of Metbolism. DACO: 6.1
- 1162748 1994, Propamocarb-HCl Water Soluble Liquid 722.0 G/L Code: CP604 Propamocarb HCl: Clearance of A Single Oral Dose From Rat Tissues. DACO: 6.4
- 1162749 1994, Propamocarb HCl:Dermal Penetration In The Rat. DACO: 6.4
- 1162750 1994, Propamocarb Hydrochloride (Code:Cr18131) Previcur N: Metabolism In The Rat. DACO: 6.4
- 1179888 1998, Propamocarb-HCl Ae B066752,1-[14C]-Propyl-Labelled Water Soluble Liquid 722.0 G/L Code:Cp406 Statement On The Metabolism Of Propamocarb Hydrochloride In Potato Tubers. DACO: 6.3
- 1179889 1998, Propamocarb Hydrochloride Ae B066752,1-[14C]-Propyl-Labelled Metabolism Of Propamocarb HCl In Cucumber Grown In Soil and Hydroculture. DACO: 6.3
- 1181351 1998, Propamocarb: Ruminant (Cow)-Metabolism, Distribution and Nature of the Residues in Milk and Edible Tissues. DACO 6.2
- 1181352 1998, Propamocarb Hydrochloride AE B066752, 1-[14C]-Propyl-Labelled, Metabolism Of Propamocarb Hydrochloride In Spinach After Seedbed Treatment And Hydroponic Uptake In Comparison To Natural Plant Products Generated By Hydroponic Uptake Of 14C-Acetate And 14CO2-ASSIMILATION. DACO 6.3
- 1181353 1998, Metabolism of [14C]-Propamocarb Hydrochloride in Spinach. DACO: 6.3
- 1184557 1998, Metabolism of Propamocarb HCl in Cucumber Grown in Soil and Hydroculture. DACO: 6.3
- 602849 2003, Validation of Method of Analysis for Propamocarb and Its Metabolites in Animal Matrices, DACO: 7.2, 7.2.3, 7.2.5
-

-
- 625684 1996, Multiresidue Method Testing For Propamocarb Hcl According to Pam I, Appendix II, as Updated January 1994. OPPTS Guideline 860.1360., DACO: 7.2.4, 7.2.5
- 625744 1999, Propamocarb Derived Residues in Tomatoes and Processed Tomato Commodities Following Five Applications of Tattoo C at the Maximum Proposed Rate-USA, DACO: 7.4.5
- 625745 1999, Propamocarb Derived Residues in Tomatoes Following Five Applications
- 625755 1991, Analytical Method for the Determination of Residues of Propamocarb X HCl in Potatoes. OPPTS Guideline 860.1340, DACO: 7.2.1, 7.2.2, 7.2.5
- 625758 1990, Stability of Propamocarb X HCl in Lettuce during Deep Freeze Storage. OPPTS Guideline Number 860.1380, DACO: 7.3
- 728428 Propamocarb: Magnitude Of Residues In Dairy Cow Milk And Tissues, DACO: 7.5
- 952686 2005, Response to Deficiencies - Supervised Residue Trial Analytical Methodology, Enforcement Analytical Methodology, DACO: 7.2.1, 7.2.2, 7.2.5
- 952689 2005, Response to Deficiencies - Residues, Enforcement Analytical Methodology, DACO: 7.2.1, 7.2.2, 7.2.5
- 953688 2002, Independent Laboratory Validation Of The Analytical Method: "Propamocarb: Analytical Method For The Determination Of Propamocarb (AE B039744) And Its Metabolites AE F155306, AE F132679 And AE F132675 In Animal Matrices Using LC/MS/MS." DACO 7.2.2
- 953707 2004, Independent Laboratory Validation of Method "02Av32853" For Propamocarb Glucuronide in Liver, DACO: 7.2.2, 7.2.3, 7.2.5
- 1067402 2005, Independent Laboratory Validation of Method "02Av32853" For Propamocarb Glucuronide in Liver. DACO: 7.2.2, 7.2.5
- 1083585 2003, Frozen Storage Stability for Propamocarb Glucuronide in Cow Milk, Kidney, Liver, Fat and Muscle Using Lc/Ms/Ms., DACO: 7.2.2, 7.3
- 1161882 1991, Propamocarb HCl: Analytical Method for the Determination Of Residues Of Propamocarb X HCl In Potatoes. DACO: 7.2.1
- 1161919 1992, Propamocarb-Hcl: Stability of Propamocarb X HCl during Deep Freeze Storage In Potatoes, DACO: 7.3
- 1162806 1990, Stability of Propamocarb X HCl in Tomatoes during Deep Freeze Storage. DACO: 7.3
-

-
- 1162807 1990, Stability Of Propamocarb \times HCl In Lettuce During Deep Freeze Storage (R144;UPSR 52/90-Pa 66752.3/16) (Tattoo C), DACO: 7.3
- 1162812 Note to Reviewer: Part 5 - Food, Feed And Residue Studies (Tattoo C), DACO: 7.1
- 1181356 1998, Rationale For Waiver of Cow Feeding Study and Analytical Method For Animal Commodities. DACO: 7.5
- 1373041 2001, Residue Decline Study Summary, DACO: 7.4.2
- 1379632 1999, Residue Data Summary, DACO: 7.4
- 1531678 1998, Analytical Methodology Xam-47; DACO URMULE D.3.1 2007-1419, Tattoo C Fungicide, DACO: 7.2.1, 7.2.5
- 1891465 1997, Propamocarb Hydrochloride Active Ingredient Analytical Grade; AE B066752; Validation of Analytical Method; Leeks, Onions, Brassicae, Tomatoes, Potatoes and Melons; Gas Chromatography Mass Selective Detection, DACO: 7.2.3
- 1901797 2010, Petition For Exemption from Additional Residue Trials For Tattoo Fungicide On Crop Group 8 (Fruiting Vegetables), Leaf Lettuce And Head Lettuce, DACO: 7.3, 7.4.1, 7.4.2
- 1525141 2001, Analytical Method of Determination of Residues of propamocarb free base in plant material, DACO: 7.2
- 602843 2003, Propamocarb: Analytical Method for the Determination of Propamocarb (AE B039744) and its Metabolites AE F155306, AE F132679, AE F132675 and Propamocarb Glucuronide in Animal Matrices Using LC/MS/MS, DACO: 7.2.1
- 602851 2003, Propamocarb: Magnitude of Residues in Dairy Cow Milk and Tissues, DACO: 7.5
- 728427 Validation of Method of Analysis for Propamocarb and its Metabolites in Animal Matrices, DACO: 7.2.2
- 1181355 1996, Propamocarb-Derived Residues in Potatoes and Processed Potato Commodities Following Application of Tattoo C-USA. DACO: 7.4.5
- 1193611 1992, Stability of Propamocarb X HCl during Deep Freeze Storage in Potatoes. DACO: 7.3
- 1193612 1999, Potato Tubers: Stability during Deep Freeze Storage Up To 26 Months. DACO: 7.3
-

1193613	1996, Addendum To Propamocarb-Derived Residues in Potatoes and Processed Potato Commodities Following Application of Tattoo C – USA. DACO: 6.4.5
1172107	1996, Summary, DACO 7.1
1172108	1997, Residue Decline Study, DACO 7.4.1, 7.4.2, 7.4.3
1172109	1997, Residue Decline Study, DACO 7.4.1, 7.4.2, 7.4.3

B. ADDITIONAL INFORMATION CONSIDERED

Published Information

PMRA Document Number	Reference
1627538	Evaluation Report for Application Number 2008-3131
2074010	Evaluation Report for Application Number 2010-1450
2267996	Evaluation Report for Application Number 2011-5929
2360133	Evaluation Report for Application Number 2013-3299
1193614	1998, Environmental Protection Agency, Federal Register Document, Propamocarb Hydrochloride; Extension of Tolerances For Emergency Exemptions.
1193615	1997, Environmental Protection Agency, Federal Register Document, Propamocarb Hydrochloride; Pesticide Tolerance for Emergency Exemptions.
1193616	1997, Environmental Protection Agency, Federal Register Document, Propamocarb Hydrochloride; Pesticide Tolerance for Emergency Exemptions.
1563673	Klonne et al., 1999. A Generic Evaluation of Homeowner Exposure Associated with Liquid Pesticide Handling and Hose-End Application to Residential Lawns. OMA004. ORETF. Submission No. 2006-4038.
1560575	Merricks, 1997a. Carbaryl Mixer/Loader/Applicator Exposure Study during Application of RP-2 Liquid (21 %), Sevin Ready to Use Insect Spray or Sevin 10 Home Dust to ~Home Vegetable Gardens. ORETF. Submission No. 2006-4038.

- 1945969 Merricks, D.L. (1998). Carbaryl Mixer/Loader/Applicator Exposure Study during Application of RP-2 Liquid (21%) to Fruit Trees and Ornamental Plants: Lab Project Number: 1518. Unpublished study prepared by Agrisearch Inc., Rhone-Poulenc Ag Co., and Morse Laboratories, Inc. 320 p. EPA MRID # 44518501.
- 1563628 Johnson, D., Thompson, R., Butterfield, B. (ORETF) October 7, 1999. Outdoor Residential Pesticide Use and Usage Survey and National Gardening Association Survey. Doane Marketing Research Inc: St. Louis, MO. Gallup Organization Inc: Burlington VT.
- 2115788 Agricultural Reentry Task Force (ARTF). 2008. Data Submitted by the ARTF to Support Revision of Agricultural Transfer Coefficients. Submission No. 2006-0257.
- 2476396 Cowell, J. and Johnson, D. (1999). Evaluation of Transferable Turf Residue
- 2476401 Techniques: Evaluation Study of Transferable Residue Techniques (OMD001) and Transferable Residue Technique Modification Study: An Evaluation of Three Turf Sampling Techniques (OMD002). October 7, 1999. Outdoor Residential Exposure Task Force. EPA MRID 44972203.
- 1619682 Klonne, D.R., Johnson, D.R. (2004) Determination of Potential Dermal Exposure to Adults and Children Reentering a Pesticide-Treated Turf Area.
- Center for Golf Course Management. 1992. Golf Course Operations: Cost of Doing Business/Profitability. Library of Congress GV975.G56.
- USEPA. 2004. Risk Assessment Guidance for Superfund. Volume 1; Human Health Evaluation Manual (Part E, Supplemental Guidance for Dermal Risk Assessment). EPA/540/R/99/005. July 2004.
- USEPA. 1995. Data Call-In Notice for Post-Application Exposure Data. U.S. Environmental Protection Agency, Washington, DC.
- USEPA. 2011. Exposure Factors Handbook 2011 Edition (Final). U.S. Environmental Protection Agency, Washington, DC, EPA/600/R-09/052F.
- USEPA 2012. Standard Operating Procedures (SOPs) for Residential Exposure Assessments.

Studies Considered in the Environmental Risk Assessment

A. LIST OF STUDIES/INFORMATION SUBMITTED BY REGISTRANT

PMRA Document Number	Reference
1162809	1995, Determination of The Acute Toxicity Of Tattoo C SC To <i>Daphnia Magna</i> . DACO: 9.3.1
1162728	1978, Toxicity test with a fungicide, in comparison with phosalone, a standard insecticide, against the honey bee (<i>Apis mellifera</i>). DACO: 9.2.4.1
1162731	1985, Determination of the LC ₅₀ of Previcur N to earthworms (<i>Eisenia foetida</i>) using an artificial soil test. DACO: 9.2.3.1.
1162811	Propamocarb Hydrochloride - The Calculation of Henry'S Law Constant (Tattoo C), DACO 8.2.1
1161903	1978, Propamocarb HCl - Mobility of propamocarb hydrochloride in four soils. DACO: 8.2.3.4.2
1161904	1979, Propamocarb HCl - Degradation of propamocarb hydrochloride in a California loamy sand, 1979. DACO 8.2.3.4.2
1161906	Propamocarb HCl - Degradation of SN 66 752 in a loamy sand under anaerobic conditions. DACO: 8.2.3.4.2
1161902	Propamocarb HCl - Mobility of SN 66 752 in two soils after ageing, 1979. DOCA 8.2.4.4
1161898	1988, Propamocarb HCl - Adsorption to and desorption from soil. DACO: 8.2.4.2
1161903	1978, Propamocarb HCl - Degradation of SN 66 752 in a loamy sand, 1978. DACO: 8.2.3.4.2
1162710	1980, Propamocarb HCl - Degradation of propamocarb hydrochloride in German Standard Soils 2.2 and 2.3 at 15°C. DACO: 8.2.3.4.2
1161905	1980, Propamocarb HCl - Degradation of propamocarb hydrochloride in a loamy sand. DACO: 8.2.3.4.2
1161910	1980, Propamocarb HCl - Degradation of propamocarb hydrochloride in a loamy sand. DACO: 8.2.3.4.2
1161907	1986, Propamocarb HCl - Degradation of propamocarb hydrochloride in a loamy sand after repeated (twofold) application. DACO: 8.2.3.4.2

- 1162709 1995, Previcur N: Column leaching in an acid soil and an alkali soil - A comparative study. DACO: 8.2.4.4
- 1162732 1988, Propamocarb HCl - Assessment of the acute toxicity of Previcur N on the cell multiplication of a pure culture of *Pseudomonas putida* bacteria (acute bacteria multiplication inhibition test). DACO: 9.2.7
- 1161931 1991, Investigation into the phytotoxic effects of propamocarb.HCl on seed germination (Tier 1). DACO: 9.8.4
- 1161932 1991, Investigation into the phytotoxic effects of propamocarb.HCl on seedling emergence (Tier 1). DACO: 9.8.4
- 1161933 1991, Investigation into the phytotoxic effects of propamocarb.HCl on vegetative vigor (Tier 1). DACO: 9.8.4
- 1161899 1988, Propamocarb HCl - The mobility of N-(3-dimethylamino-[1-14C]-propyl)-propamocarb hydrochloride in four soils determined by soil TLC. DACO: 8.2.4.2
- 1162714 Propamocarb hydrochloride: Degradation in sediment/water microcosms, DACO: 8.2.3.5.2.
- 1162734 1995, Propamocarb HCl - A laboratory assessment of the effects of Previcur® N on soil microflora respiration and nitrogen turnover according to BBA Guidelines VI 1-1 (1990). DACO: 9.2.7
- 1162720 1977, The acute toxicity of SN 66752 to the carp, *Cyprinus carpio*. DACO: 9.5.2.1.
- 1162721 1977, The acute toxicity of SN 66752 to the mirror carp (*Cyprinus carpio*) and the rainbow trout (*Salmo gairdneri*). DACO: 9.5.2.1
- 1161922 1990, Propamocarb-Hcl: An Early Life-Stage Toxicity Test with the Fathead Minnow (*Pimephales Promelas*). DACO: 9.5.3.1
- 1162719 1992, Previcur N SL: Mallard duck acute oral toxicity (LD₅₀) study. DACO: 9.6.2.1
- 1161913 1992, Previcur N SL - Bobwhite quail acute oral toxicity (LD₅₀) study. DOCA: 9.6.2.1
- 1161915 1992, Previcur N SL - Mallard duck subacute dietary toxicity (LC₅₀) study. DACO: 9.6.2.4
- 1161916 1992, Previcur N SL - Bobwhite quail subacute dietary toxicity (LC₅₀) study. DACO: 9.6.2.4

- 1161929 1990, Propamocarb-HCl: Wildlife and environment, Effect of Previcur N on the growth of the green alga *Scenedesmus quadricauda* (NEN 6506). DACO: 9.8.2
- 1161928 1991, Propamocarb HCl: A 96-hour shell deposition test with the eastern oyster (*Crassostrea virginica*). DACO: 9.4.1
- 1162735 1981, Propamocarb HCl - The effect of propamocarb hydrochloride on symbiotic dinitrogen fixation, sulphur oxidation, carbon turnover and nitrogen turnover in soils. DACO: 9.2.7
- 1162712 1980, Propamocarb HCl - Metabolism of propamocarb hydrochloride by soil microorganisms. Behaviour in sterilized and non-sterilized German Standard Soil 2.2, 1980. DACO 8.2.3.4.2
- 1161912 1980, Propamocarb HCl - Photolysis experiments with propamocarb HCl in heat sterilized aqueous solutions. DACO 8.2.3.3.2
- 1162730 1990, Propamocarb HCl - Acute toxicity of Previcur N to earthworms, *Eisenia foetida* using an artificial soil test. DACO: 9.2.3.1
- 1162808 1994, Determination of the Acute Toxicity of Tattoo C SC to Earthworms Using an Artificial Soil Test. DACO: 9.2.3.1
- 1774055 1990, Previcur N - Determination of density, Report Number 90/136, Pages 70-76, in PHY-BCJ-2 - Product Chemistry Information, Formation of Impurities, Beginning Materials and Manufacturing Process, Certification of Ingredient Limits, Verify Certified Limits, Analysis of 5 Product Samples, Colour, Physical State, Odour, Viscosity, Density, Melting Point, Identity and Disclosure of Ingredients, Solubility in Organic Solvents, Storage Stability, Solubility in Water, UV-VIS Spectrum, Dissociation Constant. DACO: 2.99
- 1162733 1989, The effect of Previcur N on soil microbial activity. DACO: 9.2.7
- 1162724 1991, An evaluation of the side-effects of Previcur® N on the staphylinid beetle (*Aleochara bilineata*). DACO: 9.2.5
- 1162723 1992, An evaluation of the side-effects of Previcur N® on the carabid beetle (*Poecilus cupreus*). DACO: 9.2.5
- 1164272 1985, The determination of the vapor pressure of SN 66 752; propyl [3-(demethylamino)propylcarbamate] hydrochloride. DACO: 8.2.4.5
- 1161930 1990, Propamocarb HCL - The temperature dependence of the vapor pressure of propamocarb HCL. DOCA: 8.2.4.5
- 625724 1991, Propamocarb HCl - The acid dissociation constant of ZK 66 752. DACO: 2.1.4.10

-
- 1164274 1979, The acute toxicity of SN 66 752 (Previcur N), 67.3 % active ingredient, to the water flea (*Daphnia magna*) Straus. DACO: 9.3.1
- 1164273 1987, Propamocarb-hydrochloride - solubility in water. DACO: 8.2.1
- 1162703 1988, Propamocarb-HCl: Solubility of Propamocarb-Hydrochloride In Water At 20°C. DACO: 8.2.1
- 1161944 1989, Propamocarb HCl: Propamocarb-Base Solubility in Water Of pH7. DACO: 8.2.1
- 1162813 1989, Propamocarb-Base: Octanol/Water Partition Coefficient (C60;89/052)(Tattoo C). DACO: 8.2.1
- 1161926 1990, Influence of Previcur N (propamocarb hydrochloride techn.) on the reproduction of *Daphnia magna*. DACO: 9.3.1.
- 1162716 Dissipation of propamocarb hydrochloride from two German standard soils, 1977. DACO : 8.3.2.3
- 1162725 Auswirkungen von Previcur N auf *Coccinella septempunctata*, Chamber of Agriculture, 1990. DACO 9.2.5
- 1161941 1976, Propamocarb - Determination of rates of hydrolysis of propamocarb base (SN 39744) at pH 5, 7, and 9 at 25°C. DACO: 8.2.3.2
- 1161945 1979, Propamocarb HCl: Determination of the Distribution Coefficient Of Propamocarb Hydrochloride (Between Octanol And Water. DACO: 8.2.1
- 1162726 1990, Previcur N Zulassungspruefung an Nutzarthropoden (*Syrphus corollae*) Im Labor. DACO: 9.2.5
- 1161914 1977, The acute oral toxicity (LD₅₀) of ZK 66 752 to the mallard duck. DACO: 9.6.2.1
- 1163775 1977, The acute oral toxicity (LD₅₀) of ZK 66 752 to the ring-necked pheasant. DACO: 9.6.2.1
- 1164263 1977, The subacute toxicity (LC₅₀) of ZK 66 752 to the mallard duck. DACO: 9.6.2.4
- 1164265 1977, The subacute toxicity (LC₅₀) of ZK 66 752 to the pheasant. DACO: 9.6.2.4
- 1163776 1978, The subacute toxicity (LC₅₀) of ZK 66 752 to the Japanese quail. DACO: 9.6.2.1
- 625724 1989, Propamocarb HCl - Base UV-VIS spectrum. DACO: 2.14.10
-

-
- 1161900 1991, Adsorption / desorption of propamocarb hydrochloride in soil. DACO: 8.2.4.2
- 1161927 1991, The static acute toxicity of propamocarb HCl to the mysid shrimp, *Mysidopsis bahia*. DACO: 9.3.1
- 1161917 1991, The static acute toxicity of propamocarb HCl to the bluegill sunfish, *Lepomis macrochirus*. DACO: 9.5.2.1
- 1161918 1991, The static acute toxicity of propamocarb HCl to the rainbow trout, *Oncorhynchus mykiss*, 1991. DACO 9.5.2.1.
- 1161920 The static acute toxicity of propamocarb HCl to the sheepshead minnow, *Cyprinodon variegatus*. DACO: 9.5.2.1
- 1161925 1992, The acute toxicity of propamocarb HCl to *Daphnia magna* in a static system. DACO: 9.3.1
- 1162727 1977, Bee toxicity of Previcur N. DACO: 9.2.4.1
- 1161943 1990, Propamocarb HCl - The photodegradation of propamocarb hydrochloride on soil surfaces. DACO: 8.2.3.3.1
- 1161912 1990, Propamocarb HCl: The dissipation of propamocarb X HCl in soil following application of BANOL - USA. DACO: 8.3.2.3
- 1161921 1990, Previcur N: 21-day prolonged toxicity study in the rainbow trout under flow-through conditions. DACO: 9.5.3.1

B. ADDITIONAL INFORMATION CONSIDERED

Published Information

Atkins EL; Kellum D; Atkins KW. 1981. Reducing pesticide hazards to honey bees: mortality prediction techniques and integrated management techniques. Univ Calif, Div Agric Sci, Leaflet 2883. 22 pp.

Harris, L.E. 1975. Guide for Estimating Toxic Residues in Animal Feeds or Diets. USEPA, Washington. EPA/540/9-75-019 (NTIS reference #: PB 243 748).

Hoerger F; Kenaga EE. 1972. Pesticide residues on plants: correlation of representative data as basis for estimation of their magnitude in the environment. In: Coulston F; Korte F. (eds). Global aspects of chemistry, toxicology and technology as applied to the environment, Vol. I. Thieme, Stuttgart, and Academic Press, New York. pp. 9-28.

- Kenaga EE. 1973. Factors to be considered in the evaluation of the toxicity of pesticides to birds in their environment. In: Coulston F; Dote F. (eds). Global aspects of chemistry, toxicology and technology as applied to the environment, Vol. II. Thieme, Stuttgart, and Academic Press, New York. pp. 166-181.
- Fletcher, J.S., Nellessen, J.E., and Pfleeger, T.G. 1994. Literature review and evaluation of the EPA food-chain (Kenaga) nomogram, an instrument for estimating pesticide residues on plants. *Environmental Toxicology and Chemistry* 13:1383-1391.
- McCall, P.J., Laskowski, D.A., Swann, R.L. and Dishburger, H.J. 1981. Measurement of sorption coefficients of organic chemicals and their use in environmental fate analysis. In: Test protocols for environmental fate & movement of toxicants. Proceedings of a symposium. Association of Official Analytical Chemists. 94th Annual Meeting, October 21-22, 1980, Washington, DC, pp 89-109.
- Spector, W.S. 1956. Handbook of Biological Data. W.B. Saunders Co., Philadelphia, PA.
- Urban DJ; Cook NJ. 1986. Hazard Evaluation Division, Standard Evaluation Procedure, Ecological Risk Assessment. EPA 540/9-85-001. US EPA, Washington, DC.
- U.S. EPA. (1988) Recommendations for and documentations of biological values for use in risk assessment. PB88 179874, EPA/600/6-87/008. Cincinnati, Ohio.
- Wauchope, R.D. 1978. The pesticide content of surface water draining from agricultural fields - a review. *J. Environ. Qual.* 7(4): 459-472.
- Willis, G.H. and McDowell, L.L. 1987. Pesticide persistence on foliage. *Rev. Environ. Contam. Toxicol.* 100:23-73.
- Wolf, T. and B.C. Caldwell, 2001. Development of a Canadian spray drift model for the determination of buffer zone distances. In Expert Committee on Weeds, Proceedings of the 2001 National Meeting, Quebec City, Sainte Anne de Bellevue, Quebec: ECW-CEM. D. Bernier, DRA Campbell, D. Cloutier, Eds.

Unpublished Information

PMRA Document Number	Reference
1311119	Ministère de l'Environnement Gouvernement du Québec. Contamination de l'eau souterraine par les pesticides et les nitrates dans les régions en culture de pommes de terre; campagne d'échantillonnage de 1999-2000-2001, DACO: 8.6
1311120	Ministère de l'Environnement Gouvernement du Québec. Annexes: contamination de l'eau souterraine par les pesticides et les nitrates dans les régions en culture de pommes de terre; campagne d'échantillonnage de 1999-2000-2001, DACO: 8.6