

April 10 to April 16, 2016 (Week 15)

Overall Summary


- Influenza activity peaked nationally in the second week of March; however, lower but sustained activity continues to be reported throughout the country.
- Many regions across Canada are reporting a greater number of influenza B detections; however, in British Columbia and the Atlantic Region, detections remain predominantly influenza A.
- Hospitalizations, ICU admissions and deaths among the pediatric population, while declining, remain above expected levels based on the past several influenza seasons.
- Despite higher pediatric hospitalizations reported, the pediatric population accounts for the smallest proportion of all deaths reported.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

Are you a primary health care practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel for the 2015-16 influenza season? Contact us at FluWatch@phac-aspc.gc.ca

Influenza/Influenza-like Illness (ILI) Activity (geographic spread)

Influenza activity continues to be reported in the majority of regions in Canada. Localized activity was reported in a total of 14 regions across seven provinces. Sporadic activity levels were reported in 31 regions.

Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, Week 15




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available in the mapping feature found in the [Weekly Influenza Reports](#).

Laboratory Confirmed Influenza Detections


In week 15, the percentage of tests positive for influenza continued to decrease from the previous week [from 23% in week 14 to 21% in week 15] (Figure 2). Compared to the previous five seasons, the percent positive (21%) reported in week 15 was above the five year average for that week and exceeded the expected levels (range 11.7%-16.9%). With the late start to the 2015-16 influenza season, these elevated levels are not unexpected.

Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2015-16



Nationally in week 15, there were 1,522 positive influenza tests reported. The number of influenza A and influenza B detections were roughly equal in week 15. Many of regions across Canada reported more influenza B detections than influenza A; however, in British Columbia and the Atlantic Region, detections remain predominantly influenza A. To date, 78% of influenza detections have been influenza A and among those subtyped, the vast majority have been influenza A(H1N1) [86% (10715/12443)].

Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province, Canada, 2015-16



Note: Specimens from NT, YT, and NU are sent to reference laboratories in other provinces. Cumulative data include updates to previous weeks.

In week 15, individuals under the age of 44 accounted for 74% of influenza B detections. To date this season, detailed information on age and type/subtype has been received for 30,052 cases. Children and teenagers (0-19) accounted for 47% of influenza B cases and approximately one third of all influenza cases. Children and teenagers (0-19), young adults (20-44) and middle-aged adults (45-64) accounted for approximately an equal proportion of influenza A(H1N1) cases (28%).

Table 1 – Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting¹, Canada, 2015-16

Age groups (years)	Weekly (Apr. 10, 2016 to Apr. 16, 2016)					Cumulative (Aug. 30, 2015 to Apr. 16, 2016)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) ³	Total	A Total	A(H1) pdm09	A(H3)	A (UnS) ³	Total	#	%
<5	80	6	<5	x	118	4396	1690	67	2639	1179	5577	19%
5-19	37	<5	0	x	160	2346	1011	97	1238	2043	4391	15%
20-44	74	17	0	57	110	5669	2699	151	2819	1726	7395	25%
45-64	121	28	<5	x	47	6165	2709	181	3275	827	6992	23%
65+	101	17	7	77	91	4608	1558	410	2640	1088	5697	19%
Total	413	72	11	330	526	23184	9667	906	12611	6863	30052	100%
Percentage²	44%	17%	3%	80%	56%	77%	42%	4%	54%	23%		

¹Table 1 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported.

²Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

³UnS: unsubtyped: The specimen was typed as influenza A, but no result for subtyping was available.


x - Suppressed to prevent residual disclosure.

For data on other respiratory virus detections see the [Respiratory Virus Detections in Canada Report](#) on the Public Health Agency of Canada website.

Influenza-like Illness Consultation Rate

The national ILI consultation rate increased from the previous week from 26.1 per 1,000 patient visits in week 14, to 56.4 per 1,000 patient visits in week 15. The highest ILI consultation rate was found in the 5-19 years age group (107.7 per 1,000) and the lowest was found in the ≥65 years age group (14.2 per 1,000) (Figure 4).

Figure 4 – Influenza-like illness (ILI) consultation rates by age group and week, Canada, 2015-16




Delays in the reporting of data may cause data to change retrospectively. In BC, AB, and SK, data are compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.

Pharmacy Surveillance

During week 15, the proportion of prescriptions for antivirals remained similar to the previous week at 50.6 antiviral prescriptions per 100,000 total prescriptions, which is lower than the five year historical average for week 15. The proportion of prescriptions for antivirals remains highest among children. In week 15, the proportion reported among children was 100.1 per 100,000 total prescriptions.

Figure 5 – Proportion of prescription sales for influenza antivirals by age group and week, Canada, 2015-16



Note: Pharmacy sales data are provided to the Public Health Agency of Canada by Rx Canada Inc. and sourced from major retail drug chains representing over 3,000 stores nationwide (excluding Nunavut) in 85% of Health Regions. Data provided include the number of new antiviral prescriptions (for Tamiflu [oseltamivir] and Relenza [zanamivir]) and the total number of new prescriptions dispensed by Province/Territory and age group.


*The average weekly proportion includes data from April 2011 to March 2015.

Influenza Outbreak Surveillance

In week 15, 12 new laboratory confirmed influenza outbreaks were reported: 11 in long-term care facilities (LTCF) and one in a hospital. Of the seven outbreaks with known strains or subtypes, four outbreaks were due to influenza B (three in LTCFs and one in a hospital), one outbreak each was due to influenza A(H1N1) and influenza A(H3N2) (both in LTCFs), and one outbreak in a LTCF was due to influenza A(UnS). Additionally, one ILLI outbreak was reported in a school.

To date this season, 387 outbreaks have been reported. At week 15 in the 2014-15 season, 1,586 outbreaks were reported and in the 2013-14 season, 210 outbreaks were reported.

Figure 6 – Overall number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2015-2016



¹All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of this report.

Sentinel Hospital Influenza Surveillance

Pediatric Influenza Hospitalizations and Deaths

In week 15, 63 laboratory-confirmed influenza-associated pediatric (≤ 16 years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network (Figure 7). The largest proportion of hospitalizations were in children aged 5-9 years, accounting for 33% of the hospitalizations. Similar to the previous week, more influenza B cases were reported than influenza A cases (41 vs. 22 respectively).

To date this season, 1,207 hospitalizations have been reported by the IMPACT network: 874 hospitalized cases (72%) were due to influenza A and 333 cases (28%) were due to influenza B. The greatest proportion of hospitalized cases were in children aged 0-2 years (40%). To date, 192 intensive care unit (ICU) admissions have been reported. Children aged 2 to 4 and 5 to 9 years accounted for 28% and 26% respectively of ICU admissions. A total of 122 ICU cases (64%) reported at least one underlying condition or comorbidity. Eight influenza-associated deaths have been reported.


Table 2 – Cumulative numbers of pediatric hospitalizations (≤ 16 years of age) with influenza reported by the IMPACT network, Canada, 2015-16*

Age Groups	Cumulative (30 Aug. 2015 to 16 Apr. 2016)					
	Influenza A				Influenza B	Influenza A and B (#(%))
	A Total	A(H1) pdm09	A(H3)	A (UnS)	B Total	
0-5m	116	32	5	79	26	142 (12%)
6-23m	270	73	7	190	66	336 (28%)
2-4y	251	79	<5	x	93	344 (29%)
5-9y	179	47	<5	x	106	285 (24%)
10-16y	58	18	<5	x	42	100 (8%)
Total	874	249	21	604	333	1207 (100%)

x - Suppressed to prevent residual disclosure

*Not included in Table 2 are two IMPACT cases that were due to co-infections of influenza A and B.

Figure 7 – Number of hospitalized cases of influenza reported by sentinel hospital networks, by week, Canada, 2015-16, pediatric and adult hospitalizations (≤ 16 years of age, IMPACT; ≥ 16 years of age, CIRN-SOS)*



*Not included in Figure 7 are two IMPACT cases that were due to co-infections of influenza A and B.

Adult Influenza Hospitalizations and Deaths

In week 15, 39 laboratory-confirmed influenza-associated adult (≥16 years of age) hospitalizations were reported by the Canadian Immunization Research Network Serious Outcome Surveillance (CIRN-SOS) (Figure 7). The largest proportion of hospitalizations was in adults 65+ years of age (54%) and due to influenza A (62%). In recent weeks, the weekly number of hospitalizations reported by CIRN has been decreasing and an increasing proportion of hospitalizations have been due to influenza B.


To date this season, 1,077 hospitalizations have been reported by CIRN-SOS (Table 3). The majority of hospitalized cases were due to influenza A (83%) and the largest reported proportion was among adults ≥65 years of age (50%). One hundred and sixty-five intensive care unit (ICU) admissions have been reported. A total of 112 ICU cases reported at least one underlying condition or comorbidity. A total of 45 deaths have been reported this season with the majority of deaths reported in adults ≥65 years of age (64%).

Table 3 – Cumulative numbers of adult hospitalizations (≥16 years of age) with influenza reported by CIRN-SOS, Canada, 2015-16

Age groups (years)	Cumulative (Nov. 1, 2015 to 16 Apr. 2016)					
	Influenza A				B	Influenza A and B
	A Total	A(H1) pdm09	A(H3)	A(UnS)	Total	# (%)
16-20	x	<5	0	<5	<5	7 (1%)
20-44	140	45	<5	x	36	176 (16%)
45-64	312	94	<5	x	39	351 (33%)
65+	434	119	23	292	103	537 (50%)
Unknown	<5	0	0	<5	<5	6 (1%)
Total	895	264	27	604	182	1077
%	83%	29%	3%	67%	17%	100%

x - Suppressed to prevent residual disclosure

Figure 8 – Percentage of hospitalizations, ICU admissions and deaths with influenza reported by age group (≥16 year of age), CIRN-SOS, Canada 2015-16



Note: The number of hospitalizations reported through CIRN-SOS and IMPACT represents a subset of all influenza-associated adult and pediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

x - Suppressed to prevent residual disclosure


Provincial/Territorial Influenza Hospitalizations and Deaths

In week 15, 182 hospitalizations were reported by participating provinces and territories*. The majority of reported hospitalizations continue to be due to influenza A, accounting for 63% of hospitalizations reported in week 15. The largest proportion of cases reported in week 15 was in adults 65+ years of age (43%).

Since the start of the 2015-16 season, 4,606 laboratory-confirmed influenza-associated hospitalizations have been reported. A total of 3,802 hospitalizations (83%) were due to influenza A and 804 (17%) were due to influenza B. Of the 486 ICU admissions reported, 258 (53%) were due to influenza A(H1N1). A total of 193 deaths have been reported; all but 24 were associated with influenza A..

Overall this season, hospitalizations have been reported more frequently among adults ≥65 years of age. The largest proportion of ICU admissions was reported in adults 45-64 years of age and the highest proportion of fatal cases was reported in adults ≥65 years of age. Pediatric (0-19 years) and young to middle-aged adults (20-44 years) accounted for 42% of all hospitalizations and 14% of all deaths reported to date this season. There have been more pediatric hospitalizations reported to date compared to the year-end totals in each of the previous [four influenza seasons](#).

Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza reported by age group, Canada 2015-16



* Note: Influenza-associated hospitalizations are not reported to PHAC by the following Provinces and Territory: BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by SK. ICU admissions are not distinguished among hospital admissions reported from ON. Data may also include cases reported by the IMPACT and CIRN-SOS networks. The number of new influenza-associated hospitalizations and deaths reported for the current week may include cases from ON that occurred in previous weeks, as a result of retrospective updates to the cumulative total. It is important to note that the hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.

x - Suppressed to prevent residual disclosure

See additional data on [Reported Influenza Hospitalizations and Deaths in Canada: 2011-12 to 2015-16](#) on the Public Health Agency of Canada website.

Influenza Strain Characterizations

During the 2015-16 influenza season, the National Microbiology Laboratory (NML) has characterized 1,901 influenza viruses [189 A(H3N2), 1,099 A(H1N1) and 613 influenza B].

Influenza A (H3N2): When tested by hemagglutination inhibition (HI) assays, 48 A(H3N2) viruses were antigenically characterized as A/Switzerland/9715293/2013-like using antiserum raised against cell-propagated A/Switzerland/9715293/2013.

Sequence analysis was done on 141 A(H3N2) viruses. All viruses belonged to a genetic group for which most viruses were antigenically related to A/Switzerland/9715293/2013. A/Switzerland/9715293/2013 is the A(H3N2) component of the 2015-16 Northern Hemisphere's vaccine.

Influenza A (H1N1): All of the 1,099 A(H1N1) viruses characterized were antigenically similar to A/California/7/2009, the A(H1N1) component of the 2015-16 influenza vaccine.

Influenza B: A total of 142 influenza B viruses characterized were antigenically similar to the vaccine strain B/Phuket/3073/2013. A total of 471 influenza B viruses were characterized as B/Brisbane/60/2008-like, one of the influenza B components of the 2015-16 Northern Hemisphere quadrivalent influenza vaccine.

The recommended components for the 2015-2016 Northern Hemisphere trivalent influenza vaccine include: an A/California/7/2009(H1N1)pdm09-like virus, an A/Switzerland/9715293/2013(H3N2)-like virus, and a B/Phuket/3073/2013 -like virus (Yamagata lineage). For quadrivalent vaccines, the addition of a B/Brisbane/60/2008-like virus (Victoria lineage) is recommended.

The NML receives a proportion of the influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Characterization data reflect the results of haemagglutination inhibition testing compared to the reference influenza strains recommended by [WHO](#).

Antiviral Resistance

During the 2015-16 season, the National Microbiology Laboratory (NML) has tested 1,178 influenza viruses for resistance to oseltamivir, 1,179 for resistance to zanamivir and 1,233 influenza viruses for resistance to amantadine. All but eight tested viruses were sensitive to oseltamivir. The eight H1N1 viruses resistant to oseltamivir had a H275Y mutation. All viruses tested for resistance were sensitive to zanamivir. All but one influenza A viruses were resistant to amantadine (Table 4).

Table 4 – Antiviral resistance by influenza virus type and subtype, Canada, 2015-16

Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	147	(0%)	147	0	193	192 (99.5%)
A (H1N1)	728	8 (1.1%)	729	0	1040	1040 (100%)
B	303	(0%)	303	0	NA ¹	NA ¹
TOTAL	1178	8 (0.7%)	1179	0	1233	1232 (99.9%)

¹NA: Not Applicable

International Influenza Reports

[World Health Organization influenza update](#)

[World Health Organization FluNet](#)

[WHO Influenza at the human-animal interface](#)

[Centers for Disease Control and Prevention seasonal influenza report](#)

[European Centre for Disease Prevention and Control - epidemiological data](#)

[South Africa Influenza surveillance report](#)

[New Zealand Public Health Surveillance](#)

[Australia Influenza Report](#)

[Pan-American Health Organization Influenza Situation Report](#)

FluWatch Definitions for the 2015-2016 Season

Abbreviations: Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

Influenza-like-illness (ILI): Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI/Influenza outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

Hospitals and residential institutions: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Workplace: Greater than 10% absenteeism on any day which is most likely due to ILI.

Other settings: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.

Influenza/ILI Activity Levels

1 = No activity: no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

2 = Sporadic: sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

3 = Localized: (1) evidence of increased ILI* ;

(2) lab confirmed influenza detection(s);

(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

4 = Widespread: (1) evidence of increased ILI*;

(2) lab confirmed influenza detection(s);

(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

** More than just sporadic as determined by the provincial/territorial epidemiologist.*

† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.

We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.

This report is available on the Government of Canada Influenza webpage under [Weekly influenza reports](#).

Ce rapport est disponible dans les deux langues officielles.