



12043041

Annotated Checklist and Bibliography of Parasites of Herring (*Clupea harengus* L.)

J.R. Arthur and H.P. Arai

PROTOZOA MONOGENEA DIGENEA CESTODA
NEMATODA ACANTHOCEPHALA ARGULOIDEA
COPEPODA AGNATHA PROTOZOA MONOGENEAE
PROTOZOA MONOGENEA DIGENEA CESTODA
NEMATODA ACANTHOCEPHALA ARGULOIDEA
COPEPODA AGNATHA PROTOZOA MONOGENEAE
PROTOZOA MONOGENEA DIGENEA CESTODA
NEMATODA ACANTHOCEPHALA ARGULOIDEA
COPEPODA AGNATHA PROTOZOA MONOGENEAE
PROTOZOA MONOGENEA DIGENEA CESTODA
NEMATODA ACANTHOCEPHALA ARGULOIDEA
COPEPODA AGNATHA PROTOZOA MONOGENEAE
PROTOZOA MONOGENEA DIGENEA CESTODA
NEMATODA ACANTHOCEPHALA ARGULOIDEA
COPEPODA AGNATHA PROTOZOA MONOGENEAE
PROTOZOA MONOGENEA DIGENEA CESTODA
NEMATODA ACANTHOCEPHALA ARGULOIDEA
COPEPODA AGNATHA PROTOZOA MONOGENEAE
PROTOZOA MONOGENEA DIGENEA CESTODA
NEMATODA ACANTHOCEPHALA ARGULOIDEA
COPEPODA AGNATHA PROTOZOA MONOGENEAE
PROTOZOA MONOGENEA DIGENEA CESTODA

QL
626
C314
no. 70
c.2



Fisheries
and Oceans

Pêches
et Océans

Canada

LIBRARY / BIBLIOTHÈQUE
FISHERIES AND OCEANS / PÊCHES ET OCÉANS
OTTAWA, ONTARIO, CANADA
K1A 0E6

Annotated Checklist and Bibliography of Parasites of Herring (*Clupea harengus* L.)

J. R. Arthur¹ and H. P. Arai

*Department of Fisheries and Oceans
Fisheries Research Branch
Pacific Biological Station
Nanaimo, B.C. V9R 5K6*

DEPARTMENT OF FISHERIES AND OCEANS
OTTAWA 1984

¹Present address: E. V. S. Consultants Ltd., 2035 Mills Road, Sidney, B.C. V8L 3S1

Published by



Fisheries and Oceans
Canada

Scientific Information
and Publications Branch
Ottawa K1A 0E6

Publié par

Pêches et Océans
Canada

Direction de l'information
et des publications scientifiques
Ottawa K1A 0E6

© Minister of Supply and Services Canada 1984
Available from authorized bookstore agents
and other bookstores
or you may send your prepaid order to the
Canadian Government Publishing Centre,
Supply and Services Canada, Ottawa, Ont. K1A 0S9

Make cheques or money orders payable in
Canadian funds
to the Receiver General for Canada.

A deposit copy of this publication is also available
for reference in public libraries across Canada.

Canada: \$4.00 Catalog No. FS 41-31/70E
Other countries: \$4.80 ISBN 0-660-11659-6
ISSN 0706-6481

Price subject to change without notice
Ottawa

Printed in Canada
by
Kromar Printing Ltd.

Correct citation for this publication:

ARTHUR, J. R., AND H. P. ARAI. 1984. Annotated checklist and bibliography of parasites of herring (*Clupea harengus* L.). Can. Spec. Publ. Fish. Aquat. Sci. 70: 26 p.

Contents

ABSTRACT/RÉSUMÉ	iv
INTRODUCTION	1
PARASITE-HOST CHECKLIST	3
HOST-PARASITE CHECKLIST	17
ERRONEOUS LISTINGS	19
ACKNOWLEDGMENTS	19
REFERENCES	20

Abstract

ARTHUR, J. R., AND H. P. ARAI. 1984. Annotated checklist and bibliography of parasites of herring (*Clupea harengus* L.). Can. Spec. Publ. Fish. Aquat. Sci. 70: 26 p.

The literature containing original observations on the parasites of herring (*Clupea harengus* L.) throughout its worldwide distribution is summarized in the form of annotated Parasite–Host and Host–Parasite checklists with accompanying bibliography. The Parasite–Host checklist, arranged on a taxonomic basis, summarizes records for over 80 identified species of parasites (9 Protozoa, 13 Monogenea, 21 Digenea, 8 Cestoda, 10 Nematoda, 12 Acanthocephala, 2 Arguloidea, 6 Copepoda, 1 Agnatha) reported from herring, and includes information on numerous other parasites for which specific identity is unknown. Included are the currently recognized scientific names of the parasites, any synonyms occurring in the literature, site of occurrence (location) in the host, subspecies of herring from which the parasite has been reported, summary of the recorded distribution of the parasites in herring, and the published source for each host and locality record. Where necessary, remarks and footnotes dealing with such topics as systematics, erroneous reports, and misidentifications are included. The Host–Parasite list summarizes the species of parasites reported from each subspecies of herring and their geographic distributions.

Résumé

ARTHUR, J. R., ET H. P. ARAI. 1984. Annotated checklist and bibliography of parasites of herring (*Clupea harengus* L.). Can. Spec. Publ. Fish. Aquat. Sci. 70: 26 p.

Les auteurs résument les ouvrages contenant les données originales sur les parasites du hareng (*Clupea harengus* L.), dans toute son aire de distribution. Ces données sont présentées sous forme de listes annotées parasite-hôte et hôte-parasite et sont accompagnées d'une bibliographie. La liste parasite-hôte, disposée selon le rang taxonomique, couvre plus de 80 espèces de parasites identifiés (9 Protozoa, 13 Monogenea, 21 Digenea, 8 Cestoda, 10 Nematoda, 12 Acanthocephala, 2 Arguloidea, 6 Copepoda, 1 Agnatha), signalés chez le hareng, et comprend des données sur un grand nombre d'autres parasites dont l'identité est inconnue. Sont inclus les noms scientifiques courants, les synonymes relevés dans les ouvrages, le lieu où on le retrouve chez l'hôte, la sous-espèce de hareng parasité, un résumé de la répartition signalée des parasites chez le hareng et la source de chaque donnée sur l'hôte et l'emplacement. Au besoin, des remarques et des notes en bas de page au sujet de la systématique, de faux signalements et d'identifications erronées sont ajoutées. La liste hôte-parasite résume les espèces de parasites signalés chez chaque sous-espèce de hareng et leur répartition géographique.

Introduction

Because of its great historical importance as a food fish and its wide distribution and great abundance throughout coastal Holarctic waters, the herring (*Clupea harengus* L.) has probably received more parasitological attention than any other species of marine fish. Since the first published account of the investigations of van Leeuwenhoek (1697), almost 300 papers and manuscripts spanning four centuries have recorded original observations on the parasites of herring. Recorded in this literature are over 80 species of parasites with numerous other reports under specific names for which the true identity cannot be determined. As no complete summary of the records of herring parasites has appeared since von Linstow (1878, 1889) and as the parasite fauna of herring has now been well documented in all major areas of its geographical distribution, it appears timely, as an aid to subsequent workers, to bring together and briefly review, in the form of an annotated checklist and bibliography, the literature reporting original observations on the parasites of herring.

The earliest published records and descriptions of parasitic helminths infecting herring were based on collections made from Atlantic herring (*Clupea harengus harengus* L.) originating from the northeastern Atlantic Ocean and Baltic Sea. These early records, by such workers as Martin (1760), Linnaeus (1767, 1789), Walch (1778), Bloch (1782), Fabricius (1794), and Zeder (1800), among others, are mainly of historical interest, the descriptions being too inadequate by today's standards to permit positive recognition of the species of parasites involved. Additional reports, contained primarily in taxonomic studies of restricted groups of helminths, were added by 19th century parasitologists, among them Bellingham (1844), Cobbold (1866), van Beneden (1871) and Olsson (1866-67, 1867-68, 1893). In the latter part of the 19th century, von Linstow (1878, 1889) was able to list 12 species of parasites (1 Monogenea, 3 Digenea, 2 Cestoda, and 6 Nematoda) occurring in herring. Since that time the parasite fauna of Atlantic herring has been well investigated throughout most of its range. In the Baltic Sea, major studies are those of Schneider (1902b, 1903), Forssell (1905), Markowski (1933), Lundström (1942), Petrushevsky (1957), Getsevichyute (1958), Arro (1964), Reimer (1970), Rokicki (1973, 1975), Grabda (1974a, b), and Gaevskaya (1977, 1979). In the Aral Sea the parasite fauna of herring transplanted from the Baltic has been examined by Osmanov (1971) and in the Barents Sea Atlantic herring have been studied by Polyansky (1955). In the northeastern Atlantic Ocean, major studies include those of Odhner (1905), Nicoll (1907, 1910, 1915), Kabata (1963), Khalil (1969), Davey (1972a), and Reimer and Jessen (1972). In the northwestern Atlantic Ocean, the parasite fauna of herring has been extensively studied, major contributions being those of Linton (1901a, 1924, 1940), Stafford (1904, 1907), Cooper (1915), Heller (1949), and Sindermann (1961a, 1963, 1965). Dollfus (1956) summarized the parasites recorded from herring from the North Atlantic Ocean and Baltic Sea, noting the occurrence of 35 species, and later (Dollfus 1970) provided a detailed and extensive review of the nematodes recorded from both Atlantic and Pacific herring.

Although the Pacific herring (*C. h. pallasi* Valenciennes) has in recent years received considerable parasitological attention, it is only since Fraser (1920) that reports on its parasite fauna have appeared in the literature. In the northeastern Pacific Ocean, the eastern Bering Sea and the Sea of Japan herring have been included in major parasitological surveys of the fishes of these regions by Soviet parasitologists, among them Layman (1930), Akhmerov (1955), Strelkov (1960), Zhukov (1960a, b, 1963, 1964), and Skrabina (1963). In the northwestern Pacific Ocean and western Bering Sea, major contributions are those of Bishop and Margolis (1955), Arai (1967a, 1969), and Arthur and Arai (1980a, b), while in the eastern Palaearctic, major reports on the parasites of White Sea herring (*C. h. pallasi n. maris-albi* Berg) are those of Shulman

and Shulman-Albova (1953), Shulman (1956), and Kulachkova (1971, 1975), among others.

Besides records occurring in restricted taxonomic studies and in parasitological surveys, a considerable volume of literature has been published in recent years dealing with the occurrence and aspects of the biology of larval *Anisakis simplex* (Rudolphi, 1809), a common and widely distributed parasite of herring which has potential significance as a human pathogen. (For reviews on this subject see Kagei 1969; Ruitenberg 1970; and Oshima 1972.) Recent studies have also dealt with aspects of the ecology of herring parasites (see, for example Shulman 1956; Polyansky and Shulman 1956; Kabata 1963; Rokicki 1973; and van Banning and Becker 1978), the zoogeography of the herring and its parasites (Dogiel 1940; Svetovidov and Shulman 1960; Svetovidov 1961), and the use of parasites as "biological tags" to study the discreteness of herring stocks, their movements, and the recruitment of young herring to adult populations (see, for example Sindermann 1957a, 1961a; MacKenzie 1974a, b, 1975a, b, 1978; MacKenzie and Johnston 1976; MacKenzie and Gallacher 1981; Kulachkova 1974b; and Arthur and Arai 1980b).

The records of parasites of *Clupea harengus* presented in this checklist were obtained primarily from the published literature but, in an effort to provide the reader with as much information as possible, records were also obtained from other sources. These included various unpublished research documents (primarily those of the International Council for the Exploration of the Sea and the International Northwest Atlantic Fisheries Commission), certain reports and pamphlets issued by various governmental departments, abstracts of scientific meetings, and university dissertations. Undoubtedly some records contained in documents of this nature have escaped the authors' attention. Additionally some records contained in foreign language papers detailing experimental studies on *Anisakis* larvae isolated from herring may have been overlooked. Reports of a very general nature and those in obvious reference to previous workers have not been included, the reports summarized being restricted to original observations only.

The basis of the system of higher classification used herein follows that given by Margolis and Arthur (1979) with modification of the "Protozoa" as given by Levine et al. (1980). Original records of parasites from herring are summarized in the form of a PARASITE-HOST LIST, a HOST-PARASITE LIST, discussion of species erroneously listed as occurring in herring, and an accompanying bibliography. For each species of parasite contained in the Host-Parasite list the following information is provided:

1) The current *scientific name*, including author(s) and date(s) followed by any recognized synonyms that have been used in establishing records from herring. No attempt has been made to evaluate systematically the validity of published reports; however, attention is drawn to obvious errors, tentative or uncertain identifications, and to cases where identifications have been questioned or material redetermined by subsequent workers.

2) The *location* or site of occurrence of the parasite in its host. In cases where the location reported is considered to be aberrant or where it was unspecified in the original report, the typical site of infection, when possible, was ascertained from records from other host species and is enclosed in square brackets.

3) The *hosts*. Records are listed by host based on the currently accepted division of the herring into two subspecies, the Atlantic herring (*Clupea harengus harengus* L.) and the Pacific herring (*C. h. pallasi* Valenciennes). In cases where only common names were given by original authors, assignment to subspecies, where possible, was based on the common name used or on collection locality. Records for which the subspecies could not be determined with confidence are listed under *Clupea harengus*. The following common names are encountered in the literature: for *C. harengus harengus*—Atlantic herring (American Fisheries Society recognized common name), Baltic herring, high vertebral count herring, herring, haring (Dutch), Hering (German), hareng (French), salaka

(Russian), sild (Norwegian), Strömling (German), and sledż (Polish); and for *C. h. pallasi*—Pacific herring (AFS recognized common name), White Sea herring, Chelsa-Pechora herring, low vertebral count herring, herring, nishin (Japanese), and sel'd (Russian). Numbers in parentheses after each host name correspond with the numbers assigned to the references establishing the particular parasite-host records.

4) The *distribution* (Dist). The geographical distribution of the herring is divided into the following regions: northeastern Atlantic Ocean (NE Atl), northwestern Atlantic Ocean (NW Atl), Baltic Sea, northeastern Pacific Ocean (NE Pac), northwestern Pacific Ocean (NW Pac), eastern Bering Sea (E Ber), western Bering Sea (W Ber), Aral Sea, eastern Palaearctic (E Pal), western Palaearctic (W Pal), eastern Nearctic (E Nea), and western Nearctic (W Nea). Where possible, records from the eastern Palaearctic region are further designated as pertaining to the White, Barents, Norwegian, or Greenland Seas. The North Pacific Ocean and Bering Sea are separated from each other by a line drawn through the Aleutian Islands to Ust Kamchatsk and are divided into eastern and western regions at 170°W longitude. The North Atlantic Ocean is similarly divided into eastern and western regions at 30°W longitude. The area north of the Arctic Circle is divided into four regions, east Palaearctic (0–90°E longitude), western Palaearctic (90–180°E longitude), eastern Nearctic (0–90°W longitude) and western Nearctic (90–180°W longitude). At present no records have been made from the western and eastern Nearctic regions or from the western Palaearctic region.

5) The *record(s)*. The authors responsible for the records of parasites are given in chronological order with each record preceded by a reference number and followed by the geographical locality or localities. This enables the reader to obtain at a glance the author(s) for a particular parasite-host-locality record, as these reference numbers correspond to those given after the appropriate host sub-

species. For example: the entries for the digenetic trematode *Lecithaster gibbosus* show that Polyansky (1955) reported this species from *C. harengus harengus* from the Barents Sea. When only one host subspecies is listed for a particular parasite the references are not numbered, and when all records are from the same geographic region it was unnecessary to list the locality after the authors' names. A special problem was encountered with the paper of Dollfus (1956), which summarizes, without citation of references, the parasites recorded from herring from the North Atlantic Ocean and Baltic Sea, including a number of original observations. References to parasites reported to occur in herring from localities that could not be traced to a previous author, but that were not clearly indicated to be original observations are listed as "Dollfus (1956) (original?)" and are accompanied by an explanatory footnote or remark.

6) *Remarks*. Under *Remarks* comments are given on such topics as systematics, possible erroneous reports, the reassignment of misidentified material, and the occurrence of species that are typically parasites of freshwater fishes.

7) *Footnotes*. Under *Footnotes* are included notes on specific items such as tentative identifications, synonyms, and ambiguous collection localities.

A *Host-Parasite list* is provided for each subspecies of *Clupea harengus*. Host synonyms given following each subspecies are only those associated with parasite records. After the name of each species of parasite its geographic distribution for the host in question is given in parentheses. An interrogation mark (?) preceding a parasite name indicates questionable validity of the parasite identification.

Under the heading *Erroneous Records* are discussed a number of species which have been incorrectly listed as occurring in *Clupea harengus*.

PARASITE-HOST CHECKLIST

PHYLUM APICOMPLEXA

CLASS SPOROZOA

ORDER EUCOCCIDIIDA

Family EIMERIIDAE

Eimeria clupearum (Thélohan, 1894) Doflein, 1909

Syn: *Coccidium clupearum* Thélohan, 1894

Location: liver

Hosts: *Clupea harengus harengus* (1, 2, 3, 4, 5, 6, 7)

C. harengus pallasi (8, 9, 10)

Dist: NE Atl, NW Atl, NE Pac, E Ber Sea

Records: 1. Thélohan 1892 (NE Atl), 2. 1894 (NE Atl); 3. Dollfus 1956 (NE Atl) (original?); 4. Sindermann 1961a (NW Atl), 5. b (NW Atl); 6. Kabata 1963 (NE Atl); 7. Reimer and Jessen 1972 (NE Atl); 8. Arthur 1978 (NE Pac, E Ber Sea); 9. Arthur and Arai 1980a (NE Pac, E Ber Sea), 10. b (NE Pac)

Eimeria nishin Fujita, 1934

Location: testis

Host: *Clupea harengus pallasi*

Dist: NE Pac, NW Pac, E Ber Sea

Records: Fujita 1934 (NW Pac); Arthur 1978 (NE Pac, E Ber Sea); Arthur and Arai 1980a (NE Pac, E Ber Sea), b (NE Pac)

Remarks: Dogiel (1940) considered the validity of *E. nishin* to be doubtful. Arthur and Arai (1980a) note that it is probably a synonym of *E. sardinae* (Thélohan, 1890).

Eimeria sardinae (Thélohan, 1890) Reichenow, 1921

Location: testis

Hosts: *Clupea harengus harengus* (3, 6, 7, 8, 9, 10, 11, 12, 13)

C. harengus pallasi (1, 2, 4, 5)

Dist: NE Atl, NW Atl, Baltic Sea, E Pal (White Sea, Barents Sea)

Records: 1. Dogiel 1940 (White Sea); 2. Shulman and Shulman-Albova 1953 (White Sea); 3. Polyansky 1955 (Barents Sea); 4. Shulman 1956 (White Sea); 5. Polyansky and Shulman 1956 (White Sea); 6. Petrushevsky 1957 (Baltic Sea); 7. Getsevichyute 1958 (Baltic Sea); 8. Sindermann 1961a (NW Atl), 9. b (NW Atl); 10. Kabata 1963 (NE Atl); 11. Gaevskaya 1977 (Baltic Sea), 12. 1979 (Baltic Sea); 13. Gaevskaya and Shapiro 1981 (Baltic Sea)

PHYLUM MICROSPORA

CLASS MICROSPOREA

ORDER MICROSPORIDA

Family POLYSPORIDAE

Pleistophora sp.

Location: cysts in musculature

Host: *Clupea harengus harengus*

¹Dollfus (1956) noted the presence of *E. clupearum* in herring from British waters. We are unaware of any previous report from this locality.

Dist: NW Atl
Records: Sindermann 1961b, 1963, 1965

Unidentified Microsporida

Microsporida gen. sp.
Location: cyst in fin
Host: *Clupea harengus pallasi*
Dist: NE Pac
Records: Arthur 1978; Arthur and Arai 1980a

PHYLUM CILIOPHORA

CLASS OLIGOHYMENOPHOREA

ORDER PERITRICHIDA

Family TRICHODINIDAE

Trichodina ploveri Zhukov, 1964
Location: gills
Host: *Clupea harengus pallasi*
Dist: W Ber Sea
Record: Zhukov 1964
Remarks: Lom and Laird (1969) regarded this species as a *nomen dubium*.

Trichodinidae gen. sp.
Location: gills
Host: *Clupea harengus harengus*
Dist: NW Atl
Record: Lom and Laird 1969

PHYLUM MYXOZOA

CLASS MYXOSPOREA

ORDER BIVALVULIDA

Family CERATOMYXIDAE

Ceratomyxa acadiensis Mavor, 1915
Location: gallbladder
Host: *Clupea harengus harengus*
Dist: NW Atl
Records: Ellis 1930,² Sindermann 1961b
Remarks: The above records are probably referable to *C. auerbachii* Kabata, 1962.

Ceratomyxa auerbachii Kabata, 1962
Syn: *Ceratomyxa sphaerulosa* of Auerbach, 1909,³ 1912
Location: gallbladder
Hosts: *Clupea harengus harengus* (1, 2, 3)
C. harengus pallasi (4, 5, 6)
Dist: NE Atl, NE Pac

²Record of Ellis (1930) is a tentative parasite identification.

³Kabata (1962) referred *C. sphaerulosa* of Auerbach, 1909 to *C. auerbachii*.

Records: 1. Auerbach 1909 (NE Atl), 2. 1912 (NE Atl);
3. Kabata 1962 (NE Atl); 4. Arthur 1978 (NE Pac);
5. Arthur and Arai 1980a (NE Pac), 6. b (NE Pac)

Ceratomyxa orientalis (Dogiel, 1948) Shulman, 1953

Location: gallbladder

Host: *Clupea harengus pallasi*

Dist: E Pal (White Sea), NW Pac

Records: Shulman and Shulman-Albova 1953 (White Sea, NW Pac); Shulman 1953 (White Sea), 1956 (White Sea); Polyansky and Shulman 1956 (White Sea)

Remarks: The occurrence of this species in *Clupea harengus* requires reassessment. *Ceratomyxa truncata orientalis*, originally described from *Sardinella melanosticta* (syn. of *Sardinops sagax melanosticta*) from the northwestern Pacific Ocean by Dogiel (1948), was first reported (as *C. orientalis*) from *Clupea harengus pallasi* by Shulman (1953) from the White Sea and was additionally recorded from this host from the northwestern Pacific Ocean by Shulman and Shulman-Albova (1953). However, substantial differences in the shape and size of spores reported from these hosts cause us to doubt their conspecificity. Spores from *Sardinella* are arcuate in shape and measure 42–45 µm in width (Dogiel 1948) whereas those from *Clupea* are only slightly curved and measure 33–72 µm in width (49–72 µm for mature spores) (Shulman 1953, 1966; Shulman and Shulman-Albova 1953). Spores of *Ceratomyxa auerbachi* reported by Arthur and Arai (1980a) from northeastern Pacific herring also differ from those of *C. orientalis* from *Sardinella* by possessing slightly curved to straight valves and by greater spore width (72–108 µm). It therefore seems possible that *C. orientalis* does not infect *Clupea harengus* and that a single species, *C. auerbachi*, may occur throughout its distribution.

Leptotheeca sp.

Location: kidney tubules

Host: *Clupea harengus pallasi*

Dist: NE Pac

Records: Arthur 1978; Arthur and Arai 1980a

Family SINUOLINEIDAE

Ortholinea orientalis (Shulman and Shulman-Albova, 1953) Shulman, 1965

Syn: *Sphaerospora orientalis* Shulman and Shulman-Albova, 1953

Location: kidney tubules

Host: *Clupea harengus pallasi*

Dist: E Pal (White Sea); NE Pac

Records: Shulman and Shulman-Albova 1953 (White Sea); Polyansky and Shulman 1956 (White Sea); Shulman 1956 (White Sea); Arthur 1978 (NE Pac); Arthur and Arai 1980a (NE Pac)

Family SPHAEROSPORIDAE

Chloromyxum sp.

Location: gallbladder

Host: *Clupea harengus pallasi*

Dist: E Pal (White Sea)

Records: Shulman and Shulman-Albova 1953; Shulman 1956

ORDER MULTIVALVULIDA

Family TETRACAPSULIDAE

Kudoa clupeidae (Hahn, 1917) Meglitsch, 1947

Syn: *Chloromyxum clupeidae* Hahn, 1917

Kudoa sp. of Laird and Bullock, 1969 (partim)⁴

Includes: Myxosporida of Tyzzer, 1900

Sporozoa of Linton, 1901, 1910

Location: cysts in musculature

Host: *Clupea harengus harengus*

Dist: NE Atl, NW Atl

Records: Tyzzer 1900 (NW Atl); Linton 1901a (NW Atl), 1910 (NW Atl); Hahn 1917 (NW Atl); Sindermann and Rosenfield 1954a (NW Atl); Sindermann and Scattergood 1954 (NW Atl); Sindermann 1957a (NW Atl), b (NW Atl), 1959 (NW Atl), 1961a (NW Atl), b (NW Atl), 1963 (NW Atl), 1965 (NW Atl); Laird and Bullock 1969 (NW Atl); MacKenzie 1983 (NE Atl).

Remarks: As *K. clupeidae* has not been reported from Pacific herring, reports of this species from rockfishes (*Sebastes* spp.) of the northeastern Pacific Ocean (Moser et al. 1976; Heckmann and Jensen 1978) are equivocal.

Unidentified Protozoa

"Sporozoa"

Location: not specified

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Record: Arro 1964

PHYLUM PLATYHELMINTHES

CLASS MONOGENEA

ORDER MONOPISTHOCTYLEA

Superfamily GYRODACTYLOIDEA

Family GYRODACTYLIDAE

Gyrodactyloides baueri Kulachkova, 1970

Location: nasal cavities

Hosts: *Clupea harengus harengus* (1, 5, 7)

C. harengus pallasi (1, 2, 3, 4, 5, 6, 7)

Dist: E Pal (White Sea), W Ber Sea

Records: 1. Kulachkova 1970 (White Sea), 2. 1971 (White Sea), 3. 1974a (White Sea), 4. b (White Sea), 5. 1975 (White Sea), 6. 1977a (White Sea), 7. b (White Sea, W Ber Sea)

Gyrodactyloides petruschewskii Bykhovsky, 1947

Location: gills

Host: *Clupea harengus pallasi*

Dist: E Pal (White Sea), NE Pac, W Ber Sea

Records: Zhukov 1960a (W Ber Sea); Kulachkova 1971 (White Sea), 1974a (White Sea), b (White Sea), 1975 (White Sea), 1977a (White Sea), b (White Sea, W Ber Sea); Arthur 1978 (NE Pac); Arthur and Arai 1980a (NE Pac)

⁴Laird and Bullock (1969) reported *Kudoa* sp. from blood films taken from the heart of several species of northwest Atlantic fishes. They noted that some if not all of their specimens were referable to *K. clupeidae*.

Gyrodactyloides spp.

Location: nasal cavities, [gills]
 Hosts: *Clupea harengus harengus*
C. harengus pallasi
 Dist: E Pal (White Sea)
 Record: Kulachkova 1968

Gyrodactylus cyclopteri Stsiborskaya, 1948

Location: gills
 Host: *Clupea harengus pallasi*
 Dist: E Pal (White Sea)
 Record: Kulachkova 1975

Gyrodactylus flesi Malmberg, 1957

Location: gills
 Host: *Clupea harengus pallasi*
 Dist: E Pal (White Sea)
 Records: Kulachkova 1971, 1974b, 1975, 1977a

Gyrodactylus gerdi Bykhovsky, 1948

Location: gills
 Host: *Clupea harengus pallasi*
 Dist: E Pal (White Sea)
 Records: Kulachkova 1974b, 1975, 1977a

Gyrodactylus groenlandicus Levinsen, 1881

Includes: *Gyrodactylus groenlandicus groenlandicus* Levinsen, 1881
 Location: gills
 Host: *Clupea harengus pallasi*
 Dist: E Pal (White Sea)
 Records: Kulachkova 1971, 1974b, 1975, 1977a

Gyrodactylus harengi Malmberg, 1957

Location: gills, fins
 Hosts: *Clupea harengus harengus* (1, 5)
C. harengus pallasi (2, 3, 4, 5, 6, 7, 8, 9)
 Dist: E Pal (White Sea), Baltic Sea, W Ber Sea, NE Pac
 Records: 1. Malmberg 1957 (Baltic Sea); 2. Zhukov 1960a (W Ber Sea); 3. Kulachkova 1971 (White Sea), 4. 1974b (White Sea), 5. 1975 (White Sea), 6. 1977a (White Sea), 7. b (W Ber Sea); 8. Arthur 1978 (NE Pac); 9. Arthur and Arai 1980a (NE Pac)
 Remarks: Malmberg (1957) also noted *G. harengi* or a closely related species on *C. h. harengus* from the northwestern Atlantic Ocean.

Gyrodactylus pterygialis Bykhovsky and Polyansky, 1953

Location: gills
 Host: *Clupea harengus pallasi*
 Dist: E Pal (White Sea)
 Records: Kulachkova 1974b, 1975, 1977a

Gyrodactylus pungitii Malmberg, 1964

Location: gills
 Host: *Clupea harengus pallasi*
 Dist: E Pal (White Sea)
 Records: Kulachkova 1974b, 1975, 1977a

Gyrodactylus robustus Malmberg, 1957

Location: [fins]
 Host: *Clupea harengus pallasi*
 Dist: E Pal (White Sea), W Ber Sea
 Records: Kulachkova 1971 (White Sea), 1974b (White Sea), 1975 (White Sea), 1977a (White Sea), b (W Ber Sea)

Gyrodactylus spp.

Location: gills
 Hosts: *Clupea harengus harengus* (1)

C. harengus pallasi (2, 4, 5, 6, 7)

C. harengus (2)

Dist: NW Atl, E Pal (White Sea)

Records: 1. Sindermann 1957a (NW Atl); 2. Kulachkova 1968 (White Sea), 3. 1971 (White Sea), 4. 1974a (White Sea), 5. b (White Sea), 6. 1975 (White Sea), 7. 1977a (White Sea)

Laminiscus dogieli (Zhukov, 1960) Pálsson and Beverley-Burton, 1983.

Syn: *Gyrodactyloides dogieli* Zhukov, 1960

Location: gills

Hosts: *Clupea harengus harengus* (5, 7)

C. harengus pallasi (1, 2, 3, 4, 5, 6, 7)

Dist: E Pal (White Sea), W Ber Sea

Records: 1. Zhukov 1960a (W Ber Sea); 2. Kulachkova 1971 (White Sea), 3. 1974a (White Sea), 4. b (White Sea), 5. 1975 (White Sea), 6. 1977a (White Sea), 7. b (White Sea, W Ber Sea)

Unidentified Gyrodactyoidea

Gyrodactyoidea gen. spp.

Location: gills, body surface

Host: *Clupea harengus pallasi*

Dist: E Pal (White Sea), NE Pac

Records: Arai 1967a (NE Pac), 1969 (NE Pac); Kulachkova 1977b (White Sea)

ORDER POLYOPISTHOCOTYLEA

Superfamily DICLIDOPHOROIDEA

Family MAZOCRAEIDAE

Mazocraeoides georgei Price, 1936

Location: gills

Host: *Clupea harengus harengus*

Dist: NW Atl

Record: Price 1961

Mazocraes harengi (van Beneden and Hesse, 1863) Baylis and Jones, 1933

Syn: *Octobothrium harengi* (van Beneden and Hesse, 1863)
Octocotyle harengi van Beneden and Hesse, 1863

Location: gills

Host: *Clupea harengus harengus*

Dist: NE Atl

Records: van Beneden and Hesse 1864; Scott 1901

CLASS TREMATODA

SUBCLASS DIGENEA

ORDER STRIGEIDA

Superfamily STRIGEOIDEA

Family DIPLOSTOMATIDAE

Diplostomum spathaceum (Rudolphi, 1819) Olsson, 1876
 (metacercaria)

Syn: *Diplostomulum spathaceum* (Rudolphi, 1819)

Location: lens of eye

Hosts: *Clupea harengus harengus* (3, 4, 5, 6, 7)

C. harengus pallasi (1, 2)

Dist: E Pal (White Sea), Baltic Sea

Records: 1. Shulman and Shulman-Albova 1953 (White Sea); 2. Shulman 1956 (White Sea); 3. Petrushevsky 1957

(Baltic Sea); 4. Getsevichyute 1958 (Baltic Sea); 5. Reimer 1962 (Baltic Sea), 6. 1970 (Baltic Sea); 7. Gaevskaia 1979 (Baltic Sea) (original?)

Remarks: *Diplostomum spathaceum* is typically a parasite of freshwater fishes.

Superfamily HEMIUROIDEA

Family DEROGENIDAE

Derogenes varicus (O. F. Müller, 1784) Looss, 1901

Location: stomach, intestine

Hosts: *Clupea harengus harengus* (1, 2, 3, 5, 8, 11)
C. harengus pallasi (4, 6, 7, 9, 10)

Dist: NE Atl, NW Atl, E Pal (Barents Sea, White Sea), NW Pac

Records: 1. Stafford 1904 (NW Atl), 2. 1907 (NW Atl); 3. Cooper 1915 (NW Atl);⁵ 4. Shulman and Shulman-Albova 1953 (White Sea); 5. Polyansky 1955 (Barents Sea); 6. Polyansky and Shulman 1956 (White Sea); 7. Shulman 1956 (White Sea); 8. Dollfus 1956 (NE Atl) (original?);⁶ 9. Zhukov 1960b (NW Pac); 10. Skrjabina 1963 (NW Pac); 11. Reimer and Jessen 1972 (NE Atl)

Family HEMIURIDAE

Brachyphallus crenatus (Rudolphi, 1802) Odhner, 1905

Syn: *Brachyphallus amuriensis* Babaskin, 1928
Hemiurus luehei of Markowski, 1933⁷

Location: stomach, intestine

Hosts: *Clupea harengus harengus* (1, 2, 3, 5, 6, 7, 10, 13, 14, 19, 22, 23, 24, 25, 26, 30)
C. harengus pallasi (4, 8, 9, 11, 12, 15, 16, 17, 18, 20, 21, 27, 28, 29)

Dist: NE Atl, NW Atl, Baltic Sea, E Pal (Barents Sea, White Sea), E Ber Sea, W Ber Sea, NE Pac, NW Pac

Records: 1. Cooper 1915 (NW Atl);⁸ 2. Manter 1925 (NW Atl), 3. 1926 (NW Atl); 4. Layman 1930 (NW Pac); 5. Markowski 1933 (Baltic Sea); 6. Linton 1940 (NW Atl); 7. Heller 1949 (NW Atl); 8. Shulman-Albova 1952 (White Sea); 9. Shulman and Shulman-Albova 1953 (White Sea); 10. Polyansky 1955 (Barents Sea); 11. Shulman 1956 (White Sea); 12. Polyansky and Shulman 1956 (White Sea); 13. Sindermann 1957a (NW Atl), 14. 1961a (NW Atl); 15. Strelkov 1960 (NW Pac); 16. Zhukov 1960b (NW Pac); 17. 1963 (W Ber Sea); 18. Skrjabina 1963 (NW Pac); 19. Reimer 1970 (Baltic Sea); 20. Ivanchenko and Grozdilova 1971 (experimental — White Sea); 21. 1981 (experimental — White Sea); 22. Reimer and Jessen 1972 (NE Atl); 23. Rokicki 1973 (Baltic Sea), 24. 1975 (Baltic Sea); 25. Gaevskaia 1977 (Baltic Sea), 26. 1979 (Baltic Sea); 27. Arthur 1978 (NE Pac, E Ber Sea); 28. Arthur and Arai 1980a (NE Pac, E Ber Sea), 29. b (NE Pac); 30. Gaevskaia and Shapiro 1981 (Baltic Sea)

⁵Cooper (1915) reported mature (egg bearing) *D. varicus* encysted on the viscera and in the musculature of *C. h. harengus*.

⁶Dollfus (1956) noted that *D. varicus* has been found very commonly in herring from the Plymouth region. We were unable to locate any previous records of *D. varicus* from herring from this locality.

⁷Rokicki (1973) referred *H. luehei* of Markowski, 1933 to *Brachyphallus*. Reimer (1970) corrected this determination to *B. crenatus*.

⁸Record of Cooper (1915) includes one specimen encysted in the musculature.

Hemiurus appendiculatus (Rudolphi, 1802) Looss, 1899

Syn: *Distoma appendiculatum* (Rudolphi, 1802)

Location: stomach, intestine

Hosts: *Clupea harengus harengus* (2, 3, 4, 5)
C. harengus (1)

Dist: NE Atl, NW Atl, Baltic Sea

Records: 1. Creplin 1839 (unspecified locality);⁹ 2. van Beneden 1871 (NE Atl); 3. Linton 1901a (NW Atl),¹⁰ 4. 1940 (NW Atl); 5. Petrushevsky 1957 (Baltic Sea)

Remarks: The report of *H. appendiculatus* from *C. h. harengus* by Stafford (1907) is referred to *H. levinseni* Odhner, 1905 following Miller (1941). Odhner (1905) referred the material reported as *Distoma appendiculatum* from *C. h. harengus* by Olsson (1867–68) to *Hemiurus luehei* Odhner, 1905.

Dr D. I. Gibson (British Museum (Natural History), London, England, personal communication) regards all records of *H. appendiculatus* from *C. harengus* to be probable misidentifications of *H. luehei*.

Hemiurus levinseni Odhner, 1905

Syn: *Hemiurus appendiculatus* of Stafford, 1907 (partim)¹¹

Location: stomach, pyloric caeca

Hosts: *Clupea harengus harengus* (1, 2, 3)
C. harengus pallasi (4, 5, 6)

Dist: NW Atl, E Pal (Barents Sea), NE Pac, NW Pac

Records: 1. Cooper 1915 (NW Atl);¹² 2. Miller 1941 (NW Atl); 3. Polyansky 1955 (Barents Sea); 4. Skrjabina 1963 (NW Pac); 5. Arthur 1978 (NE Pac); 6. Arthur and Arai 1980a (NE Pac)

Hemiurus luehei Odhner, 1905

Syn: *Fasciola ocreata* Rudolphi, 1802

Distoma ocreata (Rudolphi, 1802)

Hemiurus ocreatus (Rudolphi, 1802)

?*Hemiurus halecis* (Gmelin, 1790)

Distoma appendiculatum of Olsson, 1867–68 (partim)¹³

Includes: ?"vermiculi" of van Leeuwenhoek, 1697

Location: stomach, pyloric caeca, intestine

Hosts: *Clupea harengus harengus* (4, 6, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21)

C. harengus (1, 2, 3, 5, 7)

Dist: NE Atl, Baltic Sea

Records: 1. van Leeuwenhoek 1697 (unspecified locality); 2. Rudolphi 1802 (unspecified locality), 3. 1809 (unspecified locality), 4. 1819 (Baltic Sea); 5. Krøyer 1846–49 (unspecified locality) (original?); 6. Olsson 1867–68 (NE Atl); 7. Odhner 1905 (unspecified locality); 8. Nicoll 1907 (NE Atl), 9. 1910 (NE Atl), 10. 1915 (NE Atl); 11. Markowski 1938 (NE Atl);¹⁴ 12. Hansen 1955 (NE Atl); 13. Dollfus 1956 (NE Atl) (original?);¹⁵

⁹Specimens reported as *Distoma appendiculatum* by Creplin (1839) were probably of Baltic origin.

¹⁰The report of Linton (1901a) was a tentative parasite identification.

¹¹Miller (1941) identified material from the collection of Stafford as *H. levinseni*. He considered these specimens probably to have been reported as *H. appendiculatus* by Stafford (1907).

¹²The record of Cooper (1915) was for metacercariae of *H. levinseni* encysted in the musculature of *C. h. harengus*.

¹³Odhner (1905) referred *Distoma appendiculatum* of Olsson (1866–67) from *Clupea harengus* to *H. luehei*.

¹⁴The record of Markowski (1938) was given as "Hemiurus ocreatus (Molin 1863)."

¹⁵Dollfus (1956) noted that *H. luehei* (as *H. halecis*) occurred in herring off Belgium and northern France. We are unaware of any previous records of *H. luehei* from *C. harengus* from these localities.

14. Willemse 1968 (NE Atl); 15. Reimer 1970 (Baltic Sea); 16. Reimer and Jessen 1972 (NE Atl); 17. Rokicki 1973 (Baltic Sea), 18. 1975 (Baltic Sea); 19. Gaevskaya 1977 (Baltic Sea), 20. 1979 (Baltic Sea); 21. Gaevskaya and Shapiro 1981 (Baltic Sea).

Remarks: *Hemiuirus luehei* has an involved taxonomic history. Van Leeuwenhoek (1697) first recorded as "vermiculi" hemiurid trematodes from the stomach of herring purchased in Amsterdam. The name *Fasciola halecis* was subsequently used by Gmelin (1791) for van Leeuwenhoek's specimens. *Fasciola halecis* appears to later have been renamed *F. ocreata* by Rudolphi (1802). Unfortunately this name is a junior primary homonym of *F. ocreata* Goeze, 1782, now the type-species of *Ityogninus*. Odhner (1911) reexamined Rudolphi's specimens of *F. ocreata*, identifying them as *H. luehei*. Dollfus (1956) resurrected *Fasciola halecis* as *Hemiuirus halecis* (Gmelin, 1790). However this name is considered a questionable synonym of *H. luehei*, as the material described by van Leeuwenhoek is in fact unrecognizable. *Hemiuirus luehei* of Markowski (1933) is referred to *Brachyphallus crenatus* following Reimer (1970).

Hemiuirus raabei Slusarski, 1958

Location: stomach

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Rokicki 1973, 1975; Gaevskaya 1977

Remarks: The validity of *H. raabei* requires confirmation.

Perhaps it is a synonym of *H. luehei*.

Parahemiuirus merus (Linton, 1910) Woolcock, 1935

Location: stomach, intestine

Hosts: *Clupea harengus harengus* (5, 6)

C. harengus pallasi (1, 2, 3, 4, 7, 8)

Dist: Baltic Sea, NE Pac

Records: 1. Nahhas 1960 (NE Pac); 2. Arai 1967a (NE Pac), 3. 1969 (NE Pac); 4. Chapa 1969 (NE Pac); 5. Rokicki 1973 (Baltic Sea); 6. Gaevskaya 1977 (Baltic Sea); 7. Arthur 1978 (NE Pac); 8. Arthur and Arai 1980a (NE Pac)

Remarks: The occurrence of *P. merus* in the Baltic Sea requires substantiation. D. I. Gibson (personal communication) considers it possible that the records of Rokicki (1973) and Gaevskaya (1977) may be based on anomalous *H. luehei*.

Family LECITHASTERIDAE

Lecithaster confusus Odhner, 1905

Syn: *Distomum botryophoron* of Linton, 1901¹⁶

Location: intestine

Hosts: *Clupea harengus harengus* (1, 2, 3, 5, 8, 9)

C. harengus pallasi (4, 6, 7)

Dist: Baltic Sea, NW Atl, E Pal (White Sea, Barents Sea)

Records: 1. Linton 1901a (NW Atl); 2. 1940 (NW Atl); 3. Odhner 1905 (unspecified locality);¹⁷ 4. Shulman and Shulman-Albova 1953 (White Sea); 5. Polyansky 1955 (Barents Sea); 6. Shulman 1956 (White Sea); 7. Polyansky and Shulman 1956 (White Sea); 8. Reimer 1970 (Baltic Sea); 9. Odening 1978 (Baltic Sea)

¹⁶Linton (1940) referred his tentative identification of *Distomum botryophoron* (as "*Distomum bothryophoron* Olsson (?)") from *C. h. harengus* (Linton 1901a) to *L. confusus*.

¹⁷Odhner (1905) reported *L. confusus* as occurring in *Clupea harengus* in northern seas.

Lecithaster gibbosus (Rudolphi, 1802) Lühe, 1901

Syn: *Lecithaster salmonis* Yamaguti, 1934

Lecithaster bothryophorus of Stafford, 1904, 1907

Location: stomach, intestine

Hosts: *Clupea harengus harengus* (1, 2, 3, 4, 5, 16)
C. harengus pallasi (6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 17, 18, 19, 20)

Dist: NE Atl, NW Atl, Baltic Sea, E Pal (Barents Sea, White Sea), NE Pac, E Ber Sea, W Ber Sea

Records: 1. Stafford 1904 (NW Atl), 2. 1907 (NW Atl); 3. Odhner 1905 (unspecified locality);¹⁸ 4. Polyansky 1955 (Barents Sea); 5. Dollfus 1956 (NE Atl);¹⁹ 6. Nahhas 1960 (NE Pac); 7. Zhukov 1963 (W Ber Sea); 8. Arai 1967a (NE Pac), 9. b (NE Pac), 10. 1969 (NE Pac); 11. Barracough 1967 (NE Pac); 12. Barracough and Fulton 1967 (NE Pac); 13. Robinson et al. 1968a (NE Pac), 14. b (NE Pac); 15. Chapa 1969 (NE Pac); 16. Rokicki 1973 (Baltic Sea); 17. Arthur 1978 (NE Pac, E Ber Sea); 18. Arthur and Arai 1980a (NE Pac, E Ber Sea), 19. b (NE Pac); 20. Ivanchenko and Grozdilova 1981 (experimental — White Sea)

Lecithaster sp.

Location: digestive tract

Host: *Clupea harengus pallasi*

Dist: NW Pac

Record: Layman 1930

Superfamily FELLODISTOMATOIDEA

Family FELLODISTOMATIDAE

Pronoprymna petrowi (Layman, 1930) Bray and Gibson, 1980

Syn: *Bacciger petrowi* (Layman, 1930)

Pentagramma petrowi (Layman, 1930)

Pseudopentagramma petrowi (Layman, 1930)

Location: pyloric caeca, intestine

Host: *Clupea harengus pallasi*

Dist: NE Pac, E Ber Sea, W Ber Sea

Records: Zhukov 1963 (W Ber Sea); Margolis and Ching 1965 (NE Pac); Barracough and Fulton 1967 (NE Pac); Arai 1967a (NE Pac), 1969 (NE Pac); Arthur 1978 (NE Pac, E Ber Sea); Arthur and Arai 1980a (NE Pac, E Ber Sea), b (NE Pac)

Superfamily BUCEPHALOIDEA

Family BUCEPHALIDAE

Prosrhynchoides basargini (Layman, 1930) Margolis and Arthur, 1979 (metacercaria)

Includes: Bucephalidae gen. sp. of Arthur and Arai, 1979 (partim)

Location: encysted in fins, nasal cavities, mouth

Dist: NE Pac, E Ber Sea

Records: Arthur 1978 (NE Pac, E Ber Sea); Arthur and Arai 1979 (NE Pac), 1980a (NE Pac, E Ber Sea), b (NE Pac)

¹⁸Odhner (1905) reported *L. gibbosus* from a number of Scandinavian fishes but did not give precise localities.

¹⁹Dollfus (1956) noted that this species had been reported from Scottish herring. We have not encountered any previous report from this locality.

Prosorhynchoides gracilescens (Rudolphi, 1819) Stunkard, 1976
(metacercaria)
Syn: *Bucephalopsis gracilescens* (Rudolphi, 1819)
Location: encysted in mesenteries of heart
Host: *Clupea harengus pallasi*
Dist: NW Pac
Record: Zhukov 1960b

Rhipidocotyle sp. metacercaria
Syn: *Rhipidocotyle elongata* of Arthur, 1978
Includes: Bucephalidae gen. sp. of Arthur and Arai, 1979
(partim)
Location: encysted in fins, nasal cavities, mouth
Host: *Clupea harengus pallasi*
Dist: NE Pac
Records: Arthur 1978; Arthur and Arai 1979, 1980a, b

ORDER ECHINOSTOMATIDA

Superfamily ECHINOSTOMATOIDEA

Family ECHINOSTOMATIDAE

Stephanopora pseudoechinata (Olsson, 1876) Yamaguti, 1958
(metacercaria)
Syn: *Mesorchis pseudoechinatus* (Olsson, 1876)
Location: not specified
Host: *Clupea harengus harengus*
Dist: Baltic Sea
Record: Reimer 1970

ORDER RENICOLIDA

Superfamily RENICOLOIDEA

Family RENICOLIDAE

Renicola spp. metacercaria
Location: encysted on pyloric caeca
Host: *Clupea harengus harengus*
Dist: NE Atl
Records: MacKenzie 1974a, b, 1975a, b, c, 1976, 1978, 1983;
MacKenzie and Johnston 1976; MacKenzie and
Gallacher 1981

ORDER PLAGIORCHIIDA

Superfamily ALLOCREADIOIDEA

Family LEPOCREADIIDAE

Neophasis oculata (Levinsen, 1881) Miller, 1941
Syn: *Distoma oculatum* Levinsen, 1881
Location: [pyloric caeca, intestine]
Host: *Clupea harengus*
Dist: not specified
Record: von Linstow 1889 (original?)
Remarks: von Linstow (1889) listed this species (as "*Distomum oculatum* Rud.") from *Clupea harengus*. No collection locality or reference for this record was given.

Opechona bacillaris (Molin, 1859) Looss, 1907

Location: digestive tract
Host: *Clupea harengus harengus*
Dist: NE Atl
Record: Dollfus 1956 (original?)
Remarks: Ward and Fillingham (1934) attributed to Nicoll (publication not specified) a report of this species from *Clupea harengus* from Plymouth, England. We have not

been able to verify this report. Dollfus (1956) also noted that *O. bacillaris* was a frequently observed parasite of herring from Plymouth.

Family OPECOELIIDAE

Opecoeloides vitellous (Linton, 1900) von Wicklen, 1946

Syn: *Cymbcephalus vitellous* (Linton, 1900)
Distomum vitellous Linton, 1900

Location: [intestine]
Host: *Clupea harengus harengus*
Dist: NW Atl
Records: Linton 1901a, 1940
Remarks: Dollfus (1956) considered *Distomum vitellous* to be a "collective" name and noted that the identity of the material reported from *C. h. harengus* cannot be determined.

Podocotyle atomon (Rudolphi, 1802) Odhner, 1905

Location: digestive tract
Host: *Clupea harengus pallasi*
Dist: NW Pac
Record: Layman 1930

Podocotyle reflexa (Creplin, 1825) Odhner, 1905

Location: digestive tract
Host: *Clupea harengus pallasi*
Dist: NW Pac
Record: Skrjabina 1963

ORDER OPISTHORCHIIDA

Superfamily OPISTHORCHIOIDEA

Family HETEROPHYIDAE

Cryptocotyle lingua (Creplin, 1825) Fischöder, 1903 (metacercaria)

Location: encysted in skin
Host: *Clupea harengus harengus*
Dist: NW Atl
Records: Sindermann and Rosenfield 1954a, b; Sindermann and Scattergood 1954; Sindermann 1957a, 1963, 1965, 1966; Sindermann and Farrin 1962

Galactosomum phalacrocoracis Yamaguti, 1939 (metacercaria)

Location: encysted on pharynx
Host: *Clupea harengus pallasi*
Dist: NE Pac
Records: Arthur 1978; Arthur and Arai 1980a, b

Unidentified Digenea

Digenea gen. sp. metacercaria

Includes: Trematoda auct.
Location: encysted on skin, body wall
Hosts: *Clupea harengus harengus* (1)
C. harengus pallasi (2, 3)
Dist: Aral Sea, E Ber Sea
Records: 1. Osmanov 1971 (Aral Sea); 2. Arthur 1978 (E Ber Sea); 3. Arthur and Arai 1980a (E Ber Sea)

Digenea gen. spp.

Includes: Trematoda auct.
Location: stomach, intestine
Hosts: *Clupea harengus harengus* (1, 3, 4)
C. harengus pallasi (2, 5, 6)
Dist: NE Atl, Baltic Sea, NE Pac, NW Pac

Records: 1. Levander 1909 (Baltic Sea); 2. Layman 1930 (NW Pac); 3. Marshall et al. 1937 (NE Atl); 4. Popiel 1951 (Baltic Sea); 5. Barraclough 1967 (NE Pac); 6. Barraclough and Fulton (1967) (NE Pac)

Remarks: Hemiurid trematodes reported as "vermiculi" from the stomach of herring from Amsterdam by van Leeuwenhoek (1697) are listed under *Hemiuirus luehei*.

Class CESTOIDEA

SUBCLASS CESTODA

ORDER TRY PANORHYNCHA

Family LACISTORHYNCHIDAE

Grillotia erinaceus (van Beneden, 1858) Guiart, 1927 (plerocercoid)

Syn: *Rhynchobothrium imparispine* Linton, 1890

Location: encysted in wall of stomach

Host: *Clupea harengus harengus*

Dist: NW Atl

Records: Linton 1901a, 1924

Lacistorhynchus tenuis (van Beneden, 1858) Dollfus, 1929 (plerocercoid)

Syn: *Rhynchobothrium bulbifer* Linton, 1890

Lacistorhynchus sp. auct.

Location: encysted on viscera, in musculature, and free in intestine

Host: *Clupea harengus harengus*

Dist: NE Atl, NW Atl

Records: Linton 1924 (NW Atl); MacKenzie 1974a (NE Atl), b (NE Atl), 1975a (NE Atl), c (-), 1976 (NE Atl), 1978 (NE Atl), 1983 (NE Atl); MacKenzie and Johnston 1976 (NE Atl); MacKenzie and Gallacher 1981 (NE Atl)

Family TENTACULARIIDAE

Nybelinia surmenicola Okada in Dollfus, 1929 (plerocercoid)

Location: encysted in liver

Host: *Clupea harengus pallasi*

Dist: E Ber Sea

Records: Arthur 1978; Arthur and Arai 1980a

Unidentified Trypanorhyncha

Trypanorhyncha gen. spp. plerocercoid

Includes: *Rhynchobothrium* sp. of Linton, 1901

"Tetrahynch cestode" of Davey, 1972

Location: encysted on viscera, in liver, mesenteries, and musculature

Hosts: *Clupea harengus harengus* (1, 2, 3, 4, 5, 6, 7, 8)
C. harengus pallasi (9, 10, 11)

Dist: NE Atl, NW Atl, NE Pac

Records: 1. Linton 1901a (NW Atl), 2. b (NW Atl); 3. Sindermann 1957a (NW Atl), 4. 1959 (NW Atl), 5. 1961a (NW Atl), 6. 1963 (NW Atl), 7. 1965 (NW Atl); 8. Davey 1972a (NE Atl); 9. Arthur 1978 (NE Pac); 10. Arthur and Arai 1980a (NE Pac), 11. b (NE Pac)

ORDER PSEUDOPHYLLIDEA

Family AMPHICOTYLIDAE

Abothrium gadi van Beneden, 1870 (plerocercoid)

Location: not specified

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Record: Reimer 1970

Eubothrium crassum (Bloch, 1779) Nybelin, 1922

Syn: *Bothriotaenia proboscidea* (Batsch, 1786)

Location: intestine, pyloric caeca

Hosts: *Clupea harengus harengus* (1, 2, 3)

C. harengus pallasi (4, 5)

Dist: E Pal (White Sea), Baltic Sea, W Ber Sea

Records: 1. Schneider 1902a (Baltic Sea), 2. b (Baltic Sea);

3. Levander 1909 (Baltic Sea); 4. Shulman and

Shulman-Albova 1953 (White Sea); 5. Zhukov 1963

(W Ber Sea)

Remarks: Only immature *E. crassum* have been reported from *Clupea harengus*. *Eubothrium crassum* is typically a parasite of salmonid fishes. Records of *Bothriotaenia proboscidea* from *C. harengus* were listed as *Bothriocephalus* sp. by Jääskeläinen (1921).

Eubothrium spp.

Location: intestine, wall of intestine

Hosts: *Clupea harengus harengus* (3, 4, 5, 6)

C. harengus pallasi (1, 2)

Dist: E Pal (White Sea), Baltic Sea

Records: 1. Shulman 1956 (White Sea); 2. Polyansky and

Shulman 1956 (White Sea); 3. Petrushevsky 1957 (Baltic

Sea); 4. Gaevskaya 1977 (Baltic Sea), 5. 1979 (Baltic

Sea); 6. Gaevskaya and Shapiro 1981 (Baltic Sea)

Family BOTHRIOCEPHALIDAE

Bothriocephalus scorpii (O.F. Müller, 1776) Rudolphi, 1808

Location: pyloric caeca, intestine

Hosts: *Clupea harengus harengus* (1, 3, 4, 5)

C. harengus pallasi (2)

Dist: NW Atl, Baltic Sea, W Ber Sea

Records: 1. Linton 1941 (NW Atl); 2. Zhukov 1963 (W Ber

Sea); 3. Reimer 1970 (Baltic Sea); 4. Rokicki 1973

(Baltic Sea), 5. 1975 (Baltic Sea)

Bothriocephalus sp.

Location: not specified

Host: *Clupea harengus harengus*

Dist: Baltic Sea, NW Atl

Records: Linton 1941 (NW Atl);²⁰ Gaevskaya 1977 (Baltic Sea)

Family LIGULIDAE

Ligula intestinalis (Linnaeus, 1758) Gmelin, 1790 (plerocercoid)

Syn: *?Ligula digramma* Creplin, 1839

Location: body cavity

Host: *Clupea harengus*

Dist: not specified

Record: von Linstow 1889 (original ?)

Remarks: The authenticity of this report is suspect. *Ligula intestinalis* is typically a parasite of freshwater fishes. Dubinin (1966) lists *L. digramma* Creplin, 1839 as a synonym of *Digramma interrupta* (Rudolphi, 1810).

Unidentified Pseudophyllidea

Pseudophyllidea gen. spp.

Location: intestine, intestinal wall

Hosts: *Clupea harengus harengus* (4, 5)

C. harengus pallasi (1, 2, 3)

Dist: NW Atl, E Pal (White Sea), NW Pac

Records: 1. Shulman and Shulman-Albova 1953 (White Sea);

2. Shulman 1956 (White Sea); 3. Skrjabina 1963 (NW

Pac); 4. Sindermann 1963 (NW Atl), 5. 1965 (NW Atl)

²⁰Record of Linton (1941) was for an encysted plerocercoid.

ORDER TETRAPHYLLIDEA

Tetraphyllidea of Uncertain Position

Scolex pleuronectis O. F. Müller, 1788 (plerocercoid)

Syn: *Scolex polymorphus* Rudolphi, 1819

Phyllobothrium sp. auct.

Location: gallbladder, pyloric caeca, intestine

Hosts: *Clupea harengus harengus* (2, 3, 4, 7, 12, 13, 16)

C. harengus pallasi (5, 6, 8, 9, 10, 11, 14, 15, 17, 18, 19)

C. harengus (1)

Dist: NE Atl, NW Atl, Baltic Sea, E Pal (Barents Sea, White Sea), NE Pac, NW Pac, W Ber Sea

Records: 1. Olsson 1866–67 (unspecified locality);²¹

2. Linton 1901a (NW Atl), 3. 1924 (NW Atl); 4. Giard

1903 (NE Atl); 5. Layman 1930 (NW Pac); 6. Shulman

and Shulman-Albova 1953 (White Sea); 7. Polyansky

1955 (Barents Sea); 8. Polyansky and Shulman 1956

(White Sea); 9. Shulman 1956 (White Sea); 10. Zhukov

1960b (NW Pac), 11. 1963 (W Ber Sea); 12. Rosenthal

1966 (experimental — NE Atl);²² 13. 1967 (ex-

perimental — NE Atl); 14. Arai 1967a (NE Pac),

15. 1969 (NE Pac); 16. Reimer 1970 (Baltic Sea);

17. Ivanchenko and Grozdilova 1971 (experimental —

White Sea); 18. Arthur 1978 (NE Pac); 19. Arthur and

Arai 1980a (NE Pac)

Remarks: *Scolex pleuronectis* is a larval name under which plerocercoids of various tetraphyllidean species have been assigned.

ORDER PROTEOCEPHALIDEA

Family PROTEOCEPHALIDAE

Proteocephalus spp.

Location: intestine

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Getsevichyut 1958; Lisivnenko 1961; Gaevskaya 1977, 1979; Gaevskaya and Zhudova 1979

Remarks: *Proteocephalus* spp. are typically parasites of freshwater fishes.

Unidentified Cestoda

Cestoda gen. spp.

Includes: "Cestoscolex" of van Beneden, 1871

Scolex bothriostomplex Reimer, 1970

Location: intestine

Host: *Clupea harengus harengus*

Dist: NE Atl, Baltic Sea

Records: van Beneden 1871 (NE Atl); Marshall et al. 1937

(NE Atl); Hentschel 1950 (NE Atl); Popiel 1951 (Baltic Sea); Dollfus 1956 (NE Atl);²³ Anon. 1969 (NE Atl);

Reimer 1970 (Baltic Sea)

²¹Material reported by Olsson (1866–67) was from herring obtained at a fishmarket in Warburg [Varberg, Sweden] and was thus probably of Baltic origin.

²²The record of Rosenthal (1966) was a tentative parasite identification.

²³Dollfus (1956) noted that larval cestodes have been reported from herring from Plymouth, England. We have been unable to locate any previous report from this locality.

PHYLUM NEMATHELMINTHES²⁴

CLASS NEMATODA

SUBCLASS SECERNENTEA

ORDER ASCARIDIDA

Superfamily ASCARIDOIDEA

Family ANISAKIDAE

Anisakis simplex (Rudolphi, 1809) Dujardin, 1845 (larva)

Syn: *Anisakis marina* auct.

Acanthocheilus rotundatus auct.

Anacanthocheilus rotundatus auct.

Eustoma rotundata auct.

Pseudanisakis rotundata auct.

Includes: "haringnematoden" of Houwing, 1969

Location: body cavity, viscera, mesenteries, musculature

Hosts: *Clupea harengus harengus* (1, 2, 3, 7, 8, 9, 10, 11, 12,

14, 15, 16, 19, 20, 21, 22, 23, 24, 27, 28, 30, 31, 33, 34,

35, 36, 37, 38, 39, 40, 41)

C. harengus pallasi (6, 20, 34, 42, 43, 44, 45)

C. harengus (4, 5, 13, 17, 18, 25, 26, 29, 32)

Dist: NE Atl, NW Atl, Baltic Sea, NE Pac, NW Pac, E Ber Sea

Records: 1. Wülker 1930 (NE Atl);²⁵ 2. Kahl 1938 (NE

Atl);²⁵ 3. 1939 (NE Atl); 4. Lölicher-Müller 1955 (un-

specified locality);²⁶ 5. Chalupsky 1955 (unspecified

locality);²⁷ 6. Akhmerov 1955 (NW Pac); 7. Kreis 1958

(NE Atl); 8. van Thiel et al. 1960 (NE Atl); 9. Kuipers et

al. 1960 (NE Atl); 10. Kuipers et al. 1960 (NE Atl);

11. Roskam 1960 (NE Atl), 12. 1967 (NE Atl);

13. Kuipers et al. 1963 (unspecified locality);²⁸ 14. van

Thiel 1967 (NE Atl); 15. Polak 1967 (unspecified

locality);²⁸ 16. Mameren and Houwing 1968 (NE Atl);

17. Houwing 1969a (unspecified locality);²⁸ 18. b (un-

specified locality);²⁸ 19. Anon. 1970 (NE Atl);

20. Stankiewicz 1971 (NW Atl, unspecified locality);²⁹

21. Ruitenberg 1970 (unspecified locality),³⁰ 22. 1971

(unspecified locality);³⁰ 23. Ruitenberg et al. 1970 (un-

specified locality),³⁰ 24. 1971 (unspecified locality);³⁰

25. Priebe 1971 (unspecified locality); 26. van Banning

1971 (unspecified locality);²⁸ 27. 1972 (NE Atl);

28. Grabda 1973a (NE Atl, Baltic Sea), 29. b (un-

specified locality), 30. 1974a (Baltic Sea), 31. b (Baltic

²⁴Dollfus (1970) gave a detailed historical review of the nematodes reported from *Clupea harengus*.

²⁵The records of Wülker (1930) and Kahl (1938) of *Anacanthocheilus rotundatus* from *Clupea harengus* were referred to *Anisakis* by Dollfus (1970).

²⁶Nematodes reported as *Acanthocheilus rotundatus* by Lölicher-Müller (1955) were from herring of unspecified origin examined in Leipzig, East Germany.

²⁷Nematodes reported as *Pseudanisakis rotundata* larva by Chalupsky (1955) were from herring originating from Norway.

²⁸Nematodes reported as *Eustoma rotundatum* by Kuipers et al. (1963), as *Anisakis marina* by Polak (1967), Houwing (1969a) and van Banning (1971) and as "haringnematoden" by Houwing (1969b) probably originated from the northeastern Atlantic Ocean.

²⁹Pacific herring examined by Stankiewicz (1971) were noted to have originated from the USSR.

³⁰Records of Ruitenberg (1970, 1971) and Ruitenberg et al. (1970, 1971) were for herring obtained from the Institute for Fishery Products, IJmuiden, the Netherlands, and thus probably originated from the northeastern Atlantic Ocean.

Sea), 32. (unspecified locality), 33. 1975 (Baltic Sea), 34. 1976 (Baltic Sea, NE Atl, unspecified locality); 35. Grabda and Felinska 1975 (Baltic Sea); 36. Pippy and van Banning 1975 (NE Atl); 37. Pippy 1975 (NE Atl, NW Atl); 38. Beverley-Burton et al. 1977 (NE Atl, NW Atl); 39. Gaevskaya and Umnova 1977 (NW Atl); 40. Friess 1977 (Baltic Sea); 41. Beverley-Burton and Pippy 1977 (NE Atl, NW Atl); 42. Arthur 1978 (NE Pac, E Ber Sea); 43. Arthur and Arai 1979 (NE Pac), 44. 1980a (NE Pac, E Ber Sea), 45. b (NE Pac)

Remarks: Although reports of larval nematodes encysted on the viscera and in the body cavity of herring date from the 17th century (van Leeuwenhoek 1697), it is only recently that the true identity of the species most likely to be noted in herring, *Anisakis simplex*, has been resolved (Dollfus 1970; Pippy and van Banning 1975; Beverley-Burton et al. 1977). The brief and, by current standards, inadequate nature of descriptions of larval nematodes occurring in the older literature precludes their positive identification. These older reports are listed under Unidentified Anisakidae.

Anisakis sp. larva

Location: body cavity, viscera, mesenteries, musculature
Hosts: *Clupea harengus harengus* (1, 3, 7, 10, 11, 14, 20, 23, 30, 32, 33, 34, 40, 51, 52, 53, 54, 55, 56, 57, 58, 60, 61, 62, 63, 64, 65, 67, 68, 69, 70, 73, 74, 75, 76, 77, 80, 81, 90, 91, 92)
C. harengus pallasi (2, 4, 5, 6, 8, 9, 12, 13, 16, 18, 19, 21, 22, 24, 26, 27, 28, 29, 31, 35, 36, 37, 38, 39, 41, 42, 43, 44, 45, 65, 78, 79, 82, 83, 84, 85, 86, 87, 88, 89, 93)
C. harengus (15, 17, 26, 46, 48, 49, 50, 59, 66, 71, 72)
Dist: NE Atl, NW Atl, Baltic Sea, E Pal (Barents Sea, Greenland Sea, Norwegian Sea, White Sea), NE Pac, NW Pac
Records: 1. Heller 1949 (NW Atl); 2. Margolis 1952 (NE Pac); 3. Scott 1953 (NW Atl); 4. Gustafson 1953 (NE Pac); 5. Shulman and Shulman-Albova 1953 (White Sea); 6. Bishop and Margolis 1955 (NE Pac); 7. Polyansky 1955 (Barents Sea); 8. Polyansky and Shulman 1956 (White Sea); 9. Shulman 1956 (White Sea); 10. Sindermann 1957a (NW Atl); 11. 1963 (NW Atl); 12. Zhukov 1960b (NW Pac); 13. Strelkov 1960 (NW Pac); 14. Berland 1961 (Norwegian Sea), 15. 1962 (unspecified locality);³¹ 16. Skrjabina 1963 (NW Pac); 17. Kuipers 1964 (unspecified locality);³¹ 18. Otsuru et al. 1965 (NW Pac); 19. Otsuru et al. 1967 (NW Pac); 20. Roskam 1966 (NE Atl); 21. Kobayashi et al. 1966 (NW Pac); 22. Sano 1966 (NW Pac); 23. Vik 1966 (unspecified locality);³² 24. Oshima 1966 (unspecified locality);³³ 25. Nygard 1967 (unspecified locality);³¹ 26. Arai 1967a (NE Pac), 27. 1969 (NE Pac); 28. Oyanagi 1967 (NW Pac); 29. Okumura 1967 (NW Pac); 30. Polak 1966 (unspecified locality);³¹ 31. Yamaguchi et al. 1968 (NW Pac); 32. Khalil 1968a (NE Atl), 33. b (NE Atl), 34. 1969 (NE Atl); 35. Ichihara et al. 1968 (NW Pac); 36. Kato et al. 1968 (NW Pac); 37. Koga et al. 1968 (NW Pac); 38. Otsuru 1968a (NW Pac), 39. b (NW Pac); 40. Anon. 1969 (NE Atl); 41. Kagei 1969 (NW Pac); 42. Hagiwara et al. 1969 (NW Pac); 43. Shiraki 1969 (NW Pac), 44. 1974 (NW Pac); 45. Koyama et al. 1969 (NW Pac); 46. Ruitenberg

and Roskam 1969 (unspecified locality);³¹ 47. Young and Lowe 1969 (NE Atl); 48. Gibson 1970 (unspecified locality);³¹ 49. Andreassen 1970 (unspecified locality);³¹ 50. Andreassen and Jørring 1970 (unspecified locality);³¹ 51. Muravev 1970a (NE Atl), 52. b (Greenland Sea, Norwegian Sea); 53. Kagei et al. 1971 (NW Atl); 54. Hodder and Parsons 1971 (NW Atl); 55. Parsons and Hodder 1971a (NW Atl), 56. b (NW Atl), 57. 1974 (NW Atl); 58. Davey 1972a (NE Atl), 59. b (unspecified locality);³¹ 60. Reimer and Jessen 1972 (NE Atl); 61. Anon. 1972 (NE Atl); 62. Lubieniecki 1972 (Baltic Sea), 63. 1974 (NW Atl); 64. Rokicki 1972 (Baltic Sea), 65. 1973 (Baltic Sea, NE Atl, NW Pac); 66. Priebe et al. 1973 (unspecified locality); 67. Dornheim 1973 (NW Atl); 68. Sluiters 1973 (NE Atl), 69. 1974 (NE Atl); 70. Strzyzewska and Popiel 1974 (Baltic Sea); 71. Schulz 1974 (unspecified locality);³⁴ 72. Grabda 1974c (unspecified locality); 73. Smith 1974 (NE Atl); 74. Anon. 1974 (NE Atl); 75. Anon. 1975 (NE Atl); 76. Smith and Wootten 1975 (NE Atl); 77. MacKenzie 1975a (NE Atl); 78. Dailey 1975 (NE Pac); 79. Ono 1975 (unspecified locality);³³ 80. Möller 1975 (Baltic Sea), 81. 1978 (Baltic Sea); 82. Kulachkova 1976a (White Sea), 83. b (White Sea), 84. c (White Sea), 85. 1978 (White Sea), 86. 1980 (White Sea); 87. Hoskins et al. 1976 (NE Pac); 88. Hauck 1977 (NE Pac); 89. Hauck and May 1977 (NE Pac); 90. Meyers 1978 (NW Atl); 91. van Banning and Becker 1978 (NE Atl); 92. Jackson et al. 1978 (NW Atl); 93. Pushnikova and Pushnikov 1981 (NW Pac)

Remarks: Many of the above records were given as *Anisakis* sp. Type 1 larva, a form now recognized as *A. simplex* (see Pippy and van Banning 1975; Beverley-Burton et al. 1977). Species of *Anisakis* other than *A. simplex* have not been shown to occur in herring, although the distributions of *A. typica* (Diesing, 1860) and *A. physeteris* Baylis, 1923 in their definitive hosts overlap that of *C. harengus* (Davey 1971).

Contracaecum osculatum (Rudolphi, 1802) Baylis, 1920 (larva)

Location: liver
Host: *Clupea harengus harengus*

Dist: Baltic Sea
Records: Fagerholm 1978, 1982

Remarks: Fagerholm (1982) notes that larval *C. osculatum* have frequently been confused with *Hysterothylacium aduncum*. Records of *H. aduncum* from the liver of Baltic herring are probably based on misidentification of this species.

Contracaecum rudolphii Hartwich, 1964 (larva)

Location: not specified
Host: *Clupea harengus harengus*

Dist: NE Atl and/or Baltic Sea

Record: Hartwich 1964

Remarks: Based on studies by previous authors Hartwich (1964) considered *C. rudolphii* to occur in Atlantic herring. No original reports have been made under this name.

Contracaecum siluriglandis (von Linstow, 1883) Dogiel and Bykhovsky, 1934 (larva)

Location: wall of intestine
Host: *Clupea harengus harengus*

Dist: Aral Sea

³¹*Anisakis* sp. reported by the indicated authors were probably from herring originating from the northeastern Atlantic Ocean.

³²Locality was given only as Scando-Atlantic by Vik (1966).

³³Records of Oshima (1966) and Ono (1975) are probably from the northwestern Pacific Ocean.

³⁴*Anisakis* sp. reported by Schulz (1974) was from herring purchased from a fish vendor in Germany.

Record: Osmanov 1971

Remarks: Bykhovskaya-Pavlovskaya et al. (1962) note that this species is the larval form of *C. spiculigerum* [= *C. rudolphii*].

Contracaecum spp. larva

Includes: *Contracaecum spiculigerum* of Walton, 1928³⁵

Nematoda gen. sp. of Rosenthal, 1966³⁶

Location: viscera, body cavity, mesenteries, intestine

Hosts: *Clupea harengus harengus* (1, 5, 9, 10, 11, 16, 17, 20, 22, 23)

C. harengus pallasi (2, 3, 4, 7, 8, 12, 13, 14, 15, 18, 19, 21, 24, 25, 26, 27, 28, 29)

C. harengus (6)

Dist: NE Atl, NW Atl, E Pal (Greenland Sea, Norwegian Sea, White Sea), NE Pac, NW Pac, E Ber Sea, W Ber Sea

Records: 1. Walton 1928 (NW Atl); 2. Smedley 1934 (NE Pac);³⁷ 3. Margolis 1952 (NE Pac); 4. Gustafson 1953 (NE Pac); 5. van Thiel et al. 1960 (NE Atl); 6. Berland 1962 (unspecified locality);³⁸ 7. Zhukov 1963 (W Ber Sea);³⁹ 8. Kobayashi et al. 1966 (NW Pac); 9. Rosenthal 1966 (experimental — NE Atl), 10. 1967 (experimental — NE Atl); 11. Roskam 1967 (NE Atl); 12. Arai 1967a (NE Pac), 13. 1969 (NE Pac); 14. Koga et al. 1968 (NW Pac); 15. Kato et al. 1968 (NW Pac); 16. Khalil 1968a (NE Atl), 17. 1969 (NE Atl); 18. Hagiwara et al. 1969 (NW Pac); 19. Koyama et al. 1969 (NW Pac); 20. Anon. 1969 (NE Atl); 21. Kikuchi et al. 1970 (NW Pac); 22. Muravev 1970b (Greenland Sea, Norwegian Sea); 23. Reimer and Jessen 1972 (NE Atl); 24. Shiraki 1974 (NW Pac); 25. Dailey 1975 (NE Pac); 26. Arthur 1978 (NE Pac, E Ber Sea); 27. Arthur and Arai 1980a (NE Pac, E Ber Sea), 28. b (NE Pac); 29. Kulachkova 1980 (White Sea)

Remarks: Most of the records of *Contracaecum* sp. from *Clupea harengus* are probably referable to *Hysterothylacium aduncum* (Rudolphi, 1802).

Hysterothylacium aduncum (Rudolphi, 1802) Deardorff and Overstreet, 1981 (adult and larva)

Syn: *Contracaecum aduncum* (Rudolphi, 1802)

Contracaecum clavatum (Rudolphi, 1809)

Contracaecum hypomesi (Fujita, 1932)

Thynnascaris adunca (Rudolphi, 1802)

Location: stomach, intestine, mesenteries, viscera, body cavity

Hosts: *Clupea harengus harengus* (2, 3, 5, 10, 11, 14, 15, 16, 17, 18, 19, 21, 22, 26)

C. harengus pallasi (1, 4, 6, 8, 9, 12, 13, 20, 23, 24,

25)

C. harengus (7)

³⁵The report of Walton (1928) was based on examination of material from the Leidy collection which was labeled *Agamoneema capsularia*. Hartwich (1964) considered *Ascaris spiculigera* Rudolphi, 1809 a synonym of *Contracaecum microcephalum* (Rudolphi, 1809). However, many reports of *C. spiculigerum* were referred to *C. rudolphii*. Dollfus (1970) notes that reexamination of Leidy's specimens is necessary to determine which species is involved.

³⁶Unidentified nematodes reported from Atlantic herring by Rosenthal (1966) were later (Rosenthal 1967) assigned to *Contracaecum* sp.

³⁷Material reported as *Contracaecum* sp. by Smedley (1934) from Pacific herring apparently included specimens of *Anisakis*.

³⁸*Contracaecum* sp. reported by Berland (1962) were from salt herring which probably originated from the northeastern Atlantic Ocean.

³⁹Material listed as *Contracaecum* sp. from *Clupea harengus pallasi* in the Host-Parasite list of Zhukov (1963) was apparently reported as *Contracaecum aduncum* (syn. of *Hysterothylacium aduncum*) in his text.

Dist: NE Atl, NW Atl, Baltic Sea, E Pal (Barents Sea, Norwegian Sea, White Sea), NE Pac, NW Pac, E Ber Sea, W Ber Sea

Records: 1. Fujita 1932 (NW Pac); 2. Markowski 1933 (Baltic Sea); 3. Punt 1942 (NE Atl); 4. Shulman and Shulman-Albova 1953 (White Sea); 5. Polyansky 1955 (Barents Sea); 6. Akhmerov 1955 (NW Pac); 7. Löliger-Müller 1955 (unspecified locality);⁴⁰ 8. Shulman 1956 (White Sea); 9. Polyansky and Shulman 1956 (White Sea); 10. Petrushevsky 1957 (Baltic Sea); 11. Berland 1961 (Norwegian Sea); 12. Zhukov 1963 (W Ber Sea); 13. Popova and Valter 1965 (White Sea); 14. Muravev 1970a (NE Atl); 15. Rokicki 1973 (Baltic Sea), 16. 1975 (Baltic Sea); 17. Hartwich 1975 (NE Atl, Baltic Sea); 18. Gaevskaya 1977 (Baltic Sea); 19. Gaevskaya and Umnova 1977 (NW Atl); 20. Arthur 1978 (NE Pac, E Ber Sea); 21. Sjöblom and Kuittinen 1978a (Baltic Sea), 22. b (Baltic Sea); 23. Arthur and Arai 1979 (NE Pac), 24. 1980a (NE Pac, E Ber Sea), 25. b (NE Pac); 26. Fagerholm 1982 (Baltic Sea)

Remarks: Sheenko and Pozdnyakov (1981) reviewed the species of *Contracaecum* described by Fujita (1932), considering *C. hypomesi* a synonym of *H. aduncum*.

Phocascaris sp. larva

Location: not specified

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Sjöblom and Kuittinen 1976, 1978a, b

Remarks: Fagerholm (1982) notes that the above records are probably based on misidentification of *Contracaecum osculatum*.

Pseudoterranova decipiens (Krabbe, 1878) Gibson and Colin, 1982 (larva)

Syn: *Phocanema decipiens* (Krabbe, 1878)

Porrocaecum decipiens (Krabbe, 1878)

Terranova decipiens (Krabbe, 1878)

Location: body cavity, liver [musculature]

Hosts: *Clupea harengus harengus* (1)

C. harengus pallasi (2, 4, 5)

C. harengus (3)

Dist: NW Atl, E Pal (White Sea)

Records: 1. Walton 1928 (NW Atl);⁴¹ 2. Shulman and Shulman-Albova 1953 (White Sea); 3. Löliger-Müller 1955 (unspecified locality);⁴² 4. Polyansky and Shulman 1956 (White Sea); 5. Shulman 1956 (White Sea)

Pseudoterranova sp. larva

Syn: *Phocanema* sp. auctorum

Porrocaecum sp. auctorum

Terranova sp. auctorum

Location: musculature

Host: *Clupea harengus pallasi*

Dist: NE Pac, NW Pac, E Pal (White Sea)

Records: Strelkov 1960 (NW Pac); Dailey 1975 (NE Pac); Kulachkova 1980 (White Sea)

⁴⁰The report of Löliger-Müller (1955) of *Contracaecum clavatum* (syn. of *Hysterothylacium aduncum*) was from herring of unspecified origin examined in Leipzig, East Germany.

⁴¹Material reported from Atlantic herring as *Porrocaecum decipiens* (syn. of *Pseudoterranova* d.) from the Leidy collection by Walton (1928) was labeled *Agamoneema capsularia*.

⁴²Nematodes reported as *Porrocaecum decipiens* by Löliger-Müller (1955) were from herring of unspecified origin examined in Leipzig, East Germany.

Raphidascaris acus (Bloch, 1779) Ralliet and Henry, 1915
 Syn: *Ascaris acus* Bloch, 1779
 Location: intestine, viscera
 Host: *Clupea harengus harengus*
 Dist: NE Atl, NW Atl
 Records: Bellingham 1844 (NE Atl); Jackson et al. 1978 (NW Atl)
 Remarks: Dollfus (1956, 1970) questioned the validity of Bellingham's (1844) report of this species. The tentative identification of larval *R. acus* from the viscera of herring by Jackson et al. (1978) also requires verification. *Raphidascaris acus* is typically a parasite of freshwater fishes.

Raphidascaris sp. larva
 Location: not specified
 Host: *Clupea harengus pallasi*
 Dist: NW Pac
 Record: Kato et al. 1968

Thynnascaris sp. [larva?]
 Location: viscera and/or musculature
 Host: *Clupea harengus harengus*
 Dist: NW Atl
 Record: Jackson et al. 1978
 Remarks: The above report is probably referable to *Hysterorthyelium*.

Anisakidae gen. spp. larva
 Includes: Anisakinae auct.
Ascaris capsularia (Rudolphi, 1802)
Ascaris clupeae van Beneden, 1871
Ascaris clupearum Fabricius, 1794
Ascaris gracilescens Rudolphi, 1819
Ascaris sp. of Linton, 1901
Agamoneema capsularia (Rudolphi, 1802)
Capsularia halecis (Gmelin, 1790)
Cucullanus halecis Fabricius, 1794
Filaria capsularia Rudolphi, 1802
Filaria piscium (Rudolphi, 1809)
Filocapsularia communis Deslongchamps, 1824
Gordius harengum Bloch, 1782
Gordius marinus Linnaeus, 1767
Gordius sp. auct.
 Unidentified ascarids auct.
 Location: body cavity, mesenteries, viscera, intestine
 Hosts: *Clupea harengus harengus* (1, 13, 17, 20, 22, 24, 25, 26, 30, 31, 32, 33, 34, 37)
C. harengus pallasi (35, 36, 38)
C. harengus (2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 14, 15, 16, 18, 19, 21, 23, 27, 28, 29)
 Dist: NE Atl, NW Atl, Baltic Sea, E Pal (Norwegian Sea), NW Pac
 Records: 1. Linnaeus 1767 (NE Atl), 2. 1789 (unspecified locality);⁴³ 3. Guettard 1768 (unspecified locality);⁴⁴ 4. Beckmann 1769 (unspecified locality);⁴⁵ 5. 1770 (unspecified locality); 6. Walch 1778 (unspecified

locality);⁴⁶ 7. Bloch 1782 (unspecified locality);⁴⁷ 8. Fabricius 1794 (unspecified locality);⁴⁸ 9. Zeder 1800 (unspecified locality);⁴⁹ 10. Nöel 1807 (unspecified locality);⁵⁰ 11. Rudolphi 1802 (unspecified locality), 12. 1809 (unspecified locality), 13. 1819 (Baltic Sea);⁵¹ 14. Deslongchamps 1824 (unspecified locality);⁴⁹ 15. Creplin 1839 (unspecified locality);⁵¹ 16. 1846 (unspecified locality); 17. Bellingham 1844 (NE Atl); 18. Krøyer 1846–49 (unspecified locality) (original ?); 19. Cuvier and Valenciennes 1847 (unspecified locality) (original ?); 20. Leidy 1857 (NW Atl); 21. Cobbold 1866 (unspecified locality);⁵² 22. van Beneden 1871 (NE Atl); 23. Mühlung 1898 (unspecified locality);⁵³ 24. Linton 1901a (NW Atl); 25. Giard 1903 (NE Atl);⁵⁴ 26. Nicoll 1907 (NE Atl); 27. Baylis 1916 (unspecified locality);⁵² 28. Martin 1921 (unspecified locality);⁵⁵ 29. Jahnel 1940 (unspecified locality);⁵⁶ 30. Rees 1953 (Norwegian Sea);⁵⁷ 31. Sindermann 1957a (NW Atl), 32. 1959 (NW Atl), 33. 1961a (NW Atl), 34. 1965 (NW Atl); 35. Otsuru et al. 1967 (NW Pac); 36. Yoshimura 1966 (NW Pac); 37. Jackson et al. 1978 (NW Atl); 38. Myers 1979 (NE Pac)

Remarks: Walton (1928) identified as *Porrocaecum decipiens* (syn. of *Pseudoterranova*) and *Contracaecum spiculigerum* (syn. of *C. rudolphi*) material from the collection of Joseph Leidy which was labeled *Agamoneema capsularia*. It is uncertain whether these specimens were reported as "Agamoneema capsularia Diesing?" by Leidy (1857).

⁴⁶Records of Walch (1778) of *Gordius* and of Zeder (1800) of *Capsularia halecis* were probably from herring originating in German waters.

⁴⁷Bloch (1782) reported *Gordius harengum* from the testis of herring examined in Germany.

⁴⁸Fabricius (1794) reported the presence of *Cucullanus halecis* and *Ascaris clupearum* on the pyloric caeca and intestine of herring examined in Copenhagen, Denmark.

⁴⁹Records of Nöel (1807) of *Gordius marinus* and of Deslongchamps (1824) for *Filocapsularia communis* were probably from herring originating in French waters.

⁵⁰Rudolphi (1819) reported *Filaria capsularia* from herring from Gryphaiae. Dollfus (1970) gave the locality for this record as Griefswald, East Germany, which is situated on the Baltic Sea.

⁵¹The report of *Ascaris gracilescens* by Creplin (1839) was from herring considered by Dollfus (1970) to have been examined at Griefswald, East Germany, and thus probably of Baltic origin.

⁵²The specimens reported as *Filaria piscium* from the abdomen of *Clupea harengus* by Cobbold (1866) were contained in the Museum of the Royal College of Surgeons of England; those reported as *Ascaris capsularia* by Baylis (1916) were contained in the British Museum (Natural History). Both reports probably pertain to herring originating from the northeastern Atlantic Ocean.

⁵³The specimens reported as *Agamoneema capsularia* from *Clupea harengus* by Mühlung (1898) were contained in the Museum of the University of Königsberg and were thus probably from herring originating from the Baltic Sea.

⁵⁴Dollfus (1970) notes that *Ascaris clupeae* of Giard, 1903 was probably *Contracaecum aduncum* (syn. of *Hysterorthyelium a.*).

⁵⁵The specimens reported as ascarid larvae by Martin (1921) were from herring received from a coworker in Bergen, Norway and were thus probably from the northeastern Atlantic Ocean. Dollfus (1970) considered this report probably to be referable to *Anisakis simplex*.

⁵⁶The specimens reported as *Ascaris capsularia* by Jahnel (1940) were from herring of unknown origin examined in Vienna, Austria. Khalil (1969) considered this report probably to be referable to *Anisakis* sp. type I larvae (= *A. simplex*).

⁵⁷Specimens reported by Rees (1953) were considered to be either *Anisakis* or *Porrocaecum* [= *Pseudoterranova*] larvae.

⁴³Linnaeus (1789) reported the presence of *Gordius marinus* in *Clupea harengus* from Sweden.

⁴⁴Guettard (1768) reported the occurrence of unidentified ascarid nematodes in the testis of herring from Paris, France.

⁴⁵Beckmann (1769) reported the occurrence of *Gordius marinus* in herring from an unspecified locality and later (Beckmann 1770), probably in reference to his earlier record, noted the presence of *Gordius* in the milt of Dutch herring.

ORDER SPIRURIDA

Superfamily APROCTOIDEA

Family DESMIDOCERCIDAE

Desmidocerella numidica (Seurat, 1920) Yorke and Mapstone, 1926 (larva)

Syn: *Desmidocerca numidica* Seurat, 1920

Site: eye

Host: *Clupea harengus*

Dist: unspecified [Baltic Sea ?]

Record: Bezubik 1956

Superfamily HABRONEMATOIDEA

Family CYSTIDICOLIDAE

Ascarophis pacificus Zhukov, 1960

Location: [stomach]

Host: *Clupea harengus pallasi*

Dist: NW Pac

Record: Strelkov 1960

Cystidicola farionis Fischer, 1798

Location: swim bladder

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Arro 1964, Fagerholm 1982

Remarks: *Cystidicola farionis* is typically a freshwater parasite of salmonid fishes.

Spirurida of Undetermined Position

Spirurida gen. spp. larva

Location: stomach wall, intestine ?

Host: *Clupea harengus pallasi*

Dist: NE Pac

Records: Arthur 1978; Arthur and Arai 1980a

Remarks: Larval nematodes reported as Spirurida gen. sp. larvae type I by Arthur and Arai (1980a) have since been noted to be identical with Type X spirurid larvae of Hasegawa (1978).

ORDER OXYURIDA

Superfamily OXYUROIDEA

Oxyuroidea gen. sp. larva

Location: gill wash

Host: *Clupea harengus pallasi*

Dist: NE Pac

Records: Arthur 1978; Arthur and Arai 1980a

Nematoda of Undetermined Species

Nematoda gen. spp.

Includes: "vermiculi" of van Leeuwenhoek, 1697 (partim)
"Gordii" of Martin, 1760

Location: body cavity, viscera, mesenteries, stomach

Hosts: *Clupea harengus harengus* (3, 4, 7, 8, 9, 10, 11)
C. harengus pallasi (5, 6, 7)
C. harengus (1, 2)

Dist: NE Atl, NW Atl, Baltic Sea, NE Pac, NW Pac

Records: 1. van Leeuwenhoek 1697 (unspecified locality);⁵⁸
2. Martin 1760 (unspecified locality); 3. Linton 1901b

⁵⁸Nematodes reported by van Leeuwenhoek (1697) were from herring purchased in Amsterdam.

(NW Atl); 4. Manter 1926 (NW Atl); 5. Layman 1930 (NW Pac); 6. Robinson et al. 1968b (NE Pac); 7. Priebe 1971 (unspecified locality); 8. Molloy 1970 (NE Atl);⁵⁹
9. Lubieniecki 1972 (Baltic Sea); 10. Dornheim 1973 (NW Atl);⁵⁹ 11. Schultz 1974 (NE Atl)⁵⁹

PHYLUM ACANTHOCEPHALA

ORDER EOACANTHOCEPHALA

Family NEOECHINORHYNCHIDAE

Neoechinorhynchus rutili (O. F. Müller, 1780) Hamann, 1892

Location: [intestine]

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Gaevskaya 1977, 1979

Remarks: *Neoechinorhynchus rutili* is typically a parasite of freshwater fishes.

ORDER PALEACANTHOCEPHALA

Superfamily ECHINORHYNCHOIDEA

Family ECHINORHYNCHIDAE

Acanthocephalus clavula (Dujardin, 1845) Grabda-Kazubska and Chubb, 1968

Syn: *Echinorhynchus clavula* Dujardin, 1845

Location: intestine

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Record: Forssell 1905

Remarks: Grabda-Kazubska and Chubb (1968) indicate that the occurrence of *A. clavula* in the Baltic area has not been substantiated. Many reports under the name *Echinorhynchus clavula* from this area were referred to *E. borealis* von Linstow, 1901 by Grabda-Kazubska and Ejsymont (1969). Since Forssell's (1905) report of *E. clavula* from *Clupea harengus* was given without description or accompanying figures the identity of this material remains equivocal. *Acanthocephalus clavula* is typically a parasite of freshwater fishes.

Acanthocephalus lucii (O. F. Müller, 1776) Lühe, 1911

Location: intestine

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Record: Arro 1964

Remarks: Golvan (1969) indicated the occurrence of *A. lucii* in *Clupea harengus* to be questionable. *Acanthocephalus lucii* is typically a parasite of freshwater fishes.

Echinorhynchus gadi Zoega in O. F. Müller, 1776

Syn: *Echinorhynchus acus* Rudolphi, 1802

Echinorhynchus angustatus f. *Laborum* Olsson, 1893⁶⁰

Location: intestine

Hosts: *Clupea harengus harengus* (1, 2, 3, 4, 5, 6, 7, 8, 9, 12, 14, 15, 16)

C. harengus pallasi (10, 11, 13, 17, 18)

⁵⁹Nematodes reported by Molloy (1970) and Schultz (1974) from Atlantic herring were assumed to be *Anisakis* and/or *Contracaecum*. Those reported by Dornheim (1973) were noted to include representatives of *Anisakis*.

⁶⁰Lundström (1942) redetermined *E. angustatus* forma *Laborum* Olsson, 1893 to be *E. gadi*.

Dist: NE Atl, Baltic Sea, E Pal (White Sea), NE Pac, NW Pac
 Records: 1. Olsson 1893 (Baltic Sea); 2. Schneider 1902b (Baltic Sea); 3. Giard 1903 (NE Atl); 4. Forssell 1905 (Baltic Sea); 5. Levander 1909 (Baltic Sea); 6. Jääskelainen 1921 (Baltic Sea);⁶¹ 7. Markowski 1933 (Baltic Sea); 8. Lundström 1942 (Baltic Sea); 9. Popiel 1951 (Baltic Sea); 10. Shulman and Shulman-Albova 1953 (White Sea); 11. Shulman 1956 (White Sea); 12. Petrushevsky 1957 (Baltic Sea); 13. Strelkov 1960 (NW Pac); 14. Rokicki 1973 (Baltic Sea), 15. 1975 (Baltic Sea); 16. Gaevskaya 1977 (Baltic Sea); 17. Arthur 1978 (NE Pac); 18. Arthur and Arai 1980a (NE Pac)

Echinorhynchus salmonis O. F. Müller, 1784

Syn: *Echinorhynchus phoneix* Schneider, 1903

Location: intestine

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Schneider 1903; Nybelin 1924; Petrushevsky 1957

Remarks: *Echinorhynchus salmonis* is typically a parasite of freshwater fishes.

Family POMPHORHYNCHIDAE

Pomphorhynchus kostylewi Petrochenko, 1956

Location: intestine

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Rokicki 1973, 1975⁶²

Remarks: *Pomphorhynchus kostylewi* is typically a parasite of freshwater fishes.

Pomphorhynchus laevis (Zoega in O. F. Müller, 1776) Van Cleave, 1924

Location: intestine, body cavity [?]

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Lundström 1942; Petrushevsky 1957; Petrushevsky and Petrushevskaya 1960; Gaevskaya 1979 (original ?)

Remarks: *Pomphorhynchus laevis* is typically a parasite of freshwater fishes.

Family RHADINORHYNCHIDAE

Rhadinorhynchus trachuri Harada, 1935

Location: intestine

Host: *Clupea harengus pallasi*

Dist: NE Pac

Records: Arthur 1978; Arthur and Arai 1980a

Serrasentis socialis (Leidy, 1851) Van Cleave, 1924 (juvenile)

Location: abdomen

Host: *Clupea harengus harengus*

Dist: NW Atl

Record: Van Cleave 1924

Remarks: Van Cleave (1924) identified this species from material from the Leidy collection provided by A. Walton which was contained in a vial labeled *Agamoneema capsularia*. Leidy did not report *S. socialis* from *Clupea harengus*.

Superfamily POLYMPHOIDEA

Family POLYMPHIDAE

Corynosoma semerme (Forssell, 1904) Lühe, 1911 (juvenile)

Syn: *Echinorhynchus semermis* Forssell, 1904

Echinorhynchus sp. of Levander, 1909⁶³

Echinorhynchus strumosum of Schneider, 1902 (partim)

Location: body cavity, mesenteries, intestine

Hosts: *Clupea harengus harengus* (1, 2, 3, 5, 9, 10, 11, 12,

13, 14, 15, 16, 17)

C. harengus pallasi (6, 7, 8)

C. harengus (4)

Dist: Baltic Sea, E Pal (White Sea)

Records: 1. Schneider 1902b (Baltic Sea); 2. Forssell 1905

(Baltic Sea); 3. Levander 1909 (Baltic Sea); 4. Meyer

1931 (unspecified locality) (original?); 5. Lundström

1942 (Baltic Sea); 6. Shulman and Shulman-Albova

1953 (White Sea); 7. Shulman 1956 (White Sea);

8. Polyansky and Shulman 1956 (White Sea); 9. Arro

1964 (Baltic Sea), 10. 1967 (Baltic Sea); 11. Dubnitsky

1968 (Baltic Sea); 12. Rokicki 1973 (Baltic Sea),

13. 1975 (Baltic Sea); 14. Gaevskaya 1977 (Baltic Sea),

15. 1979 (Baltic Sea); 16. Valtonen 1979 (Baltic Sea),

17. 1980 (Baltic Sea)

Corynosoma strumosum (Rudolphi, 1802) Lühe, 1904 (juvenile)

Syn: *Echinorhynchus strumosus* Rudolphi, 1802

Echinorhynchus gibber Olsson, 1893⁶⁴

Location: mesenteries, body cavity, intestine

Hosts: *Clupea harengus harengus* (1, 2, 3, 4, 8, 11, 13, 14)

C. harengus pallasi (5, 6, 7, 9, 10, 12, 15, 16)

Dist: Baltic Sea, E Pal (White Sea), NE Pac, NW Pac, E Ber

Sea

Records: 1. Olsson 1893 (Baltic Sea); 2. Forssell 1905 (Baltic

Sea);⁶⁵ 3. Markowski 1933 (Baltic Sea); 4. Lundström

1942 (Baltic Sea); 5. Shulman and Shulman-Albova

1953 (White Sea); 6. Shulman 1956 (White Sea);

7. Polyansky and Shulman 1956 (White Sea);

8. Petrushevsky 1957 (Baltic Sea); 9. Strelkov 1960

(NW Pac); 10. Skrabina 1963 (NW Pac); 11. Dubnitsky

1968 (Baltic Sea); 12. Arthur 1978 (NE Pac, E Ber Sea);

13. Valtonen 1979 (Baltic Sea), 14. 1980 (Baltic Sea);

15. Arthur and Arai 1980a (NE Pac, E Ber Sea), 16. b

(NE Pac)

Remarks: Material reported as *E. strumosus* from *C. harengus* by Schneider (1902b) and listed as *Corynosoma strumosum* by Jääskelainen (1921) is referred to *Corynosoma semerme* following Dollfus (1956).

Corynosoma villosum Van Cleave, 1953 (juvenile)

Location: mesenteries

Host: *Clupea harengus pallasi*

Dist: NE Pac

Records: Arthur 1978; Arthur and Arai 1980a

⁶¹The record of Jääskelainen (1921) appears only in the host-parasite listing.

⁶²Record of Rokicki (1975) occurs only in his Table 22.

⁶³The report of *Echinorhynchus* sp. from *C. harengus* by Levander (1909) was referred to *Corynosoma semerme* by Jääskelainen (1921).

⁶⁴Synonymy following Lundström (1942). Other workers (Lühe 1911; Yamaguti 1963; Golvan 1969) have referred *E. gibber* in part to *Corynosoma semerme* and in part to *C. strumosum*.

⁶⁵Record of Forssell (1905) involves a tentative parasite identification.

Acanthocephala of Undetermined Position

Acanthocephala gen. sp.

Location: intestine

Host: *Clupea harengus pallasi*

Dist: NW Pac

Record: Layman 1930

PHYLUM ARTHROPODA

CLASS CRUSTACEA

SUBCLASS BRANCHIURA

ORDER ARGULOIDEA

Family ARGULIDAE

Argulus alosae Gould, 1841

Location: body surface

Host: *Clupea harengus harengus*

Dist: NW Atl

Record: Sindermann 1957a

Argulus coregoni Thorell, 1864

Location: [body surface]

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Records: Petrushevsky 1957; Gaevskaya 1979 (original ?)

Remarks: *Argulus coregoni* is typically a parasite of freshwater fishes.

SUBCLASS ENTOMOSTRACA

ORDER COPEPODA

SUBORDER POECILOSTOMATOIDA

Family BOMOLOCHIDAE

Bomolochus cuneatus Fraser, 1920

Syn: *Parabomolochus cuneatus* (Fraser, 1920)

Location: inner surface of operculum, gills

Host: *Clupea harengus pallasi*

Dist: NE Pac, NW Pac

Records: Fraser 1920 (NE Pac); Markevich 1956 (NW Pac);

Vervoort 1964 (NE Pac); Arai 1967a (NE Pac), 1969 (NE Pac); Arthur 1978 (NE Pac); Arthur and Arai 1980a (NE Pac)

Family ERGASILIDAE

Ergasilus sieboldi Nordmann, 1832

Location: gills

Host: *Clupea harengus harengus*

Dist: Baltic Sea

Record: Giesbrecht 1882⁶⁶

Remarks: *Ergasilus sieboldi* is typically a parasite of freshwater fishes.

SUBORDER SIPHONOSTOMATOIDA

Family CALIGIDAE

Caligus clemensi Parker and Margolis, 1964

Syn: *Caligus gurnardi* of Fraser, 1920 and of Bere, 1930

⁶⁶The record of *E. sieboldi* from "herring" by Giesbrecht (1882) appears in the abstract only.

Location: body surface

Host: *Clupea harengus pallasi*

Dist: NE Pac, E Ber Sea

Records: Fraser 1920 (NE Pac); Bere 1930b (NE Pac); Parker and Margolis 1964 (NE Pac); Arai 1967a (NE Pac), 1969 (NE Pac); Arthur 1978 (NE Pac, E Ber Sea); Arthur and Arai 1980a (NE Pac, E Ber Sea), b (NE Pac)

Caligus elongatus Nordmann, 1832

Syn: *Caligus rapax* auct.

Location: body surface, buccal cavity

Host: *Clupea harengus harengus*

Dist: NE Atl, NW Atl

Records: Bere 1930a (NW Atl); Dollfus 1956 (NE Atl); Sindermann 1957a (NW Atl); Rosenthal 1966 (experimental — NE Atl), 1967 (experimental — NE Atl)

Lepeophtheirus pollachius Bassett-Smith, 1896

Location: buccal cavity

Host: *Clupea harengus harengus*

Dist: NE Atl

Record: Dollfus 1956

Remarks: Dollfus (1956) considered this material to represent a separate form which he named *L. pollachius* form *harengi nobis*.

Family PENNELLIDAE

Lernaeenicus sprattae (Sowerby, 1806) Olsson, 1869

Syn: *Lernaeonema monillaris* Milne Edwards, 1840

Location: attached to eye

Host: *Clupea harengus harengus*

Dist: NE Atl

Record: Bassett-Smith 1896

Remarks: Z. Kabata (Pacific Biological Station, Nanaimo, B.C., personal communication) considers *Clupea harengus* to be an atypical host for *L. sprattae*.

Lernaeocera sp.

Location: body surface

Host: *Clupea harengus harengus*

Dist: NE Atl

Records: Rosenthal 1966 (experimental),⁶⁷ 1967 (experimental)

Remarks: Z. Kabata (personal communication) considers *Clupea harengus* to be an atypical host for this species.

PHYLUM CHORDATA

SUBPHYLUM VERTEBRATA

CLASS AGNATHA

ORDER PETROMYZONTIFORMES

Family PETROMYZONTIDAE

Lampetra ayresi (Günther, 1870) Jordan, Evermann, and Clark, 1930

Location: body surface

Host: *Clupea harengus pallasi*

Dist: NE Pac

Records: Roos et al. 1973; Beamish 1980

⁶⁷The initial report of this parasite was given by Rosenthal (1966) as a "...copepodite stage of a copepod (presumably a caligid)."

HOST-PARASITE CHECKLIST

<i>Clupea harengus harengus</i> Linnaeus	Atlantic herring	<i>Scolex pleuronectis</i> plerocercoid (NE Atl, NW Atl, Baltic Sea, Barents Sea)
Syn: <i>Clupea elongata</i> Lesueur		<i>Trypanorhyncha</i> gen. sp. plerocercoid (NE Atl, NW Atl)
<i>Clupea harengus membras</i> Linnaeus		
Includes: <i>Clypea halecis</i> auct.		
<i>Clupea harengus</i> auct.		
Apicomplexa		Nematoda
<i>Eimeria clupearum</i> (NE Atl, NW Atl)		Anisakidae gen. spp. larva (NE Atl, NW Atl, Baltic Sea, Norwegian Sea)
<i>E. sardinae</i> (NE Atl, NW Atl, Baltic Sea, Barents Sea)		<i>Anisakis simplex</i> larva (NE Atl, NW Atl, Baltic Sea)
Microspora		<i>Anisakis</i> sp. larva (NE Atl, NW Atl, Baltic Sea, Barents Sea, Greenland Sea, Norwegian Sea)
<i>Pleistophora</i> sp. (NW Atl)		<i>Contracaecum osculatum</i> larva (Baltic Sea)
Ciliophora		? <i>C. rudolphii</i> larva (NE Atl, and/or Baltic Sea)
Trichodinidae gen. sp. (NW Atl)		<i>C. siluriglandis</i> larva (Aral Sea)
"Protozoa"		<i>Contracaecum</i> sp. larva (NE Atl, NW Atl, Greenland Sea, Norwegian Sea)
Unidentified Sporozoa (Baltic Sea)		<i>Cystidicola farionis</i> (Baltic Sea)
Myxozoa		<i>Hysterothylacium aduncum</i> larva (NW Atl, NW Atl, Baltic Sea, Barents Sea, Norwegian Sea)
? <i>Ceratomyxa acadiensis</i> NW Atl)		Nematoda gen. spp. (NE Atl, NW Atl, Baltic Sea)
<i>C. auerbachii</i> (NE Atl)		? <i>Phocascaris</i> sp. larva (Baltic Sea)
<i>Kudoa clupeidae</i> (NE Atl, NW Atl)		<i>Pseudoterranova decipiens</i> larva (NW Atl)
Monogenea		? <i>Raphidascaris acus</i> (NE Atl)
<i>Gyrodactyloides baueri</i> (White Sea)		Acanthocephala
<i>Gyrodactyloides</i> spp. (White Sea)		? <i>Acanthocephalus clavula</i> (Baltic Sea)
<i>Gyrodactylus harengi</i> (Baltic Sea, White Sea)		? <i>A. lucii</i> (Baltic Sea)
<i>Gyrodactylus</i> sp. (NW Atl)		<i>Corynosoma semerme</i> juvenile (Baltic Sea)
<i>Lamniscus dogieli</i> (White Sea)		<i>C. strumosum</i> juvenile (Baltic Sea)
<i>Mazocraeoides georgei</i> (NW Atl)		<i>Corynosoma</i> sp. juvenile (Baltic Sea)
<i>Mazocraes harengi</i> (NE Atl)		<i>Echinorhynchus gadi</i> (NE Atl, Baltic Sea)
Digenea		<i>E. salmonis</i> (Baltic Sea)
<i>Brachyphallus crenatus</i> (NE Atl, NW Atl, Baltic Sea, Barents Sea)		<i>Neoechinorhynchus rutili</i> (Baltic Sea)
<i>Cryptocotyle lingua</i> metacercaria (NW Atl)		<i>Pomphorhynchus kostylewi</i> (Baltic Sea)
<i>Derogenes varicus</i> (NE Atl, NW Atl, Barents Sea)		<i>P. laevis</i> (Baltic Sea)
Digenea gen. sp. metacercaria (Aral Sea)		<i>Serrasentis socialis</i> juvenile (NW Atl)
Digenea gen. sp. (NE Atl, Baltic Sea)		
<i>Diplostomum spathaceum</i> metacercaria (Baltic Sea)		Arguloidea
? <i>Hemiurus appendiculatus</i> (NE Atl, NW Atl, Baltic Sea)		<i>Argulus alosae</i> (NW Atl)
<i>H. levinseini</i> (NW Atl, Barents Sea)		<i>A. coregoni</i> (Baltic Sea)
<i>H. leuhei</i> (NE Atl, Baltic Sea)		
<i>H. raabei</i> (Baltic Sea)		Copepoda
<i>Lecithaster confusus</i> (NW Atl, Baltic Sea, Barents Sea)		<i>Caligus elongatus</i> (NE Atl, NW Atl)
<i>L. gibbosus</i> (NE Atl, NW Atl, Baltic Sea, Barents Sea)		<i>Ergasilus sieboldi</i> (Baltic Sea)
? <i>Opechona bacillaris</i> (NE Atl)		<i>Lepophtheirus pollachius</i> (NE Atl)
<i>Opecoeloides vitellus</i> (NW Atl)		<i>Lernaeenicus sprattae</i> (NE Atl)
? <i>Parahemius merus</i> (Baltic Sea)		<i>Lernaeocera</i> sp. (NE Atl)
<i>Renicola</i> spp. metacercaria (NE Atl)		
<i>Stephanoprorapseudoechinata</i> metacercaria (Baltic Sea)		<i>Clupea harengus pallasi</i> Valenciennes
Cestoda		Syn: <i>Clupea pallasi</i> Valenciennes
<i>Abothrium gadi</i> plerocercoid (Baltic Sea)		<i>Clupea harengus pallasi</i> n. <i>maris-albi</i> Berg
<i>Bothriocephalus scorpii</i> (NW Atl, Baltic Sea)		<i>Clupea harengus pallasi</i> n. <i>suworowi</i> Rabinerson
<i>Bothriocephalus</i> sp. (Baltic Sea, NW Atl)		Includes: <i>Clupea harengus</i> auct.
Cestoda gen. sp. (NE Atl, Baltic Sea)		
<i>Eubothrium crassum</i> (Baltic Sea)		Apicomplexa
<i>Eubothrium</i> sp. (Baltic Sea)		<i>Eimeria clupearum</i> (NE Pac, E Ber Sea)
<i>Grillotia erinaceus</i> plerocercoid (NW Atl)		<i>E. nishin</i> (NE Pac, NW Pac, E Ber Sea)
<i>Lacistorhynchus tenuis</i> plerocercoid (NE Atl, NW Atl)		<i>E. sardinae</i> (White Sea)
<i>Proteocephalus</i> sp. (Baltic Sea)		
<i>Pseudophyllidea</i> gen. sp. (NW Atl)		Microspora
		<i>Microsporida</i> gen. sp. (NE Pac)
		Ciliophora
		<i>Trichodina ploveri</i> (W Ber Sea)

Myxozoa	<i>Ceratomyxa auerbachi</i> (NE Pac) <i>C. orientalis</i> (White Sea, NW Pac) <i>Chloromyxum</i> sp. (White Sea) <i>Leptotheca</i> sp. (NE Pac) <i>Ortholinea orientalis</i> (White Sea, NE Pac)	<i>Anisakis</i> sp. larva (White Sea, NE Pac, NW Pac) <i>Ascarophis pacificus</i> (NW Pac) <i>Contracaecum</i> sp. larva (NE Pac, NW Pac, E Ber, W Ber) <i>Hysterothylacium aduncum</i> (White Sea, NE Pac, NW Pac, E Ber, W Ber) Nematoda gen. spp. (NE Pac, NW Pac) <i>Oxyuroidea</i> gen. sp. larva (NE Pac) <i>Pseudoterranova decipiens</i> larva (White Sea) <i>Pseudoterranova</i> sp. larva (NE Pac, NW Pac) <i>Raphidascaris</i> sp. larva (NW Pac) <i>Spirurida</i> gen. spp. larva (NE Pac)
Monogenea	<i>Gyrodactyloidea</i> gen. spp. (White Sea, NE Pac) <i>Gyrodactyloides baueri</i> (White Sea, W Ber) <i>G. petruschewskii</i> (White Sea, NE Pac, W Ber) <i>Gyrodactyloides</i> spp. (White Sea) <i>Gyrodactylus cyclopteri</i> (White Sea) <i>G. flesi</i> (White Sea) <i>G. gerdi</i> (White Sea) <i>G. groenlandicus</i> (White Sea) <i>G. harenghi</i> (White Sea, NE Pac, W Ber) <i>G. pterygialis</i> (White Sea) <i>G. pungitii</i> (White Sea) <i>G. robustus</i> (White Sea, W Ber) <i>Gyrodactylus</i> spp. (White Sea) <i>Lamniscus dogieli</i> (White Sea, W Ber)	Acanthocephala <i>Acanthocephala</i> gen. sp. (NW Pac) <i>Corynosoma semerme</i> juvenile (White Sea) <i>C. strumosum</i> juvenile (White Sea, NE Pac, NW Pac, E Ber) <i>C. villosum</i> juvenile (NE Pac) <i>Echinorhynchus gadi</i> (White Sea, NE Pac, NW Pac) <i>Rhadinorhynchus trachuri</i> (NE Pac)
Digenea	<i>Brachyphallus crenatus</i> (White Sea, NE Pac, NW Pac, E Ber, W Ber) <i>Derogenes varicus</i> (White Sea, NW Pac) Digenea gen. sp. metacercaria (E Ber) Digenea gen. sp. (NE Pac, NW Pac) <i>Diplostomum spathaceum</i> metacercaria (White Sea) <i>Galactosomum phalacrocoracis</i> metacercaria (NE Pac) <i>Hemiurus levinseni</i> (NE Pac, NW Pac) <i>Lecithaster confusus</i> (White Sea) <i>L. gibbosus</i> (NE Pac, E Ber, W Ber) <i>Lecithaster</i> sp. (NW Pac) <i>Parahemiuirus merus</i> (NE Pac) <i>Podocotyle atomon</i> (NW Pac) <i>P. reflexa</i> (NW Pac) <i>Prosorhynchoides basargini</i> metacercaria (NE Pac, E Ber) <i>P. gracilescens</i> metacercaria (NW Pac) <i>Pronoprymna petrowi</i> (NE Pac, E Ber, W Ber) <i>Rhipidocotyle</i> sp. metacercaria (NE Pac)	Copepoda <i>Bomolochus cuneatus</i> (NE Pac, NW Pac) <i>Caligus clemensi</i> (NE Pac, E Ber)
		Agnatha <i>Lampetra ayresi</i> (NE Pac)
		<i>Clupea harengus</i> Linnaeus Includes: <i>Clupeae harengi</i> <i>Clupe hareng</i>
Monogenea		Monogenea <i>Gyrodactylus</i> sp. (White Sea)
Digenea		Digenea <i>Hemiurus appendiculatus</i> (–) <i>H. luehei</i> (–) ? <i>Neophasis oculata</i> (–)
Cestoda		Cestoda ? <i>Ligula intestinalis</i> plerocercoid (–) <i>Scolex pleuronectis</i> plerocercoid (–)
Nematoda		Nematoda Anisakidae gen. spp. (–) <i>Anisakis</i> sp. larva (–) <i>Contracaecum</i> sp. larva (–) <i>Desmidocercella numidica</i> (–) Nematoda gen. sp. (–) <i>Hysterothylacium aduncum</i> larva (–) <i>Pseudoterranova decipiens</i> larva (–)
		Acanthocephala <i>Corynosoma semerme</i> (–)

Erroneous Listings

A number of species have erroneously been listed in papers and monographs and by abstracting services as having been reported from *Clupea harengus*. The Index Catalogue of Medical and Veterinary Zoology (suppl. 18, part 7, p. 163) lists, in reference to Sanzin (1965) *Hexamita salmonis* (Moore, 1923) from *C. harengus*. Although Sanzin did examine herring for this species no infections were noted. Records of digenetic trematodes from the Black Sea region by Koval (1962) were also (suppl. 15, Parasite-Subject Catalogue: Hosts, p. 41) erroneously listed under *Clupea harengus* due to an incorrect assignment of the common name "oseledest" to this species rather than to the correct host, *Alosa kessleri pontica*, the Black Sea shad. Also attributed to *C. harengus* (Index-Catalogue Med. Vet. Zool., Subjects: Trematoda and Trematode Diseases, Part I: Supergenera and Genera A and B, p. 153) and repeated by Yamaguti (1971) is an erroneous listing attributing to Smirnova (1957) (date listed as 1958) a report of *Bucephalus polymorphus* Baer, 1827 from *C. harengus membras*. The correct host for this report is actually *Clupeonella delicatula tscharcalensis*. Records of Kulkarni (1969, 1970) of Monogenea from *Clupea rarengus* [sic] from India are considered to be the result of host misidentification. Previously mentioned are the possibly erroneous listings from herring of *Opechona bacillaris* (Molin, 1859) by Ward and Fillingham (1934) and of *Distoma oculatum* Levissen, 1881 and *Ligula digramma* Creplin, 1839 by von Linstow (1889). Additionally, the presumptive diagnosis of *Myxosoma cerebralis* (Hofer, 1903) from Baltic herring by Dannevig and Hansen (1952) is undoubtedly erroneous.

Acknowledgments

We wish to thank Dr D. I. Gibson, Parasitic Worms Section, British Museum (Natural History), London, England for assistance in tracing records, resolving questions of taxonomy, and examination of works by Gmelin (1790) and Linnaeus (1767, 1789). Dr L. W. Reimer, Pädagogische Hochschule "LiseLOTTE Herrman," Gustrow, East Germany provided information on records from herring contained in his 1962 Inaugural-Dissertation. Thanks are due to the following individuals for permission to cite restricted research documents: Dr K. MacKenzie, Department of Agriculture and Fisheries for Scotland, Aberdeen, Scotland; Dr L. F. Khalil, Commonwealth Institute of Helminthology, St. Albans, Herts, England; Dr P. Hagel, the Netherlands Institute for Fisheries Investigations, IJmuiden, the Netherlands (for permission to cite documents of the late R. Th. Roskam); L. S. Parsons, Department of Fisheries and Oceans, Ottawa, Canada; Drs C. C. Friess and H. Schultz, Institute for Deep Sea Fisheries and Fish Processing, Rostock, East Germany; Dr V. Sjöblom, Finnish Game and Fisheries Research Institute, Fisheries Division, Helsinki, Finland; Drs B. Lubieniecki and J. Popiel, Sea Fisheries Institute, Gdynia, Poland; Dr H. Rosenthal, Biologische Anstalt Helgoland, Hamburg, West Germany; Dr A. V. Gaevskaya, AtlantNIRO, Kaliningrad, USSR; and J. Molloy, Department of Agriculture and Fisheries, Fisheries Division, Dublin, Ireland. We would also like to express our appreciation to Dr Z. Kabata, R. A. Bams, K. V. Aro, F. P. J. Velsen, Ms B. Warneboldt, Ms L. Guy, Ms K. Passmans, and Ms M. Miki for assistance in the examination of foreign language publications. Thanks are also due to the staff of the Pacific Biological Station Library, under the direction of G. Miller, for assistance in obtaining literature. Drs L. Margolis and Z. Kabata of the Pacific Biological Station and Dr D. I. Gibson, British Museum (Natural History), provided critical comments on the manuscript.

References

- AKHMEROV, A. Kh. 1955. Parasite fauna of Kamchatka river fish. Izv. Tikhookean. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr. 43: 99–137. (In Russian)
- ANDREASSEN, J. 1970. The first human case of anisakiosis in Denmark. Nytt Mag. Zool. (Oslo) 18: 105.
- ANDREASSEN, J., AND K. JØRRING. 1970. Anisakose i Danmark. Infektion med nematodlarver fra marine fisk. Nord. Med. 84: 1492–1495.
- ANON. 1969. Natural environment research council. Report of the council for the period 1 April, 1968–31 March, 1969. Fisheries Helminthological Unit, London, 58–59.
1970. Natural environment research council. Report of the council for the period 1 April 1969–31 March 1970. Fisheries Helminthological Unit, London, 58.
1972. Scotland Department of Agriculture and Fisheries Report for 1971: 33–44.
1974. Scotland Department of Agriculture and Fisheries Report for 1972: 39–45.
1975. Scotland Department of Agriculture and Fisheries Report for 1974: 30–31.
- ARAI, H. P. 1967a. A preliminary report on a study of the parasites of marine fishes of Burke Channel, British Columbia. Fish. Res. Board Can. MS Rep. 925: 26 p.
- 1967b. Ecological specificity of parasites of some embiotocid fishes. J. Fish. Res. Board Can. 24: 2161–2168.
1969. Preliminary report on the parasites of certain marine fishes of British Columbia. J. Fish. Res. Board Can. 26: 2319–2337.
- ARRO, I. V. 1964. Kalade invasiooni karusloomadele ohtlike parasiitide vastsetega. Eesti Pollumajanduse Akad. Tead. Tööde Kogumik Vet.-Alased Tööd 38: 216–221.
1967. Helminth larvae of fish, dangerous for fur-bearing animals. Veterinariya (Moscow) 44: 65–66. (In Russian)
- ARTHUR, J. R. 1978. Studies on the parasites of Pacific herring (*Clupea harengus pallasi*) in North American waters. Ph.D. thesis, Univ. Calgary, Calgary, Alta.: 206 p.
- ARTHUR, J. R., AND H. P. ARAI. 1979. A preliminary analysis of the value of parasites as biological indicators for stocks of Pacific herring. Proc. Annu. Meet. Can. Soc. Zool., May 13–16, 1979, Univ. Laval, Quebec City, p. 97.
- 1980a. Studies on the parasites of Pacific herring (*Clupea harengus pallasi* Valenciennes): survey results. Can. J. Zool. 58: 64–70.
- 1980b. Studies on the parasites of Pacific herring (*Clupea harengus pallasi* Valenciennes): a preliminary evaluation of parasites as indicators of geographical origin for spawning herring. Can. J. Zool. 58: 521–527.
- AUERBACH, M. 1909. Bemerkungen über Myxosporidien. Zool. Anz. 34: 65–82.
1912. Studien über die Myxosporidien der Norwegischen See-fische und ihre Verbreitung. Zool. Jahrb. Abt. Syst. Oekol. Geogr. Tiere 34: 1–50.
- BANNING, P. VAN. 1971. Some notes on the successful rearing of the herring-worm, *Anisakis marina* L. (Nematoda: Heterocheilidae). J. Cons. Int. Explor. Mer 34: 84–88.
1972. De haringworm nader verkend. Vakbl. Biol. 52: 24–33.
- BANNING, P. VAN, AND H. B. BECKER. 1978. Long-term survey data (1965–1972) on the occurrence of *Anisakis* larvae (Nematoda: Ascaridida) in herring, *Clupea harengus* L., from the North Sea. J. Fish Biol. 12: 25–33.
- BARRACLOUGH, W. E. 1967. Data record. Number, size composition and food of larval and juvenile fish caught with a two-boat surface trawl in the Strait of Georgia, June 6–8, 1966. Fish. Res. Board Can. MS Rep. 928: 58 p.
- BARRACLOUGH, W. E., AND J. D. FULTON. 1967. Data record. Number, size composition and food of larval and juvenile fish caught with a two-boat surface trawl in the Strait of Georgia, July 4–8, 1966. Fish. Res. Board Can. MS Rep. 940: 82 p.
- BASSETT-SMITH, P. W. 1896. A list of the parasitic Copepoda of fish obtained at Plymouth. J. Mar. Biol. Assoc. U.K. 4: 155–163.
- BAYLIS, H. A. 1916. Some ascarids in the British Museum (Natural History). Parasitology 8: 360–378.
- BEAMISH, R. J. 1980. Adult biology of the river lamprey (*Lampetra ayresi*) and Pacific lamprey (*Lampetra tridentata*) from the Pacific coast of Canada. Can. J. Fish. Aquat. Sci. 37: 1906–1923.
- BECKMANN, J. 1769. Beitrag Naturgeschichte, des sogenannten Fiecks in Fischen. Hannöverches Magazin 1769: 665–672.
1770. Caroli a Linné. Amoenitates academicae seu dissertationes variae physicae, medicæ, botanicæ antehac seorsim editæ, nunc collectæ et auctæ cum tabulis aeneis. Volumen septimum. Holmiae 1769. 8, pagg 508. p. 197–200. In J. Beckmann. Physikalisch-Ökonomische Bibliothek worin von den neuesten Büchern, welche die Naturgeschichte, Naturlehre und die Land-und Stadtwirthschaft betreffen, zuverlässige und vollständige Nachrichten ertheilet werden. Vol. 1, Part 2. Wittwe Vandenhoeck. Göttingen.
- BELLINGHAM, O. 1844. Catalogue of Irish Entozoa, with observations. Ann. Mag. Nat. Hist. 13: 101–105, 167–174.
- BENEDEN, P.-J. VAN. 1871. Les poissons des côtes de Belgique, leurs parasites et leurs commensaux. Mem. Acad. R. Sci. Belg. 38: 1–100.
- BENEDEN, P.-J. VAN, AND C.-E. HESSE. 1864. Recherches sur les bdelloses (hirudinées) et les trématodes marins. Mem. Acad. R. Sci. Belg. 34: 1–142.
- BERE, R. 1930a. The parasitic copepods of the fish of the Passamaquoddy region. Contrib. Can. Biol. Fish. (New Ser.) 5: 423–430.
- 1930b. Parasitic copepods from the Vancouver Island region. Fish. Res. Board Can. MS Rep. (Biol.) 259: 3 p.
- BERLAND, B. 1961. Nematodes from Norwegian marine fishes. Sarsia 2: 1–50.
1962. Rå sild og markellarver. Naturen (1962) No. 2: 106–111.
- BEVERLEY-BURTON, M., O. L. NYMAN, AND J. H. C. PIPPY. 1977. The morphology, and some observations on the population genetics of *Anisakis simplex* larvae (Nematoda: Ascaridata) from fishes of the North Sea. J. Fish. Res. Board Can. 34: 105–112.
- BEVERLEY-BURTON, M., AND J. H. C. PIPPY. 1977. Morphometric variations among larval *Anisakis simplex* (Nematoda: Ascaridoidea) from fishes of the North Atlantic and their use as biological indicators of host stocks. Environ. Biol. Fishes 2: 309–314.
- BEZUBIK, B. 1956. Materiały do helmintofauny ptaków wodnych Polski. Acta Parasitol. Pol. 4: 59–88. (English summary).
- BISHOP, Y. M. M., AND L. MARGOLIS. 1955. A statistical examination of *Anisakis* larvae (Nematoda) in herring (*Clupea pallasi*) of the British Columbia coast. J. Fish. Res. Board Can. 12: 571–592.
- BLOCH, M. E. 1782. Abhandlung von der Erzeugung der Eingeweidewürmer und den Mitteln wider dieselben. Siegmund Friedrich Hesse, Berlin. 54 p.
- BYKHOVSKAYA-PAVLOVSKAYA, I. E., ET AL. 1962. Key to parasites of freshwater fish of the USSR. Izdat. Akad. Nauk SSSR, Moscow and Leningrad. (Transl. from Russian by Israel Program for Sci. Transl. Jerusalem, No. 1136, 1964)
- CIALUPSKÝ, J. 1955. Parazitické červi v mrazeňích a uzených morských rybách. Cesk. Parasitol. 2: 83–88.
- CHAPA, E. 1969. Studies on digenetic trematodes of marine fishes from Monterey and Tomales Bays, California. M.Sc. thesis, Univ. Pacific, Stockton, CA. 85 p.
- COBBOLD, T. S. 1866. Catalogue of the specimens of Entozoa in the Museum of the Royal College of Surgeons of England. R. Hardwicke, London. 24 p.
- COOPER, A. R. 1915. Trematodes from marine and fresh-water fishes, including one species of ectoparasitic turbellarian. Trans. R. Soc. Can., Sect. 4: 181–205.
- CREPLIN, F. C. H. 1839. Eingeweidewürmer, Binnenwürmer, Thierwürmer, p. 277–302. In Allgemeine Encyklopädie der Wissenschaften und Künste, Vol. 32. J. S. Ersch and J. G. Gruber, Leipzig.
1846. Filaria, p. 154–178. In Allgemeine Encyklopädie der Wissenschaften und Künste, Vol. 44. J. S. Ersch and J. G. Gruber, Leipzig.
- CUVIER, G., AND A. VALENCIENNES. 1847. p. 72. In Histoire naturelle des poissons. Vol. 20. P. Bertrand, Paris.
- DAILEY, M. D. 1975. Investigations on the viability of larval helminths after freezing. Aquat. Mammals 3: 22–25.
- DANNEVIG, A., AND S. HANSEN. 1952. Faktorer av betydning for fiskeegegenes og fiskeyeengelens oppvekst. Fiskeridirekt. Skrift., Ser. Havundersokser 10: 5–36.
- DAVEY, J. T. 1971. A revision of the genus *Anisakis* Dujardin, 1845 (Nematoda: Ascaridata). J. Helminthol. 45: 51–72.
- 1972a. The incidence of *Anisakis* sp. larvae (Nematoda: Ascaridata) in the commercially exploited stocks of herring (*Clupea harengus* L., 1758) (Pices: Clupeidae) in British and adjacent waters. J. Fish Biol. 4: 535–554.
- 1972b. The public health aspects of a larval roundworm from the herring. Community Health (Bristol) 4: 104–106.

- DESLONGCHAMPS, J. A. E. 1824. Filocapsulaire; *Filocapsularia*, p. 398–400. In J. V. F. Lamouroux et al. [ed.] Encyclopédie Méthodique, Histoire naturelle des Zoophytes, Part 2, Livraison 98 (published 1827) Ve. Agasse, Paris.
- DOGIEL, V. A. 1940. Coccidia from the testes of Clupeidae and their zoogeographical significance. Tr. Leningr. Obshch. Estestvoispyt. 68: 32–39. (Transl. from Russian by Fish. Res. Board Can. Transl. Ser. No. 3750, 1976)
1948. Parasitic Protozoa of fishes of Peter the Great Bay. Izv. Vses. Nauchno-Issled. Inst. Ozern. Rech. Rybn. Khoz. 27: 17–66. (In Russian)
- DOLLFUS, R. P. 1956. Liste des parasites animaux du hareng de l'Atlantique Nord et de la Baltique. J. Cons. Int. Explor. Mer 22: 58–65.
1970. Les vers nematodes du hareng. C. R. Congr. Natl. Soc. Savantes Sect. Sci. (94th), Pau, 1969, 3: 351–426.
- DORNHEIM, H. 1973. Nematodenlarven im Hering der nordamerikanischen Ostküste in den Jahren 1969 bis 1972. Inf. Fischwirtsch. 20: 3–5.
- DUBININA, M. N. 1966. Tapeworms (Cestoda, Ligulidae) of the fauna of the USSR. Izdat. "Nauka," Moscow and Leningrad. 261 p. (Transl. from Russian by Amerind Publ. Co., New Delhi, 1980)
- DUBNITSKY, A. A. 1968. Use of low temperatures for the prophylaxis of *Corynosoma* infections of fur-bearing animals p. 171–173. In V. S. Ershev et al. [ed.] Gelminty Cheloveka, Zhivotnykh i Rastenii i Mery Borby s Nimi (K 90-Letyu Akademika K. I. Skrjabina). Izdat. "Nauka," Moscow. (In Russian)
- ELLIS, M. F. 1930. Investigations on the protozoan fish parasites of the St. Andrew's region. Trans. Nova Scotian Inst. Sci. 17: 268–275.
- FABRICIUS, O. 1794. Bidrag til snylte-ormenes historie. Skr. Naturh.-Selsk., Copenhagen, 3(2): 1–45.
- AGERHOLM, H.-P. 1978. New implications of the nematode infection of Baltic cod liver. Short Commun. 4th Int. Congr. Parasitol. Sect. C: p. 192.
1982. Parasites of fish in Finland. VI. Nematodes. Acta Acad. Abo. 40: 1–128.
- FORSSELL, A. L. 1905. Bidrag till kännedom om Echinorhyncherna i Finlands fishar. Acta. Soc. Fauna Flora Fenn. 27: 1–30.
- FRASER, C. M. 1920. Copepods parasitic on fish from the Vancouver Island region. Trans. R. Soc. Can., Ser. 3, 13(5): 45–67.
- FRIESS, C. C. 1977. Investigations on the spring spawning Rügen herring infested by *Anisakis* in 1975–1977. Int. Counc. Explor. Sea, C.M. 1977/P:16, Baltic Fish Comm.: 2 p.
- FUJITA, T. 1932. On new nematodes (*Contracaecum*) in fishes of Japan. Bull. Sch. Fish., Hokkaido Imp. Univ. 2: 36–40.
1934. Note on *Eimeria* of herring. Proc. 5th Pac. Sci. Congr. 5 (for 1933): 4135–4139.
- GAEVSKAYA, A. V. 1977. Parasitofauna of Baltic herring (*Clupea harengus membras*) from the southeast Baltic Sea. Int. Counc. Explor. Sea, C.M. 1977/H:7, Pelagic Fish (Northern) Comm.: 11 p.
1979. Parasites of Baltic herring of the southeastern part of the Baltic Sea as indicators of their biological characteristics, p. 21–22. In I. N. Verbitskaya et al. [ed.] Tez. Dokl. Vses. Soveshch. Parazitol. Bolezn. Ryb, Leningrad. (In Russian)
- GAEVSKAYA, A. V., AND L. S. SHAPIRO. 1981. On the problem of the distinctness of Baltic herring (*Clupea harengus membras* L.) of Vistula Bay of the Baltic Sea, p. 11–19. In E. Bakova. [ed.] The status of stocks and essentials of rational fisheries in the Atlantic Ocean. Atl. Nauchno-Issled. Inst. Rybn. Khoz. Okeanogr., Izdat. "Kaliningradskaya Prava," Kaliningrad. (In Russian)
- GAEVSKAYA, A. V., AND B. A. UMNOVA. 1977. On the parasite fauna of the principal commercial fishes of the northwest Atlantic. Biol. Morya (Vladivost.) 4: 40–48. (In Russian)
- GAEVSKAYA, A. V., AND A. M. ZHUDOVA. 1979. Infection of young Baltic herring from Vislinsk Bay with plerocercoids of *Proteocephalus* (Cestoda). Gidrobiol. Zh. 15: 98. (In Russian)
- GETSEVICHYUTE, S. Yu. 1958. Parasite fauna of fish in Kurshy Mares bay. Liet. TSR Mokslu Akad. Biol. Inst. Darb. Ser. B, 3: 101–139. (In Russian)
- GIARD, M. A. 1903. Exuviations métamorphiques chez les ascarides des poissons (group de l'*Ascaris adunca* Rud.). C. R. Soc. Biol. 55: 627–630.
- GIBSON, D. I. 1970. Aspects of the development of "herringworm" (*Anisakis* sp. larva) in experimentally infected rats. Nytt Mag. Zool. (Oslo) 18: 176–187.
- RIESBRECHT, W. 1882. Die freilebenden Copepoden der Kieler Föhrde. Jahresber. Comm. Wissensch. Untersuch. Deutsch. Meere in Kiel (1877–81) 4: 85–168.
- GMELIN, J. F. 1791. Caroli à Linné. . . Systema naturae per regna tria naturae, secundum classes ordines genera, species cum characteribus differentiis, synonymis, locis. 13th ed. Vol. 1, Part 6 (Vermes), J. B. Delamoliere, Leipzig, p. 3021–3910. (Not seen)
- GOLVAN, Y. J. 1969. Systématique des Acanthocéphales (Acanthocephala Rudolphi 1801) Première partie. L'ordre des Palaeacanthocephala Meyer 1931. Premier fascicule. La super-famille des Echinorhynchoidea (Cobbold 1876) Golvan et Houin 1963. Mém. Mus. Natl. Hist. Nat. Ser. A Zool. 57: 1–373.
- GRABDA, J. 1973a. Z inwajzjologii *Anisakis simplex*. Ann. Acad. Med. Stetin. Supl. 10: 39–45.
- 1973b. Badania nad przezywalnością larw nicieni *Anisakis simplex*. Ann. Acad. Med. Stetin. Supl. 10: 47–52.
- 1974a. The dynamics of the nematode larvae, *Anisakis simplex* (Rud.) invasion in the south-western Baltic herring (*Clupea harengus* L.). Acta Ichthyol. Piscatoria 4: 3–21.
- 1974b. Apparition saisonnière des larves d'*Anisakis simplex* chez le hareng de la Baltique du sud-ouest. Proc. Int. Congr. Parasitol. 3: 1625.
- 1974c. Znaczenie pasozytów z rodziny Anisakidae (Nematoda) dla przewrotwora rybnego i zdrowia człowieka. Zesz. Nauk. Akad. Roln. Szczecinie 43: 97–103.
- 1974d. Technologia przetworów rybnych, p. 221. In Informator o wynikach badań naukowych zakończonych w 1972. Part 2 (Sect. 10) Fisheries.
1975. Studies on the life-history and morphogenesis of *Anisakis simplex* (Rudolphi, 1809) cultured *in vitro*. 2nd Eur. Multicoll. Parasitol., Summaries: 47–48.
1976. Studies on the life cycle and morphogenesis of *Anisakis simplex* (Rudolphi, 1809) (Nematoda: Anisakidae) cultured *in vitro*. Acta Ichthyol. Piscatoria 6: 119–141.
- GRABDA, J., AND C. FELINSKA. 1975. Studies on the toxicity of larval *Anisakis simplex* (Rud.) Part I. Biological test on white mice. Acta Ichthyol. Piscatoria 5: 3–11.
- GRABDA-KAZUBSKA, B., AND J. C. CHUBB. 1968. *Acanthocephalus* — the correct generic name for *Echinorhynchus clavula* Dujardin, 1845. Acta Parasitol. Pol. 15: 305–311.
- GRABDA-KAZUBSKA, B., AND L. EJSYMONT. 1969. Studies on the morphology, variability and systematic status of *Echinorhynchus borealis* Linstow, 1901 (Acanthocephala, Echinorhynchidae). Acta Parasitol. Pol. 17: 65–87.
- GUETTARD, J.-E. 1768. Sur des vers ascarides des harengs, p. 85–86. In J. E. Guettard. Mémoirs sur différentes parties des Sciences et Arts, Vol. 1, Laurent Prault, Paris.
- GUSTAFSON, P. V. 1953. The effect of freezing on encysted *Anisakis* larvae. J. Parasitol. 39: 585–588.
- HAGIWARA, Y., S. TAKAHATA, T. SHINKURA, T. ONO, Y. KITAMURA, AND J. HOSHIHARA. 1969. Investigations for the presence of anisakid nematodes in marine fishes from Hokkaido. J. Jpn. Vet. Med. Assoc. 22: 397. (In Japanese)
- HAHN, C. W. 1917. On the sporozoan parasites of the fishes of Woods Hole and vicinity. III. On the *Chloromyxum clupeidae* of *Clupea harengus* (young), *Pomolobus pseudoharengus* (young) and *P. aestivalis* (young). J. Parasitol. 4: 13–20.
- HANSEN, V. 1955. The food of herring on the Blöden Ground (North Sea) in 1953. J. Cons. Int. Explor. Mer 21: 61–64.
- HARTWICH, G. 1964. Revision der Vogelparasitischen Nematoden Mittel-europas. II. Die Gattung *Contracaecum* Ralliet and Henry, 1912 (Ascaridoidea). Mitt. Zool. Mus. Berlin 40: 15–53.
1975. Schlachtwürmer, Nemathelminthes Rund-oder Faden-würmer, Nematoda parasitische Rundwürmer von Wirbeltieren. I. Rhabditida und Ascaridida. Die Tierwelt Deutschlands 62: 256 p.
- HASEGAWA, H. 1978. Larval nematodes of the superfamily Spiruroidea — a description, identification and examination of their pathogenicity. Acta Med. Biol. 26: 79–116.
- HAUCK, A. K. 1977. Occurrence and survival of the larval nematode *Anisakis* sp. in the flesh of fresh, frozen, brined, and smoked Pacific herring, *Clupea harengus pallasi*. J. Parasitol. 63: 515–519.
- HAUCK, A. K., AND E. B. MAY. 1977. Histopathologic alterations associated with *Anisakis* larvae in Pacific herring from Oregon. J. Wildl. Dis. 13: 290–293.
- HECKMANN, R. A., AND L. A. JENSEN. 1978. The histopathology and prevalence of *Henneguya sebasta* and *Kudoa clupeidae* in the rockfish, *Sebastodes paucispinis* of southern California. J. Wildl. Dis. 14: 259–262.

- HELLER, A. F. 1949. Parasites of cod and other marine fish from the Baie de Chaleur region. Can. J. Res., Sect. D, 27: 243–264.
- HENTSCHEL, E. 1950. Die Nahrung der Heringslarven. Helgol. Wiss. Meeresunters. 3: 59–81.
- HODDER, V. M., AND L. S. PARSONS. 1971. Preliminary comparison of Scotian Shelf and southwest Newfoundland herring taken during the winter of 1971. Int. Comm. Northwest Atl. Fish. Res. Doc. 71/120: 1–8.
- HOSKINS, G. E., G. R. BELL, AND T. P. T. EVELYN. 1976. The occurrence, distribution and significance of infectious diseases and neoplasms observed in fish in the Pacific region up to the end of 1974. Environ. Can. Fish. Mar. Serv. Res. Dev. Tech. Rep. 609: 37 p.
- HOUWING, H. 1969a. Het inactiveren van de haringnematode door keukenzout en azijnzuur. Visserijwereld, year 28(20): 19, 22–23.
- 1969b. Het inactiveren van haringnematoden tijdens het rookproces. Visserijwereld, year 28(20): 23.
- ICHIHARA, A., M. MACHIDA, T. KOGA, AND T. ABE. 1968. Investigation for the presence of *Anisakis* of marine fishes and others. Jpn. J. Parasitol. 17(6, Suppl): 582–583. (In Japanese)
- IVANCHENKO, O. F., AND T. A. GROZDIOLOVA. 1971. Parasites of young White Sea herring (*Clupea harengus pallasi maris-albi*) reared under artificial conditions. Parazitologiya 5: 233–236. (In Russian)
1981. Infection of larval White Sea herring reared under experimental conditions with the trematode *Lecithaster gibbosus*. Parazitologiya 15: 132–134. (In Russian, English summary)
- JÄÄSKELÄINEN, V. 1921. Über die Nahrung und die Parasiten der Fische im Ladogasee nebst einem Verzeichnis der in Finnland bisher konstatierten Fischhelminthen nach ihren Wirten geordnet. Ann. Acad. Sci. Fenn., Ser. A. 14: 1–55.
- JACKSON, G. J., J. W. BIER, W. L. PAYNE, T. A. GERDING, AND W. G. KNOLLENBERG. 1978. Nematodes in fresh market fish of the Washington D.C. area. J. Food Prot. 41: 613–620.
- JAHNEL, J. 1940. Lebende Nematodenlarven in einem marinierter Hering. Wien. Tierärztl. Monatschr. 27: 442–444.
- KABATA, Z. 1962. Five new species of Myxosporidia from marine fishes. Parasitology 52: 177–186.
1963. Incidence of coccidioses in Scottish herring (*Clupea harengus* L.). J. Cons. Int. Explor. Mer 28: 201–210.
- KAGEI, N. 1969. Life history of nematodes of the genus *Anisakis*. Saishin Igaku 24: 389–400. (Transl. from Japanese by Fish. Res. Board Can. Transl. Ser. No. 1324, 1969)
- KAGEI, N., G. IMAE, AND T. TANAKA. 1971. Studies on anisakid Nematoda (Anisakinac). III. On *Anisakis*- larvae found in the kazu-no-ko (herring roes). Bull. Inst. Public Health (Tokyo) 20: 202–206. (In Japanese)
- KAHL, W. 1938. Nematoden in Seefischen. II. Erhebungen über den Befall von Seefischen mit Larven von *Anacanthocheilus rotundatus* (Rudolphi) und die durch diese Larven hervorgerufenen Reaktionen des Wirtsgewebes. Z. Parasitenkd. 10: 513–534.
1939. Nematoden in Seefischen. III. Statistische Erhebungen über den Nematodenbefall von Seefischen. Z. Parasitenkd. 11 (for 1940): 16–41.
- KATO, T., M. UMINUMA, K. ITO, AND K. MIURA. 1968. On Anisakinae from the marine fishes at the Central Market of Japan. Shokuhin Eisei Kenkyu 18: 784–793. (In Japanese)
- KHALIL, L. F. 1968a. Larval nematodes in juvenile and adult herring. Int. Counc. Explor. Sea, Symp. "The biology of early stages and recruitment mechanisms of herring," No. 12: 10 p.
- 1968b. Anisakiasis — "herring-worm disease" — The *Anisakis*-type larvae in the herring of the North Sea. Trans. R. Soc. Trop. Med. Hyg. 62: 11.
1969. Larval nematodes in the herring (*Clupea harengus*) from British coastal waters and adjacent territories. J. Mar. Biol. Assoc. U.K. 49: 641–659.
- KIKUCHI, S., K. KOSUGI, H. HIRABAYASHI, AND S. HAYASHI. 1970. Six types of *Contracaecum* larvae (Nematode) found in the sea fishes in Japan. Yokohama Igaku 21: 421–427. (In Japanese)
- KOBAYASHI, A., T. KOYAMA, M. KUMADA, Y. KOMIYA, T. OSHIMA, N. KAGEI, T. ISHII, AND M. MACHIDA. 1966. A survey of marine fishes and squids for the presence of Anisakinae larvae. Jpn. J. Parasitol. 15: 348–349. (In Japanese)
- KOGA, A., A. ICHIHARA, M. MACHIDA, AND M. ABE. 1968. Survey of anisakid nematodes infecting fishes. Min. Fish. Res. Study Group, Japan, Pamphlet: 11 p. (In Japanese)
- KOVAL, V. P. 1962. Parasite fauna of fishes of the Dnieper delta. Visn. Kiv. Univ., Ser. Biol. 5: 98–104. (In Ukrainian)
- KOYAMA, T., A. KOBAYASHI, M. KUMADA, Y. KOMIYA, T. OSHIMA, N. KAGEI, T. ISHII, AND M. MACHIDA. 1969. Morphological and taxonomical studies on Anisakidae larvae found in marine fishes and squids. Jpn. J. Parasitol. 18: 466–487. (Transl. from Japanese by Fish. Res. Board Can. Transl. Ser. No. 3082, 1974)
- KREIS, H. A. 1958. Parasitic Nematoda. Zool. Iceland 2, Part 15b: 1–24.
- KRØYER, H. 1846–49. Danmark's fiske. Vol. 3. Part 1. S. Triers, Copenhagen, p. 174.
- KUIPERS, F. C. 1964. Eosinophilic phlegmonous inflammation of the alimentary canal caused by a parasite from the herring. Pathol. Microbiol. 27: 925–930.
- KUIPERS, F. C., E. H. KAMPELMACHER, AND F. STEENBERGEN. 1963. Onderzoeken over haringwormziekte bij konijnen. Ned. Tijdschr. Geneesk. 107: 990–995.
- KUIPERS, F. C., P. H. VAN THIEL, W. RODENBURG, W. J. WIELINGA, AND R. T. ROSKAM. 1960. Eosinophilic phlegmon of the alimentary canal caused by a worm. Lancet, Year 1960, 2: 1171–1173.
- KUIPERS, F. C., P. H. VAN THIEL, AND R. T. ROSKAM. 1960. Eosinofiele flegmone van de dunne darm, veroorzaakt door een niet aan het lichaam van de mens aangepaste worm. Ned. Tijdschr. Geneesk. 104: 422–427.
- KULACHKOVA, V. G. 1968. Monogenetic trematodes of the White Sea herring (*Clupea harengus pallasi n. maris-albi* Berg), p. 62. In O. N. Bauer et al. [ed.] Ref. Dokl. 5th Vses. Soveshch. Bolezn. Parazit. Ryb Vodnykh Bespozvonochnykh. Izdat. "Nauka," Leningrad. (In Russian)
1970. *Gyrodactyloides baueri* sp. n. (Monogeneida: Gyrodactylidae) from the nasal cavities of White Sea herrings. Parazitologiya 4: 544–546. (In Russian)
1971. Some data on the biology and variability of the Monogenea of White Sea herring. Tez. Dokl. Otchetnaya Nauchn. Sess. Itog. Rabot 1970 G., Akad. Nauk SSSR, Zool. Inst., Leningrad. 25–27. (In Russian)
- 1974a. Monogeneans of the genus *Gyrodactyloides* Bychowsky, 1947 — their biology and distribution. Proc. 3rd Int. Congr. Parasitol. 1: 329–330.
- 1974b. Monogenetic trematodes as indicators of local stocks of White Sea herring, p. 128–134. In I. N. Verbitskaya et al. [ed.] Tez. Dokl. 6th Vses. Soveshch. Bolezn. Parazit. Ryb, Moscow. (In Russian)
1975. Monogenea of the White Sea herring. Rabot. Belomorsk. Biol. Sta. Issled. Fauny Morei 16: 134–168. (In Russian)
- 1976a. Some data on the biology and life cycle of larval *Anisakis* sp. (nematodes), p. 57–58. In Problem. Parazitol. Izdat. "Nauka," Leningrad. (In Russian)
- 1976b. On life of *Anisakis* sp. larvae in herring (*Clupea harengus pallasi n. maris-albi* Berg) of the White Sea. In Tez. Dokl. 3rd Int. Symp. Helm. Inst. Kosice, 12–15 Oct., 1976, High Tatra Mountains, Czech. Helmintol. Ustav Slovensk. Akad. Vied, Kosice. 2 p.
- 1976c. Infection of White Sea herring (*Clupea harengus pallasi n. maris-albi* Berg) with larvae of *Anisakis* sp. Kratk. Tez. Dokl. 2 Vses. Simp. Parazit. Bolezn. Morsk. Zhivotn. AtlantNIRO, Kaliningrad. 36–37. (In Russian)
- 1977a. Monogeneans (Gyrodactylidae) as indicators of local herring stocks (*Clupea harengus pallasi n. maris-albi* Berg) in the White Sea. Parazitol. Sb. 27: 27–34 (Transl. from Russian by Dep. Agric. Fish. Scotland Mar. Lab. Aberdeen Transl. No. 2078.)
- 1977b. Ecological and geographical analysis of the monogenean genus *Gyrodactyloides* Bychowsky, 1947 (Monogeneida), p. 124–130. In O. A. Skarlato et al. [ed.] Issled. Monogenei v SSSR (Mat. Vses. Simp. Monogen.) Zool. Inst. Akad. Nauk SSSR, Leningrad. (In Russian)
1978. The role of herring in the life-cycle of *Anisakis* sp. (Nematoda). Short Commun. 4th Int. Congr. Parasitol., Sect. C. p. 195.
1980. Infection of White Sea herring with *Anisakis* sp. larvae (Nematoda: Anisakinac). Parazitol. Sb. 29: 126–142. (In Russian, English summary.)
- KULKARNI, T. 1969. Studies on the monogenetic trematodes of fishes found in Hyderabad, Andhra Pradesh (India). Part II. Riv. Parassitol. 30: 263–282.
1970. Studies on the monogenetic trematodes of fishes found in Hyderabad, Andhra Pradesh (India). Part III. Riv. Parassitol. 31: 15–28.
- LAIRD, M., AND W. L. BULLOCK. 1969. Marine fish haematozoa from New Brunswick and New England. J. Fish. Res. Board Can. 26: 1075–1102.

- LAYMAN, E. M. 1930. Parasitic worms from the fishes of Peter the Great Bay. Izv. Tikhookean. Nauchno-Prom. Sta. 3: 1–120. (In Russian)
- LEEUWENHOEK, A. VAN. 1697. Continuatio arcanorum naturae detectorum, qua continetur quicquid hactenus ab auctore lingua vernacula editum and in linguam Latinam transfusum non fuit. Epistola 97. Delphis Batavorum. 45–55.
- LEIDY, J. 1857. A synopsis of Entozoa and some of their ecto-congeners observed by the author. Proc. Acad. Nat. Sci. Philadelphia 8 (for 1856): 42–58.
- LEVANDER, K. M. 1909. Beobachtungen über die Nahrung und die Parasiten der Fische des Finnischen Meerbusens. Finnland. Hydrogr. Biol. Untersuch. 5: 1–44.
- LEVINE, N. D., J. O. CORLISS, F. E. G. COX, G. DEROUX, J. GRAIN, B. M. HONIGBERG, G. F. LEEDALE, A. R. LOEBLICH, J. LOM, D. LYNN, E. G. MERINFELD, F. C. PAGE, G. POLJANSKY, V. SPRAGUE, J. VAVRA, AND F. G. WALLACE. 1980. A newly revised classification of the Protozoa. J. Protozool. 27: 37–58.
- LINNAEUS, C. 1767. Systema naturae per regna tria naturae. Secundum classes ordines genera, species, cum characteribus, differentiis, synonymis, locis. 12th ed. reformata, Vol. 1, Part 2, Holmiae. 533–1327. (Not seen)
1789. CXXXI. Lepra quam prae side D. D. Car. V. Linné pro posuit Isacus Uddman ostrobotniensis. Upsaliae 1763. Junii 17. In Caroli A. Linné. Amoenitates Academicae, 2nd ed., Vol. 7, Io. Iacobi Palm, Erlangen. p. 94–108. (Not seen)
- LINSTOW, O. VON. 1878. Compendium der Helminthologie. Hahn'sche, Hannover. 382 p.
1889. Compendium der Helminthologie. Nachtrag. Die Litterature de Jahre 1878–1889. Hahn'sche, Hannover. 151 p.
- LINTON, E. 1901a. Parasites of fishes of the Woods Hole region. Bull. U.S. Fish. Comm. 19 (for 1899): 405–492.
- 1901b. Fish parasites collected at Woods Hole in 1898. Bull. U.S. Fish. Comm. 19 (for 1899): 267–304.
1910. Notes on the flesh parasites of marine food fishes. Bull. U.S. Bur. Fish. 28, part 2 (for 1908): 1195–1209.
1924. Notes on cestode parasites of sharks and skates. Proc. U.S. Natl. Mus. 64 (No. 2511): 1–114.
1940. Trematodes from fishes mainly from the Woods Hole region, Massachusetts. Proc. U.S. Natl. Mus. 88 (No. 3078): 1–172.
1941. Cestode parasites of teleost fishes of the Woods Hole region, Massachusetts. Proc. U.S. Natl. Mus. 90 (No. 3112): 417–442.
- LISIVNENKO, L. N. 1961. Plankton and the food of larval Baltic herring, in the Gulf of Riga. Tr. Nauchno-Issled. Inst. Rybn. Khoz. Soveta Narodn. Khoz. Lat. SSR 3: 105–138. (Transl. from Russian by Fish. Res. Board Can. Transl. Ser. No. 444. 36 p.)
- LÖLIGER-MÜLLER, B. 1955. Nematoden-Larven aus der Bauchhöhle des Herings. Monatsh. Veterinaermed. 10: 486–489.
- LOM, J., AND M. LAIRD. 1969. Parasitic Protozoa from marine and euryhaline fish of Newfoundland and New Brunswick. I. Peritrichous ciliates. Can. J. Zool. 47: 1367–1380.
- LUBIENIECKI, B. 1972. The occurrence of *Anisakis* sp. larvae (Nematoda) in herring from the southern Baltic. Int. Counc. Explor. Sea, C.M. 1972/H: 21, Pelagic Fish (Northern) Comm.: 5 p.
1974. Note on the occurrence of larval *Anisakis* in adult herring and mackerel from Long Island to Chesapeake Bay. Int. Comm. Northwest Atl. Fish. Res. Bull. 10: 79–81.
- LÜHE, M. 1911. Acanthocephalen. Register der Acanthocephalen und parasitischen Plattwürmer geordnet nach ihren Wirten. In A. Brauer [ed.] Die Süßwasserfauna Deutschlands, Eine Exkursionfauna, No. 16. Gustav Fischer, Jena. 116 p.
- LUNDSTRÖM, A. 1942. Die Acanthocephalen Schwedens mit Ausnahme der Fischacanthocephalen von Süßwasserstandorten. Ph.D. thesis, Lund Univ., Lund, Sweden: 238 p.
- MACKENZIE, K. 1974a. The use of parasites in tracing herring recruitment migrations. Int. Counc. Explor. Sea, C.M. 1974/H:31, Pelagic Fish (Northern) Comm.: 5 p.
- 1974b. Parasites as biological tags for herring. Proc. 3rd Int. Congr. Parasitol. 3: 1724.
- 1975a. Parasites as indicators of herring migrations in the North Sea and to the north and west of Scotland. Int. Counc. Explor. Sea, C.M. 1975/H:42, Pelagic Fish (Northern) Comm.: 7 p.
- 1975b. *Renicola* metacercariae (Digenea: Renicolidae) in clupeoid fish: new host records. J. Fish Biol. 7: 359–360.
- 1975c. *Renicola* spp. and *Lacistorhynchus* sp. as biological tags for herring. Parasitology 71(2): iii.
1976. Parasites as biological tags for herring. Norweg. J. Zool. 24: 464–465.
1978. Parasites as biological tags for herring, *Clupea harengus* L. Proc. 4th Int. Congr. Parasitol., Sect. H: p. 6.
1983. Parasites as biological tags in fish population studies. Adv. Mar. Biol. 20: 251–331.
- MACKENZIE, K., AND J. GALLACHER. 1981. Further studies of parasites as biological tags for herring. Int. Counc. Explor. Sea, C.M. 1981/H:58, Pelagic Fish Comm.: 11 p.
- MACKENZIE, K., AND C. JOHNSTON. 1976. Recruitment to the Minch herring population, as determined by the use of parasites as biological tags and a new meristic character. Int. Counc. Explor. Sea, C.M. 1976/H:34, Pelagic Fish (Northern) Comm.: 9 p.
- MALMBERG, G. 1957. Om förekonsten av *Gyrodactylus* på Svenska fiskar. Skr. Sod. Sver. Fiskeriför., Årsskr. 1956: 19–76.
- MAMEREN, J. VAN, AND H. HOUWING. 1968. Effect of irradiation of *Anisakis* larvae in salted herring. Eliminat. Harmful Organisms Food Feed Irradiat., Vienna 68: 73–80.
- MANTER, H. W. 1925. Some marine fish trematodes of Maine. J. Parasitol. 12: 11–18.
1926. Some North American fish trematodes. Ill. Biol. Monogr. 10: 133–264.
- MARGOLIS, L. 1952. Studies on parasites and diseases of marine and anadromous fish from the Canadian Pacific coast. Ph.D. thesis, McGill Univ., Montreal, Que.: 231 p.
- MARGOLIS, L., AND J. R. ARTHUR. 1979. Synopsis of the parasites of fishes of Canada. Bull. Fish. Res. Board Can. 199: 269 p.
- MARGOLIS, L., AND H. L. CHING. 1965. Review of the trematode genera *Bacciger* and *Pentagramma* (Fellodistomatidae) and description of *P. petrowi* (Layman, 1930) n. comb. from marine fishes from the Pacific coast of Canada. Can. J. Zool. 43: 381–405.
- MARKEVICH, A. P. 1956. Parasitic copepods on the fishes of the USSR. Izdat. Akad. Nauk Ukr. SSR, Kiev. 209 p. (Transl. from Russian by Indian Nat. Sci. Doc. Centre, New Delhi, 1976).
- MARKOSWKI, S. 1933. Die Eingeweiderwürmer der Fische des Polnischen Baltikums (Trematoda, Cestoda, Nematoda, Acanthocephala). Arch. Hydrobiol. Rybact. 7: 1–58.
1938. Contribution à la connaissance des helminthes des poissons de la côte Belge. Bull. Mus. R. Hist. Nat. Belg. 14: 10 p.
- MARSHALL, S. M., A. G. NICHOLLS, AND A. P. ORR. 1937. On the growth and feeding of the larval and post-larval stages of the Clyde herring. J. Mar. Biol. Assoc. U.K. 22: 245–267.
- MARTIN, A. R. 1760. Anmarkningar over den sa kallade spitelska fisk och boskap i Norrige. K. Vetenskapsacad. Handl. 21: 306–311.
- MARTIN, O. 1921. Über Ascaridenlarven aus dem Fleisch von Seefischen. Z. Infektionskr. Parasit. Krankh. Hyg. Haustiere 22: 13–36.
- MEYER, A. 1931. Die Acanthocephalen des Arktischen Gebietes. Fauna Arctica 6: 9–20.
- MEYERS, T. R. 1978. Prevalence of fish parasitism in Raritan Bay, New Jersey. Proc. Helminthol. Soc. Wash. 45: 120–128.
- MILLER, M. J. 1941. A critical study of Stafford's report on "Trematodes of Canadian Fishes" based on his trematode collection. Can. J. Res., Sect. D, 19: 28–52.
- MÖLLER, H. 1975. Der Einfluss von Temperatur und Salzgehalt auf Entwicklung und Verbreitung von Fischparasiten. Diss. Univ. Kiel. 109 p.
1978. The effects of salinity and temperature on the development and survival of fish parasites. J. Fish Biol. 12: 311–323.
- MOLLOY, J. 1970. Nematodes in Irish herrings. Int. Counc. Explor. Sea, C.M. 1970/H:9, Pelagic Fish (Northern) Comm.: 5 p.
- MOSER, M., M. S. LOVE, AND L. A. JENSEN. 1976. Myxosporida (Protozoa) in California rockfish, *Sebastodes* spp. J. Parasitol. 62: 690–692.
- MÜHLING, P. 1898. Die Helminthen-fauna der Wirbeltiere Ostpreussens. Arch. Naturgesch. 1: 1–66.
- MURAVEV, V. I. 1970a. On the biology of parasitic nematodes of herring and blue whiting in the North Atlantic. Mater. Rybokhoz. Issled. Sev. Baseina 14: 86–95. (Transl. from Russian by Fish. Res. Board Can. Transl. Ser. No. 1888, 1971).
- 1970b. Parasitic nematodes in commercial fishes of the North Atlantic, p. 75–76. In A. M. Parukhin [ed.] Vopr. Morsk. Parazitol. (Mater. I Vses. Simp. Parazitol. Bolez. Morsk. Zhivotnykh, Sevastopol, 1970), Izdat. "Naukova Dumka," Kiev. (In Russian)
- MYERS, B. J. 1979. Anisakine nematodes in fresh commercial fish from waters along the Washington, Oregon and California coasts. J. Food Protect. 42: 380–384.

- NAHHAS, F. M. 1960. Some hemiurid trematodes of marine fishes of California. M.Sc. thesis, Univ. Pacific, Stockton, CA.
- NICOLL, W. 1907. A contribution towards a knowledge of the Entozoa of British marine fishes. Part 1. Ann. Mag. Nat. Hist. Ser. 7, 19: 66–94.
1910. On the Entozoa of fishes from the Firth of Clyde. Parasitology 3: 322–359.
1915. A list of trematode parasites of British marine fishes. Parasitology 7: 339–378.
- NÖEL, S. B.-J. 1807. Observations sur l'ascaride du clupé hareng, (*gordius marinus* de Linné), (*Ascaris halecis* de Gmelin). Précis Analyt. Trav. Acad. Sci. Belles-Lett. Arts Rouen 1804: 81–82.
- NYBELIN, O. 1924. Zur postembryonalen Entwicklungsgeschichte der Acanthocephalen II. Zool. Anz. 61: 190–193.
- NYGARD, J. J. 1967. Levende *Anisakis*-larver i spekessild. Nord. Veterinaermed. 19: 411–414.
- ODENING, K. 1978. Trematodenliste der DDR. Angew. Parasitol. 19: 168–186.
- ODHNER, T. 1905. Die Trematoden des artischen Gebietes, p. 291–372. In F. Römer and F. Schaudinn [ed.] Fauna Arctica, Vol. 4. Gustav Fischer, Jena.
1911. Zum naturlichen System der digenen Trematoden IV. Zool. Anz. 38: 513–531.
- OKUMURA, T. 1967. Experimental studies on anisakiasis. Med. J. Osaka Univ. 16: 465–499. (In Japanese)
- OLSSON, P. 1866–67, 1867–68. Entozoa, iakttagna hos skandinaviska hasfiskar. I. Platyhelminthes. Lunds Univ. Arsskr., Math. Naturv.-Vetensk. 3(Art. 3): 1–59; 4(Art. 8): 1–64.
1893. Bidrag till Skandinaviens helminthfauna. II. K. Svenska Vetenskapsakad. Handl. 25: 1–41.
- ONO, Y. 1975. Anisakiasis as a parasitic zoonosis and its prevention. I. Chikusan No Kenkyu 29: 605–610 (Transl. from Japanese by Fish. Mar. Serv. Transl. Ser. No. 3872, 1976)
- OSHIMA, T. 1966. Biological aspects on the anisakiasis. Jpn. J. Parasitol. 15: 286–290. (In Japanese)
1972. *Anisakis* and anisakiasis in Japan and adjacent areas. Prog. Med. Parasitol. Jpn. 4: 301–393.
- OSMANOV, S. O. 1971. Parasites of the fishes of Uzbekistan. Izdat. "FAN" Uzbek. SSR, Tashkent. 532 p. (In Russian)
- OTSURU, M. 1968a. Anisakiasis. Niigata Igakkai Zasshi 82: 295–298. (In Japanese)
- 1968b. Anisakiasis. Mod. Media 7: 361–370. (In Japanese)
- OTSURU, M., T. HATSUKANO, T. OYANAGI, and M. KENMOTSU. 1965. Visceral migrans of the gastro-intestinal tract and its vicinity caused by some larval nematodes. Jpn. J. Parasitol. 14: 542–552. (In Japanese)
- OTSURU, M., T. HOTTA, T. HATSUKANO, T. OYANAGI, T. SHIRAKI, AND M. KENMOTSU. 1967. Observations on *Anisakis*-like larvae in marine fishes. Jpn. J. Parasitol. 16: 288. (In Japanese)
- OYANAGI, T. 1967. Experimental studies on the visceral migrans of gastro-intestinal walls due to *Anisakis* larvae. Jpn. J. Parasitol. 16: 470–493.
- PARKER, R. R., AND L. MARGOLIS. 1964. A new species of parasitic copepod, *Caligus clemensi* sp. nov. (Caligidae: Caligidae), from pelagic fishes in the coastal waters of British Columbia. J. Fish. Res. Board Can. 21: 873–889.
- PARSONS, L. S., AND V. M. HODDER. 1971a. Variation in the incidence of larval nematodes in herring from Canadian Atlantic waters. Int. Comm. Northwest Atl. Fish. Res. Bull. 8: 5–14.
- 1971b. Variation in the incidence of larval nematodes in herring from Canadian Atlantic waters. Int. Comm. Northwest Atl. Fish. Res. Doc. 71/6, Ser. No. 2495: 1–21.
1974. Some biological characteristics of the Fortune Bay, Newfoundland, herring stock, 1966–71. Int. Comm. Northwest Atl. Fish. Res. Bull. 10: 15–22.
- PETRUSHEVSKY, G. K. 1957. On the parasites of the herring (*Clupea harengus membras*). Izv. Vses. Nauchno-Issled. Inst. Ozern. Rechn. Rybn. Khoz. 42: 332. (Transl. from Russian by Israel Prog. Sci. Transl. Jerusalem, 1961).
- PETRUSHEVSKY, G. K., AND M. G. PETRUSHEVSKAYA. 1960. The accuracy of quantitative indices relating to the study of parasite faunas of fishes. Parazitol. Sb. 19: 333–353. (In Russian)
- PIPPY, J. H. C. 1975. Larval *Anisakis simplex* in Atlantic salmon (*Salmo salar*) and their value as biological indicators. Ph.D. thesis, Univ. Guelph, Guelph, Ont.: 118 p.
- PIPPY, J. H. C., AND P. VAN BANNING. 1975. Identification of *Anisakis* larva (I) as *Anisakis simplex* (Rudolphi, 1809, det. Krabbe 1878) (Nematoda: Ascaridata). J. Fish. Res. Board Can. 32: 29–32.
- POLAK, M. F. 1966. Haringwormbesmetting bij de mens, 1965. Rijks Inst. voor de Volksgezondheid, Rap. No. U 41/66 Epid, 16 p.
1967. Medische en biologische aspecten van haringwormziekte (Anisakiasis). Rijks Inst. voor de Volksgezondheid, 13 p.
- POLYANSKY, YU. I. 1955. Parasites of the fish of the Barents Sea. Tr. Zool. Inst. Akad. Nauk SSSR 19: 5–170. (Transl. from Russian by Israel Prog. Sci. Transl., Jerusalem, 1966).
- POLYANSKY, YU. I., AND S. S. SHULMAN. 1956. Age alteration in the parasite fauna of fish. Tr. Karelo-Finsk. Fil. Akad. Nauk SSSR, Ser. Parasitol. 4: 3–26. (In Russian)
- POPIEL, J. 1951. Pokarm i odzywianie sie sledzia (*Clupea harengus* L.) naterenie zatoki Gdańskiej i wod przyleglych. Pr. Morsk. Inst. Ryback Gdyni 6: 29–56.
- POPOVA, T. I., AND E. D. VALTER. 1965. On the elucidation of the life cycle of the fish nematode *Contracaecum aduncum* (Rudolphi, 1802) Baylis, 1920 (Ascaridata). Mater. Nauchn. Konf. Vses. Obshch. Gel'mintol., Part 1: 175–178. (Transl. from Russian by Fish. Res. Board Can. Transl. Ser. No. 1797, 1971).
- PRICE, E. W. 1961. North American monogenetic trematodes IX. The families Mazocraeidae and Plectanocotylidae. Proc. Biol. Soc. Wash. 74: 127–156.
- PRIEBE, K. 1971. Die lebensmittelhygienische Bedeutung des Nematodenlarven-befalls bei Seefischen. Arch. Lebensmittelhyg. 22: 193–200. (Transl. from German by Fish. Mar. Serv. Transl. Ser. No. 361, 1976).
- PRIEBE, K., H. JENDRUSCH, AND U. HAUSTEDT. 1973. Problematik und Experimentaluntersuchungen zum erlöschen der einborhrpotenz von *Anisakis*-larven des Herings bei der herstellung von Kaltmarinaden. Arch. Lebensmittelhyg. 24: 217–222. (Transl. from German by Fish. Mar. Serv. Transl. Ser. No. 3303, 1974.)
- PUNT, A. 1942. Recherches sur quelques nématodes parasites de poissons de la mer du Nord. Mem. Mus. Hist. Nat. Belg., Mem. No. 98: 1–110.
- PUSHNIKOVA, G. M., AND V. V. PUSHNIKOV. 1981. Infection of herring with larval nematodes in the waters of Sakhalin. Biol. Morya, 1981, (Vladivostok.) 5: 71–73. (In Russian)
- REES, G. 1953. Some parasitic worms from fishes off the coast of Iceland III. Monogenea, Nematoda, Acanthocephala. Parasitology 43: 193–198.
- REIMER, L. W. 1962. Die digenetischen Trematoden der Wasservögel der Insel Hiddensee und ihre Larval-stadien aus den die Insel umgebenden Brackgewässern. Inaug. Diss., Greifswald. 128 p.
1970. Digene Trematoden und Cestoden der Ostseefische als natürliche Fischmarken. Parasitol. Schriftenr. 20: 3–144.
- REIMER, L. W., AND O. JESSEN. 1972. Parasitenbefall der Nordseeheringe. Angew. Parasitol. 13: 65–71.
- ROBINSON, D. G., W. E. BARRACLOUGH, AND J. D. FULTON. 1968a. Data record. Number, size composition, weight and food of larval and juvenile fish caught with a two-boat surface trawl in the Strait of Georgia May 1–4, 1967. Fish. Res. Board Can. MS Rep. 964: 105 p.
- 1968b. Data record. Number, size composition, weight and food of larval and juvenile fish caught with a two-boat surface trawl in the Strait of Georgia June 5–9, 1967. Fish. Res. Board Can. MS Rep. 972: 109 p.
- ROKICKI, J. 1972. Lawy *Anisakis* sp. u śledzi *Clupea harengus* L. w Bałtyku. Wiad. Parazytol. 18: 89–96.
1973. Helminths of certain Clupeidae mainly of the herring *Clupea harengus* L., in south Baltic. Acta Parasitol. Pol. 21: 443–464.
1975. Helminth fauna of fishes of the Gdańsk Bay (Baltic Sea). Acta Parasitol. Pol. 23: 37–84.
- ROOS, J. F., P. GILHOUSEN, S. R. KILICK, AND E. R. ZYBLUT. 1973. Parasitism on juvenile Pacific salmon (*Oncorhynchus*) and Pacific herring (*Clupea harengus pallasi*) in the Strait of Georgia by the river lamprey (*Lampetra ayresi*). J. Fish. Res. Board Can. 30: 565–568.
- ROSENTHAL, H. 1966. Parasites in herring larvae (Observations in aquaria). Int. Counc. Explor. Sea, C.M. 1966/H:2, Herring Comm.: 3 p.
1967. Parasites in larvae of herring (*Clupea harengus* L.) fed with wild plankton. Mar. Biol. (Berl.) 1: 10–15.
- ROSKAM, R. T. 1960. A human disease caused by a nematode from herring. Int. Counc. Explor. Sea, C.M. 1960, Herring Comm. No. 98: 4 p.
1966. *Anisakis* larvae in North Sea herring. Int. Counc. Explor. Sea, C.M. 1966/H:13, Herring Comm.: 3 p.
1967. *Anisakis* and *Contracaecum* larvae in North Sea herring. Int. Counc. Explor. Sea, C.M. 1967/H:19, Pelagic Fish (Northern) Comm.: 7 p.
- RUDOLPHI, C. A. 1802. Fortsetzung der Beobachtungen über die Einweidewürmer. Arch. Zool. u. Zoot. 2: 1–67; 3: 61–125.

1809. Entozoorum, seve vermium intestinalium historia naturalis. Vol. 2, Part 1. Amsterdam. 1–457.
1819. Entozoorum synopsis cui accedunt mantissa duplex et indices locupletissimi. S. A. Rucker, Berlin. 881 p.
- RUITENBERG, E. J. 1970. Anisakiasis. Pathogenesis, serodiagnosis and prevention. Thesis. Rijksuniversiteit, Utrecht. 138 p.
1971. Experimental anisakiasis in rabbits, p. 359–366. In S. M. Gaafar [ed.] Pathology of parasitic diseases. Purdue Univ. Studies, Lafayette, IN.
- RUITENBERG, E. J., J. M. BERKVENS, AND M. J. M. DUYZINGS. 1970. Experimentele Anisakis-marina-infecties bij het konijn. Ned. Tijdschr. Geneesk. 114: 1904–1909.
1971. Experimental *Anisakis marina* infections in rabbits. J. Comp. Pathol. 81: 157–163.
- RUITENBERG, E. J., AND R. T. ROSKAM. 1969. Eenvoudige controlemethode op de inboringscapaciteit van *Anisakis*-larven. Conserva 17: 236–237.
- SANO, M. 1966. Research on helminths in the human body carried by marine fish in Japan, and its progress. Shokuhin Eisei Kenkyu 16: 543–551. (In Japanese)
- SANZIN, W.-D. 1965. Untersuchungen zur Ökologie, Artspezifität und Variabilität von *Hexamita salmonis* Moore. Inaug. -Diss., Ludwig-Maximilians-Univ., Munich. 103 p.
- SCHNEIDER, G. 1902a. Über das Vorkommen von Larven des Bandwurms *Bothriotaenia proboscidea* Batsch im Magen und Darm von Ostseeheringen (*Clupea harengus membras* L.). Sitzungsber. Ges. Naturforsch. Freunde, Berlin 1: 28–30.
- 1902b. Ichthyologische Beiträge III. Über die in den Fischen des Finnischen Meerbusens vorkommenden Endoparasiten. Acta Soc. Fauna Flora Fenn. 22: 1–83.
1903. Beiträge zur Kenntnis der Helminthenfauna des Finnischen Meerbusens. Acta Soc. Fauna Flora Fenn. 26: 1–35.
- SCHULZ, H. P. 1974. *Anisakis* larvae from the herring: in vitro development and morphology. Proc. 3rd Int. Congr. Parasitol. 3: 1627.
- SCHULTZ, H. 1974. Investigations on the occurrence of nematodes in herring off the western British and Irish coasts. Int. Counc. Explor. Sea, C.M. 1974/H:24, Pelagic Fish (Northern) Comm.: 9 p.
- SCOTT, D. M. 1953. Experiments with the harbour seal, *Phoca vitulina*, a definitive host of a marine nematode, *Porrocaecum decipiens*. J. Fish. Res. Board Can. 10: 539–547.
- SCOTT, T. 1901. Notes on some parasites of fishes. 19th Annu. Rep. Fish. Board Scotland (for 1900), Part 3: 120–153.
- SHEENKO, P. S., AND S. E. POZDNYAKOV. 1981. Systematics and nomenclature of some nematodes of the genus *Contracaecum* sensu lato (Ascaridida: Anisakidae) of the northwestern Pacific, p. 82–85. In Yu. L. Mamaev and N. I. Demshin, [ed.] Biology and systematics of helminths of animals of the Far-East. Akad. Nauk SSSR; Dalnevost. Nauchn. Tsentr, Vladivostok, USSR. (In Russian)
- SHIRAKI, T. 1969. On the histopathological diagnosis of the disease caused by migrating larval nematodes in the gastro-intestinal tract (mainly disease caused by *Anisakis*). Saishin-Igaku 24: 378–389. (Transl. from Japanese by Fish. Res. Board Can. Transl. Ser. No. 1323, 1969).
1974. Larval nematodes of family Anisakidae (Nematoda) in the northern Sea of Japan — as a causative agent of eosinophilic phlegmone or granuloma in the human gastro-intestinal tract. Acta Med. Biol. (Niigata) 22: 57–98.
- SHULMAN, S. S. 1953. New and little known myxosporidians from the White Sea. Zool. Zh. 32: 384–393. (In Russian)
1956. Parasite fauna of *Clupea haengus*, *Osmerus eperlanus*, and *Eleginus navaga* in the White Sea. Tr. Karelo-Finsk. Fil. Akad. Nauk SSSR, Ser. Parasitol. 4: 50–67. (In Russian)
1966. Myxosporidia in the fauna of the USSR. Izdat. "Nauka," Moscow and Leningrad. 507 p. (In Russian)
- SHULMAN, S. S., AND R. E. SHULMAN-ALBOVA. 1953. Parasites of fishes of the White Sea. Izdat. Akad. Nauk SSSR, Moscow and Leningrad. 200 p. (In Russian)
- SHULMAN-ALBOVA, R. E. 1952. Parasites of fishes of Gridinsk Gulf of the White Sea. Part I. Uchen. Zap. Karelo-Finsk. Gos. Univ. 4: 78–97. (In Russian)
- SINDERMANN, C. J. 1957a. Diseases of fishes of the western North Atlantic. V. Parasites as indicators of herring movements. Maine Dep. Sea Shore Fish. Res. Bull. 27: 30 p.
- 1957b. Diseases of fishes of the western North Atlantic. VI. Geographic discontinuity of myxosporidiosis in immature herring from the Gulf of Maine. Maine Dep. Sea Shore Fish. Res. Bull. 29: 17 p.
1959. Zoogeography of sea herring parasites. J. Parasitol. 45 (4, Sect. 2): 34.
- 1961a. Parasite tags for marine fish. J. Wildl. Manage. 25: 41–47.
- 1961b. Sporozoan parasites of sea herring. J. Parasitol. 47 (4, Sect. 2): 34.
1963. Disease in marine populations. Trans. North Am. Wildl. Nat. Resour. Conf.: 336–356.
1965. Effects of environment on several diseases of herring from the western North Atlantic. Int. Comm. Northwest Atl. Fish. Spec. Publ. 6: 603–610.
1966. Larval ecology of the trematode *Cryptocotyle lingua*, p. 12–13. In A. Corradetti [ed.] Proc. 1st Int. Congr. Parasitol. Vol. 1. Pergamon press, Milano.
- SINDERMANN, C. J., AND A. E. FARRIN. 1962. Ecological studies of *Cryptocotyle lingua* (Trematoda: Heterophyidae) whose larvae cause "pigment spots" of marine fish. Ecology 43: 69–75.
- SINDERMANN, C., AND A. ROSENFIELD. 1954a. Diseases of fishes of the western North Atlantic I. Diseases of the sea herring (*Clupea harengus*). Maine Dep. Sea Shore Fish. Res. Bull. 18: 23 p.
- 1954b. Diseases of fishes of the western North Atlantic III. Mortalities of sea herring (*Clupea harengus*) caused by larval trematode invasion. Maine Dep. Sea Shore Fish. Res. Bull. 21: 16 p.
- SINDERMANN, C. J., AND L. W. SCATTERGOOD. 1954. Diseases of fishes of the western North Altantic II. *Ichthyosporidium* disease of the sea herring (*Clupea harengus*). Maine Dep. Sea Shore Fish. Res. Bull. 19: 40 p.
- SIÖBLOM, V., AND E. KUUTTINEN. 1976. *Phocascaris* sp. (Nematoda) larvae in Baltic herring, a new parasite for the Baltic Sea. Finn. Fish. Res. 2: 1–3.
- 1978a. Further finds of *Phocascaris* sp. (Nematoda) in Baltic herring off the coast of Finland. Int. Counc. Explor. Sea., C.M. 1978/J:19, Baltic Fish Comm.: 7 p.
- 1978b. Further finds of larvae of *Phocascaris* sp. (Nematoda) in Baltic herring off the coast of Finland. Finn. Fish. Res. 2: 19–21.
- SKRJABINA, E. S. 1963. On the helminth fauna of marine fish of Kamchatka. Tr. Gel'mintol. Lab. Akad. Nauk SSSR 13: 313–329. (In Russian)
- SLUITERS, J. F. 1973. Study on the life-cycle of *Anisakis* sp. Parasitology 67: xxviii.
1974. *Anisakis* sp. larvae in the stomachs of herrings (*Clupea harengus* L.). Z. Parasitenkd. 44: 279–288.
- SMEDLEY, E. M. 1934. Some parasitic nematodes from Canadian fishes. J. Helminthol. 12: 205–220.
- SMIRNOVA, K. V. 1957. On the parasite fauna of *Clupeonella delicatula tscharcalensis* and some other fishes of lake Charkhal. Izv. Vses. Nauchno-Issled. Inst. Ozern. Rechn. Rybn. Khoz. 42: 332–333. (Transl. from Russian by Israel Prog. Sci. Transl. Jerusalem, 1961.)
- SMITH, J. W. 1974. Experimental transfer of *Anisakis* sp. larvae (Nematoda: Ascaridida) from one fish host to another. J. Helminthol. 48: 229–234.
- SMITH, J. W., AND R. WOOTTEN. 1975. Experimental studies on the migration of *Anisakis* sp. larvae (Nematoda: Ascaridida) into the flesh of herring, *Clupea harengus* L. Int. J. Parasitol. 5: 133–136.
- STAFFORD, J. 1904. Trematodes from Canadian fishes. Zool. Anz. 27: 481–495.
1907. Preliminary report on the trematodes of Canadian marine fishes. Contrib. Can. Biol., 1901–05: 91–94.
- STANKIEWICZ, E. B. 1971. Paszozytnicze larwy nicieni *Anisakis marina* (L. 1767) w aspekcie schorzeń człowieka. Med. Weter. 27: 419–421.
- STRELKOV, YU. A. 1960. Endoparasitic worms of marine fishes of northern Kamchatka. Tr. Zool. Inst. Akad. Nauk SSSR 28: 147–196. (In Russian)
- STRZYZEWSKA, K., AND J. POPIEL. 1974. Characteristic of herring infested by *Anisakis* larvae caught in the Gulf of Gdańsk. Int. Counc. Explor. Sea, C.M. 1974/H:15, Pelagic Fish (Northern) Comm.: 4 p.
- SVETOVIDOV, A. N. 1961. On the origin of certain amphiboreal gadoid and clupeoid fishes in the light of the species composition of their parasites. Zool. Zh. 40: 1335–1344. (Transl. from Russian by Fish. Res. Board Can. Transl. Ser. No. 593, 1965)
- SVETOVIDOV, A. N., AND S. S. SHULMAN. 1960. An association between the species composition of the parasites of certain groups of fishes and parallels in their ecology and distribution. Zool. Zh. 39: 1866–1870. (Transl. from Russian by Fish. Res. Board Can. Transl. Ser. No. 592, 1965)
- THÉLOHAN, P. 1892. Sur quelques coccidies nouvelles. Parasites des poissons. J. Anat. Physiol. 28: 151–162.
1894. Nouvelles recherches sur les coccidies. Arch. Zool. Exp. Gen. 2, Ser. 3: 541–573.

- THIEL, P. H. VAN. 1967. Le phénomène de la "formation de la calotte" chez la larve d'*Anisakis marina* (Linnaeus 1767) dans du sérum humain. *Ann. Parasitol. Hum. Comp.* 42: 633-642.
- THIEL, P. H. VAN, F. C. KUIPERS, AND R. T. ROSKAM. 1960. A nematode parasitic to herring, causing acute abdominal syndromes in man. *Trop. Geogr. Med.* 2: 97-113.
- TYZZER, E. E. 1900. Tumors and Sporozoa in fishes. *J. Boston Soc. Med. Sci.* 5: 63-68.
- VALTONEN, E. T. 1979. *Corynosoma* species in the fish of the Bothnian Bay, p. 34-35. In G. Bylund [ed.] *Proc. 9th Symp. Scandinavian Soc. Parasitol.*, Abo Akad., Finland No. 15. Inform Inst. Parasitol., Turku.
1980. On the occurrence of acanthocephalans in the fish of the Bothnian Bay. *Proc. 3rd Eur. Multicoll. Parasitol.*: p. 102.
- VAN CLEAVE, H. J. 1924. A critical study of the Acanthocephala described and identified by Joseph Leidy. *Proc. Acad. Nat. Sci. Philadelphia* 73: 279-334.
- VERVOORT, W. 1964. Notes on Bomolochidae (Copepoda), I. A redescription of *Parabomolochus cuneatus* (Fraser, 1920) and notes on its synonymy. *Crustaceana* (Leiden) 6: 291-302.
- VIK, R. 1966. *Anisakis* larvae in Norwegian food fishes, p. 568-569. In A. Corradetti [ed.] *Proc. 1st Int. Congr. Parasitol.* Vol. 1. Pergamon Press, Milano.
- WALCH, J. E. J. 1778. Beiträge zur Insecten-Geschichte. Drittes Stück. Der Naturforscher, Halle, 12: 56-67.
- WALTON, A. C. 1928. A revision of the nematodes of the Leidy collections. *Proc. Acad. Nat. Sci. Philadelphia* 79 (for 1927): 49-163.
- WARD, H. B., AND J. FILLINGHAM. 1934. A new trematode in a toadfish from southeastern Alaska. *Proc. Helminthol. Soc. Wash.* 1: 25-31.
- WILLEMSE, J. J. 1968. Helminth and sporozoan parasites of fishes in the Netherlands. *Bull. Zool. Mus. Univ. Amsterdam* 1: 83-87.
- WÜLKER, G. 1930. Über Nematoden aus Nordseetieren. II. *Zool. Anz.* 88: 1-16. (Transl. from German by Fish. Res. Board Can. Transl. Ser. No. 2479, 1973)
- YAMAGUCHI, T., N. KUDO, S. KAWADA, Y. NAKADE, AND N. TAKADA. 1968. Studies on larva migrans. (24). The incidence of infection of *Anisakis* larvae in marine fishes. *Jpn. J. Parasitol.* 17: 262. (In Japanese)
- YAMAGUTI, S. 1963. *Systema helminthum*. V. *Acanthocephala*. Interscience, New York, N.Y. 432 p.
1971. *Synopsis of digenetic trematodes of vertebrates*. Vol. 1. Keigaku Publ. Co., Tokyo. 1074 p.
- YOSHIMURA, H. 1966. Migrations of *Anisakis*-like larvae into the human alimentary tract with special reference to clinical pathology. *Nihon Iji Shinpo* 2204: 10-16.
- YOUNG, P. C., AND D. LOWE. 1969. Larval nematodes from fish of the subfamily Anisakinae and gastro-intestinal lesions in mammals. *J. Comp. Pathol.* 79: 301-313.
- ZEDER, J. G.-H. 1800. Erster Nachtrag zur Naturgeschichte der Eingeiedewürmer von Johann August Ephraim Goeze mit Zusätzen und Anmerkungen herausgegeben von D. Johann Georg Heinrich Zeder. Siegfried Lebrecht Crusius, Leipzig. 320 p.
- ZHUKOV, E. V. 1960a. Parasite fauna of fish in Chukotka. I. Monogenetic trematodes of marine and freshwater fish. *Parazitol. Sb.* 19: 308-332. (In Russian)
- 1960b. Endoparasitic worms of fishes from the Sea of Japan and South-Kurile shallow-waters. *Tr. Zool. Inst. Akad. Nauk SSSR* 28: 3-146. (In Russian, English summary)
1963. On the parasite fauna of fishes of the Chukotsk Peninsula and adjoining seas. II. Endoparasitic worms of marine and freshwater fishes. *Parazitol. Sb.* 21: 96-139. (In Russian, English summary)
1964. On the parasite fauna of fishes of the Chukotsk Peninsula and the adjoining seas. III. The parasitic Protozoa of marine and freshwater fishes. *Parazitol. Sb.* 22: 224-262. (In Russian, English summary)

QL 626 C314 no.70 c.2
Arthur, J.R.
Annotated checklist and
bibliography of parasite...
69952 12043041 c.2

