

# **Photo-identification Catalogue and Status of the Northern Resident Killer Whale Population in 2014.**

J.R. Towers, G.M. Ellis and J.K.B. Ford

Fisheries and Oceans Canada  
Science Branch, Pacific Region  
Pacific Biological Station  
3190 Hammond Bay Road  
Nanaimo, BC  
V6T 6N7

2015

## **Canadian Technical Report of Fisheries and Aquatic Sciences 3139**

## **Canadian Technical Report of Fisheries and Aquatic Sciences**

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of Fisheries and Oceans Canada, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in the data base *Aquatic Sciences and Fisheries Abstracts*.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

## **Rapport technique canadien des sciences halieutiques et aquatiques**

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais qui ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques de Pêches et Océans Canada, c'est-à-dire les sciences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications à part entière. Le titre exact figure au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la base de données *Résumés des sciences aquatiques et halieutiques*.

Les rapports techniques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre.

Les numéros 1 à 456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Canadian Technical Report of  
Fisheries and Aquatic Sciences 3139

2015

PHOTO-IDENTIFICATION CATALOGUE AND STATUS OF THE NORTHERN  
RESIDENT KILLER WHALE POPULATION IN 2014.

by

J.R. Towers, G.M. Ellis and J.K.B Ford

Fisheries and Oceans Canada  
Science Branch, Pacific Region  
Pacific Biological Station  
3190 Hammond Bay Road  
Nanaimo, BC  
V9T 6N7

E-mail: [Jared.Towers@dfo-mpo.gc.ca](mailto:Jared.Towers@dfo-mpo.gc.ca)

© Her Majesty the Queen in Right of Canada, 2015  
Cat. No. Fs97-6/3139E-PDF ISBN 978-0-660-03209-2 ISSN 1488-5379

Correct citation for this publication:

Towers, J.R., Ellis, G.M. and Ford, J.K.B. 2015. Photo-identification catalogue and status of the northern resident killer whale population in 2014. Can. Tech. Rep. Fish. Aquat. Sci. 3139: iv + 75 p.



## TABLE OF CONTENTS

|   |    |
|---|----|
| ABSTRACT.....   | iv |
| RÉSUMÉ .....  | iv |
| 1.0 INTRODUCTION .....  | 1  |
| 2.0 MATERIALS AND METHODS .....   | 3  |
| 2.1 DATA COLLECTION AND ANALYSIS .....                                  | 3  |
| 2.2 DATA PRESENTATION .....   | 3  |
| 3.0 RESULTS AND DISCUSSION .....  | 4  |
| 4.0 ACKNOWLEDGEMENTS .....  | 7  |
| 5.0 REFERENCES .....  | 7  |
| APPENDIX.....   | 14 |
| PHOTO-IDENTIFICATION CATALOGUE OF NORTHERN RESIDENT KILLER WHALES ..... | 14 |

## **ABSTRACT**

Towers, J.R., Ellis, G.M. and Ford, J.K.B. 2015. Photo-identification catalogue and status of the northern resident killer whale population in 2014. Can. Tech. Rep. Fish. Aquat. Sci. 3139: iv + 75 p.

Field studies of the life history and ecology of killer whale populations off Canada's west coast have been conducted annually since 1973. These studies are based on the identification of individual whales from photographs of permanent, natural markings. In this report, we summarize abundance trends in the northern resident killer whale population between 1974 and 2014, and provide an updated photo-identification catalogue of individuals. In 2014, the northern resident population was composed of 290 whales. The population has grown at an average mean annual rate of 2.2% since 1974. Continued population monitoring by photo-identification is a key research activity in the recovery strategy for this threatened population.

## **RÉSUMÉ**

Towers, J. R., Ellis, G. M. et Ford, J. K. B. 2015. Catalogue de photographies d'identification et état de la population d'épaulards résidents du nord en 2014. Rapp. tech. can. sci. halieut. aquat. 3139: iv + 75 p.

Depuis 1973, des études sur le terrain sont menées chaque année afin d'observer le cycle de vie et les caractéristiques écologiques des populations d'épaulards qui vivent le long de la côte ouest du Canada. Ces études reposent sur l'identification d'individus à partir de photographies de leurs marques naturelles permanentes. Dans le présent rapport, nous présentons la courbe évolutive de la population d'épaulards résidents du nord de 1974 à 2014, ainsi qu'une mise à jour du catalogue de photographies d'identification des individus qui forment cette population. En 2014, la population d'épaulards résidents du nord était composée de 290 individus. Depuis 1974, la croissance de la population a suivi un taux annuel moyen de 2,2 %. La surveillance continue à partir de photographies d'identification est une activité de recherche essentielle à la stratégie de rétablissement de cette population menacée.

## 1.0 INTRODUCTION

Killer whales off the west coast of Canada were among the first cetaceans found to be individually recognizable from photographs of natural markings (Bigg et al. 1976). The use of photo-identification to study killer whale populations in British Columbia began in 1973 and continues to the present (Ford 2011). Data collected using this method have been fundamental to the description of killer whale populations in the region and assessments of their abundance, life history traits and social organization (Bigg 1982; Bigg et al. 1990; Olesiuk et al. 1990, 2005; Ford and Ellis 1999; Ford et al. 1994, 2000, 2007, 2013, 2014; Williams and Lusseau 2006; Ellis et al. 2007, 2008, 2011; Foster et al. 2012; Towers et al. 2012a). The photo-identification technique has also facilitated a wide range of studies on the behaviour (e.g., Jacobsen 1986; Morton 1990; Baird and Dill 1995), acoustics (Ford and Fisher 1983; Ford 1991; Barrett-Lennard et al. 1996; Deecke et al. 2010), foraging ecology (Nichol and Shackleton 1996; Ford et al. 1998, 2005, 2010a,b; Ford and Ellis 2006) and genetics (Barrett-Lennard 2000) of killer whale populations in British Columbia.

Three separate ecotypes of killer whales have been found to share coastal waters off the west coast of Canada – residents, Bigg's (transients) and offshores. These ecotypes do not mix despite living in the same habitats, and as a result they are reproductively isolated and genetically distinct (Ford et al. 2000; Krahn et al. 2007a; Ford 2014). Each ecotype specializes on different prey resources. Resident killer whales feed on fish, with a strong preference for salmon, especially Chinook (Ford and Ellis 2006; Ford et al. 2010b). Bigg's killer whales prey on marine mammals (Ford et al. 1998) and offshore killer whales feed on sharks and other large fish species (Ford et al. 2011, 2014).

Resident killer whales in British Columbia are represented by two distinct populations, northern and southern, which have overlapping ranges but do not intermingle. The northern resident population, which is the subject of this report, ranges between southern Washington and southeastern Alaska (Dahlheim et al. 1997; Ford et al. 2000; Ellis et al. 2011). Within this overall range, the majority of encounters with northern resident killer whales have taken place in the coastal waters of British Columbia (Ford 2006; Ellis et al. 2011). The smaller southern resident killer whale population ranges from central California to southeastern Alaska (Ford 2012), but is found most frequently during summer and fall in the transboundary waters of the Salish Sea, between northern Washington and southern Vancouver Island (Ford et al. 1994).

Resident killer whales travel in family groups that remain cohesive over long periods of time (Bigg et al. 1987). These social groups, referred to as pods (Bigg et al. 1976), subpods (Bigg et al. 1987) or matriline (Ford et al. 2000) are typically composed of one or more maternally-related reproductive females and their offspring. Maternally-related individuals that have descended from a common ancestor share a common

acoustic dialect and are thus referred to as belonging to the same clan (Ford 1991). Different clans have no calls in common and sound distinct. The northern resident population is composed of 3 clans – A, G and R. Southern resident killer whales all belong to J clan (Bigg et al. 1987).

Killer whales from both the northern and southern resident populations were exploited in British Columbia and Washington by a live-capture fishery that began in 1964 to supply demand for killer whales in aquariums. At least fifty individuals were removed from the northern and southern resident populations in the following decade (Bigg and Wolman, 1975; Bigg et al. 1976; Ford et al. 1994). Many other individuals were caught and released, sometimes on more than one occasion. Removals by this fishery and mortalities likely due to intentional shootings resulted in depleted resident killer whale populations by the early 1970s. Preliminary results of photo-identification studies by Bigg et al. (1976) showed that resident killer whale population sizes were too small to sustain this fishery. Combined with public opposition to the captures, this led to the end of the fishery in 1976 (Ford 2011).

Annual censuses of resident killer whales in both British Columbia and Washington have been undertaken each year since 1974, coordinated by researchers with the Pacific Biological Station, Nanaimo, BC (primarily northern residents) and the Center for Whale Research, Friday Harbor, WA (southern residents). Both resident populations showed a long-term positive growth trend between 1974 and the mid 1990s, increasing at average rates of 2.5 to 3% per annum (Bigg 1982; Olesiuk et al. 1990, 2005). Northern and southern resident populations both experienced sharp increases in mortality rates and a decline in abundance during the latter half of the 1990s and early 2000s, which coincided with a coast-wide decline in availability of their primary prey, Chinook salmon (Ford et al. 2005, 2010a). In 2001, the conservation status of killer whales off the west coast was assessed by the Committee on the Status of Endangered Wildlife in Canada (COSEWIC). The northern residents were designated as threatened and southern residents as endangered due to their small population sizes, low reproductive rate and the existence of a variety of anthropogenic threats (Ford 2006; COSEWIC 2008; Fisheries and Oceans Canada 2011). These populations were subsequently legally listed as such under Canada's Species at Risk Act (SARA), and the process of recovery strategy development and identification of critical habitat was initiated (Fisheries and Oceans Canada 2011). Effort to annually census resident killer whales by photo-identification is recognized in the resident killer whale recovery strategy as an integral component of on-going population monitoring in Canada.

Results of the annual photo-identification censuses of resident killer whales have been presented in various scientific reports as well as catalogues intended for use by managers, field researchers, the whale-watch community and the interested public. Technical reports describing the population status and dynamics of resident killer whales include Bigg et al. (1976, 1990), Bigg (1982), Balcomb et al. (1980, 1982), Olesiuk et al. (1990, 2005), Ford et al. (2005, 2010b), and Ellis et al. (2011). Photo-

identification catalogues for northern and southern resident populations were provided in Bigg et al. (1987), and Ford et al. (1994, 2000). Updated catalogues of southern and northern residents have been periodically provided by the Center for Whale Research (whaleresearch.com) and the Cetacean Research Program at the Pacific Biological Station respectively (Ellis et al. 2007; 2011).

In this report, we present a summary of the current population status of the northern resident killer whale population through 2014, and an updated photo-identification catalogue of individuals. This catalogue includes demographic and genealogical data for all whales in the northern resident population. These data were gathered from annual photo-identification population censuses and associated long-term visual monitoring. This catalogue is intended to provide colleagues with a tool to facilitate their work. Furthermore, it is hoped that it will enrich observations of this population by marine wildlife viewers off the west coasts of Canada and the US.

## **2.0 MATERIALS AND METHODS**

### **2.1 DATA COLLECTION AND ANALYSIS**

Methods for the collection of identification photos of killer whales were first described in Bigg et al. (1976, 1986). Details outlining the use of updated field methods and equipment are described in Ellis et al. (2011). Photo-identification data analysis and management techniques are provided in Towers et al. (2012b).

### **2.2 DATA PRESENTATION**

The appendix includes identification photographs of whales known or presumed to be alive at the end of the 2014 field season. In most cases, individuals laid out in clusters represent socially cohesive groups. However, some maternally-related individuals (primarily adult females with their own offspring, orphans and mature bulls without living mothers) may or may not exhibit social cohesion with others in their matriline (Ford and Ellis 2002; Ford 2006; DFO unpubl. data). Matrilines are typically named after the eldest living reproductive female within a group of maternally-related individuals, although if she has a surviving brother or uncle, the matriline is named after the deceased mother of the eldest male. Pod names are also provided and refer to the most distinctive whale documented in each social group when they were first identified in the early 1970s. Tabs are provided on the outside edge of each page to show the acoustic clan to which each group belongs (see introduction).


The appendix also includes schematic diagrams of the different northern resident killer whale lineages. The structure of each lineage is portrayed to represent matrilineal-based genealogical relationships that have been inferred from long-term observations of social organization (Bigg et al. 1987, 1990; Ford et al. 1994, 2000; Ford and Ellis 2002; Ellis et al. 2011). Diagrams include every individual documented over the course of the

study. Clear boxes represent individuals known or presumed to be alive in 2014 and shaded boxes represent deceased individuals (see key on page 15). Lines linking mothers and offspring are solid if the relationship is positive (i.e., known with certainty). This includes individuals that have been documented since birth. Relationships of whales born before the study began in the early 1970s are either probable, indicated by a dashed line, or possible, indicated by a dotted line.

### **3.0 RESULTS AND DISCUSSION**


Photo-identification censuses of the northern resident killer whale population have been conducted annually since 1974. The last population update and associated photo-identification catalogue was based on the 2010 field season and published in Ellis et al. (2011). During 2011–2014, we travelled up to 10,000 km per year in search of northern residents and averaged 58 encounters per year (range: 52-67) with this population. This is roughly consistent with the effort and number of encounters with northern residents during previous years (Bigg 1982; Ellis et al. 2011). Between 2011 and 2014 we analyzed a total of 28,012 identification images of whales in this population.

Although the majority of whales were photographed each year, it was not always possible to locate every animal during each field season. As a result, there was occasionally some uncertainty in the number of animals alive each year. For example, 33 individuals from the northern resident population could not be accounted for in 2014. The status of these individuals was confirmed when 28 of them were photo-identified in 2015 and five were not. Three other animals that disappeared from their matriline part way through 2014 are included in the population count but identification images of them have been excluded from the appendix because they are now confirmed to be dead. In total, the northern resident killer whale population numbered 290 individuals in 2014.



*Figure 1. Abundance trend of the northern resident killer whale population, 1974–2014. In years with uncertainty, the minimum and maximum population sizes are represented with shading.*

The abundance trend of the northern resident population during 1974–2014 is shown in Figure 1. From the mid 1970s to mid 1990s, the population grew steadily at an average mean annual rate of 2.6%, from approximately 122 individuals in 1974 to 218 in 1997. The population then declined by about 7% during 1998–2001, a period that coincided with a significant reduction in the availability of the whales’ primary prey, Chinook salmon (Ford et al. 2010b). Starting in 2002, the growth trend became positive once again, with the population increasing at an average mean rate of 2.9% per year until 2014 (range = -0.4 – 8.6% per annum). This represents an average annual mean increase of 2.2% over the 40 year time series. All three northern resident clans experienced growth during 1974–2014, but at different rates (Figure 2). The G clan grew at the greatest average mean rate of 2.9% per annum followed by R clan at 2.1% per annum, and A clan at 1.9% per annum. By contrast, the southern resident population (J clan) has not maintained similar growth over the same time period (Bigg et al. 1976; Ellis et al. 2011; Cogan 2014).



*Figure 2. Abundance trends of A, G and R clans, 1974–2014. In years with uncertainty, the minimum and maximum population sizes are represented with shading.*

Several factors may influence population trends for resident killer whales. Among these, prey availability (Ford et al. 2005, 2010a,b; Ayres et al. 2012), disturbance (Erbe 2002; Morton and Symonds 2002; Williams et al. 2002a,b; Williams and Ashe 2007; Lusseau et al. 2009) and pollution (Ross et al. 2000; Rayne et al. 2004; Ross 2006; Krahn et al. 2007b; 2009; Lachmuth et al. 2010, 2011; Williams et al. 2014) are of primary conservation concern (Fisheries and Oceans Canada 2011). Several other threats such as vessel strikes (Williams and O'Hara 2009), entanglement (Bigg and Wolman 1975; Fisheries and Oceans Canada 2011) and intentional shootings also have the potential to directly affect mortality rates of the northern and southern resident killer whale populations.

With each year of field research, our understanding of the distribution, demographics, behaviour and population trends of northern resident killer whales improves, as does our ability to detect subtle changes in population parameters that may result from anthropogenic threats. On-going photo-identification monitoring will continue to be a key component of recovery efforts focused on the northern resident population and other communities of killer whales that live off the west coast of North America.



## 4.0 ACKNOWLEDGEMENTS

During the course of this long-term study, numerous people, especially research colleagues and marine tourism operators, have shared data and provided logistical support essential to our work. We thank them all. We also thank the following individuals who have provided images used in this report. They are: Robin Abernethy, Lance Barrett-Lennard, Archie Dundas, Brian Falconer, Brian Gisborne, Mark Malleson, James Pilkington, Nick Sinclair and Eva Stredulinsky. Other individuals who have contributed to this work in recent years include: Adrienne Akmajian, Erin Ashe, Ken Balcomb, Caitlin Birdsall, Jim and Mary Borrowman, Kate Brauer, Troy Bright, Randy Burke, John Calambokidis, Miray Campbell, Ed Chutter, Volker Deecke, John Durban, Dave Ellifrit, Rachel Elliott, Holly Fearnbach, Bev Ford, Marie Fournier, Wayne Garton, Karen Hansen, Jackie Hilderling, Stan Hutchings, Andrew Jones, Bill and Donna Mackay, Susan MacKay, Dena Matkin, Yvonne Maximchuk, Roger McDonell, Christie McMillan, Hermann Mueter, Linda Nichol, Stef Olcen, Rod Palm, Bruce Paterson, Kathy Peavey, Leah Robinson, Nicole Robinson, Angela Smith, Lisa Spaven, Paul Spong, Jack Springer, Jan Straley, Helena Symonds, Dave and Maureen Towers, Jane Watson, Aaron Webber, Rob Williams, Janie Wray and Brianna Wright. We are also grateful to our colleagues with the Canadian Coast Guard for their hospitality and efforts in the field. Additionally, we appreciate support and cooperation from the following organizations: BC Cetacean Sightings Network, BC Marine Mammal Response Network, Cascadia Research Collective, Center for Whale Research, Cetacealab, Cetus Research and Conservation Society, Gitga'at First Nation, Langara Fishing Lodge, MERS Marine Education and Research Society, Marine Mammal Research Unit, North Island Marine Mammal Stewardship Association, Orcalab, Raincoast Conservation Society, Salmon Coast Field Station and the Vancouver Aquarium. Finally, we thank Brianna Wright for conducting a thorough review of this report and preparing the figures. Since 2001, major funding for this research has been provided by the Species at Risk program of Fisheries and Oceans Canada.

## 5.0 REFERENCES

- Ayres, K.L., Booth, R.K., Hemplemann, J.A., Koski, K.L., Emmons, C.K., Baird, R.W., Balcomb-Bartok, K., Hanson, B.M., Ford, M.J., and Wasser, S.K. 2012. Distinguishing the impacts of inadequate prey and vessel traffic on an endangered killer whale (*Orcinus orca*) population. PLoS ONE. 7(6): e36842. doi: 10.1371/journal.pone.0036842
- Baird, R.W. and Dill, L.M. 1995. Occurrence and behaviour of transient killer whales: seasonal and pod-specific variability, foraging behaviour, and prey handling. Can. J. Zool. 73(7): 1300-1311.

- Balcomb, K.C.III, Boran, J.R. and Heimlich, S.L. 1982. Killer whales in greater Puget Sound. Rep. Int. Whal. Commn. 32: 681-685.
- Balcomb, K.C.III, Boran, J.R., Osborne, R.W. and Haenel, N.J. 1980. Observations of killer whales (*Orcinus orca*) in greater Puget Sound, State of Washington. Report MM1300731-7, U.S. Marine Mammal Commission, Washington, D.C.
- Barrett-Lennard, L.G. 2000. Population structure and mating patterns of killer whales (*Orcinus orca*) as revealed by DNA analysis. Ph.D. Thesis, University of British Columbia, Vancouver, BC. xi + 97 p.
- Barrett-Lennard, L.G., J.K.B. Ford and K. Heise. 1996. The mixed blessing of echolocation: Differences in sonar use by fish-eating and mammal-eating killer whales. Anim. Behav. 51: 553-565.
- Bigg, M.A. 1982. An assessment of killer whale (*Orcinus orca*) stocks off Vancouver Island, British Columbia. Rep. Int. Whal. Commn. 32: 655-666.
- Bigg, M.A. and Wolman, A.A. 1975. Live-capture killer whale (*Orcinus orca*) fishery, British Columbia and Washington, 1962-73. J. Fish Res. Bd Can. 32(7): 1213-1221.
- Bigg, M.A., Ellis, G.M. and Balcomb, K.C. 1986. The photographic identification of individual cetaceans. Whalewatcher J. Amer Cetacean Soc. 20(2): 10-12.
- Bigg, M.A., Ellis, G.M., Ford, J.K.B. and Balcomb, K.C. 1987. Killer Whales: A Study of their Identification, Genealogy and Natural History in British Columbia and Washington State. Phantom Press and Publishers, Nanaimo, BC. 79 p.
- Bigg, M.A., MacAskie, I.B. and Ellis, G. 1976. Abundance and movements of killer whales off eastern and southern Vancouver Island with comments on management. Unpubl. Rep., Arctic Biological Station, Dept. of Fisheries and Environment, Ste Anne-de-Bellevue, Quebec.
- Bigg, M.A., Olesiuk, P.F., Ellis, G.M., Ford, J.K.B. and Balcomb, K.C.III. 1990. Social organization and genealogy of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. Rep. Int. Whal. Commn. Special Issue 12: 383-405.
- Cogan, J. 2014. Southern Resident State of the Population – 2014. Center for Whale Research. Friday Harbour, WA. 13 p.
- COSEWIC. 2008. COSEWIC assessment and update status report on the Killer Whale *Orcinus orca*, Southern Resident population, Northern Resident population,

West Coast Transient population, Offshore population and Northwest Atlantic / Eastern Arctic population, in Canada. Committee on the Status of Endangered Wildlife in Canada. Ottawa. viii + 65 pp.  
([www.sararegistry.gc.ca/status/status\\_e.cfm](http://www.sararegistry.gc.ca/status/status_e.cfm)).

- Dahlheim, M., Ellifrit, D.K. and Swenson, J.D. 1997. Killer whales of southeast Alaska: a catalogue of photo-identified individuals. National Marine Mammal Laboratory, National Marine Fisheries Service, U.S. Department of Commerce, Seattle, Washington.
- Deecke, V.B., Barrett-Lennard, L.G., Spong, P. and Ford, J.K.B. 2010. The structure of stereotyped calls reflects kinship and social affiliation in resident killer whales (*Orcinus orca*). *Naturwissenschaften*. DOI 10.1007/s00114-010-0657-z.
- Ellis, G.M., Ford, J.K.B. and Towers, J.R., 2007. Northern Resident Killer Whales in British Columbia: Photo-Identification Catalogue 2007. Pacific Biological Station, Nanaimo. ii + 38 p.
- Ellis, G.M., Towers, J.R. and Ford, J.K.B. 2008. Transient Killer Whales of British Columbia and Southeast Alaska: Photo-Identification Catalogue 2008. Pacific Biological Station, Nanaimo. 48 p.
- Ellis, G.M., Towers, J.R. and Ford, J.K.B. 2011. Northern resident killer whales of British Columbia: Photo-identification catalogue and population status to 2010. *Can. Tech. Rep. Fish. Aquat. Sci.* 2942: v + 71 p.
- Erbe, C. 2002. Underwater noise of whale-watching boats and potential effects on killer whales (*Orcinus orca*), based on an acoustic impact model. *Mar. Mamm. Sci.* 18:394-418.
- Fisheries and Oceans Canada. 2011. Recovery Strategy for the Northern and Southern Resident Killer Whales (*Orcinus orca*) in Canada. Species at Risk Act Recovery Strategy Series, Fisheries & Oceans Canada, Ottawa, ix + 80 p.
- Ford, J.K.B. 1991. Vocal traditions among resident killer whales (*Orcinus orca*) in coastal waters of British Columbia. *Can. J. Zool.* 69: 1454-1483.
- Ford, J.K.B. 2006. An assessment of critical habitats of resident killer whales in waters off the Pacific coast of Canada. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2006/072. iv + 34 p.
- Ford, J.K.B. 2011. Killer whales of the Pacific northwest coast: From pest to paragon. *Whalewatcher J. Amer. Cetacean Soc.* 40(1): 15-23.

- Ford, J.K.B. 2012. Assessment of Potential Food Limitation in Resident Killer Whales: How, When and Where. In Evaluating the Effects of Salmon Fisheries on Southern Resident Killer Whales: Workshop 2, March 13-15, 2012. NOAA Fisheries and DFO (Fisheries and Oceans Canada), Vancouver, BC.
- Ford, J.K.B. 2014. Marine Mammals of British Columbia. Royal BC Museum Handbook. Royal British Columbia Museum. Victoria, BC. 464 p.
- Ford, J.K.B. and G.M. Ellis. 1999. Transients: Mammal-Hunting Killer Whales. UBC Press, Vancouver, British Columbia. 96 p.
- Ford, J.K.B. and Ellis, G.M. 2002. Reassessing the social organization of resident killer whales in British Columbia. Pages 72-75 in the Fourth International Orca Symposium and Workshop, September 23-28, 2002. CEBC-CNRS, France.
- Ford, J.K.B. and Ellis, G.M. 2005. Prey selection and food sharing by fish-eating 'resident' killer whales (*Orcinus orca*) in British Columbia. DFO Can. Sci. Advis. Sec. Res. Doc. 2005/041. ii + 30 p.
- Ford, J.K.B. and Ellis, G.M. 2006. Selective foraging by fish-eating killer whales *Orcinus orca* in British Columbia. Mar. Ecol. Prog. Ser. 316: 185-199.
- Ford, J.K.B., and Fisher, H.D. 1983. Group-specific dialects of killer whales (*Orcinus orca*) in British Columbia. pp. 129-61. In: R. Payne (ed.) Communications and Behaviour of whales. AAAS Selected Symposium 76. Boulder, Co. 643 p.
- Ford, J.K.B., Ellis, G.M. and Balcomb, K.C. 1994. Killer Whales: The Natural History and Genealogy of *Orcinus orca* in British Columbia and Washington State. UBC Press and University of Washington Press, Vancouver, BC and Seattle, WA. 102 p.
- Ford, J.K.B., Ellis, G.M. and Balcomb, K.C. 2000. Killer Whales: The Natural History and Genealogy of *Orcinus orca* in British Columbia and Washington, Second edition. UBC Press and University of Washington Press, Vancouver, BC and Seattle, WA. 104 p.
- Ford, J.K.B., Ellis, G.M. and Durban, J.W. 2007. An assessment of the potential for recovery of West Coast Transient Killer Whales using coastal waters of British Columbia. DFO Can. Sci. Advis. Sec. Res. Doc. 2007/088.
- Ford, J.K.B., Ellis, G.M. and Olesiuk, P.F. 2005. Linking prey and population dynamics: did food limitation cause recent declines of 'resident' killer whales (*Orcinus orca*) in British Columbia? DFO Can. Sci. Advis. Sec. Res. Doc. 2005/042.

- Ford, J.K.B., Ellis, G.M., Barrett-Lennard, L.G., Morton, A.B., Palm, R.S. and Balcomb, K.C.III. 1998. Dietary specialization in two sympatric populations of killer whales (*Orcinus orca*) in coastal British Columbia and adjacent waters. *Can. J. Zool.* 76: 1456-1471.
- Ford, J.K.B., Ellis, G.M., Matkin, C.O., Wetklo, M.H., Barrett-Lennard, L.G. and Withler, R.E. 2011. Shark predation and tooth wear in a population of northeastern Pacific killer whales. *Aquat. Biol.* 11: 213-224.
- Ford, J.K.B., Ellis, G.M., Olesiuk, P.F. and Balcomb, K.C. 2010a. Linking killer whale survival and prey abundance: food limitation in the oceans' apex predator? *Biol. Letters*, 6:139-142.
- Ford, J.K.B., Stredulinsky, E.H., Ellis, G.M., Durban, J.W. and Pilkington, J.F. 2014. Offshore Killer Whales in Canadian Pacific Waters: Distribution, Seasonality, Foraging Ecology, Population Status and Potential for Recovery. DFO Can. Sci. Advis. Sec. Res. Doc. 2014/088. vii + 55 p.
- Ford, J.K.B., Stredulinsky, E.H., Towers, J.R. and Ellis, G.M. 2013. Information in Support of the Identification of Critical Habitat for Transient Killer Whales (*Orcinus orca*) off the West Coast of Canada. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/155: v + 46 p.
- Ford, J.K.B., Wright, B.M., Ellis, G.M. and Candy, J.R. 2010b. Chinook salmon predation by resident killer whales: seasonal and regional selectivity, stock identity of prey, and consumption rates. DFO Can. Sci. Advis. Sec. Res. Doc. 2009/101. iv + 43 p.
- Foster, E.A., Franks, D.W., Mazzi, S., Darden, S.K., Balcomb, K.C., Ford, J.K.B., and Croft, D.P. 2012. Adaptive prolonged postreproductive life span in killer whales. *Science*. 337(6100): 1313.
- Jacobsen, J.K. 1986. The behavior of *Orcinus orca* in the Johnstone Strait, British Columbia, pp.135-85. In: B.C. Kirkevold and J.S. Lockard (ed.), *The Behavioral Biology of Killer Whales*. Alan R. Liss, New York. 457 p.
- Krahn, M.M., Hanson, B.M., Baird, R.W., Boyer, R.H., Burrows, D.G., Emmons, C.K., Ford, J.K.B., Jones, L.L., Noren, D.P., Ross, P.S., Schorr, G.S. and Collier, T.K. 2007a. Persistent organic pollutants and stable isotopes in biopsy samples (2004/2006) from southern resident killer whales. *Mar. Pollute. Bull.* 54: 1903-1911.
- Krahn, M.M., Hanson, B.M., Schorr, G.S., Emmons, C.K., Burrows, D.G., Bolton, J.L., Baird, R.W. and Ylitalo, G.M. 2009. Effects of age, sex and reproductive status

on persistent organic pollutant concentrations in “Southern Resident” killer whales. Mar. Pollute. Bull. 58: 1522-1529.

Krahn M.M., Herman, D.P., Matkin, C.O., Durban, J.W., Barrett-Lennard, L., Burrows, D.G., Dahlheim, M.E., Black, N., LeDuc, R.G. and Wade, P.R. 2007b. Use of chemical tracers in assessing the diet and foraging regions of eastern North Pacific killer whales. Mar. Env. Res. 63(2): 91-114.

Lachmuth, C.L., Alava, J.J., Hickie, B.E., Johannessen, S.C., Macdonald, R.W., Ford, J.K.B., Ellis, G.M., Gobas, F.A.P.C. and Ross, P.S. 2010. Ocean Disposal in Resident Killer Whale (*Orcinus orca*) Critical Habitat: Science in Support of Risk Management. DFO Can. Sci. Advis. Sec. Res. Doc. 2010/116. x + 172 p.

Lachmuth, C.L., Barrett-Lennard, L.G., Steyn, D.Q. and Milsom, W.K. 2011. Estimation of southern resident killer whale exposure to exhaust emissions from whale-watching vessels and potential adverse health effects and toxicity thresholds. Mar. Pollute. Bull. 62(4): 792-805.

Lusseau, D., Bain, D.E., Williams, R. and Smith, J.C. 2009. Vessel traffic disrupts the foraging behavior of southern resident killer whales *Orcinus orca*. Endanger. Species Res. 6, 211–221.

Morton, A.B. 1990. A quantitative comparison of the behavior of resident and transient forms of the killer whale off the central British Columbia coast. Rep. Int. Whal. Commn. Special Issue 12: 245–248.

Morton, A.B. and Symonds, H.K. 2002. Displacement of *Orcinus orca* (L.) by high amplitude sound in British Columbia, Canada. ICES J. Mar. Sci. 59: 71-80.

Nichol, L.M. and Shackleton, D.M. 1996. Seasonal movements and foraging behaviour of northern resident killer whales (*Orcinus orca*) in relation to the inshore distribution of salmon (*Oncorhynchus* spp.) in British Columbia. Can. J. Zool. 74: 983-991.

Olesiuk, P.F., Bigg, M.A. and Ellis, G.M. 1990. Life history and population dynamics of resident killer whales (*Orcinus orca*) in the coastal waters of British Columbia and Washington State. Rep. Int. Whal. Commn. Special Issue 12: 209-242.

Olesiuk, P.F., Ellis, G.M. and Ford, J.K.B. 2005. Life history and population dynamics of northern resident killer whales (*Orcinus orca*) in British Columbia. DFO Can. Sci. Advis. Sec. Res. Doc. 2005/045. iv + 75 p.

Rayne, S., Ikonomou, M.G., Ross, P.S., Ellis, G.M. and Barrett-Lennard, L.G. 2004. PBDEs, PBBs, and PCNs in three communities of free-ranging killer whales

- (*Orcinus orca*) from the northeastern Pacific Ocean. Environ. Sci. Technol. 38: 4293-4299.
- Ross, P.S. 2006. Fireproof killer whales (*Orcinus orca*): Flame retardant chemicals and the conservation imperative in the charismatic icon of British Columbia, Canada. Can. J. Fish. Aquat. Sci. 63: 224-234.
- Ross, P.S., Ellis, G.M., Ikonumou, M.G., Barrett-Lennard, L.G. and Addison, R.F. 2000. High PCB concentrations in free-ranging Pacific Killer Whales, *Orcinus orca*: effects of age, sex and dietary preference. Mar. Pollute. Bull. 40: 504-515.
- Towers, J.R., Ellis, G.M., and Ford, J.K.B. 2012a. Photo-identification catalogue of Bigg's (Transient) killer whales from coastal waters of British Columbia, northern Washington, and southeastern Alaska. Can. Data Rep. of Fish. and Aquat. Sci. 1241: v + 127 p.
- Towers, J.R., Ford, J.K.B., and Ellis, G.M. 2012b. Digital photo-identification dataset management and analysis: Testing protocols using a commercially available application. Can. Tech. Rep. Fish. Aquat. Sci. 2978: iv + 16 p.
- Williams, R. and Ashe, E. 2007. Killer whale evasive tactics vary with boat number. J. Zool. 272: 390–397.
- Williams, R. and Lusseau, D. 2006. A killer whale social network is vulnerable to targeted removals. Biol. Letters. 2: 497-500.
- Williams, R. and O'Hara, P. 2010. Modelling ship strike risk to fin, humpback and killer whales in British Columbia, Canada. J. Cetacean Res. Manag. 11: 1–8.
- Williams, R., Bain, D.E., Ford, J.K.B. and Trites, A.W. 2002b. Behavioural responses of male killer whales to a 'leapfrogging' vessel. J. Cetacean Res. Manag. 4(3): 305-310.
- Williams, R., Clark, C.W., Ponirakis, D. and Ashe, E. 2014. Acoustic quality of critical habitats for three threatened whale populations. Anim. Conserv. 17:174-185.
- Williams, R., Trites, A.W. and Bain, D.E. 2002a. Behavioural responses of killer whales (*Orcinus orca*) to whale-watching boats; opportunistic observations and experimental approaches. J Zool. 256: 255-270.

## **APPENDIX**

### **PHOTO-IDENTIFICATION CATALOGUE OF NORTHERN RESIDENT KILLER WHALES**



# A1 Pod, A36 Matriline

A46 ♂ 1982



A-Clan

## CATALOGUE KEY


A1 ♀  
1930 Name, sex, and birth year of deceased whale

A46 ♂ Name and sex of live whale

— Positive Relationship


..... Probable Relationship

..... Possible Relationship



A1 Pod,  
A30 Matriline

A38 ♂ 1971



A50 ♀ 1984



A54 ♀ 1989



A72 ♀ 1999



A84 2005



A99 2011



A75 ♀ 2002



A86 2006





A106 2013









A108 2014



A101 2012




# A-Clan





A4 Pod,  
A35 Matriline







A4 Pod,  
A56 Matriline




A4 Pod,  
A24 Matriline


A73 and her offspring  
normally travel with the  
A35 and A56 matriline.



A4 Pod,  
A24 Matriline  
continued





A5 Pod,  
A42 Matriline





A-Clan


A-Clan



A5 Pod,  
A23 Matriline




A5 Pod,  
A25 Matriline






B1 Pod,  
B7 Matriline


A-Clan




A-Clan





C1 Pod,  
C6 Matriline



C8, her offspring, and C27 are not always documented with C14 and C18.







D1 Pod,  
D12 Matriline

D12 ♀ 1982



D20 ♀ 1999



D29 2014



D1 Pod,  
D9 Matriline

D9 ♀ 1972



Since the death of C19,  
C31 has been travelling  
with D9.

D1 Pod,  
D13 Matriline

D13 ♀ 1984




D19 ♀ 1998



D23 2005





D27 2012





D1 Pod,  
D11 Matriline



The four matriline in  
D pod are usually found  
together.





H1 Pod,  
H5 Matriline





I1 Pod,  
I19 Matriline


A-Clan



A-Clan


I1 Pod,  
I40 Matriline

A-Clan



I18 Pod,  
I17 Matriline

A-Clan



I18 Pod,  
**I17 Matriline**  
continued

A-Clan

I50 and her offspring are  
not always found with the  
rest of the I17 matriline.

I50 ♀ 1983



I89 ♂ 1996



I100 ♀ 2003



I130 2010



I147 2014



I17 ♀ 1960



I57 ♀ 1989





I99 ♀ 2003



I124 2007



I149 2014




A-Clan




I18 Pod,  
I18 Matriline



A-Clan





A-Clan



I18 Pod,  
I18 Matriline  
continued



I24, her offspring, and I53 are not often found with others in the I18 matriline.



I2 Pod,  
I22 Matriline

