


RESIDUAL TOTAL MAGNETIC FIELD



This aeromagnetic survey and the production of this map were funded by phase 2 of the Geo-Mapping for Energy and Minerals program (GEM-2) of the Earth Sciences Sector, Natural Resources Canada.


GEOLOGICAL SURVEY OF CANADA OPEN FILE 7673

RESIDUAL TOTAL MAGNETIC FIELD


AEROMAGNETIC SURVEY OF SOMERSET ISLAND AREA

Authors: M. Coyle and D. Oneschuk
Data acquisition and data compilation by
EON Geosciences Inc., Montréal, Québec.
Contract and project management and map production
by the Geological Survey of Canada, Ottawa, Ontario.

Parts of NTS 58-C/11, 12 and 13
NUNAVUT



Universal Transverse Mercator Projection
North American Datum 1983
©Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources Canada, 2015
Digital topographic data from Natural Resources Canada




MAP LOCATION

Residual Total Magnetic Field

This map of the residual total magnetic field was derived from data acquired during an aeromagnetic survey carried out by EON Geosciences Inc. from May 30 to July 8, 2014. The data were recorded using split-beam cesium vapour magnetometers (sensitivity = 0.005 nT) mounted in each of the tail booms of a Piper Cheyenne II aircraft (C-GFON) and a Cessna 206 aircraft (C-FTPN). The nominal traverse line spacings were 400 m and 500 m. The nominal control line spacing was 2400 m. The aircraft flew at a nominal terrain clearance of 150 m. Traverse lines were oriented E-W with orthogonal control lines. The flight path was recovered following post-flight differential corrections to the raw Global Navigational Satellite System (GNSS) data and inspection of ground images recorded by a vertically-mounted video camera. The survey was flown on a pre-determined flight surface to minimize differences in magnetic values at the intersections of control and traverse lines. These differences were computer-analysed to obtain a mutually levelled set of flight-line magnetic data. The levelled values were then interpolated to a 100 m grid. The International Geomagnetic Reference Field (IGRF), defined at the average GPS altitude of 399.88 m for the year 2014.465 was then removed. Removal of the IGRF, representing the magnetic field of the Earth's core, produces a residual component related almost entirely to magnetizations within the Earth's crust.

A digital version of this map can be downloaded, at no charge, from Natural Resources Canada's GeoGratis portal at <http://geogratis.gc.ca/>. Corresponding digital profile and gridded data as well as similar data for adjacent airborne geophysical surveys are available from Natural Resources Canada's Geoscience Data Repository for Aeromagnetic data at http://nrcan.nrcan.gc.ca/index_e.html. The same products are also available, for a fee, from the Geophysical Data Centre, Geological Survey of Canada, 615 Booth Street, Ottawa, Ontario K1A 0E9. Telephone: (613) 995-5326, email: infoqdc@gsg.nrcan.gc.ca.



AEROMAGNETIC SURVEY OF SOMERSET ISLAND AREA

OPEN FILE
DOSSIER PUBLIC
7673
GEOLOGICAL SURVEY OF CANADA
COMMISSION GÉOLOGIQUE DU CANADA
2015

Publications in this series
have not been edited;
they are released as
submitted by the author.

Les publications de cette
série ne sont pas révisées;
elles sont publiées telles
que soumises par l'auteur.

Recommended citation:

Coyle, M. and Oneschuk, D., 2015.
Residual total magnetic field,
Aeromagnetic survey of Somerset Island Area,
Parts of NTS 58-C/11, 12 and 13, Nunavut;
Geological Survey of Canada, Open File 7673;
Scale 1:100 000.