

June 18 to July 22, 2017 (Weeks 25-29)


## Overall Summary

- Influenza activity is at interseasonal levels across the country, with a few regions reporting sporadic or localized activity.
- In weeks 25-29, influenza A and B viruses circulated at interseasonal levels in Canada. The majority of subtyped influenza A viruses were A(H3N2).
- FluWatch will publish monthly reports over the summer. The next report will be published on August 25, 2017. We continue to monitor influenza and other respiratory infections via the [RVDSS report](#), published every Thursday.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

## Influenza/Influenza-like Illness (ILI) Activity (geographic spread)

In weeks 25-29, influenza or influenza-like illness activity levels continued to decline and the majority of regions reported no activity. In week 29, one region in Ontario reported localized activity, and seven regions across five provinces reported sporadic activity. For more details on a specific region, click on the map.

**Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, Week 29**




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available in the [mapping feature](#) found in the [Weekly Influenza Reports](#).

## Laboratory Confirmed Influenza Detections

In weeks 25-29, the number of tests positive for influenza remained at interseasonal levels, although slightly higher than in the previous six seasons. The percentage of tests positive was low, around 2% during weeks 25-29. Detections of influenza A and B were nearly equal in weeks 25-29, and the majority of influenza A viruses subtyped were A(H3N2). For data on other respiratory virus detections, see the [Respiratory Virus Detections in Canada Report](#) on the Public Health Agency of Canada (PHAC) website.


**Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2016-17, Week 29**



The shaded area indicates weeks where the positivity rate was at least 5% and a minimum of 15 positive tests were observed, signalling the period of [seasonal influenza activity](#).

To date this season, 39,235 laboratory-confirmed influenza detections have been reported, of which 89% have been influenza A. Influenza A(H3N2) has been the most common subtype detected this season, representing over 99% of influenza A detections. For more detailed weekly and cumulative influenza data, see the text descriptions for Figures 2 and 3 or the [Respiratory Virus Detections in Canada Report](#).

**Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province/territory, Canada, 2016-17, Week 29**



To date this season, detailed information on age and type/subtype has been received for 27,258 laboratory-confirmed influenza cases (Table 1). Among cases with reported age and type/subtype information, adults aged 65+ accounted for nearly half of the reported influenza cases. Adults aged 65+ have predominantly been affected by influenza A accounting for 51% of influenza A detections. Influenza B, while much smaller in number, is mainly affecting individuals less than 65 years of age.

**Table 1 – Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting<sup>1</sup>, Canada, 2016-17, Week 29**

Age groups (years)	Weeks 25-29 (June 18 to July 22, 2017)					Cumulative (August 28, 2016 to July 22, 2017)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) <sup>3</sup>		A Total	A(H1) pdm09	A(H3)	A (UnS) <sup>3</sup>		Total	#
0-4	<5	<5	<5	<5	9	2253	22	833	1398	298	2551	9%
5-19	5	0	<5	<5	9	2225	18	1080	1127	555	2780	10%
20-44	>10	<5	5	5	8	3460	43	1810	1607	551	4011	15%
45-64	16	<5	10	5	9	3952	34	1971	1947	740	4692	17%
65+	34	<5	18	15	24	12141	21	5470	6650	1083	13224	49%
<b>Total</b>	<b>71</b>	<b>6</b>	<b>36</b>	<b>29</b>	<b>59</b>	<b>24031</b>	<b>138</b>	<b>11164</b>	<b>12729</b>	<b>3227</b>	<b>27258</b>	<b>100%</b>
<b>Percentage<sup>2</sup></b>	<b>55%</b>	<b>8%</b>	<b>51%</b>	<b>41%</b>	<b>45%</b>	<b>88%</b>	<b>1%</b>	<b>46%</b>	<b>53%</b>	<b>12%</b>		

<sup>1</sup>Table 1 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported. Cumulative data include updates to previous weeks.

<sup>2</sup>Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

<sup>3</sup>UnS: unsorted: The specimen was typed as influenza A, but no result for subtyping was available.

x - Supressed to prevent residual disclosure

Specimens from NT, YT, and NU are sent to reference laboratories in the provinces


## Syndromic/Influenza-like Illness Surveillance

### Healthcare Professionals Sentinel Syndromic Surveillance

In weeks 25-29, overall, the proportion of visits to healthcare professionals due to influenza-like illness was low. The weekly percentage varied between 0.5% and 1.9% due to a large proportion of visits for ILI reported by a single sentinel practitioner.

**Figure 4 – Percentage of visits for ILI reported by sentinels by report week, Canada, 2016-17**

Number of Sentinels Reporting Week 29: 94



Delays in the reporting of data may cause data to change retrospectively. In BC, AB, and SK, data are compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.


**Are you a primary healthcare practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel? Please visit our [Influenza Sentinel page](#) for more details.**

## Influenza Outbreak Surveillance

During the period of weeks 25-29, four laboratory-confirmed influenza outbreaks were reported, of which three were in long-term care facilities, and one was in an institutional or community (other) setting.

To date this season, 1,194 outbreaks have been reported and the majority (66%) have occurred in LTC facilities. Fewer outbreaks were reported this season compared to the same period in the most recent previous A(H3N2)-predominant season (2014-15) when 1,732 outbreaks were reported, of which 74% occurred in LTC facilities.

**Figure 5 – Number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2016-17, Week 29**



<sup>1</sup>All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of this report.

## Provincial/Territorial Influenza Hospitalizations and Deaths

During the period of weeks 25-29, the number of weekly influenza-associated hospitalizations reported by participating provinces and territories\* decreased to low levels. In weeks 25-29, 37 hospitalizations were reported, of which 21 were associated with influenza A and 65% occurred in adults 65+. Four intensive care unit (ICU) admissions and two deaths were reported.

To date this season, 6,527 hospitalizations have been reported, of which 88% were due to influenza A. Among cases for which the subtype of influenza A was reported, 99% were influenza A(H3N2). Adults 65+ accounted for 67% of the hospitalizations. A total of 273 ICU admissions and 387 deaths have been reported. The majority of deaths (88%) were reported in adults aged 65+ years.

**Table 2 – Cumulative number of hospitalizations, ICU admissions and deaths by age and influenza type reported by participating provinces and territories, Canada, 2016-17, Week 29**

Age Groups (years)	Cumulative (August 28, 2016 to July 22, 2017)						
	Hospitalizations			ICU Admissions		Deaths	
	Influenza A Total	Influenza B Total	Total [# (%)]	Influenza A and B Total	%	Influenza A and B Total	%
0-4	447	94	541 (8%)	20	7%	<5	x%
5-19	241	99	340 (5%)	20	7%	<5	x%
20-44	296	52	348 (5%)	27	10%	5	1%
45-64	765	143	908 (14%)	83	30%	37	10%
65+	3964	426	4390 (67%)	123	45%	340	88%
<b>Total</b>	<b>5713</b>	<b>814</b>	<b>6527 (99%)</b>	<b>273</b>	<b>99%</b>	<b>387</b>	<b>100%</b>

x: Suppressed to prevent residual disclosure

\*Note: Influenza-associated hospitalizations are not reported to PHAC by BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by SK. ICU admissions are not distinguished among hospital admissions reported from ON. The hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.


# Sentinel Hospital Influenza Surveillance

## Pediatric Influenza Hospitalizations and Deaths


In weeks 25-29, nine laboratory-confirmed influenza-associated pediatric ( $\leq 16$  years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network. Six of the nine hospitalizations were associated with influenza B.

To date this season, 595 laboratory-confirmed influenza-associated pediatric hospitalizations were reported by the IMPACT network. Children aged 0-23 months accounted for approximately 37% of hospitalizations and influenza A accounted for 78% of the reported hospitalizations. Among the 132 hospitalizations due to influenza B, 69 (52%) were in children 5 to 16 years of age. In comparison, children over the age of 5 years accounted for 33% of influenza A hospitalizations. Additionally, 100 intensive care unit (ICU) admissions have been reported. A total of 67 ICU cases (67%) reported at least one underlying condition or comorbidity. Less than five deaths have been reported this season.

**Figure 6 – Cumulative numbers of pediatric hospitalizations ( $\leq 16$  years of age) with influenza by age-group reported by the IMPACT network, Canada, 2016-17, Week 29**



**Figure 7 – Number of pediatric hospitalizations ( $\leq 16$  years of age) with influenza reported by the IMPACT network, by week, Canada, 2016-17, Week 29**



The shaded area represents the maximum and minimum number of cases reported by week from seasons 2010-11 to 2015-16


The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated pediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

## Adult Influenza Hospitalizations and Deaths


Surveillance for the 2016-2017 influenza season ended on April 30<sup>th</sup>, 2017 (week 20).

This season, 1,535 laboratory-confirmed influenza-associated adult ( $\geq 20$  years of age) hospitalizations have been reported by the Canadian Immunization Research Network (CIRN). Influenza A accounted for 92% of hospitalizations. Adults aged 65+ accounted for 78% of hospitalizations. A total of 143 intensive care unit (ICU) admissions have been reported. Among ICU cases with available data, 126 cases (88%) reported at least one underlying condition or comorbidity. The median age of patients admitted to the ICU was 71 years. Approximately 86 deaths have been reported this season, the majority in adults aged 65+. The median age of reported deaths was 85 years.

**Figure 8 - Cumulative numbers of adult hospitalizations ( $\geq 20$  years of age) with influenza by type and age-group reported by CIRN, Canada, 2016-17, Week 20**



**Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza by age-group ( $\geq 20$  years of age) reported by CIRN, Canada 2016-17, Week 20**



The number of hospitalizations reported through CIRN represents a subset of all influenza-associated adult hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.



## Influenza Strain Characterizations

During the 2016-17 influenza season, the National Microbiology Laboratory (NML) has characterized 2,307 influenza viruses [1,628 A(H3N2), 59 A(H1N1), 620 influenza B]. All seasonal influenza A viruses and 20% of influenza B viruses characterized were antigenically or genetically similar to the vaccine strains included in both the trivalent and quadrivalent vaccines. Eighty percent of influenza B viruses characterized were similar to the strain which is only included in the quadrivalent vaccine.

**Table 3 – Influenza strain characterizations, Canada, 2016-17, Week 29**

Strain Characterization Results <sup>1</sup>	Count	Description
<b>Influenza A (H3N2)</b>		
Antigenically A/Hong Kong/4801/2014-like	395	Viruses antigenically similar to A/Hong Kong/4801/2014, the A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine.
Genetically <sup>2</sup> A/Hong Kong/4801/2014-like	1232	Viruses belonging to genetic group 3C.2a. A/Hong Kong/4801/2014-like virus belongs to genetic group 3C.2a and is the influenza A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine.  Additionally, one virus belonged to genetic group 3C.3a. Genetic characterization of the 395 influenza A (H3N2) viruses that underwent HI testing determined that 329 viruses belonged to genetic group 3C.2a and 65 viruses belonged to genetic group 3C.3a. The majority of viruses belonging to genetic group 3C.3a are inhibited by antisera raised against A/Hong Kong/4801/2014 <sup>3</sup> .
Antigenically A/Indiana/10/2011-like <sup>4</sup>	1	Viruses antigenically similar to A/Indiana/10/2011, a candidate H3N2v vaccine virus.
<b>Influenza A (H1N1)</b>		
A/California/7/2009-like	59	Viruses antigenically similar to A/California/7/2009, the A(H1N1) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
<b>Influenza B</b>		
B/Brisbane/60/2008-like (Victoria lineage)	125	Viruses antigenically similar to B/Brisbane/60/2008, the influenza B component of the 2016-17 Northern Hemisphere's <b>trivalent</b> and <b>quadrivalent</b> influenza vaccine.
B/Phuket/3073/2013-like (Yamagata lineage)	495	Viruses antigenically similar to B/Phuket/3073/2013, the additional influenza B component of the 2016-17 Northern Hemisphere <b>quadrivalent</b> influenza vaccine.

<sup>1</sup>The NML receives a proportion of the influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Strain characterization data reflect the results of hemagglutination inhibition (HI) testing compared to the reference influenza strains recommended by [WHO](#).

<sup>2</sup>Determined by sequence analysis

<sup>3</sup>[WHO](#) - Recommended composition of the influenza virus vaccines for use in the 2016-17 northern hemisphere influenza season.

<sup>4</sup>Detected in epidemiological week 50. For more details, see [Week 50 report](#)

## Antiviral Resistance

During the 2016-17 season, the National Microbiology Laboratory (NML) has tested 1,250 influenza viruses for resistance to oseltamivir, 1,250 influenza viruses for resistance to zanamivir and 272 influenza viruses for resistance to amantadine. All but two influenza A(H3N2) viruses and one of the A(H1N1) viruses were sensitive to oseltamivir and all viruses were sensitive to zanamivir. All 272 influenza A viruses were resistant to amantadine (Table 4).

**Table 4 – Antiviral resistance by influenza virus type and subtype, Canada, 2016-17, Week 24**

Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
<b>A (H3N2)</b>	757	2 (0.3%)	756	0 (0%)	218	218 (100%)
<b>A (H3N2v)</b>	1	0 (0%)	1	0 (0%)	1	1 (100%)
<b>A (H1N1)</b>	52	1 (1.9%)	51	0 (0%)	53	53 (100%)
<b>B</b>	440	0 (0%)	442	0 (0%)	NA <sup>1</sup>	NA <sup>1</sup>
<b>TOTAL</b>	1250	3 (0.2%)	1250	0 (0%)	272	272 (100%)

<sup>1</sup>NA: Not Applicable

## Provincial and International Influenza Reports

- [World Health Organization influenza update](#)
- [World Health Organization FluNet](#)
- [WHO Influenza at the human-animal interface](#)
- [Centers for Disease Control and Prevention seasonal influenza report](#)
- [European Centre for Disease Prevention and Control - epidemiological data](#)
- [South Africa Influenza surveillance report](#)
- [New Zealand Public Health Surveillance](#)
- [Australia Influenza Report](#)
- [Pan-American Health Organization Influenza Situation Report](#)
- [Alberta Health – Influenza Surveillance Report](#)
- [BC - Centre for Disease Control \(BCCDC\) - Influenza Surveillance](#)
- [New Brunswick – Influenza Surveillance Reports](#)
- [Newfoundland and Labrador – Surveillance and Disease Reports](#)
- [Nova Scotia - Flu Information](#)
- [Public Health Ontario – Ontario Respiratory Pathogen Bulletin](#)
- [Manitoba – Epidemiology and Surveillance – Influenza Reports](#)
- [Saskatchewan – influenza Reports](#)
- [PEI – Influenza Summary](#)



## FluWatch Definitions for the 2016-2017 Season

**Abbreviations:** Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

**Influenza-like-illness (ILI):** Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

### ILI/Influenza outbreaks

**Schools:** Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

**Hospitals and residential institutions:** two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

**Workplace:** Greater than 10% absenteeism on any day which is most likely due to ILI.

**Other settings:** two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

*Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.*

### Influenza/ILI Activity Levels

**1 = No activity:** no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

**2 = Sporadic:** sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

**3 = Localized:** (1) evidence of increased ILI\* ;  
(2) lab confirmed influenza detection(s);  
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region**†

**4 = Widespread:** (1) evidence of increased ILI\*;  
(2) lab confirmed influenza detection(s);  
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region**†

*Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.*

*\* More than just sporadic as determined by the provincial/territorial epidemiologist.*

*† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.*

*We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.*

This [report](#) is available on the Government of Canada Influenza webpage. Ce rapport est disponible dans les deux langues officielles.