

January 15 to 21, 2017 (Week 03)


Overall Summary

- In week 03, activity from several indicators including laboratory detections, outbreaks and hospitalizations declined from the previous week indicating that nationally the influenza season may have reached its peak in week 02.
- A total of 2,667 positive influenza detections were reported in week 03, a decrease from the previous week.
- A(H3N2) continues to be the most common type of influenza affecting Canadians.
- The majority of cases, hospitalizations and deaths have been among adults aged 65+ years.
- Sixty-six confirmed influenza outbreaks were reported in week 03, with the majority occurring in long-term care facilities and due to influenza A.
- A total of 417 hospitalizations were reported by participating provinces and territories, down from 467 hospitalizations reported in the previous week.
- For more information on the flu, see our [Flu\(influenza\)](#) web page.

Influenza/Influenza-like Illness (ILI) Activity (geographic spread)

In week 03, all but two reporting regions are reporting influenza or influenza-like illness activity. One region in NS and one region in NB reported no activity. Sporadic influenza activity was reported in 23 regions across ten provinces and territories. Localized activity was reported in twenty regions across seven provinces and territories. Widespread activity was reported in five provinces (three regions in AB, one region in ON, one region in QC, one region in PE and one region in BC). For more details on a specific region, click on the map.

Figure 1 – Map of overall influenza/ILI activity level by province and territory, Canada, Week 03




Note: Influenza/ILI activity levels, as represented on this map, are assigned and reported by Provincial and Territorial Ministries of Health, based on laboratory confirmations, sentinel ILI rates and reported outbreaks. Please refer to detailed definitions at the end of the report. Maps from previous weeks, including any retrospective updates, are available in the mapping feature found in the [Weekly Influenza Reports](#).

Laboratory Confirmed Influenza Detections

The percentage of tests positive for influenza decreased from 27% in week 02 to 23% in week 03. This is an indication that we may have reached the peak in the percent positive and number of detections for influenza in week 02. In 2014-15, the previous influenza A(H3N2)-predominant season, the peak occurred in week 52 with 34% of tests positive for influenza. For data on other respiratory virus detections, see the [Respiratory Virus Detections in Canada Report](#) on the Public Health Agency of Canada (PHAC) website.


Figure 2 – Number of positive influenza tests and percentage of tests positive, by type, subtype and report week, Canada, 2016-17, Week 03



The shaded area indicates weeks where the positivity rate was at least 5% and a minimum of 15 positive tests were observed, signalling the period of [seasonal influenza activity](#).

Nationally in week 03, 2,667 positive influenza tests were reported, down from 3,477 tests reported in week 02. To date, a total of 15,231 laboratory confirmed influenza detections have been reported, of which 98% have been influenza A. Influenza A(H3N2) is the most common subtype detected, representing over 99% of subtyped influenza A detections (8664/8703). For more detailed weekly and cumulative influenza data, see the text descriptions for Figures 2 and 3 or the [Respiratory Virus Detections in Canada Report](#).

Figure 3 – Cumulative numbers of positive influenza specimens by type/subtype and province/territory, Canada, 2016-17, Week 03



To date, detailed information on age and type/subtype has been received for 10, 808 laboratory confirmed influenza cases. Among cases with reported age and type/subtype information, adults aged 65+ accounted for over half of the reported influenza cases and the largest proportion (45%) of influenza A (H3N2) cases.

Table 1 – Weekly and cumulative numbers of positive influenza specimens by type, subtype and age-group reported through case-based laboratory reporting¹, Canada, 2016-17, Week 03

Age groups (years)	Weeks (Jan. 15 to Jan. 21, 2017)					Cumulative (Aug. 28, 2016 to Jan. 21, 2017)						
	Influenza A				B	Influenza A				B	Influenza A and B	
	A Total	A(H1) pdm09	A(H3)	A (UnS) ³		A Total	A(H1) pdm09	A(H3)	A (UnS) ³		Total	#
0-4	147	0	46	101	5	>844	<5	404	440	44	>888	x%
5-19	139	0	40	99	5	>1032	<5	593	439	37	>1069	x%
20-44	193	0	66	127	<5	>1661	<5	944	717	34	>1695	x%
45-64	248	0	94	154	9	1833	6	960	867	36	1869	17%
65+	691	0	209	482	6	>5227	<5	2336	2891	52	>5279	x%
Total	1418	0	455	963	>25	10605	14	5237	5354	203	10808	100%
Percentage²	98%	0%	32%	68%	2%	98%	0%	49%	50%	2%		

¹Table 1 includes specimens for which demographic information was reported. These represent a subset of all positive influenza cases reported. Cumulative data include updates to previous weeks.

²Percentage of tests positive for sub-types of influenza A are a percentage of all influenza A detections.

³UnS: unsubtyped: The specimen was typed as influenza A, but no result for subtyping was available.

x: Suppressed to prevent residual disclosure

Specimens from NT, YT, and NU are sent to reference laboratories in the provinces


Syndromic/Influenza-like Illness Surveillance

Healthcare Professionals Sentinel Syndromic Surveillance

In week 03, 2.1% of visits to healthcare professionals were due to influenza-like illness.

Figure 4 – Percentage of visits for ILI reported by sentinels by report week, Canada, 2016-17

Number of Sentinels Reporting Week 03: 128



Delays in the reporting of data may cause data to change retrospectively. In BC, AB, and SK, data are compiled by a provincial sentinel surveillance program for reporting to FluWatch. Not all sentinel physicians report every week.


Are you a primary healthcare practitioner (General Practitioner, Nurse Practitioner or Registered Nurse) interested in becoming a FluWatch sentinel? Please visit our [Influenza Sentinel page](#) for more details.

Influenza Outbreak Surveillance

In week 03, sixty-six laboratory confirmed influenza outbreaks were reported: 39 in long-term care (LTC) facilities, 19 in hospitals and 8 in institutional or community (other) settings. Of the outbreaks with known strains or subtypes, all outbreaks were due to influenza A, of which 17 were due to influenza A(H3N2) and 35 were due to influenza A(UnS).

To date this season, 536 outbreaks have been reported and the majority (68%) have occurred in LTC facilities. In comparison, at week 03 in the 2014-15 season, the previous influenza A(H3N2)-predominant season, 1,036 outbreaks were reported, of which 77% occurred in LTC facilities.

Figure 5 – Number of new laboratory-confirmed influenza outbreaks by report week, Canada, 2016-17, Week 03



¹All provinces and territories except NU report influenza outbreaks in long-term care facilities. All provinces and territories with the exception of NU and QC report outbreaks in hospitals. Outbreaks of influenza or influenza-like-illness in other facilities are reported to FluWatch but reporting varies between jurisdictions. Outbreak definitions are included at the end of this report.

Provincial/Territorial Influenza Hospitalizations and Deaths

In week 03, 417 influenza-associated hospitalizations were reported by participating provinces and territories*. Influenza A accounted for all but six reported hospitalizations. A total of ten intensive care unit (ICU) admissions and 24 deaths were reported in week 03. Adults aged 65+ accounted for the largest proportion of hospitalizations (69%). All ICU admissions and deaths in week 03 were reported in adults.

To date this season, 2,479 hospitalizations have been reported, of which 99% were due to influenza A. Among cases for which the subtype of influenza A was reported, almost all (1446/1450) were influenza A(H3N2). Adults 65+ accounted for 70% of the hospitalizations. Eighty-nine ICU admissions and 79 deaths have been reported. The majority of deaths (81%) were reported in adults aged 65+ years.

Table 2 – Cumulative number of hospitalizations, ICU admissions and deaths by age and influenza type reported by participating provinces and territories, Canada, 2016-17, Week 03

Age Groups (years)	Cumulative (Aug. 28, 2016 to Jan. 21, 2017)						
	Hospitalizations			ICU Admissions		Deaths	
	Influenza A Total	Influenza B Total	Total [# (%)]	Influenza A and B Total	%	Influenza A and B Total	%
0-4	168	5	173 (7%)	7	8%	0	0%
5-19	100	5	105 (4%)	6	7%	<5	x%
20-44	137	<5	>137 (x%)	7	8%	0	0%
45-64	319	<5	>319 (x%)	28	31%	14	x%
65+	1725	15	1740 (70%)	41	46%	64	x%
Total	2449	30	2479 (100%)	89	100%	79	100%

x: Suppressed to prevent residual disclosure

*Note: Influenza-associated hospitalizations are not reported to PHAC by BC, NU, and QC. Only hospitalizations that require intensive medical care are reported by SK. ICU admissions are not distinguished among hospital admissions reported from ON. The hospitalization or death does not have to be attributable to influenza, a positive laboratory test is sufficient for reporting.

Sentinel Hospital Influenza Surveillance

Pediatric Influenza Hospitalizations and Deaths

In week 03, 36 laboratory-confirmed influenza-associated pediatric (≤ 16 years of age) hospitalizations were reported by the Immunization Monitoring Program Active (IMPACT) network. All but two cases were due to influenza A. The number of hospitalizations reported in week 03 is below the six year average for the same time period (Figure 7).

To date this season, 236 laboratory-confirmed influenza-associated pediatric hospitalizations were reported by the IMPACT network. Children aged 0-2 years accounted for approximately 41% of hospitalizations. Influenza A accounted for 93% (n=219) of the reported hospitalizations, of which 44% (n=96) were influenza A(H3N2) and the remainder were A(UnS). Additionally, 40 intensive care unit (ICU) admissions have been reported, of which the largest proportion (27%) was reported in children 10-16 years. A total of 25 ICU cases reported at least one underlying condition or comorbidity. No deaths have been reported this season.

In 2014-15, the previous influenza A(H3N2)-predominant season, there were 446 hospitalizations, 50 ICU admissions and three deaths reported as of week 03.

Figure 6 – Cumulative numbers of pediatric hospitalizations (≤ 16 years of age) with influenza by age-group reported by the IMPACT network, Canada, 2016-17, Week 03





Figure 7 – Number of pediatric hospitalizations (≤ 16 years of age) with influenza reported by the IMPACT network, by week, Canada, 2016-17, Week 03



The shaded area represents the maximum and minimum number of cases reported by week from seasons 2010-11 to 2015-16

The number of hospitalizations reported through IMPACT represents a subset of all influenza-associated pediatric hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Adult Influenza Hospitalizations and Deaths

In week 03, a total of 104 laboratory-confirmed influenza-associated adult (≥ 20 years of age) hospitalizations were reported by the Canadian Immunization Research Network (CIRN). All cases were due to influenza A and the greatest proportion of cases (74%) occurred in adults aged 65+.

To date this season, 607 laboratory-confirmed influenza-associated adult (≥ 20 years of age) hospitalizations have been reported by CIRN. All but four hospitalized cases were due to influenza A. Adults aged 65+ accounted for 78% of hospitalizations. To date, greater than 37 intensive care unit (ICU) admissions have been reported. A total of 26 ICU cases reported at least one underlying condition or comorbidity. A total of 12 deaths have been reported this season, all in adults aged 65+. The median age of reported deaths was 72 years.

Figure 8 - Cumulative numbers of adult hospitalizations (≥ 20 years of age) with influenza by type and age-group reported by CIRN, Canada, 2016-17, Week 03





Figure 9 – Percentage of hospitalizations, ICU admissions and deaths with influenza by age-group (≥ 20 years of age) reported by CIRN, Canada 2016-17, Week 03



The number of hospitalizations reported through CIRN represents a subset of all influenza-associated adult hospitalizations in Canada. Delays in the reporting of data may cause data to change retrospectively.

Influenza Strain Characterizations

During the 2016-17 influenza season, the National Microbiology Laboratory (NML) has characterized 412 influenza viruses [372 A(H3N2), 10 A(H1N1), 30 influenza B]. All but one influenza A virus (n=371) and all (n=30) influenza B viruses characterized were antigenically or genetically similar to the vaccine strains included in both the trivalent and quadrivalent vaccines. Sixteen influenza B viruses were similar to the strain which is included only in the quadrivalent vaccine.

Table 3 – Influenza strain characterizations, Canada, 2016-17, Week 03

Strain Characterization Results ¹	Count	Description
Influenza A (H3N2)		
Antigenically A/Hong Kong/4801/2014-like	132	Viruses antigenically similar to A/Hong Kong/4801/2014, the A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine.
Genetically ² A/Hong Kong/4801/2014-like	239	Viruses belonging to genetic group 3C.2a. A/Hong Kong/4801/2014-like virus belongs to genetic group 3C.2a and is the influenza A(H3N2) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent vaccine. Additionally, genetic characterization of the 132 influenza A (H3N2) viruses that underwent HI testing determined that 92 viruses belonged to genetic group 3C.2a and 21 viruses belonged to genetic group 3C.3a. The majority of viruses belonging to genetic group 3C.3a are inhibited by antisera raised against A/Hong Kong/4801/2014 ³ .
Antigenically A/Indiana/10/2011-like ⁴	1	Viruses antigenically similar to A/Indiana/10/2011, a candidate H3N2v vaccine virus.
Influenza A (H1N1)		
A/California/7/2009-like	10	Viruses antigenically similar to A/California/7/2009, the A(H1N1) component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
Influenza B		
B/Brisbane/60/2008-like (Victoria lineage)	14	Viruses antigenically similar to B/Brisbane/60/2008, the influenza B component of the 2016-17 Northern Hemisphere's trivalent and quadrivalent influenza vaccine.
B/Phuket/3073/2013-like (Yamagata lineage)	16	Viruses antigenically similar to B/Phuket/3073/2013, the additional influenza B component of the 2016-17 Northern Hemisphere quadrivalent influenza vaccine.

¹The NML receives a proportion of the influenza positive specimens from provincial laboratories for strain characterization and antiviral resistance testing. Strain characterization data reflect the results of hemagglutination inhibition (HI) testing compared to the reference influenza strains recommended by [WHO](#).

²Determined by sequence analysis

³[WHO](#) - Recommended composition of the influenza virus vaccines for use in the 2016-17 northern hemisphere influenza season.

⁴Detected in epidemiological week 50. For more details, see [Week 50 report](#)

Antiviral Resistance

During the 2016-17 season, the National Microbiology Laboratory (NML) has tested 313 influenza viruses for resistance to oseltamivir, 312 to zanamivir and 127 to amantadine. All viruses were sensitive to oseltamivir and zanamivir. All 127 influenza A viruses were resistant to amantadine (Table 4).

Table 4 – Antiviral resistance by influenza virus type and subtype, Canada, 2016-17, Week 03

Virus type and subtype	Oseltamivir		Zanamivir		Amantadine	
	# tested	# resistant (%)	# tested	# resistant (%)	# tested	# resistant (%)
A (H3N2)	273	0 (0%)	273	0 (0%)	118	118 (100%)
A (H3N2v)	1	0 (0%)	1	0 (0%)	1	1 (100%)
A (H1N1)	10	0 (0%)	9	0 (0%)	8	8 (100%)
B	29	0 (0%)	29	0 (0%)	NA ¹	NA ¹
TOTAL	313	0 (0%)	312	0 (0%)	127	127 (100%)

¹NA: Not Applicable

Provincial and International Influenza Reports

- [World Health Organization influenza update](#)
- [World Health Organization FluNet](#)
- [WHO Influenza at the human-animal interface](#)
- [Centers for Disease Control and Prevention seasonal influenza report](#)
- [European Centre for Disease Prevention and Control - epidemiological data](#)
- [South Africa Influenza surveillance report](#)
- [New Zealand Public Health Surveillance](#)
- [Australia Influenza Report](#)
- [Pan-American Health Organization Influenza Situation Report](#)
- [Alberta Health – Influenza Surveillance Report](#)
- [BC - Centre for Disease Control \(BCCDC\) - Influenza Surveillance](#)
- [New Brunswick – Influenza Surveillance Reports](#)
- [Newfoundland and Labrador – Surveillance and Disease Reports](#)
- [Nova Scotia - Flu Information](#)
- [Public Health Ontario – Ontario Respiratory Pathogen Bulletin](#)
- [Manitoba – Epidemiology and Surveillance – Influenza Reports](#)
- [Saskatchewan – influenza Reports](#)
- [PEI – Influenza Summary](#)

FluWatch Definitions for the 2016-2017 Season

Abbreviations: Newfoundland/Labrador (NL), Prince Edward Island (PE), New Brunswick (NB), Nova Scotia (NS), Quebec (QC), Ontario (ON), Manitoba (MB), Saskatchewan (SK), Alberta (AB), British Columbia (BC), Yukon (YT), Northwest Territories (NT), Nunavut (NU).

Influenza-like-illness (ILI): Acute onset of respiratory illness with fever and cough and with one or more of the following - sore throat, arthralgia, myalgia, or prostration which is likely due to influenza. In children under 5, gastrointestinal symptoms may also be present. In patients under 5 or 65 and older, fever may not be prominent.

ILI/Influenza outbreaks

Schools: Greater than 10% absenteeism (or absenteeism that is higher (e.g. >5-10%) than expected level as determined by school or public health authority) which is likely due to ILI. Note: it is recommended that ILI school outbreaks be laboratory confirmed at the beginning of influenza season as it may be the first indication of community transmission in an area.

Hospitals and residential institutions: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case. Residential institutions include but not limited to long-term care facilities (LTCF) and prisons.

Workplace: Greater than 10% absenteeism on any day which is most likely due to ILI.

Other settings: two or more cases of ILI within a seven-day period, including at least one laboratory confirmed case; i.e. closed communities.

Note that reporting of outbreaks of influenza/ILI from different types of facilities differs between jurisdictions.

Influenza/ILI Activity Levels

1 = No activity: no laboratory-confirmed influenza detections in the reporting week, however, sporadically occurring ILI may be reported

2 = Sporadic: sporadically occurring ILI and lab confirmed influenza detection(s) with **no outbreaks** detected within the influenza surveillance region†

3 = Localized: (1) evidence of increased ILI* ;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring in **less than 50% of the influenza surveillance region†**

4 = Widespread: (1) evidence of increased ILI*;
(2) lab confirmed influenza detection(s);
(3) **outbreaks** in schools, hospitals, residential institutions and/or other types of facilities occurring **in greater than or equal to 50% of the influenza surveillance region†**

Note: ILI data may be reported through sentinel physicians, emergency room visits or health line telephone calls.

** More than just sporadic as determined by the provincial/territorial epidemiologist.*

† Influenza surveillance regions within the province or territory as defined by the provincial/territorial epidemiologist.

We would like to thank all the Fluwatch surveillance partners who are participating in this year's influenza surveillance program.

This [report](#) is available on the Government of Canada Influenza webpage. Ce rapport est disponible dans les deux langues officielles.