

Environment Canada

Water Science and Technology Directorate

**Direction générale des sciences
et de la technologie, eau**

Environnement Canada

Graphical Representation of Primary Outputs from Energy
Balance Computations Conducted on Lake Erie Central Basin
(1950-1997)

TD
226
N87
no.08-556

Oskar Resler and William Schertzer

WSTD Contribution No. 08-556

**Graphical Representation of Primary Outputs
from Energy Balance Computations
Conducted on Lake Erie Central Basin
(1950-1997)**

**EC Library
Burlington**

Oskar Resler¹ and William Schertzer²

WSTD Contribution # 08-556

Prepared - March 2008

¹ 1610-44 Queen St. N.
Hamilton, ON, Canada, L8R 3H8

² Project Leader, Research Scientist
WS&TD, National Water Research Institute
Canada Centre for Inland Waters
867 Lakeshore Rd.
Burlington, ON, Canada, L7R 4A6

Abstract

As a contribution to the Canada Climate Program, a major research effort was undertaken to examine the long-term heat balance for the lower Great Lakes (Sawchuk and Schertzer, 1988). A new model for evaluating the radiation and surface energy balance components for Lake Ontario and Lake Erie was developed. Improvements included the evaluation of solar radiation from a true radiative transfer model that treats multiple scattering effects explicitly and the evaluation of incoming longwave radiation that incorporated effects of aerosols and clouds. The model derived lake evaporation from a mass transfer formulation that includes dependency on wind speed and atmospheric stability. Accurate evaluation of the lake evaporation component is particularly important since it is a major term in the lake hydrology and longer-term variability in lake levels. A major difficulty in all energy budget studies of lakes over historical periods is the lack of water surface temperatures which is an important variable in evaluation of many of the radiative fluxes and turbulent exchanges at the air-water interface. In this model, a method for estimating surface temperature based on a hysteresis with lake heat content was applied. The relationship represents a strong negative feedback mechanism and provides a constraint that prevents numerical instabilities from amplifying (see Sawchuk and Schertzer, 1988). The model was applied to estimate daily surface radiation and energy balance components for Lake Erie and Lake Ontario over the period 1950 – 1997. This report provides graphical representations of the primary radiative and turbulent exchanges and selected meteorological/limnological components at the daily time scale (1950-1997) for Lake Erie (central basin). The graphical output is particularly relevant for current and future comparisons of the magnitude, variability and interannual variability of key computed variables derived in a consistent method over the 48 year period. In particular, the graphical output documents the limitations in deriving the energy balance of large deep lakes without directly observed water temperatures.

Résumé

En tant que contribution au Programme sur le climat du Canada, un important effort de recherche a été entrepris pour examiner le bilan thermique à long terme des Grands Lacs inférieurs (Sawchuk et Schertzer, 1988). Un nouveau modèle a été créé pour évaluer les éléments du bilan radiatif et du bilan énergétique de surface des lacs Ontario et Érié. Les améliorations comprennent l'évaluation du rayonnement solaire à partir d'un modèle de transfert radiatif vrai qui traite explicitement les effets de diffusion multiple et l'évaluation du rayonnement incident de grandes longueurs d'onde qui a intégré les effets des aérosols et des nuages. Le modèle a calculé l'évaporation des lacs à partir d'une formule de transfert de masse qui inclut la dépendance de la vitesse du vent et de la stabilité atmosphérique. L'évaluation exacte de la composante d'évaporation des lacs est particulièrement importante puisqu'il s'agit d'un élément essentiel de l'hydrologie des lacs et de la variabilité à plus long terme des niveaux des lacs. Une grande difficulté dans toutes les études des bilans énergétiques des lacs sur des périodes historiques est le manque de données sur la température de l'eau de surface, qui est une variable importante dans l'évaluation de nombreux flux radiatifs et échanges turbulents à l'interface air-eau. Le présent modèle a utilisé une méthode d'estimation de la température de surface à partir de l'hystéresis du contenu en chaleur des lacs. La relation représente un solide mécanisme de rétroaction négative et constitue une contrainte qui empêche les instabilités numériques d'augmenter (voir Sawchuk et Schertzer, 1988). Le modèle a servi à estimer les éléments du rayonnement de surface quotidien et du bilan énergétique des lacs Érié et Ontario au cours de la période de 1950 à 1997. Le présent rapport fournit des représentations graphiques des échanges radiatifs et turbulents primaires et de composantes météorologiques et limnologiques sélectionnées sur une échelle quotidienne (1950-1997) pour le lac Érié (bassin central). Les produits graphiques sont particulièrement pertinents pour les comparaisons courantes et futures de l'ampleur, des variations et de la variabilité interannuelle des principaux paramètres calculés par une méthode constante sur la période de 48 ans. Plus particulièrement, les produits graphiques documentent les limites du calcul du bilan énergétique de grands lacs profonds en l'absence de températures de l'eau directement observées.

Table of Contents

| | |
|---|-----|
| Abstract | i |
| Table of Contents | iii |
| Introduction | 1 |
| Elements of the Energy Budget Model Approach | 1 |
| Data File Processing | 3 |
| Table 1: Lake Erie (central basin) Flux and Meteorological Output Parameters | 4 |
| Figure 1: Sample Ordinal Summary | 5 |
| Figure 2: Sample Yearly Summary | 5 |
| Computations Involving Surface Water Temperature | 5 |
| Figure 3: Relationship between surface water temperature and total lake heat content for Lake Erie (central basin). Solid line represents warming period and dashed line represents cooling period (based on Sawchuk and Schertzer, 1988) | 6 |
| Table 2: Root mean square errors of differences between predicted and observed water surface temperatures for Lake Erie (central basin) (Sawchuk and Schertzer, 1988) | 6 |
| Implications Regarding Surface Temperature | 7 |
| References | 9 |
| Appendix 1 Summary Plots | 11 |
| Appendix 2 Heat Fluxes | 31 |
| Appendix 3 Other Fluxes | 79 |
| Appendix 4 Radiative | 127 |
| Appendix 5 Meteorological | 175 |
| Appendix 6 Evaporation | 223 |
| Appendix 7 MATLAB Scripts | 272 |

Introduction

Accurate quantification of the energy exchange across the air-water interface of a lake is crucial for understanding processes such as thermal stratification and the potential impacts of climate / climate change on the lake and aquatic ecosystem components (Schertzer and Sawchuk, 1990). The surface heat exchange can be defined as the following:

$$Q^* - Q_H - Q_{LE} - Q_s - Q_i - Q_M = 0$$

where, Q^* is net radiation, Q_H is sensible heat flux, Q_{LE} is latent heat flux, Q_s is the surface heat flux, Q_i is ice heat flux, and Q_M represents minor heat flux terms such as exchanges resulting from hydrological exchanges. The surface heat flux may then be evaluated as a residual. Details of the component formulations and computational techniques can be found in Sawchuk and Schertzer (1988), Schertzer and Sawchuk (1990).

Evaluation of the energy balance components for a large lake requires measurement of individual components at representative locations over the lake. However, other than for specialized research studies, such data is not usually available and recourse is to model mean meteorological conditions at the lake surface using meteorological stations at the lake periphery. Computational techniques can then be applied to estimate radiation and energy balance components. Such approaches have been applied to long-term studies conducted on the lower Great Lakes.

Derecki (1975) estimated monthly totals of energy budget components for Lake Erie (1950-1968) using shoreline meteorology and water intake temperatures to estimate lake surface temperature. Schertzer (1987) computed daily heat flux components, and heat storage of Lake Erie (lakewide and for the west, central and east basins) which formed boundary conditions for Lake Erie water quality model development (Lam et al., 1983, Lam and Schertzer, 1987). Few comparable long-term studies exist for Lake Ontario, although intensive measurements and calculations of energy balance components were performed during the international Field Year for the Great Lakes (IFYGL) from April 1972 to March 1973 (Pinsak and Rodgers, 1981). Quinn and den Hartog (1983) compared latent and sensible heat fluxes evaluated from solving the energy balance components with other methods such as estimates from the water balance equation and mass transfer models with values recommended by the IFYGL Energy Balance Panel.

This report is focused on the model results of Sawchuk and Schertzer (1998), who developed and applied a new energy budget model approach to Lake Erie and Lake Ontario. The objective here is to provide detailed graphical output of the primary input and output variables from the Sawchuk and Schertzer (1988) investigation.

Elements of the Energy Budget Model Approach

Sawchuk and Schertzer (1988) conducted an investigation to develop a model for evaluating daily values of surface radiation and energy balance components for large deep lakes. The focus of the study was to replace empirical and site specific formulations of the energy budget technique with physical models (where possible) in an effort to develop a numerical algorithm with general application. Considerable effort was expended in ensuring the algorithm was numerically stable, provided accurate flux estimates, required little computer time and performed in all environmental conditions. The model was applied to Lake Erie and its three sub-basins and Lake Ontario for the climatological period 1953 to 1983 to evaluate the magnitude and variability of surface energy

balance components and to compare these estimates, where possible, with estimates from other long-term studies.

Model development focused on three aspects: radiation receipt, evaporation rates and ice cover. Net radiation was estimated from the sum of net shortwave and net longwave radiation fluxes. The shortwave model is a true radiative transfer model and marked the first time that such a model was applied to evaluate shortwave radiation receipts in a boundary layer study. Comparison of model estimates with measured values showed that the model could estimate daily totals of incoming shortwave radiation within, or close to, the range of values measured during IFYGL. Largest differences between model estimates and measured values were observed on days with predominately thin, high-altitude clouds, e.g. cirro-stratus combinations with maximum differences up to 30 percent of measured values. The impact of high altitude clouds on surface radiation receipt was much less when low clouds were also present.

Estimates of longwave radiation were obtained from a variant of Beer's law that incorporated effects of clouds, aerosols, and atmospheric gases on incoming longwave radiation. Estimates of daily totals of incoming longwave radiation were usually within 2 to 3 MJ m⁻² day⁻¹ of measured values which is within the uncertainty of incoming longwave radiation measurements on most days. In general, largest errors in estimates of incoming longwave radiation were observed for mostly clear days resulting in model underestimates. These underestimates may have resulted from several causes, (a) underestimated over-lake air temperatures, (b) inaccurate specification of reference heights for mean atmospheric temperature for emission of radiation and/or inaccurate specification of vertical temperature profile for a mostly clear atmosphere, and (c) neglect of multiple scattering effects. Specification of reference height and vertical temperature profile are likely the most important considerations. Sawchuk and Schertzer (1988) assumed a reference height ranging from 200 m (winter) to 300 m (summer) as recommended by Partridge and Platt (1976). If the reference height was changed to 50 m as recommended by Ulden and Holtslag (1985), mean atmospheric temperature for emission would increase by approximately 2.5 °C. This temperature increase would be sufficient to bring clear sky longwave radiation estimates to within instrumental uncertainty of measured values on most days.

The evaporation model used in Sawchuk and Schertzer (1988) is a variant of the mass transfer approach. The primary improvement is the replacement of the Lake Hefner value for the mass transfer coefficient with a formulation that includes a dependency on wind speed (Quinn, 1978; Quinn and den Hartog, 1981). Although this formulation was strictly applicable to neutral conditions, atmospheric stability is incorporated through scaling of over-lake wind speeds. This approach resulted in evaporation estimates for Lake Ontario during IFYGL that compared favourably with values recommended by the IFYGL Energy Balance Panel. Estimated Bowen ratios were found to compare favourably with values tabulated by Pinsak and Rodgers (1981).

A simple model of ice extent, based on the cumulative freezing day concept provided acceptable estimates of ice heat flux due to ice formation and decay. Although the heat transfer associated with ice formation and decay is small, ice extent is important because of the high albedo for shortwave radiation and the effect of ice in reducing evaporation rates. Comparison of mean ice extent derived from a 31 year simulation for Lake Ontario and Lake Erie with tabulated values of Assel et al. (1983) showed good agreement.

The model was applied to produce 31 year simulations of daily radiation and energy balance components for Lake Erie and Lake Ontario. Daily lake heat contents were obtained from the product of estimated surface heat flux and lake surface area. Heat content estimates were compared

with measured heat contents for both lakes for the period 1967-1982 (Schertzer and Sawchuk, 1985; Schertzer, 1987). Estimated and measured heat contents showed excellent agreement on most days, usually within the accuracy of heat content measurements.

Examination of annual surface heat flux values, obtained from summing daily estimates, reveals that annual surface heat fluxes are not balanced on an annual basis. For Lake Erie, surface heat flux appears to be balanced over periods ranging from three to six years. Lake Ontario appeared to be gaining heat, although gains were small and within the uncertainty of measured heat contents.

One of the important recommendations resulting from the Sawchuk and Schertzer (1988) study was that future research was required for improvement of estimates of surface water temperature. Surface water temperature is the crucial variable in energy balance studies over lakes. It arises in estimation of over-lake meteorology variables, Bowen ratio, and incoming and emitted longwave radiation, etc. Consequently, almost all components of the model require knowledge of surface water temperature. In this investigation, a first approximation was estimated as the hysteresis between surface temperature and lake / basin heat content.

The computations conducted by Sawchuk and Schertzer (1988) for Lake Erie, Lake Erie West, Central and East Basin and Lake Ontario resulted in an enormous input and output database. The final report summarized the results for the 1953 – 1983 period as monthly and annual means. This report provides a detailed graphical summary of primary variables of the lake / basin heat flux, meteorological components and computed evaporation. While the graphs and tables of Sawchuk and Schertzer (1988) served to provide the essential overview of the energy budget conducted over these lakes and basins, the detailed summary plots in this report document the magnitude and variability in the key components at a finer temporal scale. The figures in this report documents the results of the current stage of computations and provides a solid basis for future advanced research on the development of improved techniques and applications of energy budget research for climate change impacts research on large deep lakes.

Data File Processing

The detailed meteorological data supplied for Lake Erie (Central Basin) consisted of plain ASCII text files. Initial processing of the file was required to allow efficient processing by MATLAB and Excel. Two conditions need to be corrected before they could be processed.

Originally, these file were formatted for efficient processing by FORTRAN programs. At first glance, the files looked like they could be processed as space delimited files but, where missing data values occurred in adjacent columns the value “-999.99M” did not leave a separating space. These values were replace with the simpler “-999.99” along with a blank space.

It was also found that some entries consisted of the character string “*****”. These would cause MATLAB to fail, expecting only numeric entries. Since no documentation existed as to the meaning of this string, they were also replaced with the generic “missing value” value.

The parameters present in these meteorological files (see table 1 for a detailed description) were grouped into five categories with detailed daily time series plots produced:

1. Heat Fluxes (Appendix 2),
2. Other Fluxes (Appendix 3),
3. Radiative (Appendix 4),

4. Meteorological (Appendix 5),
5. Evaporation (Appendix 6).

Plotting routines were written using MATLAB to generate yearly plots and to produce some basic statistics. All routines used to process the data file are listed in Appendix 7.

Table 1: Lake Erie (central basin) Flux and Meteorological Output Parameters

| | | |
|----|--------------------------------------|---------------------------------|
| Kd | MJ m ⁻² day ⁻¹ | incoming solar radiation |
| Ku | MJ m ⁻² day ⁻¹ | reflected solar radiation |
| Ld | MJ m ⁻² day ⁻¹ | incoming long wave radiation |
| Lu | MJ m ⁻² day ⁻¹ | emitted long wave radiation |
| Q* | MJ m ⁻² day ⁻¹ | net radiation |
| Uk | m/s | over-lake wind speed |
| Ta | °C | over-lake air temperature |
| Td | °C | over-lake dew point temperature |
| Rh | percent | over-lake relative humidity |
| Tw | °C | surface water temperature |
| As | fractional value | surface albedo |
| Qh | MJ m ⁻² day ⁻¹ | sensible heat flux |
| Qe | MJ m ⁻² day ⁻¹ | latent heat flux |
| Ev | Mm | daily evaporation |
| Qi | MJ m ⁻² day ⁻¹ | ice heat flux |
| Ic | fraction surface area | lake ice extent |
| Qm | MJ m ⁻² day ⁻¹ | minor heat flux terms |
| Qt | MJ m ⁻² day ⁻¹ | total surface heat flux |
| Qc | Exa Joules | lake heat content (computed) |

In addition to the detailed plots, two types of summary plots were also produced (see Appendix 1 for the complete set of summary plots). The first averages the above listed parameters for each ordinal (also sometimes referred to as Julian) day over the study period, in this case 1950 through 1997. The Maximum, Mean and Minimum values are plotted.




Figure 1: Sample Ordinal Summary

The second summary plot consists of simple yearly averages. Each year is represented by a vertical line, Maximum to Minimum (Absolute), along with a horizontal bar representing the Mean.




Figure 2: Sample Yearly Summary

Computations Involving Surface Water Temperature

Sawchuk and Schertzer (1988) modeled surface water temperature as surface temperatures were generally not available till 1966 and from then on there were different databases involving water intake temperatures, airborne radiometer measurements, lake surveillance observations and finally remote sensing from satellite observations. Schertzer and Sawchuk (1985) summarized all available water temperature measurements for the lower Great Lakes. Schertzer computed one dimensional temperature profiles for the lakes and estimated the lake and basin heat contents.

The algorithm used to estimate surface water temperature is based on the relationship between surface water temperature and the total heat content of the lake or basin. The relationships were derived by using paired surface water temperatures and heat content values. Figure 3 illustrates this "hysteresis" relationship for two periods of the year: a warming phase (15 March to 10 September) and a cooling phase (11 September to 14 March) (Sawchuk and Schertzer, 1988). The relationship between surface temperature and heat content was approximated by a linear combination of Legendre Polynomials of order 0 to 5 shown as dashed and solid curves on Figure 3. Higher order approximations did not appreciably reduce the root mean square error of the estimates or improve statistics of the fit (Table 2).




Figure 3: Relationship between surface water temperature and total lake heat content for Lake Erie (central basin). Solid line represents warming period and dashed line represents cooling period (based on Sawchuk and Schertzer, 1988)

Table 2: Root mean square errors of differences between predicted and observed water surface temperatures for Lake Erie (central basin) (Sawchuk and Schertzer, 1988).

| | Warming Phase | Cooling Phase |
|------|---------------|---------------|
| RMSE | 1.12 °C | 0.82 °C |

As indicated in Table 2, the approximations to water surface temperature in the present approach perform better during the cooling phase of the lake. Figure 3 also illustrates that there is larger variability during the warming period of the year, especially during late summer especially when heat content is at or near the maximum. Procedures used to initialize the heat content for a particular year and the computation of the daily mean surface temperature is described in detail in Sawchuk and Schertzer (1988) and also applied in Schertzer and Sawchuk (1990).

Implications Regarding Surface Temperature

Appendix 1 shows a summary of the magnitude and variability of the long-term (45 year) daily computations of the radiative and turbulent fluxes and selected meteorological and limnological variables and succeeding Appendices show the detailed computations for each year. Many of the variables are dependent on the accuracy of water surface temperature; however, water temperature is not often available in historical databases. The application of hysteresis functions, in this case, the relation between surface temperature and lake heat content has advantages and limitations. The advantage is that root mean square errors in the overall temperature approximations are not dissimilar to comparisons between surface temperature observations from different measurement types (e.g. ART vs. satellite vs. buoy observations). However, there are times of the year when variability is larger as noted during the warming phase and, in particular, at the period of maximum heat content. The effect of this variability can be seen in examination of the summary graphs for the surface water temperature in Appendix 1 and also in the Detailed Appendices. In particular, surface water temperature in many cases is constrained at the time of peak heat content. The inaccuracy in the surface water temperature at this time of year will affect computation of the temperature dependent radiative flux and turbulent exchange components.


Future research is required in order to develop new algorithms / approaches that can be applied to approximate water surface temperature in this lake as well as many lakes in Canada that do not have detailed water temperature observations, especially over climatologically significant periods. Such research is required especially in developing baseline conditions for future climate impacts analyses on lakes.

References


- Assel, R.A., Quinn, F.H., Leshkevich, G.A., and Bolsenga, S.J. 1983. Great Lakes Ice Atlas, Great Lakes Environmental Research Laboratory, Ann Arbor, MI, USA. 48104, 115p.
- Sawchuk, A.M., and Schertzer, W.M. (Project Leader), 1988. Climatology of Lake Ontario and Lake Erie Surface Heat Exchanges 1953-1983. NWRI Contribution 88-37, National Water Research Institute, Canada Centre for Inland Waters, 169p.
- Schertzer, W.M. and Sawchuk A.M. 1985. Summary of Water Surface Temperature Observations for Lake Ontario (1966-1984) and Lake Erie (1967 – 1982), NWRI Contribution Number 85-159, Canada Centre for Inland Waters, Burlington, ON, Canada, 49p.
- Schertzer, W.M. Heat balance and heat storage estimates for Lake Erie, *J. Great Lakes Research*, 13(4): 454-4567.
- Schertzer, W.M. and Sawchuk, A.M. 1990. Thermal structure of the lower Great Lakes in a warm year: Implications for the occurrence of hypolimnion anoxia, *Transactions of the American Fisheries Society*, 119(2): 1950-209.
- Lam, D.C.L., Schertzer, W.M., and Fraser, A.S. 1983. Simulation of Lake Erie Water Quality Responses to Loading and Weather Variations, National Water Research Institute, Inland Waters Directorate, Canada Centre for Inland Waters, Burlington, ON, Canada, IWD-Scientific Series Number 134, 310p.
- Lam, D.C.L and Schertzer, W.M. 1987 Lake Erie thermocline model results: Comparison with 1967-1982 Data and relation to anoxic occurrences, *J. Great Lakes Research*, 13(4): 757-769.
- Paltridge, G.W. and Platt, C.M.R. 1976. Radiative Processes in Meteorology and Climatology, Elsevier, New York, 318 p.
- Pinsak, A.P. and Rodgers, G.K. 1981. Energy Balance, Ch. 6 of IFYGL– The International Field Year for the Great Lakes, ed. E.J. Aubert and T.L. Richards, NOAA Great Lakes Environmental Research Laboratory, US Department of Commerce, Ann Arbor, Michigan, 48104: pp. 169-198.
- Quinn, F.H. 1978. An improved aerodynamic evaporation technique for large lakes with application to IFYGL, *Water Resources research*, 15(4): 935-940.
- Quinn, F.H. and Den Hartog, G. 1981. Evaporation Synthesis, Ch. 7 of IFYGL – The International Field Year for the Great Lakes, ed. E.J. Aubert and T.L. Richards, NOAA Great Lakes Environmental Research Laboratory, Ann Arbor, Michigan, 1981: pp. 241-246.
- van Ulden, A.P. and Holtslag, A.A. 1985. Estimation of Atmospheric Boundary Layer Parameters for Diffusion Applications, *J. Applied Meteorology*, 24(11): 1196-1207.

Appendix 1


Summary Plots





Latent Heat Flux (1950-1997)




Latent Heat Flux







Total Surface Heat Flux (1950-1997)




Total Surface Heat Flux





Surface Albedo (1950-1997)




Surface Albedo







Minor Heat Flux Terms (1950-1997)




Minor Heat Flux Terms




Lake Heat Content - Computed (1950-1997)




Lake Heat Content - Computed




Incoming Solar Radiation (1950-1997)




Incoming Solar Radiation




Reflected Solar Radiation (1950-1997)




Reflected Solar Radiation




Incoming Long Wave Radiation (1950-1997)




Incoming Long Wave Radiation




Emitted Long Wave Radiation (1950-1997)




Emitted Long Wave Radiation






Over-Lake Air Temperature (1950-1997)




Over-Lake Air Temperature









Over-Lake Wind Speed (1950-1997)




Over-Lake Wind Speed





Surface Water Temperature (1950-1997)




Surface Water Temperature







Daily Evaporation (1950-1997)




Daily Evaporation




Cumulative Evaporation (1950-1997)





Cumulative Evaporation




Appendix 2 Heat Fluxes










Lake Erie (Central Basin), ON, 1953 Daily Climate Data




Lake Erie (Central Basin), ON, 1954 Daily Climate Data










Lake Erie (Central Basin), ON, 1957 Daily Climate Data







Lake Erie (Central Basin), ON, 1959 Daily Climate Data












Lake Erie (Central Basin), ON, 1963 Daily Climate Data




Lake Erie (Central Basin), ON, 1964 Daily Climate Data




Lake Erie (Central Basin), ON, 1965 Daily Climate Data




Lake Erie (Central Basin), ON, 1966 Daily Climate Data








Lake Erie (Central Basin), ON, 1968 Daily Climate Data




Lake Erie (Central Basin), ON, 1969 Daily Climate Data












Lake Erie (Central Basin), ON, 1973 Daily Climate Data




Lake Erie (Central Basin), ON, 1974 Daily Climate Data




Lake Erie (Central Basin), ON, 1975 Daily Climate Data




Lake Erie (Central Basin), ON, 1976 Daily Climate Data




Lake Erie (Central Basin), ON, 1977 Daily Climate Data







Lake Erie (Central Basin), ON, 1979 Daily Climate Data




Lake Erie (Central Basin), ON, 1980 Daily Climate Data




Lake Erie (Central Basin), ON, 1981 Daily Climate Data




Lake Erie (Central Basin), ON, 1982 Daily Climate Data




Lake Erie (Central Basin), ON, 1983 Daily Climate Data




Lake Erie (Central Basin), ON, 1984 Daily Climate Data




Lake Erie (Central Basin), ON, 1985 Daily Climate Data







Lake Erie (Central Basin), ON, 1987 Daily Climate Data




Lake Erie (Central Basin), ON, 1988 Daily Climate Data




Lake Erie (Central Basin), ON, 1989 Daily Climate Data







Lake Erie (Central Basin), ON, 1991 Daily Climate Data





Lake Erie (Central Basin), ON, 1992 Daily Climate Data




Lake Erie (Central Basin), ON, 1993 Daily Climate Data









Lake Erie (Central Basin), ON, 1996 Daily Climate Data







Appendix 3 Other Fluxes




Lake Erie (Central Basin), ON, 1951 Daily Climate Data




Lake Erie (Central Basin), ON, 1952 Daily Climate Data




Lake Erie (Central Basin), ON, 1953 Daily Climate Data




Lake Erie (Central Basin), ON, 1954 Daily Climate Data




Lake Erie (Central Basin), ON, 1955 Daily Climate Data




Lake Erie (Central Basin), ON, 1956 Daily Climate Data




Lake Erie (Central Basin), ON, 1957 Daily Climate Data




Lake Erie (Central Basin), ON, 1958 Daily Climate Data




Lake Erie (Central Basin), ON, 1959 Daily Climate Data




Lake Erie (Central Basin), ON, 1960 Daily Climate Data




Lake Erie (Central Basin), ON, 1961 Daily Climate Data




Lake Erie (Central Basin), ON, 1962 Daily Climate Data




Lake Erie (Central Basin), ON, 1963 Daily Climate Data




Lake Erie (Central Basin), ON, 1964 Daily Climate Data




Lake Erie (Central Basin), ON, 1965 Daily Climate Data




Lake Erie (Central Basin), ON, 1966 Daily Climate Data




Lake Erie (Central Basin), ON, 1967 Daily Climate Data




Lake Erie (Central Basin), ON, 1968 Daily Climate Data




Lake Erie (Central Basin), ON, 1969 Daily Climate Data




Lake Erie (Central Basin), ON, 1970 Daily Climate Data




Lake Erie (Central Basin), ON, 1971 Daily Climate Data




Lake Erie (Central Basin), ON, 1972 Daily Climate Data




Lake Erie (Central Basin), ON, 1973 Daily Climate Data




Lake Erie (Central Basin), ON, 1974 Daily Climate Data




Lake Erie (Central Basin), ON, 1975 Daily Climate Data




Lake Erie (Central Basin), ON, 1976 Daily Climate Data




Lake Erie (Central Basin), ON, 1977 Daily Climate Data




Lake Erie (Central Basin), ON, 1978 Daily Climate Data




Lake Erie (Central Basin), ON, 1979 Daily Climate Data




Lake Erie (Central Basin), ON, 1980 Daily Climate Data




Lake Erie (Central Basin), ON, 1981 Daily Climate Data




Lake Erie (Central Basin), ON, 1982 Daily Climate Data




Lake Erie (Central Basin), ON, 1983 Daily Climate Data




Lake Erie (Central Basin), ON, 1984 Daily Climate Data




Lake Erie (Central Basin), ON, 1985 Daily Climate Data




Lake Erie (Central Basin), ON, 1986 Daily Climate Data




Lake Erie (Central Basin), ON, 1987 Daily Climate Data




Lake Erie (Central Basin), ON, 1988 Daily Climate Data




Lake Erie (Central Basin), ON, 1989 Daily Climate Data




Lake Erie (Central Basin), ON, 1990 Daily Climate Data




Lake Erie (Central Basin), ON, 1991 Daily Climate Data




Lake Erie (Central Basin), ON, 1992 Daily Climate Data




Lake Erie (Central Basin), ON, 1993 Daily Climate Data




Lake Erie (Central Basin), ON, 1994 Daily Climate Data




Lake Erie (Central Basin), ON, 1995 Daily Climate Data



Lake Erie (Central Basin), ON, 1996 Daily Climate Data




Lake Erie (Central Basin), ON, 1997 Daily Climate Data





Appendix 4

Radiative




Lake Erie (Central Basin), ON, 1951 Daily Climate Data







Lake Erie (Central Basin), ON, 1953 Daily Climate Data




Lake Erie (Central Basin), ON, 1954 Daily Climate Data




Lake Erie (Central Basin), ON, 1955 Daily Climate Data




Lake Erie (Central Basin), ON, 1956 Daily Climate Data




Lake Erie (Central Basin), ON, 1957 Daily Climate Data




Lake Erie (Central Basin), ON, 1958 Daily Climate Data




Lake Erie (Central Basin), ON, 1959 Daily Climate Data




Lake Erie (Central Basin), ON, 1960 Daily Climate Data




Lake Erie (Central Basin), ON, 1961 Daily Climate Data




Lake Erie (Central Basin), ON, 1962 Daily Climate Data




Lake Erie (Central Basin), ON, 1963 Daily Climate Data




Lake Erie (Central Basin), ON, 1964 Daily Climate Data.




Lake Erie (Central Basin), ON, 1965 Daily Climate Data




Lake Erie (Central Basin), ON, 1966 Daily Climate Data




Lake Erie (Central Basin), ON, 1967 Daily Climate Data




Lake Erie (Central Basin), ON, 1968 Daily Climate Data




Lake Erie (Central Basin), ON, 1969 Daily Climate Data




Lake Erie (Central Basin), ON, 1970 Daily Climate Data




Lake Erie (Central Basin), ON, 1971 Daily Climate Data




Lake Erie (Central Basin), ON, 1972 Daily Climate Data




Lake Erie (Central Basin), ON, 1973 Daily Climate Data




Lake Erie (Central Basin), ON, 1974 Daily Climate Data




Lake Erie (Central Basin), ON, 1975 Daily Climate Data




Lake Erie (Central Basin), ON, 1976 Daily Climate Data




Lake Erie (Central Basin), ON, 1977 Daily Climate Data




Lake Erie (Central Basin), ON, 1978 Daily Climate Data




Lake Erie (Central Basin), ON, 1979 Daily Climate Data




Lake Erie (Central Basin), ON, 1980 Daily Climate Data




Lake Erie (Central Basin), ON, 1981 Daily Climate Data





Lake Erie (Central Basin), ON, 1982 Daily Climate Data







Lake Erie (Central Basin), ON; 1984 Daily Climate Data







Lake Erie (Central Basin), ON, 1986 Daily Climate Data






Lake Erie (Central Basin), ON, 1988 Daily Climate Data







Lake Erie (Central Basin), ON, 1990 Daily Climate Data




Lake Erie (Central Basin), ON, 1991 Daily Climate Data




Lake Erie (Central Basin), ON, 1992 Daily Climate Data




Lake Erie (Central Basin), ON, 1993 Daily Climate Data




Lake Erie (Central Basin), ON, 1994 Daily Climate Data







Lake Erie (Central Basin), ON, 1996 Daily Climate Data





Lake Erie (Central Basin), ON, 1997 Daily Climate Data





Appendix 5 Meteorological








































Lake Erie (Central Basin), ON, 1963 Daily Climate Data










































Lake Erie (Central Basin), ON, 1977 Daily Climate Data








Lake Erie (Central Basin), ON, 1979 Daily Climate Data



















































Lake Erie (Central Basin), ON, 1996 Daily Climate Data








Appendix 6 Evaporation



















Lake Erie (Central Basin), ON, 1956 Daily Climate Data









































Lake Erie (Central Basin), ON, 1970 Daily Climate Data





Lake Erie (Central Basin), ON, 1971 Daily Climate Data





Lake Erie (Central Basin), ON, 1972 Daily Climate Data


























Lake Erie (Central Basin), ON, 1981 Daily Climate Data




Lake Erie (Central Basin), ON, 1982 Daily Climate Data





Lake Erie (Central Basin), ON, 1983 Daily Climate Data






Lake Erie (Central Basin), ON, 1985 Daily Climate Data








Lake Erie (Central Basin), ON, 1987 Daily Climate Data




Lake Erie (Central Basin), ON, 1988 Daily Climate Data













Lake Erie (Central Basin), ON, 1992 Daily Climate Data







Lake Erie (Central Basin), ON, 1994 Daily Climate Data









Appendix 7

MATLAB Scripts

ConvertErie_CB.m

Reads data from the original data files and generates a MATLAB compatible data files (one per year). These files are formatted to be compatible the "textread" function (comma delimited ASCII files). This script also generates a Cumulative Evaporation value. This conversion is done to simplify the process of reading in information and to greatly speed up the plotting of the data.

```
%For plotting Lake Erie (Central Basin) Data
%   Years 1950 - 1997

close all
clear all

year=0;
for PlotYear = 1950: 1: 1997
close all

%Internal parameters
%PlotYear = 1965;

if ((PlotYear >= 1950) & (PlotYear <= 1995))
    InFile = 'cbf5095.dat';
else
    InFile = 'cbf9697.dat';
end

OutFile = sprintf('L_Erie_CB_%d.txt', PlotYear);
PrintFile = sprintf('L_Erie_CB_%d', PlotYear);

%Read the file; reformat it; save it in its new textread friendly format.
fid=fopen(InFile,'r');
fid_out=fopen(OutFile,'w');
%Print a header row
fprintf(fid_out,'year, ordinal, Kd, Ku, Ld, Lu, Q*, Uk, Ta, Td, Rh, Tw, As, Qh, Qe, Ev, Qi,
Ic, Qm, Qt, Qc, cumEv\r\n');

count=0;
linecount=0;
SkipLine=0;
clear cumEv;
cumEv=0;
while (~feof(fid) & year <= PlotYear)
```

```

line=fgetl(fid);

clear matches;
matches=findstr(line,'*****');
MatchCount=length(matches);
if (MatchCount > 0)
    SkipLine=1;
end

count=count+1;
if (count>3)
    clear c1 c2 c3 c4 c5 c6 c7 c8
    linecount=linecount+1;
    switch linecount
        case 1
            [year,month,day,ordinal,id]=strread(line,' %d %d %d %d %d');
        case 2
            if ((MatchCount == 0) & (year == PlotYear))
                [Kd,Ku,Ld,Lu,Q,Uk,Ta,Td]=strread(line,' %f %f %f %f %f %f %f %f %f');
            end
        case 3
            if ((MatchCount == 0) & (year == PlotYear))
                [Rh,Tw,As,Qh,Qe,Ev,Qi,Ic]=strread(line, ' %f %f %f %f %f %f %f %f %f');
                if (Ev ~= -999.99)
                    if (count == 6)
                        cumEv = Ev;
                    else
                        cumEv = cumEv + Ev;
                    end
                else
                    Ev = NaN;
                end
            end
        otherwise
            if ((MatchCount == 0) & (year == PlotYear))
                [Qm,Qt,Qc]=strread(line,' %f %f %f');
            end

            linecount=0;
            if (SkipLine==0)
                if (year==PlotYear)
                    fprintf(fid_out,'%d, %d, %f, %f\r\n',...

```

```

year,ordinal,Kd,Ku,Ld,Lu,Q,Uk,Ta,Td,Rh,Tw,As,Qh,Qe,Ev,Qi,Ic,Qm,Qt,Qc,cumEv);
    end
end
SkipLine=0;
end
else
linecount=0;
end
end
fclose(fid);
fclose(fid_out);

end

```

ErieMonthly_CB.m

This script reads data from the converted yearly files and generates three comma delimited (.csv) files containing monthly Max, Min and Mean values (1950 through 1997).

```
% Script To generate Monthly summaries.
```

```

StartYear = 1950;
EndYear = 1997;
OutFile_Mean = 'L_Erie_CB_Monthly_Mean.csv';
OutFile_Max = 'L_Erie_CB_Monthly_Max.csv';
OutFile_Min = 'L_Erie_CB_Monthly_Min.csv';

% Define a data structure.
ds = struct('Year',{},'Data',{});

% Read in the data,
for PlotYear = StartYear: 1: EndYear
close all

InFile = sprintf('L_Erie_CB_%d.txt', PlotYear);

[year,ordinal,Kd,Ku,Ld,Lu,Q,Uk,Ta,Td,Rh,Tw,As,Qh,Qe,Ev,Qi,Ic,Qm,Qt,Qc,cumEv]=textread(InFile
'...
' %d %d %f %f',...
'headerlines',1,'delimiter','','');

data=[year,ordinal,Kd,Ku,Ld,Lu,Q,Uk,Ta,Td,Rh,Tw,As,Qh,Qe,Ev,Qi,Ic,Qm,Qt,Qc,cumEv];

ds(PlotYear).Year=PlotYear;
ds(PlotYear).Data=data;

```

```

clear year ordinal Kd Ku Ld Lu Q_Uk Ta Td Rh Tw As Qh Qe Ev Qi Ic Qm Qt Qc cumEv;
end

%Initialize the results file.
fid_mean=fopen(OutFile_Mean,'w');
fid_max=fopen(OutFile_Max,'w');
fid_min=fopen(OutFile_Min,'w');

%Print a header row
fprintf(fid_mean,'year, month, Kd, Ku, Ld, Lu, Q*, Uk, Ta, Td, Rh, Tw, As, Qh, Qe, Ev, Qi,
Ic, Qm, Qt, Qc, cumEv\r\n');
fprintf(fid_max,'year, month, Kd, Ku, Ld, Lu, Q*, Uk, Ta, Td, Rh, Tw, As, Qh, Qe, Ev, Qi,
Ic, Qm, Qt, Qc, cumEv\r\n');
fprintf(fid_min,'year, month, Kd, Ku, Ld, Lu, Q*, Uk, Ta, Td, Rh, Tw, As, Qh, Qe, Ev, Qi,
Ic, Qm, Qt, Qc, cumEv\r\n');

%Calculate the Monthly means.
for PlotYear = StartYear: 1: EndYear %Loop through the years.
    clear MMean MMax MMin;

    %Determine the start and ends ordinal date for each month.
    IsLeap = mod(PlotYear,4);
    if (IsLeap == 0) %0 represents a leap year.
        Month_Start = [1 32 61 92 122 153 183 214 245 275 306 336];
        Month_End = [31 60 91 121 152 182 213 244 274 305 335 366];
    else
        Month_Start = [1 32 60 91 121 152 182 213 244 274 305 335];
        Month_End = [31 59 90 120 151 181 212 243 273 304 334 365];
    end

    for i = 1: 1: 12 %Loop through the months.
        %Kd
        clear FD Filtered ii
        FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),3);
        [ii] = find(FD ~= -999.99);
        Filtered = FD(ii);
        if (size(Filtered) > 0)
            MMean(i,3) = mean(Filtered);
            MMax(i,3) = max(Filtered);
            MMin(i,3) = min(Filtered);
        else
            MMean(i,3) = -999.99;
            MMax(i,3) = -999.99;
        end
    end

```

```

    MMin(i,3) = -999.99;
end
%Ku
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),4);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,4) = mean(Filtered);
    MMax(i,4) = max(Filtered);
    MMin(i,4) = min(Filtered);
else
    MMean(i,4) = -999.99;
    MMax(i,4) = -999.99;
    MMin(i,4) = -999.99;
end
%Ld
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),5);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,5) = mean(Filtered);
    MMax(i,5) = max(Filtered);
    MMin(i,5) = min(Filtered);
else
    MMean(i,5) = -999.99;
    MMax(i,5) = -999.99;
    MMin(i,5) = -999.99;
end
%Lu
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),6);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,6) = mean(Filtered);
    MMax(i,6) = max(Filtered);
    MMin(i,6) = min(Filtered);
else
    MMean(i,6) = -999.99;
    MMax(i,6) = -999.99;
    MMin(i,6) = -999.99;

```

```

end
%Q
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),7);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,7) = mean(Filtered);
    MMax(i,7) = max(Filtered);
    MMin(i,7) = min(Filtered);
else
    MMean(i,7) = -999.99;
    MMax(i,7) = -999.99;
    MMin(i,7) = -999.99;
end
%Ük
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),8);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,8) = mean(Filtered);
    MMax(i,8) = max(Filtered);
    MMin(i,8) = min(Filtered);
else
    MMean(i,8) = -999.99;
    MMax(i,8) = -999.99;
    MMin(i,8) = -999.99;
end
%Ta
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),9);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,9) = mean(Filtered);
    MMax(i,9) = max(Filtered);
    MMin(i,9) = min(Filtered);
else
    MMean(i,9) = -999.99;
    MMax(i,9) = -999.99;
    MMin(i,9) = -999.99;
end

```

```

%Td
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),10);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,10) = mean(Filtered);
    MMax(i,10) = max(Filtered);
    MMin(i,10) = min(Filtered);
else
    MMean(i,10) = -999.99;
    MMax(i,10) = -999.99;
    MMin(i,10) = -999.99;
end
%Rh
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),11);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,11) = mean(Filtered);
    MMax(i,11) = max(Filtered);
    MMin(i,11) = min(Filtered);
else
    MMean(i,11) = -999.99;
    MMax(i,11) = -999.99;
    MMin(i,11) = -999.99;
end
%Tw
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),12);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,12) = mean(Filtered);
    MMax(i,12) = max(Filtered);
    MMin(i,12) = min(Filtered);
else
    MMean(i,12) = -999.99;
    MMax(i,12) = -999.99;
    MMin(i,12) = -999.99;
end
%As

```

```

clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),13);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,13) = mean(Filtered);
    MMax(i,13) = max(Filtered);
    MMin(i,13) = min(Filtered);
else
    MMean(i,13) = -999.99;
    MMax(i,13) = -999.99;
    MMin(i,13) = -999.99;
end
%Qh
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),14);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,14) = mean(Filtered);
    MMax(i,14) = max(Filtered);
    MMin(i,14) = min(Filtered);
else
    MMean(i,14) = -999.99;
    MMax(i,14) = -999.99;
    MMin(i,14) = -999.99;
end
%Qe
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),15);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,15) = mean(Filtered);
    MMax(i,15) = max(Filtered);
    MMin(i,15) = min(Filtered);
else
    MMean(i,15) = -999.99;
    MMax(i,15) = -999.99;
    MMin(i,15) = -999.99;
end
%Ev
clear FD Filtered ii

```

```

FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),16);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,16) = mean(Filtered);
    MMax(i,16) = max(Filtered);
    MMin(i,16) = min(Filtered);
else
    MMean(i,16) = -999.99;
    MMax(i,16) = -999.99;
    MMin(i,16) = -999.99;
end
%Qi
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),17);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,17) = mean(Filtered);
    MMax(i,17) = max(Filtered);
    MMin(i,17) = min(Filtered);
else
    MMean(i,17) = -999.99;
    MMax(i,17) = -999.99;
    MMin(i,17) = -999.99;
end
%Ic
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),18);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,18) = mean(Filtered);
    MMax(i,18) = max(Filtered);
    MMin(i,18) = min(Filtered);
else
    MMean(i,18) = -999.99;
    MMax(i,18) = -999.99;
    MMin(i,18) = -999.99;
end
%Qm
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),19);

```

```

[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,19) = mean(Filtered);
    MMax(i,19) = max(Filtered);
    MMin(i,19) = min(Filtered);
else
    MMean(i,19) = -999.99;
    MMax(i,19) = -999.99;
    MMin(i,19) = -999.99;
end
%Qt
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),20);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,20) = mean(Filtered);
    MMax(i,20) = max(Filtered);
    MMin(i,20) = min(Filtered);
else
    MMean(i,20) = -999.99;
    MMax(i,20) = -999.99;
    MMin(i,20) = -999.99;
end
%Qc
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),21);
[ii] = find(FD ~= -999.99);
Filtered = FD(ii);
if (size(Filtered) > 0)
    MMean(i,21) = mean(Filtered);
    MMax(i,21) = max(Filtered);
    MMin(i,21) = min(Filtered);
else
    MMean(i,21) = -999.99;
    MMax(i,21) = -999.99;
    MMin(i,21) = -999.99;
end
%cumEv
clear FD Filtered ii
FD = ds(PlotYear).Data(Month_Start(i):Month_End(i),22);
[ii] = find(FD ~= -999.99);

```

```

        Filtered = FD(ii);

        if (size(Filtered) > 0)
            MMean(i,22) = mean(Filtered);
            MMax(i,22) = max(Filtered);
            MMin(i,22) = min(Filtered);
        else
            MMean(i,22) = -999.99;
            MMax(i,22) = -999.99;
            MMin(i,22) = -999.99;
        end

        %Save the results to the output file.

        fprintf(fid_mean,' %d, %d, %f, %f,
        %f, %f, %f, %f, %f\r\n',...
        PlotYear, i, MMean(i,3:22));
        fprintf(fid_max,' %d, %d, %f, %f,
        %f, %f, %f, %f, %f, %f\r\n',...
        PlotYear, i, MMax(i,3:22));
        fprintf(fid_min,' %d, %d, %f, %f,
        %f, %f, %f, %f, %f, %f\r\n',...
        PlotYear, i, MMin(i,3:22));
    end
end

fclose all;

```

Erie_CB_Summary.m

This script generate a comma delimited (.csv) file containing daily Max, Min, Mean and Median values for the 1950 through 1997 years (i.e., maximum temperature for ordinal day 1, over the 48 year span).

% Script for stats gathering of daily met data.

```

StartYear = 1950;
EndYear = 1997;
OutFile = 'L_Erie_CB_Summary.csv';

% Define a data structure.
ds = struct('Year',{},'Data', {});

% Read in the data,
for PlotYear = StartYear: 1: EndYear
    close all

    InFile = sprintf('L_Erie_CB_%d.txt', PlotYear);

```



```

Filtered = Kd(ii);
DataMax(Day,3) = max(Filtered);
DataMin(Day,3) = min(Filtered);
DataMean(Day,3) = mean(Filtered);
DataMedian(Day,3) = median(Filtered);

%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Ku(1,:) ~= -999.99);
Filtered = Ku(ii);
DataMax(Day,4) = max(Filtered);
DataMin(Day,4) = min(Filtered);
DataMean(Day,4) = mean(Filtered);
DataMedian(Day,4) = median(Filtered);

%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Ld(1,:) ~= -999.99);
Filtered = Ld(ii);
DataMax(Day,5) = max(Filtered);
DataMin(Day,5) = min(Filtered);
DataMean(Day,5) = mean(Filtered);
DataMedian(Day,5) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Lu(1,:) ~= -999.99);
Filtered = Lu(ii);
DataMax(Day,6) = max(Filtered);
DataMin(Day,6) = min(Filtered);
DataMean(Day,6) = mean(Filtered);
DataMedian(Day,6) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Q(1,:) ~= -999.99);
Filtered = Q(ii);
DataMax(Day,7) = max(Filtered);
DataMin(Day,7) = min(Filtered);
DataMean(Day,7) = mean(Filtered);
DataMedian(Day,7) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Uk(1,:) ~= -999.99);
Filtered = Uk(ii);

```

```

DataMax(Day,8) = max(Filtered);
DataMin(Day,8) = min(Filtered);
DataMean(Day,8) = mean(Filtered);
DataMedian(Day,8) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Ta(1,:) ~= -999.99);
Filtered = Ta(ii);
DataMax(Day,9) = max(Filtered);
DataMin(Day,9) = min(Filtered);
DataMean(Day,9) = mean(Filtered);
DataMedian(Day,9) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Td(1,:) ~= -999.99);
Filtered = Td(ii);
DataMax(Day,10) = max(Filtered);
DataMin(Day,10) = min(Filtered);
DataMean(Day,10) = mean(Filtered);
DataMedian(Day,10) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Rh(1,:) ~= -999.99);
Filtered = Rh(ii);
DataMax(Day,11) = max(Filtered);
DataMin(Day,11) = min(Filtered);
DataMean(Day,11) = mean(Filtered);
DataMedian(Day,11) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Tw(1,:) ~= -999.99);
Filtered = Tw(ii);
DataMax(Day,12) = max(Filtered);
DataMin(Day,12) = min(Filtered);
DataMean(Day,12) = mean(Filtered);
DataMedian(Day,12) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(As(1,:) ~= -999.99);
Filtered = As(ii);
DataMax(Day,13) = max(Filtered);
DataMin(Day,13) = min(Filtered);
DataMean(Day,13) = mean(Filtered);

```

```

DataMedian(Day,13) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Qh(1,:) ~= -999.99);
Filtered = Qh(ii);
DataMax(Day,14) = max(Filtered);
DataMin(Day,14) = min(Filtered);
DataMean(Day,14) = mean(Filtered);
DataMedian(Day,14) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Qe(1,:) ~= -999.99);
Filtered = Qe(ii);
DataMax(Day,15) = max(Filtered);
DataMin(Day,15) = min(Filtered);
DataMean(Day,15) = mean(Filtered);
DataMedian(Day,15) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Ev(1,:) ~= -999.99);
Filtered = Ev(ii);
DataMax(Day,16) = max(Filtered);
DataMin(Day,16) = min(Filtered);
DataMean(Day,16) = mean(Filtered);
DataMedian(Day,16) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Qi(1,:) ~= -999.99);
Filtered = Qi(ii);
DataMax(Day,17) = max(Filtered);
DataMin(Day,17) = min(Filtered);
DataMean(Day,17) = mean(Filtered);
DataMedian(Day,17) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Ic(1,:) ~= -999.99);
Filtered = Ic(ii);
DataMax(Day,18) = max(Filtered);
DataMin(Day,18) = min(Filtered);
DataMean(Day,18) = mean(Filtered);
DataMedian(Day,18) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;

```

```

[ii] = find(Qm(1,:) ~= -999.99);
Filtered = Qm(ii);
DataMax(Day,19) = max(Filtered);
DataMin(Day,19) = min(Filtered);
DataMean(Day,19) = mean(Filtered);
DataMedian(Day,19) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Qt(1,:) ~= -999.99);
Filtered = Qt(ii);
DataMax(Day,20) = max(Filtered);
DataMin(Day,20) = min(Filtered);
DataMean(Day,20) = mean(Filtered);
DataMedian(Day,20) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(Qc(1,:) ~= -999.99);
Filtered = Qc(ii);
DataMax(Day,21) = max(Filtered);
DataMin(Day,21) = min(Filtered);
DataMean(Day,21) = mean(Filtered);
DataMedian(Day,21) = median(Filtered);
%Only look at non "missing number" numbers
clear Filtered;
[ii] = find(cumEv(1,:) ~= -999.99);
Filtered = cumEv(ii);
DataMax(Day,22) = max(Filtered);
DataMin(Day,22) = min(Filtered);
DataMean(Day,22) = mean(Filtered);
DataMedian(Day,22) = median(Filtered);

clear Kd Ku Ld Lu Q Uk Ta Td Rh Tw As Qh Qe Ev Qi Ic Qm Qt Qc cumEv;
end

%Save the results to a text file

fid_out=fopen(OutFile,'w');
%Print a header row
fprintf(fid_out,'ordinal, Kd, Ku, Ld, Lu, Q*, Uk, Ta, Td, Rh, Tw, As, Qh, Qe, Ev, Qi, Ic,
Qm, Qt, Qc, cumEv\r\n');
fprintf(fid_out,'Max (%d-%d)\r\n',StartYear,EndYear)
for Day = 1: 1: 365

```

```

fprintf(fid_out,'%d, %f, %f,
%f, %f, %f\r\n',...
    Day,...

DataMax(Day,3),DataMax(Day,4),DataMax(Day,5),DataMax(Day,6),DataMax(Day,7),DataMax(Day,8),Da
taMax(Day,9),...

DataMax(Day,10),DataMax(Day,11),DataMax(Day,12),DataMax(Day,13),DataMax(Day,14),DataMax(Day,
15),DataMax(Day,16),...

DataMax(Day,17),DataMax(Day,18),DataMax(Day,19),DataMax(Day,20),DataMax(Day,21),DataMax(Day,
22));
end

fprintf(fid_out,'Min (%d-%d)\r\n',StartYear,EndYear);
for Day = 1: 1: 365
    fprintf(fid_out,'%d, %f, %f,
%f, %f, %f\r\n',...
        Day,...

DataMin(Day,3),DataMin(Day,4),DataMin(Day,5),DataMin(Day,6),DataMin(Day,7),DataMin(Day,8),Da
taMin(Day,9),...

DataMin(Day,10),DataMin(Day,11),DataMin(Day,12),DataMin(Day,13),DataMin(Day,14),DataMin(Day,
15),DataMin(Day,16),...

DataMin(Day,17),DataMin(Day,18),DataMin(Day,19),DataMin(Day,20),DataMin(Day,21),DataMin(Day,
22)');
end

fprintf(fid_out,'Mean (%d-%d)\r\n',StartYear,EndYear);
for Day = 1: 1: 365
    fprintf(fid_out,'%d, %f, %f,
%f, %f, %f\r\n',...
        Day,...

DataMean(Day,3),DataMean(Day,4),DataMean(Day,5),DataMean(Day,6),DataMean(Day,7),DataMean(Day
,8),DataMean(Day,9),...

DataMean(Day,10),DataMean(Day,11),DataMean(Day,12),DataMean(Day,13),DataMean(Day,14),DataMea
n(Day,15),DataMean(Day,16),...

DataMean(Day,17),DataMean(Day,18),DataMean(Day,19),DataMean(Day,20),DataMean(Day,21),DataMea
n(Day,22));
end

fprintf(fid_out,'Median (%d-%d)\r\n',StartYear,EndYear);
for Day = 1: 1: 365
    fprintf(fid_out,'%d, %f, %f,
%f, %f, %f\r\n',...
        Day,...

DataMedian(Day,3),DataMedian(Day,4),DataMedian(Day,5),DataMedian(Day,6),DataMedian(Day,7),Da
taMedian(Day,8),DataMedian(Day,9),...

```

```

DataMedian(Day,10),DataMedian(Day,11),DataMedian(Day,12),DataMedian(Day,13),DataMedian(Day,1
4),DataMedian(Day,15),DataMedian(Day,16),...

DataMedian(Day,17),DataMedian(Day,18),DataMedian(Day,19),DataMedian(Day,20),DataMedian(Day,2
1),DataMedian(Day,22));
end
fclose(fid_out);

```

PlotErie_CB.m

This script generates daily time series plots for all parameters contained within the converted data files. One plot is generated per year.

```

%For plotting Lake Erie (Central Basin) Data
% Years 1950 - 1997

close all
clear all

for PlotYear = 1950: 1: 1997
close all

OutFile = sprintf('L_Erie_CB_%d.txt', PlotYear);
PrintFile = sprintf('L_Erie_CB_%d', PlotYear);

PlotTitle=sprintf('Lake Erie (Central Basin), ON, %d Daily Climate Data',PlotYear);

%Read the Data
[year,ordinal,Kd,Ku,Ld,Lu,Q,Uk,Ta,Td,Rh,Tw,As,Qh,Qe,Ev,Qi,Ic,Qm,Qt,Qc,cumEv]=textread(OutFil
e, ...
    '%d %d %f %f',...
    'headerlines',1,'delimiter','','');
data=[year,ordinal,Kd,Ku,Ld,Lu,Q,Uk,Ta,Td,Rh,Tw,As,Qh,Qe,Ev,Qi,Ic,Qm,Qt,Qc,cumEv];
clear year ordinal Kd Ku Ld Lu Q Uk Ta Td Rh Tw As Qh Qe Ev Qi Ic Qm Qt Qc cumEv

%Remove "Missing Data" values.
for i=1:22 %Number of columns
    [ii]=find(data(:,i) == -999.99);
    data(ii,i) = NaN;
end

%Plot the data
TickLabelFontSize = 8;

```

```

AxisLabelFontSize = 10;
TitleSize = 12;

IsLeap = mod(PlotYear,4);
if (IsLeap == 0) %0 represents a leap year.
    X_Ticks = [1 32 61 92 122 153 183 214 245 275 306 336];
    X_Labels = 'J|F|M|A|M|J|J|A|S|O|N|D';
else
    X_Ticks = [1 32 60 91 121 152 182 213 244 274 305 335];
    X_Labels = 'J|F|M|A|M|J|J|A|S|O|N|D';
end
X_Label_Ticks = [15 47 75 106 136 168 197 228 259 289 320 350];

%Set the data for the x-axis
x=data(:,2);
XMin=1;
XMax=366;

%=====
% Page 1 Plots
%
% Radiative Exchanges
% Kd (MJ m^-2 day^-1)
% Ku (MJ m^-2 day^-1)
% Ld (MJ m^-2 day^-1)
% Lu (MJ m^-2 day^-1)
%
%-----

close all
clear Y1 Y2 Y3 Y4

Y1 = data(:,3);
Y2 = data(:,4);
Y3 = data(:,5);
Y4 = data(:,6);

%'-k' = Solid Black Line
subplot(4,1,1),plot(x,Y1,'-k');
orient tall;
title(PlotTitle,'fontsize',TitleSize);
set(gca,'fontsize',TickLabelFontSize);
ylabel('Kd (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.

```

```

axis([XMin XMax 0 35]);
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,2),plot(x,Y2,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Ku (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 0 14]);
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,3),plot(x,Y3,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Ld (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 15 40]);
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,4),plot(x,Y4,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Lu (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 25 40]);
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

% Plot a second Axis to place Months between tick marks.
ax1 = gca;
ax2 = axes('Position',get(ax1,'Position'), ...
    'XAxisLocation','bottom', 'YAxisLocation', 'left', ...
    'Color', 'none', ...
    'XColor', 'k', 'YColor', 'k');
set(ax2,'fontsize',TickLabelFontSize);

```

```

set(ax2,'XLim',get(ax1,'XLim'));
set(ax2,'YLim',get(ax1,'YLim'));
set(ax2,'XTick',X_Label_Ticks);
set(ax2,'XTickLabel',X_Labels);
set(ax2,'YTickLabel',[]);
set(ax2,'TickLength',[0 0]);

%Save the plot to a file.
PlotFile = ['Radiative ', PrintFile];
print('-dmeta', '-r600', PlotFile);

%=====
% Page 2 Plots
%
% Heat Fluxes
% Q* (MJ m^-2 day^-1)
% Qe (MJ m^-2 day^-1)
% Qh (MJ m^-2 day^-1)
% Qt (MJ m^-2 day^-1)
%-----

close all
clear Y1 Y2 Y3 Y4

Y1 = data(:,7);
Y2 = data(:,15);
Y3 = data(:,14);
Y4 = data(:,20);

%'-k' = Solid Black Line
subplot(4,1,1),plot(x,Y1,'-k');
orient tall;
title(PlotTitle,'fontsize',TitleSize);
set(gca,'fontsize',TickLabelFontSize);
ylabel('Q* (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax -5 30])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'YTickLabel',[]);
%YLim('Auto');

```

```

    subplot(4,1,2),plot(x,Y2,'-k')
    set(gca,'fontsize',TickLabelFontSize);
    ylabel('Qe (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
    %set the X and Y axis limits. Set Y axis to auto scaling.
    axis([XMin XMax -10 50])
    XLim('Manual');
    set(gca,'XTick',X_Ticks);
    set(gca,'XTickLabel',[]);
    %YLim('Auto');

    subplot(4,1,3),plot(x,Y3,'-k')
    set(gca,'fontsize',TickLabelFontSize);
    ylabel('Qh (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
    %set the X and Y axis limits. Set Y axis to auto scaling.
    axis([XMin XMax -80 40])
    XLim('Manual');
    set(gca,'XTick',X_Ticks);
    set(gca,'XTickLabel',[]);
    %YLim('Auto');

    subplot(4,1,4),plot(x,Y4,'-k')
    set(gca,'fontsize',TickLabelFontSize)
    ylabel('Qt (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
    %set the X and Y axis limits. Set Y axis to auto scaling.
    axis([XMin XMax -80 40])
    XLim('Manual');
    set(gca,'XTick',X_Ticks);
    set(gca,'XTickLabel',[]);
    %YLim('Auto');

    % Plot a second Axis to place Months between tick marks.
    ax1 = gca;
    ax2 = axes('Position',get(ax1,'Position'), ...
        'XAxisLocation','bottom', 'YAxisLocation', 'left', ...
        'Color', 'none', ...
        'XColor', 'k', 'YColor', 'k');
    set(ax2,'fontsize',TickLabelFontSize);
    set(ax2,'XLim',get(ax1,'XLim'));
    set(ax2,'YLim',get(ax1,'YLim'));
    set(ax2,'XTick',X_Label_Ticks);
    set(ax2,'XTickLabel',X_Labels);
    set(ax2,'YTickLabel',[]);
    set(ax2,'TickLength',[0 0]);

```

```

%Save the plot to a file.
PlotFile = ['HeatFluxes_', PrintFile];
print('-dmeta', '-r600', PlotFile);

%=====
% Page 3 Plots
%
% Meteorology
% Ta (Deg C)
% Td (Dec C)
% Rh (%)
% Uk (m/s)
%=====

close all
clear Y1 Y2 Y3 Y4

Y1 = data(:,9);
Y2 = data(:,10);
Y3 = data(:,11);
Y4 = data(:,8);

%-k' = Solid Black Line
subplot(4,1,1),plot(x,Y1,'-k');
orient tall;
title(PlotTitle,'fontsize',TitleSize);
set(gca,'fontsize',TickLabelFontSize);
ylabel('Ta (^oC)', 'FontSize', AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax -20 30])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,2),plot(x,Y2,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Td (^oC)', 'FontSize', AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax -30 30])
XLim('Manual');

```

```

set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,3),plot(x,Y3,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Rh (%)','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 0 100])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,4),plot(x,Y4,'-k')
set(gca,'fontsize',TickLabelFontSize)
ylabel('Uk (m/s)','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 0 16])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

% Plot a second Axis to place Months between tick marks.
ax1 = gca;
ax2 = axes('Position',get(ax1,'Position'), ...
    'XAxisLocation','bottom', 'YAxisLocation', 'left',...
    'Color', 'none',...
    'XColor', 'k', 'YColor', 'k');
set(ax2,'fontsize',TickLabelFontSize);
set(ax2,'XLim',get(ax1,'XLim'));
set(ax2,'YLim',get(ax1,'YLim'));
set(ax2,'XTick',X_Label_Ticks);
set(ax2,'XTickLabel',X_Labels);
set(ax2,'YTickLabel',[]);
set(ax2,'TickLength',[0 0]);

%Save the plot to a file.
PlotFile = ['Meteorology_', PrintFile];
print('-dmeta', '-r600', PlotFile);

```

```

=====
% Page 4 Plots
%
% Other fluxes
% As (fraction)
% Qi (MJ/m^2/day)
% Qm (MJ/m^2/day)
% Qc (Exa Joules)
-----

close all
clear Y1 Y2 Y3 Y4

Y1 = data(:,13);
Y2 = data(:,17);
Y3 = data(:,19);
Y4 = data(:,21);

%'-k' = Solid Black Line
subplot(4,1,1),plot(x,Y1,'-k');
orient tall;
title(PlotTitle,'fontsize',TitleSize);
set(gca,'fontsize',TickLabelFontSize);
ylabel('As (fractional value)', 'FontSize', AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 0 0.7])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,2),plot(x,Y2,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Qi (MJ m^{-2} day^{-1})', 'FontSize', AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax -6 8])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,3),plot(x,Y3,'-k')
set(gca,'fontsize',TickLabelFontSize);

```

```

ylabel('Qm (MJ m^{-2} day^{-1})','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax -0.5 0.5])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,4),plot(x,Y4,'-k')
set(gca,'fontsize',TickLabelFontSize)
ylabel('Qc (x10^{18} Joules)','FontSize',AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax -10 30])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

% Plot a second Axis to place Months between tick marks.
ax1 = gca;
ax2 = axes('Position',get(ax1,'Position'), ...
'XAxisLocation','bottom', 'YAxisLocation', 'left', ...
'Color', 'none', ...
'XColor', 'k', 'YColor', 'k');
set(ax2,'fontsize',TickLabelFontSize);
set(ax2,'XLim',get(ax1,'XLim'));
set(ax2,'YLim',get(ax1,'YLim'));
set(ax2,'XTick',X_Label_Ticks);
set(ax2,'XTickLabel',X_Labels);
set(ax2,'YTickLabel',[]);
set(ax2,'TickLength',[0 0]);

%Save the plot to a file.
PlotFile = ['OtherFluxes_', PrintFile];
print('-dmeta', '-r600', PlotFile);

% =====
% Page 5 Plots
%
% Evaporation
% Tw (Deg C)
% Ic (fraction surface area)
% Ev (mm)

```

```

% cumEv (mm)
% -----
%
close all
clear Y1 Y2 Y3 Y4

Y1 = data(:,12);
Y2 = data(:,18);
Y3 = data(:,16);
Y4 = data(:,22);

%'-k' = Solid Black Line
subplot(4,1,1),plot(x,Y1,'-k');
orient tall;
title(PlotTitle,'fontsize',TitleSize);
set(gca,'fontsize',TickLabelFontSize);
ylabel('Tw (^oC)', 'FontSize', AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 0 25])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,2),plot(x,Y2,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Ic (fraction surface area)', 'FontSize', AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax 0 100])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

subplot(4,1,3),plot(x,Y3,'-k')
set(gca,'fontsize',TickLabelFontSize);
ylabel('Ev (mm)', 'FontSize', AxisLabelFontSize);
%set the X and Y axis limits. Set Y axis to auto scaling.
axis([XMin XMax -5 20])
XLim('Manual');
set(gca,'XTick',X_Ticks);
set(gca,'XTickLabel',[]);
%YLim('Auto');

```

```

    subplot(4,1,4),plot(x,Y4,'-k')
    set(gca,'fontsize',TickLabelFontSize)
    ylabel('cumEv (mm)', 'FontSize', AxisLabelFontSize);
    %set the X and Y axis limits. Set Y axis to auto scaling.
    axis([XMin XMax 0 1200])
    XLim('Manual');
    set(gca,'XTick',X_Ticks);
    set(gca,'XTickLabel',[]);
    %YLim('Auto');

    % Plot a second Axis to place Months between tick marks.
    ax1 = gca;
    ax2 = axes('Position',get(ax1,'Position'), ...
    'XAxisLocation','bottom', 'YAxisLocation', 'left', ...
    'Color', 'none', ...
    'XColor', 'k', 'YColor', 'k');
    set(ax2,'fontsize',TickLabelFontSize);
    set(ax2,'XLim',get(ax1,'XLim'));
    set(ax2,'YLim',get(ax1,'YLim'));
    set(ax2,'XTick',X_Label_Ticks);
    set(ax2,'XTickLabel',X_Labels);
    set(ax2,'YTickLabel',[]);
    set(ax2,'TickLength',[0 0]);

    %Save the plot to a file.
    PlotFile = ['Evaporation_', PrintFile];
    print('-dmeta', '-r600', PlotFile);

end

```

ENVIRONMENT CANADA LIBRARY, BURLINGTON



3 9055 1016 5803 6



**Environment
Canada**

**Environnement
Canada**

Canada

Canada Centre for Inland Waters
P.O. Box 5050
867 Lakeshore Road
Burlington, Ontario
L7R 4A6 Canada

National Hydrology Research Centre
11 Innovation Boulevard
Saskatoon, Saskatchewan
S7N 3H5 Canada

St. Lawrence Centre
105 McGill Street
Montreal, Québec
H2Y 2E7 Canada

Place Vincent Massey
351 St. Joseph Boulevard
Gatineau, Quebec
K1A 0H3 Canada

Centre canadien des eaux intérieures
Case postale 5050
867, chemin Lakeshore
Burlington (Ontario)
L7R 4A6 Canada

Centre national de recherche en hydrologie
11, boul. Innovation
Saskatoon (Saskatchewan)
S7N 3H5 Canada

Centre Saint-Laurent
105, rue McGill
Montréal (Québec)
H2Y 2E7 Canada

Place Vincent-Massey
351 boul. St-Joseph
Gatineau (Québec)
K1A 0H3 Canada