



Natural Resources  
Canada

Ressources naturelles  
Canada

2nd  
EDITION

# **CANADIAN GEOSCIENCE MAP 212**

## **SURFICIAL GEOLOGY**

# **TANQUARY FIORD**

Nunavut  
NTS 340-D



**Map Information  
Document**

**Preliminary**

**Geological Survey of Canada  
Canadian Geoscience Maps**

**2016**

**Canada** 



## **MAP NUMBER**

Natural Resources Canada, Geological Survey of Canada  
Canadian Geoscience Map 212 (Preliminary, 2nd Edition)

## **TITLE**

Surficial geology, Tanquary Fiord, Nunavut, NTS 340-D

## **SCALE**

1:250 000

## **CATALOGUE INFORMATION**

Catalogue No. M183-1/212-2016E-PDF  
ISBN 978-0-660-05372-1  
doi:10.4095/298701

## **COPYRIGHT**

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources, 2016

Information contained in this publication or product may be reproduced, in part or in whole, and by any means, for personal or public non-commercial purposes, without charge or further permission, unless otherwise specified.

You are asked to:

- exercise due diligence in ensuring the accuracy of the materials reproduced;
- indicate the complete title of the materials reproduced, and the name of the author organization; and
- indicate that the reproduction is a copy of an official work that is published by Natural Resources Canada (NRCan) and that the reproduction has not been produced in affiliation with, or with the endorsement of, NRCan.

Commercial reproduction and distribution is prohibited except with written permission from NRCan. For more information, contact NRCan at [nrcan.copyrightdroitdauteur.nrcan@canada.ca](mailto:nrcan.copyrightdroitdauteur.nrcan@canada.ca).

## **RECOMMENDED CITATION**

Bednarski, J.M., 2016. Surficial geology, Tanquary Fiord, Nunavut, NTS 340-D; Geological Survey of Canada, Canadian Geoscience Map 212 (2nd edition, preliminary), scale 1:250 000. doi:10.4095/298701

## **ABSTRACT**

The Quttinirpaaq National Park region is mountainous including the highest peak in eastern North America. Glaciers cover about half of the map area, including unique floating ice shelves along the north coast, which have largely broken up in the last few

decades but many fiord and valley glaciers still contact the sea. Sedimentary rock outcrops form the dominant surficial unit, including large areas of frost shattered rubble mantling broad summits and slopes. Glacial debris is also widespread normally forming a thin discontinuous veneer of till or as isolated erratic boulders. In the past, as now, the main source of run-off and sediment is supplied by glacial meltwater. Consequently most of the unconsolidated sediment lies within formerly glaciated valleys and coastlines where there are complex associations of moraines, glaciofluvial and glaciomarine deposits related to the advance and retreat of valley glaciers and ice caps, coupled with falling sea levels caused by postglacial crustal uplift.

## **RÉSUMÉ**

La région du Parc national Quttinirpaaq est une zone montagneuse, avec le plus haut sommet de la partie est de l'Amérique du Nord. Les glaciers couvrent environ la moitié de la région de la carte et des plateformes uniques de glace flottante, qui se sont en grande partie disloquées au cours des dernières décennies, sont présentes le long de la côte nord, mais des glaciers de vallée sont encore en contact avec la mer à la tête de nombreux fjords. Des affleurements de roches sédimentaires constituent l'unité superficielle dominante, et comportent de grandes étendues de blocaille résultant de la gélifraction de la roche qui couvrent les larges sommets et les versants. Des débris glaciaires non triés, déposés lors de la dernière glaciation, sont également répandus, mais se présentent habituellement sous forme d'un mince placage de till discontinu ou de blocs erratiques isolés, dispersés à la surface du substratum rocheux. À l'heure actuelle, comme dans le passé, les eaux de fonte glaciaire représentent la principale source de ruissellement et d'apport de sédiments. Ainsi, la plupart des sédiments meubles reposent dans d'anciennes vallées glaciaires et sur les côtes, là où se trouvent des associations complexes de sédiments de moraines, de dépôts fluvioglaciaires et glaciomarins associés à l'avancée et au retrait des glaciers de vallée, en combinaison avec la baisse du niveau de la mer causée par le relèvement postglaciaire de la croûte terrestre.

## **LICENCE AGREEMENT**

View the license agreement at

<http://open.canada.ca/en/open-government-licence-canada>

## **ACCORD DE LICENCE**

Voir l'accord de licence à

<http://ouvert.canada.ca/fr/licence-du-gouvernement-ouvert-canada>

## **SHEET 1 OF 1, SURFICIAL GEOLOGY**

### **GENERAL INFORMATION**

Author: J.M. Bednarski

Geology based on aerial photograph interpretation and field work by J.M. Bednarski in 1979 to 1988. Additional geological field data by R.L. Christie in 1954, 1957, 1958.

Geology conforms to Surficial Data Model v. 2.0.1

Geological compilation by D.E. Kerr, 2013–2014

Data conversion by F. Fortin, 2013 and S. Eagles, 2013, 2014

Geomatics by F. Fortin

Cartography by T. Konopelky

Initiative of the Geological Survey of Canada as part of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) Program, with participation from Parks Canada Agency – Nunavut Field Unit.

Map projection Universal Transverse Mercator, zone 18.  
North American Datum 1983

Base map at the scale of 1:250 000 from Natural Resources Canada, with modifications.

Elevations in feet above mean sea level

Proximity to the North Magnetic Pole causes the magnetic compass to be erratic in this area.

Mean magnetic declination 2016, 56°46'W, decreasing 75.3' annually. Readings vary from 54°10'W in the SE corner to 59°23'W in the NW corner of the map.

This map is not to be used for navigational purposes.

Title photograph: Glaciofluvial sediments, Tanquary Fiord, Ellesmere Island, Nunavut.  
Photograph by J.M. Bednarski. 2013-083

The Geological Survey of Canada welcomes corrections or additional information from users.

Data may include additional observations not portrayed on this map. See map info document accompanying the downloaded data for more information about this publication.

This publication is available for free download through GEOSCAN (<http://geoscan.nrcan.gc.ca/>).

Preliminary publications in this series have not been scientifically edited.

### **MAP VIEWING FILES**

The published map is distributed as a Portable Document File (PDF), and may contain a subset of the overall geological data for legibility reasons at the publication scale.

### **DESCRIPTIVE NOTES**

The 1:250 000 scale map of the Quttinirpaaq National Park of Canada is based on information compiled from aerial photographs of a scale of approximately 1:64 000. This represents a fourfold reduction in scale which necessitated some generalization and combining of units. The reduction process involved an intermediate step of plotting the information on base maps of 1:125 000 scale. Plotting and scale reduction was accomplished with a projection table on which corrections for distortion could be made.

The most substantial amalgamation was for glaciofluvial sediments (unit GF). These deposits are commonly concentrated in narrow valleys where they form terraces (unit GFt), active outwash plains (unit GFp), outwash fans (unit GFf), and combinations of the three. Except where large terraces are found (unit GFt), some of these units were combined on the final map. Melting and undercutting of permafrost in these sediments produces thermokarst and slumping.

The 1:250 000 scale map also includes the amalgamation of bedrock units: weathered bedrock (felsenmeer; unit W) and unaltered bedrock (unit R). Different tills were also amalgamated. On slopes, both R and T units were transported downslope by periglacial processes (e.g. C units).

The glacier cover on the topographic base map is generalized and not very accurate in some areas, therefore, minor supraglacial features interpreted on the airphotos were not retained on the final map.

During the scale reduction process the units become progressively more generalized as the boundaries between adjacent units become smoothed out. This means that actual delineations on the airphotos, are not entirely 'real' on the final map. Nonetheless, an attempt was made to maintain the character of the original landscape.

Ice shelves mapped along the north coast were originally included as a basic unit, but large changes have occurred along the margins in recent years which may make some of the delineations on the airphotos taken in the late 1950's inaccurate. Due to this difficulty, the ice shelf designation represents generalized polygons based on extent of glaciers between 1959 and 2011; data vary locally depending on date of glacier ice coverage. Some valley glaciers were also classed as Isn when they float in deep fiords and display ice shelf morphology.

### **REFERENCES**

- Christie, R.L., 1966. Surficial geology, northeastern Ellesmere Island, District of Franklin and northwestern Greenland; Geological Survey of Canada, Map 1192A, scale 1:506 880. doi:10.4095/107408
- Deblonde, C., Plouffe, A., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Kerr, D.E., Moore, A., Parent, M., Robertson, L., Smith, I.R., St-Onge, D.A., and Weatherston, A., 2014. Science language for an integrated Geological Survey of Canada data model for surficial geology maps, version 2.0; Geological Survey of Canada, Open File 7631, 464 p. doi:10.4095/294225

### **AUTHOR CONTACT**

Questions, suggestions, and comments regarding the geological information contained in the data sets should be addressed to:

J.M. Bednarski  
Geological Survey of Canada

9860 West Saanich Road, P.O. Box 6000  
Sidney, British Columbia  
V8L4B2  
[ian.bednarski@canada.ca](mailto:ian.bednarski@canada.ca)

### **COORDINATE SYSTEM**

Projection: Universal Transverse Mercator  
Units: metres  
Zone: 18  
Horizontal Datum: NAD83  
Vertical Datum: mean sea level

### **BOUNDING COORDINATES**

Western longitude: 80°00'00"W  
Eastern longitude: 72°00'00"W  
Northern latitude: 82°00'00"N  
Southern latitude: 81°00'00"N

### **SOFTWARE VERSION**

Data has been originally compiled and formatted for use with ArcGIS™ desktop version 10.2.2 developed by ESRI®.

### **DATA MODEL INFORMATION**

#### **Surficial**

The Geological Survey of Canada (GSC) through the Geo-mapping for Energy and Minerals Program (GEM) has undertaken the Geological Map Flow to develop protocols for the collection, management (compilation, interpretation), and dissemination of surficial and bedrock geology data and map information. To this end, a data model has been created.

The Surficial Data Model (SDM) was designed using ESRI geodatabase architecture. The XML workspace document provided can be imported into a geodatabase, and the geodatabase will then be populated with the feature datasets, feature classes, tables, relationship classes, subtypes, and domains.

Shapefile and table (.dbf) versions of the data are included within the data. Column names have been simplified and the text values have been maintained within the shapefile attributes. The direction columns are numerical, to display rotation for points, and the symbol fields will hold the correct values to be matched to the appropriate style file.

For a more in depth description of the data model please refer to the official publication:

Deblonde, C., Plouffe, A., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Kerr, D.E., Moore, A., Parent, M., Robertson, L., Smith, I.R., St-Onge, D.A., and Weatherston, A., 2014. Science language for an integrated Geological Survey of Canada data model for surficial geology maps, version 2.0; Geological Survey of Canada, Open File 7631, 464 p. doi:10.4095/294225