


Figure 3. 3.5 kHz seismic reflection profile showing acoustically transparent postglacial mud adjacent to an area of seafloor winnowing and erosion in which acoustically stratified glaciomarine sediments are exposed at the sea floor. Location indicated on the principal map.

This map depicts the geomorphology of the Kitimat fiord Cette carte illustre la géomorphologie du système de system, British Columbia, and is based on bathymetry fjords de Kitimat, en Colombie-Britannique, et elle est and backscatter data from multibeam sonar surveys, fondée sur des données bathymétriques et des complemented by 3.5 kHz sub-bottom profiler data. données de rétrodiffusion acquises lors de levés au sonar multifaisceaux, auxquelles s'aioutent des grab samples, cores, and bottom photographs. The coverage extends from the head of Kitimat Arm to données recueillies à l'aide d'un sondeur de sédiments Hecate Strait, and includes Douglas Channel. The à fréquence d'émission de 3,5 kHz, des échantillons seafloor is divided into classes based on morphology prélevés par benne, des carottes et des photographies and texture. The classes include glacial landforms that du fond marin. La zone couverte s'étend du fond du record the advance and retreat of grounded ice, and a bras Kitimat au détroit d'Hecate et comprend le chenal wide range of submarine slope failures. The map also Douglas. Le fond marin est divisé en classes selon la shows that postglacial sediment distribution on the fiord morphologie et la texture. Les classes englobent des floors is highly variable: large areas are current- reliefs glaciaires qui témoignent de l'avancée et du recul scoured, so that underlying glaciomarine sediments are de la glace ancrée, ainsi qu'une vaste gamme de exposed. More than a hundred submarine fans deltas ruptures sous-marines de versant. La carte illustre are present within the fiords. également la répartition très variable des sédiments postglaciaires sur le fond des fjords : de vastes zones ont été érodées par les courants, ce qui a permis d'exposer les sédiments glaciomarins sous jacents. Plus d'une centaine d'éventails deltaïques sous-marins sont présents dans les fjords.





© Her Majesty the Queen in Right of Catalogue No. M183-1/275-1-2016E-PDF ISBN 978-0-660-05580-0 Canada, as represented by the doi:10.4095/298793 Minister of Natural Resources, 2016

CANADIAN GEOSCIENCE MAP 275 MARINE GEOLOGY **GEOMORPHOLOGY OF THE** KITIMAT FIORD SYSTEM

Natural Resources Ressources naturelles du Canada

Parts of NTS 103-A, NTS 103-H, and NTS 103-I





Authors: J. Shaw and D.G. Lintern Geological compilation by J. Shaw

Map projection Universal Transverse Mercator, zone 9.

North American Datum 1983

Base map at the scale of 1:250 000 from Natural Resources Shaded relief image derived from the Canadian Digital Elevation Data Illumination: azimuth 345°, altitude 45°, vertical factor 10x

CANADIAN GEOSCIENCE MAP 275 MARINE GEOLOGY GEOMORPHOLOGY OF THE KITIMAT FIORD SYSTEM British Columbia Parts of NTS 103-A, NTS 103-H, and NTS 103-I

Columbia. Photograph by J. Shaw. 2016-001

Mean magnetic declination 2016, 18°22'E, decreasing 12.8' annually Readings vary from 18°37'E in the NW corner to 18°06'E in the SE This map is not to be used for navigational purposes. Title photograph: Snow covered hills near Douglas Channel, British

Preliminary Preliminary The Geological Survey of Canada welcomes corrections or additional Data may include additional observations not portrayed on this map. This publication is available for free download through GEOSCAN (http://geoscan.nrcan.gc.ca/).

associated grab sample consisted of dark olive gray sandy mud.

and irregular morphology interpreted as rockfall. Photographs

show a rippled gravelly sandy seafloor littered with shell

instance, a sea pen. The associated grab sample consisted of

gravelly sand with shell fragments.

fragments, scattered anemones and sponges, and, in this

Figure 7. Representative seafloor images from survey 2015002PGC, each covering an area approximately 1.25 x 0.85 m.

103-H, and NTS 103-I; Geological Survey of Canada, Canadian Geoscience Map 275 (2nd edition, preliminary), scale 1:200 000.

scientifically edited.

Canada

Preliminary

Canadian Geoscience Maps

Geological Survey of Canada

fiord floors and in basins on the inner throughout the fiord system.

STREAMLINED GLACIAL

on the shoals of the inner continental shelf. throughout the fiord system.

Station 01, Photo 106: fiord- postglacial mud / sandy mud.

Station 04, Photo 03: fiord-mass transport complex. Depth

Station 09, Photo 21: fiord-postglacial mud / sandy mud.

Station 10, Photo 13: fiord-floor-winnowed / scoured. Depth of 673 m near the margin of an area of low backscatter. The seafloor is fine-grained, with scattered shell fragments, and a sediment with high backscatter strength. The pavement of subsediment with high backscatter strength and subsediment with high backscatter strength. The pavement of subsediment with high backscatter strength and subsediment with high backscatter strength. The pavement of subsediment with high backscatter strength and subsediment with high backscatter strength. The pavement of subsediment with high backscatter strength and subsed mottled appearance, with areas of dark blue and light brown. The angular/sub-rounded gravel has fine sediment in the interstices.

The associated grave semile separated gravel as flow could be appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness, and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness are roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a mottled (dark blue/light brown) appearance, with a roughness and a roughness and a roughness and a roughness and a roughn

Station 13, Photo 08: fiord-bioherm. Depth 125 m on the slope

Station 14, Photo 07: fiord-bioherm. Depth 85 m on top of a

Station 65, Photo 08: fiord – sidewall – unconsolidated. Depth

Station 68, Photo 07: fiord – sidewall – bedroe

anemones, brachiopods, urchins. The associated grab sample mostly dead, with several sea stars visible. The grab sample sand grading to clayey mud with depth.

of a submarine moraine in Devastation Channel. The seafloor is sponge reef — a ridge of acoustically transparent sediment with 58 m on a section of fiord sidewall with high backscatter, showing on steeply sloping sidewall with high backscatter. Colonies of covered by disarticulated tests of the giant acorn barnacle low backscatter on the crest of the submarine moraine in sub-angular/sub-rounded pebbles strewn on a fine-grained zooanthid anemones are present on the bedrock surface. Other Balanus nubilus, with a profusion of other biota including sponges, Devastation Channel. The seafloor is covered by glass sponges, seafloor. The associated grab sample showed gravel over coarse photographs show glass sponges and sea stars. In the

Station 115, Photo 13: fiord – bedrock slump / rock avalanche.

Station 141, Photo 13: Shelf – glaciomarine sediment. Depth

Station 142, Photo 11: Shelf – postglacial mud / sandy mud.

Station 145, Photo 11: Shelf – winnowed / scoured seafloor.

encrusted on top and clean below, suggesting the presence of an associated grab sample consisted of dark olive sandy mud.

The associated grab sample revealed a surface gravel lag animal burrows and tracks, and bivalve remains. The associated

252 m in an area of high backscatter with iceberg furrows and pits. Depth 53 m in an area of smooth seafloor Depth 53 m in an area of high backscatter, close to submarine

have encrusting fauna. A sea star and fish are also visible. The sediment. The relatively smooth seafloor is marked by large pavement, with sandy interstices. In this image a large anemone

grab sampler failed to penetrate but collected a single cobble, burrows. Other photographs show numerous animal tracks. The is present. Other photographs at this site show sponges, sea

grab sample contained sandy mud.

Bedrock is exposed on fiord sidewalls and Unconsolidated fiord sidewalls are found Postglacial mud/sandy mud is found on Winnowed and scoured seafloors are found Glaciomarine sediment is found on the

Submarine moraine crests are located Streamlined glacial landforms are located Deltas are found mostly in the inner fiords. Mass transport deposits occur throughout Bioherms have been mapped in several transverse to fiord axes; on the inner on the inner continental shelf. areas, but only those in Devastation the fiord system continental shelf they trend NW to SE. Channel (arrowed) have been confirmed by Figure 8. Distribution of seascape units. For this figure, distributions of bedrock inside and outside the fiord have been amalgamated, as have the distributions of postglacial mud.

> Shaw, J. and Lintern, G., 2016. Marine geology, Geomorphology of the Kitimat fiord system, British Columbia, Parts of NTS 103-A, NTS

background the seafloor is fine-grained sediment. Other

stars, and a bed of scallops. The associated grab sample showed

Preliminary publications in this series have not been

CANADIAN GEOSCIENCE MAP 275 MARINE GEOLOGY **GEOMORPHOLOGY OF THE** KITIMAT FIORD SYSTEM British Columbia

Parts of NTS 103-A, NTS 103-H, and NTS 103-I