References and additional ice flow and field data from:

Arsenault, L., Aylesworth, J.M., Cunningham, C.M., Kettles, I.M., and Shilts, W.W., 1981. Surficial geology, Eskimo Point, District of Keewatin, Geological Survey of Canada, Map 8-1980, scale 1:125 000. https://doi.org/10.4095/119389

Cocking, R.B., Deblonde, C., Kerr, D.E., Campbell, J.E., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Lavoilette, A., Parent, M., Plouffe, A., Robertson, L., Smith, I.R., and Weatherston, A., 2016. Surficial Data Model, version 2.2.0: Revisions to the science language of the integrated Geological Survey of Canada data model for surficial geology maps; Geological Survey of Canada, Open File 8041, 45 p. https://doi.org/10.4095/298767

Wright, G.M., 1967. Surficial geology, southeastern Barren Grounds, District of Keewatin and District of Mackenzie; Geological Survey of Canada, Map 1217A, scale 1:1 000 000. https://doi.org/10.4095/108855

Abstract

This new surficial geology map product represents the conversion of Map 8-1980 and its legend only, using the Geological Survey of Canada's Surficial Data Model (SDM version 2.2) which can be found in Open File 8041. All geoscience knowledge and information from Map 8-1980 that conformed to the current SDM were maintained during the conversion process. Additional material such as marginal notes or figures which may exist on the original map, are not included here. Supplementary legacy information was added to complement the converted geoscience data. This consists of a few glacial striations from Wright, 1967.It is identified in the accompanying geodatabase. The purpose of converting legacy map data to a common science language and common legend is to enable and facilitate the efficient digital compilation, interpretation, management, and dissemination of geologic map information in a structured and consistent manner. This provides an effective knowledge management tool designed around a geodatabase which can expand following the type of information to appear on new surficial geology maps.

Résumé

Ce nouveau produit cartographique de la géologie des formations superficielles correspond à la conversion de la Carte 8-1980 et de sa légende uniquement, en se servant du Modèle de données pour les formations superficielles (MDFS version 2.2) de la Commission géologique du Canada, lequel peut être consulté dans le Dossier public 8041. Toutes les connaissances et l'information de nature géoscientifique de la Carte 8-1980 qui sont en conformité avec le modèle de données ont été conservées pendant le processus de conversion. Des éléments additionnels tels que des notes marginales ou des figures qui pourraient être présents sur la carte originale ne sont pas inclus ici. Des données complémentaires ont été ajoutées pour compléter les données géoscientifiques converties. Ce sont quelques stries glaciaires de Wright, 1967. Elles sont identifiées dans la base de données géospatiales. Le but de la conversion de cartes publiées antérieurement suivant un langage scientifique commun et une légende commune est de permettre et de faciliter la compilation, l'interprétation, la gestion et la diffusion efficaces de l'information géologique cartographique en mode numérique de façon structurée et cohérente. Cette façon de faire offre un outil efficace de gestion des connaissances élaboré à l'aide d'une géodatabase qui pourra évoluer suivant le type d'information à paraître sur les nouvelles cartes des formations superficielles.



National Topographic System reference and index to adjoining published Geological Survey of Canada maps

Catalogue No. M183-1/240-2016E-PDF
ISBN 978-0-660-04097-4
https://doi.org/10.4095/300687

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources, 2017

Natural Resources Canada | Ressources naturelles Canada

CANADIAN GEOSCIENCE MAP 240

RECONNAISSANCE SURFICIAL GEOLOGY

ESKIMO POINT

Nunavut

NTS 55-E

1:125 000



QUATERNARY

NONGLACIAL ENVIRONMENT

ALLUVIAL SEDIMENTS: stream-deposited material within modern active drainage systems; "modern" is defined as the period since retreat of the sea, proglacial lakes, or glacial ice.

Ap — **Floodplain sediments:** silt, sand, and gravel; variable thickness; deposited in channels and on floodplains; may include alluvial sediments in terraces which formed as streams cut to present level in glacial and marine sedimentary fill.

A — **Alluvial sediments, undifferentiated:** silt, sand and gravel; variable thickness; when associated with marine sediments it occurs as flat areas consisting of modern alluvium mixed with silt and sand that were washed from slopes by wave action or deposited in the sea by meltwater streams; surface generally covered by 40 cm to more than 1 m of fibrous peat on which mosses, sedges, and grasses grow; maximum active layer is 15 to 50 cm thick; surface characterized by frost polygons and thaw ponds related to vertical ice wedges extending to a depth of 2 to 3 m.

L — **Lacustrine sediments, undifferentiated:** silt and sand; variable thickness; associated with permanently drained postglacial lake basins; may include deposit of silty sediment with up to 15% organic carbon.

NONGLACIAL AND GLACIAL ENVIRONMENT

MARINE SEDIMENTS: materials deposited in the Tyrrell Sea and during marine regression, and glacial deposits modified by marine processes.

Mr — **Beach sediments:** sand, gravel, cobbles, or boulders; variable thickness; generally well sorted; deposited as beaches, bars, spits, and ice-pushed ridges in littoral and nearshore environments; surface characterized by sparse vegetation and orthogonal frost cracks.

Md — **Deltaic sediments:** sand, gravel, and boulders; variable thickness; deposited where modern streams enter Hudson Bay.

Mi — **Intertidal sediments:** stony silt and sand; variable thickness; poorly sorted; coastal plain complex includes wind-blown sand and marine clayey silt associated with tidal flats along Hudson Bay, with pockets of sorted nearshore sand and gravel; probably a till plain levelled by filling of depressions and planation by wave action during emergence; forms a flat plain extending as far as 20 km inland; a generally complete organic cover and high density of shallow lakes make determining the nature of underlying sediments difficult; periglacial features include mudboils on muddy substrate to frost cracks or polygons on organic, gravelly or sandy deposits.

Mn — **Nearshore sediments:** sand; variable thickness; thin sheet deposited by a migrating shoreline; thought to be a lag developed by wave reworking of marine clayey sand or silty sand, generally associated with Mi; characterized on some aerial photos by a mottled pattern resulting from periglacial processes or gullying by streams or tidal currents.

Mo — **Offshore sediments:** clay, silt and silty sand; variable thickness; deposited in a deep water environment; may occur anywhere below marine limit but distribution is patchy above 60 m a.s.l.; thickest deposits generally occur in major river valleys or valleys with major eskers; may include prominent striped or mottled pattern on airphotos; difficult to differentiate on airphotos from till; surface vegetation is dominantly shrubs in turf circles around mudboils; striped pattern consists of dark and light stripes, 10 to 20 m wide, running downslope and reflects vegetation differences; mottled pattern due to shrub-and mudboil-covered mounds, 5 to 10 m in diameter and less than 0.5 m high, surrounded by light coloured grass-sedge vegetation; thought to have high ground ice content.

M — **Marine sediments, undifferentiated:** clay, silt and sand; variable thickness; deposited in various marine environments.

GLACIAL ENVIRONMENT

GLACIOFLUVIAL SEDIMENTS: water-sorted sediments deposited in, or around, or near a glacier, largely as a result of meltwater stream flow.

GFz2 — **Subaqueous outwash fan sediments:** silt, sand, fine gravel; variable thickness; terraced, hummocky, and kettled surfaces; distal fine-sediment debouching from the subaqueous mouth of an ice tunnel into the sea; vegetation is commonly sedge, grass, and moss growing on thin, fibrous peat; surface characterized by thaw ponds and tundra polygons; difficult to differentiate on aerial photos from unit A.

GFz — **Ice-contact sediments:** sand and gravel; variable thickness; stratified, deposited near ice margins in, over, or around ice or in ice tunnels, commonly as eskers but includes isolated hummocky deposits of uncertain origin; many eskers are beaded or interrupted at irregular intervals by major bulges marking positions where deltaic or subaqueous fan deposition occurred during periods of slowed ice retreat.

GLACIAL SEDIMENTS (TILL): poorly sorted till (diamicton) with distinctive forms deposited directly by glacial ice.

Th — **Hummocky till:** till; variable thickness; without significant boulder cover occurring as hummocks; includes ridges of till that are minor end moraines or erosion remnants between subparallel meltwater channels; extensive areas are present in the region of the Keewatin Ice Divide and in some places adjacent to areas of ribbed moraines (Tr); exact origin unknown; may have formed in association with stagnant ice and in some places from erosion of till surfaces by streams in anastomosing meltwater channels; vegetation and periglacial features similar to those on till plain (Tp).

Tr — **Ridged moraine:** generally bouldery till, in places sand and gravel; variable thickness; ribbed (Ridge) moraine, forming hummocks and straight to sinuous ridges, generally less than 1 km long and 2 to 10 m high; ridges generally oriented at right angles and form trains parallel to direction of ice flow; individual ribs may be asymmetric in cross-section with steep side facing down-ice; trains of ribs pass laterally into drumlin fields and may be drumlinized in the transition zone; surfaces have sparse vegetation and generally a heavy cover of large boulders and mudboils where composed of till, and are marked by frost cracks where composed of gravel.

Tp — **Till plain:** generally sandy, silty till with less than 25% clay sized particles; noncalcareous; grey; variable thickness; forms till plains; includes areas of clay-rich red till; vegetation (shrubs, mosses, grasses) grows in elevated peaty rings around bare or lichen-covered mudboils 1–2 m diameter; may exhibit striped pattern on clay-rich red till as observed on aerial photos.

Tv — **Till veneer:** generally sandy, silty diamicton; less than 1 m thick. Occurs in secondary unit only.

T — **Till, undifferentiated:** diamicton; variable thickness; associated with complex map units of till-cored landforms blanketed by marine sediments; based solely on aerial photo interpretation. Occurs in secondary unit only.

PRE-QUATERNARY

R — **Bedrock, undifferentiated:** Precambrian intrusive igneous and metamorphic rocks, red volcanic rocks, and unmetamorphosed sediments; surface comprises more than 80% outcrop; vegetation sparse; surface may be glacially rounded or covered by felsenmeer; bedrock (20 to 80% outcrop) is mantled with less than 1 m of the surficial deposit indicated in complex polygons.

Complex units: two map-unit designators are used in cases where the surficial cover forms a complex area and the map units are too small to be mapped individually, yet constitute a significant areal extent of the total polygon (e.g. R.Tv designates an area of bedrock with numerous small deposits of till veneer). In such instances a dot (.') is used to separate the map-unit designator.

Stratigraphic relationship: a stratigraphic relationship is shown with two map-unit designators separated by a slash ('/') (e.g. M/T designates marine sediments overlying till).

- Geological contact, defined
- Terrace scarp
- Beach crest, bar, or ice-shoved ridge
- Limit of submergence, marine, defined
- Meltwater channel, major, direction unknown
- Moraine ridge, minor, unspecified, De Geer
- Esker, paleoflow known
- Drumlinoid ridge:
  - *Buried, may be obscured by solifluction processes, water-laid sediments or wave reworking*
  - *Drumlinoid, fluting*
- Palsa or lithalsa
- Hummock, hummocky moraine
- Striation:
  - *Ice flow direction unknown*
  - *Ice flow direction known*
  - *Crossed. 1 = older, 2 = younger*
- Small bedrock outcrop
- Dated sample location, with number (*see* Table 1)

Map ID	Sample ID	Latitude	Longitude	Elevation (m a.s.l.)	Material	Radiocarbon Age
1	GSC-1626	61.961764	95.687028	61 ± 3	Shells	5980 ± 180
2	GSC-1427	61.111203	94.065655	2	Shells	3200 ± 220

Table 1. Radiocarbon age.

Recommended citation
Geological Survey of Canada, 2017, Reconnaissance surficial geology, Eskimo Point, Nunavut, NTS 55-E; Geological Survey of Canada, Canadian Geoscience Map 240 (preliminary, Surficial Data Model v. 2.2 conversion of Map 8-1980), scale 1:125 000. https://doi.org/10.4095/300687

Preliminary Preliminary Preliminary

Celebrating 175 yrs

Geological Survey of Canada
Canadian Geoscience Maps

Canada

Author: Geological Survey of Canada

Geology based mainly on airphoto interpretation and ground observations by L. Arsenault, J.M. Aylesworth, C.M. Cunningham, I.M. Kettles, and W.W. Shilts, 1971, 1973, 1977.

Geology conforms to Surficial Data Model v. 2.2

Data conversion by D.E. Kerr, 2014, 2015, 2016

Geomatics by GSP Geographics Inc. and S. Eagles

Cartography by N. Côté

Initiative of the Geological Survey of Canada, conducted under the auspices of the Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) program

Map projection Universal Transverse Mercator, zone 15. North American Datum 1983

CANADIAN GEOSCIENCE MAP 240
RECONNAISSANCE SURFICIAL GEOLOGY
ESKIMO POINT
Nunavut
NTS 55-E
1:125 000

2 0 2 4 6 8 10 km

Preliminary Preliminary Preliminary

Base map at the scale of 1:250 000 from Natural Resources Canada, with modifications.
Elevations in metres above mean sea level

Mean magnetic declination 2017, 2°24'E, increasing 0.8' annually. Readings vary from 0°53'W in the SW corner to 4°00'W in the NE corner of the map.

This map is not to be used for navigational purposes.

The Geological Survey of Canada welcomes corrections or additional information from users.

Data may include additional observations not portrayed on this map. See map info document accompanying the downloaded data for more information about this publication.

This publication is available for free download through GEOSCAN (http://geoscan.nrcan.gc.ca/).

Preliminary publications in this series have not been scientifically edited.

CANADIAN GEOSCIENCE MAP 240
RECONNAISSANCE SURFICIAL GEOLOGY
ESKIMO POINT
Nunavut
NTS 55-E