



Health
Canada Santé
Canada

Your health and
safety... our priority.

Votre santé et votre
sécurité... notre priorité.

Proposed Maximum Residue Limit

PMRL2017-08

Ethaboxam

(publié aussi en français)

11 April 2017

This document is published by the Health Canada Pest Management Regulatory Agency. For further information, please contact:

Publications
Pest Management Regulatory Agency
Health Canada
2720 Riverside Drive
A.L. 6607 D
Ottawa, Ontario K1A 0K9

Internet: pmra.publications@hc-sc.gc.ca
healthcanada.gc.ca/pmra
Facsimile: 613-736-3758
Information Service:
1-800-267-6315 or 613-736-3799
pmra.infoserv@hc-sc.gc.ca

Canada 

ISSN: 1925-0835 (print)
1925-0843 (online)

Catalogue number: H113-24/2017-8E (print version)
H113-24/2017-8E-PDF (PDF version)

© Her Majesty the Queen in Right of Canada, represented by the Minister of Health Canada, 2017

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of the Minister of Public Works and Government Services Canada, Ottawa, Ontario K1A 0S5.

Under the authority of the *Pest Control Products Act*, Health Canada's Pest Management Regulatory Agency (PMRA) has concluded that the addition of new uses for seed treatment on field peas, cowpeas and Crop Subgroup 20B (sunflower subgroup) to the product label of Intego™ Solo Fungicide, containing technical grade ethaboxam, is acceptable. The specific uses approved in Canada are detailed on the label of Intego™ Solo Fungicide, *Pest Control Products Act* Registration Number 31324.

The evaluation of this ethaboxam application indicated that the end-use product has value and the human health and environmental risks associated with the new uses are acceptable.

Before registering a pesticide for food use in Canada, the PMRA must determine the quantity of residues that are likely to remain in or on the food when the pesticide is used according to label directions and that such residues will not be a concern to human health. This quantity is then legally established as a maximum residue limit (MRL). An MRL applies to the identified raw agricultural food commodity as well as to any processed food product that contains it, except where separate MRLs are specified for the raw agricultural commodity and a processed product made from it.

Consultation on the proposed MRLs for ethaboxam is being conducted via this document (see Next Steps, the last section of this document). A summary of the field trial data used to support the proposed MRLs can be found in Appendix I.

To comply with Canada's international trade obligations, consultation on the proposed MRLs is also being conducted internationally by notifying the World Trade Organization, as coordinated by the Canada's Notification Authority and Enquiry Point.

The proposed MRLs, to be added to the MRLs already established for ethaboxam, are as follows.

Table 1 Proposed Maximum Residue Limits for Ethaboxam

Common Name	Residue Definition	MRL (ppm) ¹	Food Commodity
Ethaboxam	N-(cyano-2-thienylmethyl)-4-ethyl-2-(ethylamino)-5-thiazolecarboxamide	0.01	Sunflowers (crop subgroup 20B), dry cowpea seeds, dry field peas, succulent shelled cowpeas

¹ ppm = parts per million

MRLs are proposed for each commodity included in the listed crop groupings in accordance with the Residue Chemistry Crop Groups webpage in the Pesticides and Pest Management section of Health Canada's website.

MRLs established in Canada may be found using the Maximum Residue Limit Database on the Maximum Residue Limits for Pesticides webpage. The database allows users to search for established MRLs, regulated under the *Pest Control Products Act*, both for pesticides or for food commodities.

International Situation and Trade Implications

The MRLs proposed for ethaboxam in Canada do not have corresponding tolerances in the US. Ethaboxam is registered for use as a seed treatment in the US on legume vegetables, cereal grains and rapeseeds. These uses are considered “non-food” by the US, as residues are not expected in any subsequent RACs. As such, no MRLs were established on these crops. The only American tolerance for ethaboxam is for imported grapes as listed in the Electronic Code of Federal Regulations, 40 CFR Part 180, by pesticide. Currently, there are no Codex MRLs¹ listed for ethaboxam in or on any commodity on the Codex Alimentarius Pesticide Residues in Food and Feed webpage.

Next Steps

The PMRA invites the public to submit written comments on the proposed MRLs for ethaboxam up to 75 days from the date of publication of this document. Please forward your comments to Publications (see the contact information on the cover page of this document). The PMRA will consider all comments received before making a final decision on the proposed MRLs. Comments received will be addressed in a separate document linked to this PMRL. The established MRLs will be legally in effect as of the date that they are entered into the Maximum Residue Limit Database.

¹ The Codex Alimentarius Commission is an international organization under the auspices of the United Nations that develops international food standards, including MRLs.

Appendix I

Summary of Field Trial Data Used to Support the Proposed Maximum Residue Limits

Residue data for ethaboxam seed treatment in soybean and sunflower were submitted to support the domestic use of Intego™ Solo Fungicide on field pea, cowpea and sunflowers (crop subgroup 20B).

Maximum Residue Limits

The recommendation for maximum residue limits (MRLs) for ethaboxam was based upon the submitted field trial data, and the guidance provided in the OECD MRL Calculator. Table A1 summarizes the residue data used to calculate the proposed MRLs for field pea, cowpea and sunflowers (crop subgroup 20B).

Table A1 Summary of Field Trial Data Used to Support MRLs

Commodity	Application Method/ Total Application Rate	Preharvest Interval (days)	Lowest Average Field Trial Residues (ppm)	Highest Average Field Trial Residues (ppm)	Experimental Processing Factor
Soybean seeds	Seed treatment 75 g a.i./100 kg seed ¹	130-138	<0.01	<0.01	No quantifiable residues were observed at exaggerated rates
Sunflower	Seed treatment 0.0945-0.202 mg a.i./seed ²	105	<0.005	<0.005	No quantifiable residues were observed at exaggerated rates
	Seed Treatment 0.185-0.192 mg a.i./seed ²	105	<0.005	0.005	

¹ g a.i./100 kg seed = grams of active ingredient per 100 kilogram of seed

² mg a.i./seed = milligrams of active ingredient per seed

Following the review of all available data, MRLs as proposed in Table 1 are recommended to cover residues of ethaboxam. Residues of ethaboxam in these crop commodities at the proposed MRLs will not pose an unacceptable risk to any segment of the population, including infants, children, adults and seniors.