0.3.

CANADIAN
SICKNESS SURVEY
1950-51

No. 8

VOLUME OF HEALTH CARE
(NATIONAL ESTIMATES)

Prepared jointly

bу

THE DOMINION BUREAU OF STATISTICS and

THE DEPARTMENT OF NATIONAL HEALTH AND WELFARE

OTTAWA, CANADA OCTOBER, 1955


Price 25 cents

ACKNOWLEGEMENT

The Canadian Sickness Survey, the first nationwide study of illness in the general population of Canada, was carried out during a twelve-month period commencing in the autumn of 1950.

The Survey was initiated by the Department of National Health and Welfare and carried out by the ten provincial health departments with federal funds made available to the provinces through the National Health Program.

The planning and organization of the Survey was a joint undertaking of the Dominion Bureau of Statistics and the Department of National Health and Welfare in consultation with the provinces. Every provincial health department co-operated fully in gathering the extensive body of information which makes these publications possible.



016

1950-51

100.8

312.3

Note: This is part 8 of D.B.S. Reference Paper No. 51. The first five bulletins in this series were referred to as Special Compilations.

TABLE OF CONTENTS

				Page
1 1	in tr c Ti	odu he	ction: Canadian Sickness Survey 1950-51	5
п			ets, Definitions, and Methods:	5
	34	op.	e of present bulletin	
	M	eas	sures of health care	6
*** *			ulation and presentation of data	8
Ш			nary	9
			TABLES AND CHARTS	
			Total Health Care	
Tabl	e	1.	Persons reporting health care during survey year	14
Tabl Char	e	2.	Complaint periods under health care during survey year Estimated number of persons reporting various classes of health care	15 13
			Physicians' Services	
Tabl	e	3.	Physicians' services during survey year (summary table)	17
Tabl	e	4.	Persons reporting physicians' care during survey year.	17
Tabl	_	5	Complaint periods under physicians' care during survey year	19
Tabl	0	e.	Number of doctors' calls and clinic visits during survey year	20
	•	7	Programmy of doctors' calls and clinic visits during survey year	
Tabl Char			Frequency of doctors' calls and clinic visits during survey year Estimated average number of doctors' calls and clinic visits reported per 1,000 persons of each age-sex group	25 22
Char		9	Estimated number of persons with various dumbers of destart's calls and	44
Clian	L	J.	Estimated number of persons with various numbers of doctors' calls and clinic visits	26
			In-Patient Hospital Care	
Tabl	e :	8.	Persons reporting in-patient hospital care during survey year	29
Tabl	e !	9.	Periods of in-patient hospital care during survey year	30
Tabl	e 1	0.	Number of person-days of hospitalization during survey year	30
Tabl	e 1	1.	Duration of hospitalization of persons during survey year	32
Char	t	4.	Estimated average number of days of hospitalization reported per 1,000 persons of each age-sex group	31
Char	t	5.	Estimated number of persons with various number of days in hospital	31
	_	_	Home Nursing Services ,	
			Home nursing services during survey year (summary table)	35
Tabl	e 1	3.	Persons reporting home nursing care during survey year	35
Tabl	le 1	4.	Complaint periods under home nursing care during survey year	37
Tabl	e 1	5.	Number of visits or days of home nursing care during survey year	38
	-	_	Operations	
Tabl	e 1	6.	Operations during survey year	41
Char	t	6.	Estimated average number of operations reported per 1,000 persons of each age group	42
			Dental Care	
Tabl	e 1	7.	Dental care during survey year	42
Char	t	7.	Estimated average number of dental visits reported per 1,000 persons of each age group	43
			Examinations for Glasses	
Tabl	e 1	8.	Examinations for glasses during survey year (other than by physicians)	43
			Estimated average number of examinations for glasses reported per 1,000 persons of each age group	44
			Miscellaneous Health Care	
Tahl	1 ما	Q	Miscellaneous health care during survey year	44
~ 401		٠.	MIDOUTOMOVED HEATIN ONE CHAINE DATACT LEGIS TO THE MIDOUTOMOVED HEATING ONE CHAINED DATACT.	77

	•
•	
•	
•	
	, ·
	•
	•
	·
•	
	•
•	
·	•
·	
·	
	÷

INTRODUCTION

THE CANADIAN SICKNESS SURVEY 1950-51

The object of the Canadian Sickness Survey was to obtain estimates of the incidence and prevalence of illness and accidents of all kinds; the amount of medical, nursing and other health care received; the volume of family expenditures for the various types of health services; and the prevalence of permanent physical disabilities. The present bulletin is one of a series intended to provide a means of releasing interim information on the more important of the results, pending the preparation of a comprehensive publication containing more detailed information together with a fuller description of the methods used.

The survey method consisted of personal visits by trained lay enumerators - in British Columbia the enumerators were public health nurses - to a sample of approximately 10,000 households distributed throughout the ten provinces in metropolitan, small urban, and rural areas. Less than five per cent of these households refused to participate in the Survey. Of the remaining households over 80 per cent of the individuals involved remained in the sample throughout the survey period. All information, including particulars of income, housing, and environment, was obtained by direct interview of a household informant, usually the housewife. While the starting dates for the Survey varied somewhat in different provinces, in most cases a total of 14 monthly visits were made to each household in the sample. In the first visit the enumerator introduced the Survey and left a special calendar designed to help the informant keep a detailed day-to-day record of current sickness and of expenditures on health care and services for each member of the household. During each of the succeeding twelve months the enumerator interviewed the informant and recorded the sickness experienced by each person since the previous visit. The final visit was made to review the information recorded throughout the whole survey period. Uniformity of practice in the ten provinces was maintained by frequent consultation among the agencies involved, by uniform instructions to the enumerators, and by the use of three standard record forms - a Household Record, an Individual Sickness Record, and an Expenditures Form. Auxiliary schedules, also standardized, were used to

record permanent physical disabilities and also health services which were desired but notobtained.

The sample was designed to obtain estimates with a sampling error of 20 per cent.² Indications are that for almost all of the estimates the error is substantially smaller. Area sampling was used for the Survey. As a first stage the following six domains (regions) of study were established consisting of four single provinces and two groups of three provinces each:

- 1. Newfoundland
- Maritimes (includes Prince Edward Island, Nova Scotia, and New Brunswick)
- 3. Quebec
- 4. Ontario
- Prairies (includes Manitoba, Saskatchewan, and Alberta)
- 6. British Columbia

Within each domain three types of area were considered - metropolitan, urban, and rural. Within these areas multi-stage sampling was adopted. In metropolitan centres, all of which were included in sample, and in some of the sampled urban areas, the first stage of sampling was the block, the second stage being the household. In other urban areas systematic sampling from a list of households was used. Rural areas were divided into primary sampling units and grouped into strata. Within each stratum one primary sampling unit was selected and multi-stage sampling applied. The first stage was the selection of clusters or segments within the primary sampling unit while the second stage was the selection of households within the chosen clusters.

In designing the sample extensive use was made of population, social, and economic data obtained from 1941 Census material. The results of the 1951 Census, which was taken at about mid-point of the survey period, provided the necessary distributions concerning persons and families for the calculation of weights used to inflate figures to national and provincial totals. The basic survey units for data on illness were individual persons, while the units for expenditures on health services included families, as defined in the Census, together with certain single persons living alone or with other families as roomers or relatives.

CONCEPTS, DEFINITIONS, AND METHODS

SCOPE OF PRESENT BULLETIN

This bulletin is the eighth in the series of preliminary publications of the Canadian Sickness Survey. Family expenditures for health services were dealt with in the first four reports. The fifth report concerned the volume of sickness, the sixth permanent physical disabilities, and the seventh the incidence and prevalence of illness. Illness statistics classified by diagnostic group will be considered in bulletins to follow. The purpose of the present report is to provide estimates by age group and sex of the volume of health care received by Canadians. The major categories of health care recorded are physicians' services, hospital care, and nursing care. In addition information is presented regarding operations, dental care, examinations for glasses, and miscellaneous other health care.

2. With a 95 per cent confidence limit.

^{1.} The population sampled did not include residents of institutions, military establishments, Indian reservations, and remote areas.

The volume of health care has been estimated, broadly, according to three different measures. In the first place, estimates are given of the number of persons who received various types of health care, e.g. the number of persons who were attended by a physician during the survey year and the number of persons who received hospital treatment during the survey year.

Secondly, estimates are presented of the number of "complaint periods" for which various types of health care was received. A complaint period in this care is the same as defined in bulletin No. 5. viz. a series of days, ranging from a single day to a period covering the entire survey year, throughout which time a person was reported as continuously experiencing a disturbance in his state of health, and was at no time free of symptoms. Health disturbances covered by complaint periods included injuries and confinements as well as diagnosed diseases and undiagnosed symptoms. A complaint period might have a single diagnostic or symptomatic cause, in which case it would correspond to a single illness as defined in bulletin No. 7, or it might be due to a number of overlapping or concurring illnesses. In relating health care to complaint periods, only those services were counted which were provided during the survey year. In effect, therefore, complaint periods current at the beginning of the survey year were considered to commence on the first day of the Survey and complaint periods current at the end of the survey year were considered to terminate on the last day of the Survey. It bears emphasis that complaint periods cover every indisposition - no matter how slight - which was reported by informants, and the high number of recorded complaint periods should be interpreted accordingly.

In the third place, this bulletin presents estimates of the number of units provided of various categories of health care, e.g. the total number of doctors' calls and the total number of person-days of hospitalization. In addition, estimates are given of the rates at which various types of care were provided per thousand persons or per complaint period.

The estimates of health care presented in this bulletin are based on information which was supplied directly by the Sickness Survey informants, i.e. by the persons who actually received the health care concerned or by their household spokesmen. The categories of health care recorded for the Survey, on the whole, were objective in nature and clearly definable. A reasonably accurate and uniform interpretation by informants of questions concerning health care may therefore be expected. Only in relating health care to the subjectively determined complaint periods was objective definition of health care categories impaired.

It should be noted that residents of institutions were excluded from the Sickness Survey sample (see page 5, n. 1). While the number of persons thus excluded was very small relative to the total population, the excluded persons may be expected to

have received a proportionately large amount of health care. Residents of institutions would include, for instance, most persons staying in sanatoria, in mental institutions, or in homes for the aged, as well as those long-term hospital patients who were no longer reported living at their household addresses. Some downward bias may therefore be expected in a few of the estimates given in this bulletin, principally where the older age groups are concerned and in the case of long-term hospital stays.

MEASURES OF HEALTH CARE

Physicians' Services

Physicians' services cover care given by qualified medical doctors. For the Sickness Survey information was collected on three classes of physicians' services - office calls, home calls, and clinic visits. An office call consisted of a visit by a patient to the office of his physician for treatment, examination, or prescription by the latter. A home call consisted of a visit by a physician to the home of a patient for treatment, examination, or prescription. A clinic visit was a visit by a patient for treatment, examination, or prescription to an outpatient hospital clinic, dispensary, or other special diagnostic or treatment centre where the services of qualified medical doctors were regularly available, even though in some instances patients visiting the clinic were attended by a nurse rather than by a physician. Visits to public health clinics or school health clinics where service was preventive in . nature - rather than therapeutic or diagnostic - were not included. Visits to medical group-practice clinics were counted as ordinary visits to a doctor's office. Visits by physicians to their patients in hospital were not recorded.

Where a person had a doctor's call or clinic visit for two or more illnesses at the same time, the call or visit was counted only once. However, when more than one doctor was called in consultation at the same time and for the same patient, a separate call or visit was counted for each physician.

Usually doctors' calls and clinic visits were made in connection with current illness. In some instances, however, doctors' calls or clinic visits were only for general medical examination or for routine check-up, in which case they could not be related to specific complaint periods. The number of doctors' calls and clinic visits recorded for complaint periods (Table 6 F) is therefore a little lower than the total number (Table 6 A).

In-Patient Hospital Care

Every person who was admitted as a patient to a hospital for one day or longer was considered to have received in-patient hospital care. A person who visited a hospital for medical treatment, examination, or prescription but did not occupy a hospital bed overnight was credited with a clinic visit instead (see above—Physicians' Services).

Hospitals were considered to consist of all those public and private institutions which were listed as hospitals by provincial hospital inspection authorities. This excluded most places known as rest homes or homes for the aged.

Two types of periods were considered in relation to hospital care. Estimates were prepared of the number of complaint periods for which in-patient hospital care was provided during the survey year. In this calculation each complaint period was counted once even though in some cases a person was admitted to hospital on two or more separate occasions in connection with the same complaint period. Separate estimates were prepared of the number of 'hospital periods' i.e. unbroken periods of in-patient hospital care. These hospital periods also included certain contingencies which were not covered by complaint periods. Such was the case when persons stayed in hospital not in connection with any active illness but only for observation or medical check-up. The initial stay in hospital of a newborn infant - not sick - was also counted as a hospital period but not as a complaint period. The number of hospital periods (Table 9, column (e)) was, of course, somewhat larger than the number of complaint periods with in-patient hospital care (Table 9, column (b)). The total number of persondays of hospitalization (Table 10, column (a)) was also a little larger than the number of person-days of hospitalization for complaint periods (Table 10, column (f)).

Home Nursing Services

This heading covers bedside nursing care obtained in the home. Nursing services provided in the hospital or other institution are not included, nor are visits by public health nurses who did not give bedside care. Information on the amount of care given by graduate nurses was recorded separately from that given by persons without graduate nurses' qualifications.

Home nursing care given by graduate nurses most commonly consisted of short visits to the homes of patients. In a small number of cases fultime bedside care in the home was given by graduate nurses. Graduate nursing care was measured in terms of the number of visits made, each full day of care also being counted as a visit.

Non-graduate nursing care was recorded in cases where a "practical nurse" or other outsider came into the home for the specific purpose of giving bedside care to a sick person. No account was taken of care given by members of the sick person's family or of the services of outsiders who did other housework in the home in order to relieve a member of the family who looked after the sick person. Non-graduate nursing care was measured in terms of eight-hour days of work. Where the daily hours of work were clearly different, they were totalled and divided by eight to determine the number of days of nursing care recorded for the Survey.

Operations

An operation was defined as any cutting or piercing procedure performed by a surgeon or—as occurred *less frequently—by a physician. The application of sutures, the setting of fractures, and the removal of splinters or foreign bodies from the eye or ear as such were not considered operations.

Dental Care

For the Sickness Survey a record was kept of care received by patients from qualified dentists. The amount of dental care received was measured in terms of the number of visits made to dentists' offices or clinics. Dental treatment received at hospital out-patient clinics was counted but treatment received at school dental clinics was excluded.

Examinations for Glasses

Under this caption information was recorded concerning eye examinations performed by optomethists or opticians. Examinations by eye specialists with medical qualifications were not included here but were counted as doctors' calls under the heading 'Physicians' Services'.

Examinations for glasses (other than by physicians) were measured in terms of the number of visits made to the offices of optometrists and opticians for the purpose of eye examination.

Miscellaneous Health Care

A record was also kept of health care which was not given in a hospital and which was not given by a qualified medical doctor, nurse, dentist, optometrist, or optician. Such miscellaneous health care included services performed by persons with formal qualifications as chiropodists, chiropractors, herbalists, homoeopaths, naturopaths, osteopaths, and physiotherapists, as well as by other practitioners such as faith healers, "bone doctors", etc.

Miscellaneous health care treatments were measured by the number of visits for treatment, examination, or prescription made by patients to practitioners or vice versa.

Total Health Care

Combined information concerning the various categories of health care are shown in Tables 1 and 2.

The category "medical care" in Table 1 is meant to cover all cases attended by medical doctors. Persons reporting any medical care are those reporting doctors' calls or clinic visits and/or in-patient hospital care. "Miscellaneous health care" in Table 1 refers to the categories listed as miscellaneous health care above and in Table 19.

In Table 2 estimates are given of the number of complaint periods with various combinations of

doctors' calls or clinic visits, in-patient hospital care, and home nursing care. "Other health care" in Table 2 covers dental care, examinations for glasses (other than by physicians), and the miscellaneous categories of health care listed in Table 19.

CALCULATION AND PRESENTATION OF DATA

Population Universe

The population universe from which the sample for the Canadian Sickness Survey was drawn consisted of the total population of Canada minus persons residing in institutions, military establishments, Indian reservations, and remote areas. This population universe, estimated at 13,540,000, was calculated from the total population of Canada, as recorded in the 1951 Census, with appropriate adjustments for the excluded sections of the population.

Sample

The estimates given in this bulletin were calculated from data obtained from a sample of about 33,000 persons, most of whom were reported on for the entire survey year. A few persons who were absent from the Survey for less than 31 days were also included in the sample. Persons who were born or who died during the survey year, but were otherwise not absent from the Survey for more than 30 days, were included in the sample.

An appropriate small downward adjustment was made to estimates of numbers of persons recorded in the tables of this bulletin, in order to compensate for partial absence from the Survey of some persons. In other words, all estimates pertain to person-years.

Estimating

The total sample was broken down into cells. by a division into provinces, areas (metropolitan and non-metropolitan) within provinces, and age-sex groups (five male and five female age groups) within areas. The total population of each cell was taken from the 1951 Census, with appropriate adjustments for excluded sections of the population. For each cell the ratio of total population to sample population was used as a weight and applied to all sample data pertaining to the cell. Weighted figures for the various cells were appropriately combined to provide the published estimates. These estimates are provisional in the sense of being subject to minor adjustments arising from further analysis of the sample material and from study and comparison of more detailed tabulations.

Age and Sex Groups

In most tables of this publication separate estimates are given for the two sexes combined and separately. For each of the three groups of figures thus obtained a further breakdown is given by five age groups. In the case of hospital care statistics

the youngest age group (under 15) is shown both in its entirety and exclusive of the newborn (i.e. those under one year of age or not yet born on June 1, 1951). The latter estimate was made to remove the undue influence of time spent in hospital by many newly born babies.

Frequency

The maximum margin of sampling error allowed for estimates published in this bulletin was 20 per cent.² In order to achieve this standard each published figure had to be based on a certain minimum frequency of occurrence in the Sickness Survey sample. Where the required frequency was not reached, estimates were omitted from the tables (indicated by a double dot) or else age and sex groups or other categories were combined to a sufficient total frequency. Low frequencies occurred most often with the smaller age groups—the 15-24 age group and the 65 and over age group.

Arrangement of Tables and Charts

For each of the major categories of health care considered in this bulletin tabular material is presented in a separate table or set of tables, together with appropriate charts. Combined information is shown as "total health care" in the first two tables. The contents of the tables are self-explanatory.

Rounding

In the tables, absolute numbers are given in millions with two decimal places, i.e. they are rounded to the nearest ten thousand. Percentage figures are given with one decimal place. Rates of occurrence per thousand persons are given in integral numbers. Averages per complaint period are given with three decimals.

As each figure was rounded individually, additions of rounded figures may differ slightly from the rounded totals shown.

Rates and averages were calculated from original full estimates, not from rounded figures shown in the published tables.

Analysis -

It is the purpose of these bulletins to release certain basic data as soon as possible and to leave more extensive tabulations and interpretations to a more comprehensive report (to be issued at a later date). For that reason no attempt is made in the present series of bulletins to analyse the data in detail. However, a summary of the tabular material of this bulletin is given below.

^{1.} The remote areas consisted of Labrador, the Northwest Territories, the Yukon, and the northernmost stretches of Quebec, Ontario, Manitoba, Saskatchewan, Alberta, and British Columbia.

^{2.} With a 95 per cent confidence limit, i.e. no individual figure was allowed a chance greater than one in 20 of exceeding the 20 per cent margin of error.

FINDINGS

SUMMARY

Physicians' Services

Tables 3-7 and Charts 2-3

More than two-fifths (43.2 per cent) of the surveyed Canadian population of 13,540,000 persons reported doctors' calls or clinic visits during the survey year (see Table 4C). About the same proportion of males (42.9 per cent) and females (42.1 per cent) under 15 reported doctors' calls or clinic visits, but in all other age groups females reporting physicians' services outnumbered males by a wide margin. The 15-24 age group showed a smaller proportion of males (29.2 per cent) and of females (41.1 per cent) reporting physicians' services than any other age group. The proportion generally increased successively with older age groups. The exception was for females in the 25-44 year bracket -covering most of the child bearing age - with the highest proportion of all (54.7 per cent) reporting physicians' services.

One quarter (25.3 per cent) of all complaint periods were attended by a physician through doctors' calls or clinic visits (see Table 5C). Females received physicians' services for a slightly higher proportion (25.6 per cent) of their complaint periods than males (24.9 per cent). The proportion of complaint periods attended to increased sharply and consistently with the advance of age. For males the proportion ranged from 20.5 per cent (under 15) to 35.5 per cent (65 and over) and for females it ranged from 19.1 per cent (under 15) to 39.5 per cent (65 and over).

A total of 24,180,000 doctors' calls and clinic visits were reported in the year 1950-51 (see Table 6A). Most numerous were doctors' office calls, accounting for 63.3 per cent of the total of calls and visits (see Table 6C). Doctors' home calls made up 28.7 per cent of the total, and clinic visits accounted for the remaining 8.0 per cent. Clinic visits made up a substantially larger portion of total calls and visits for males (10.4 per cent) than for females (6.3 per cent). For females, conversely, doctors' office calls and doctors' home calls accounted for a larger share of total calls and visits. For persons in the oldest (65 and over) and youngest (under 15) age groups doctors' home calls made up a larger than average proportion of total calls and visits (45.7 per cent and 34.5 per cent respectively against an average for all age groups of 28.7 per cent), while doctors' office calls made up a smaller than average proportion of the total (46.6 per cent and 57.5 per cent against an average of 63.3 per cent).

The average number of doctors' calls and clinic visits reported per 1,000 persons amounted to 1,786, less than two per person (see Table 6D). For males the average was 1,452 per 1,000 persons and for females almost half as many again, viz.

2.124 per 1,000. Males reported on the average more clinic visits than females (152 per 1.000 as against 135 per 1,000) but far fewer doctors' office calls (901 per 1,000 as against 1,364 per 1,000) and home calls (400 per 1,000 as against 626 per 1,000). The average number of calls and visits reported increased generally with age from 1,210 per 1,000 persons in the age group under 15 to 2,751 per 1,000 persons in the 65 and over age group (see Chart 2). For the separate sexes there were exceptions to this rule: males in the 15-24 age group reported fewer calls and visits (895 per 1,000 persons) than males under 15 (1,261 per 1,000 persons) and females in the 25-44 age group reported more calls and visits (2,775 per 1,000 persons) than females 45-64 years old (2.542 per 1.000 persons).

The average number of calls or visits per 1,000 persons, counting only those who did report at least one call or visit during the survey year, amounted to 4,132, or over four per person (see Table 6E). The total number of calls or visits reported per 1,000 persons reporting calls or visits increased consistently from 2,848 for the under 15 to 5,541 for the 65 and over age group. The count for each female age group was higher than for the corresponding male age group, except for the age group under 15.

Less than one (0.678) doctor's call or clinic visit was reported per complaint period (see Table 6G). The average number of calls or visits reported per complaint period increased consistently from 0.342 for the under 15 to 1.470 for the 65 and over age group. Except in the under 15 age group the count was higher for females than for males.

Counting only those complaint periods which did have physicians' attendance, close to three (2,682) doctors' calls or clinic visits were reported per complaint period (see Table 6H). The number increased consistently from 1.726 for the under 15 to 3.904 for the 65 and over age group. The total average for females was 2,797 and for males 2,547.

Well over half of the population (56.8 per cent) reported no doctors' calls or clinic visits during the survey year (see Table 7C). Persons reporting one call or visit amounted to 15.1 per cent of the population, persons reporting two calls or visits amounted to 8.1 per cent of the population, and successively smaller proportions reported larger number of doctors' calls or clinic visits (see Chart 3). The age group under 15 made up a comparatively large proportion (36.7 per cent) of persons reporting one call or visit and a successively smaller proportion of persons reporting increasing numbers of calls (only 14.4 per cent of persons reporting 10-14 calls or visits-see Table 7B). The reverse trend was evident for the age groups 25 and over. The proportion of persons reporting various numbers of calls and visits who were in the 15-24 year age group was fairly constant.

In-Patient Hospital Care

Tables 8-11 and Charts 4-5

One tenth (10.2 per cent) of the surveyed population reported hospital care during the year 1950-51 (see column (d) of Table 8). For both sexes combined the proportion of persons in each age group reporting hospitalization did not vary greatly, except that it was rather low (6.8 per cent) for persons under 15 if the newborn are excluded from that age group. There was a very great difference, however, between the proportion of males in the 15-24 and 25-44 age groups reporting hospital care (5.0 per cent and 6.1 per cent respectively) and the proportion of females in the same age categories hospital care (13.3 per cent and 16.2 per cent respectively). For the other age groups the difference between the sexes was small.

In-patient hospital care was given for almost one twentieth (4.7 per cent) of all complaint periods reported (see column (d) of Table 9). The proportion of complaint periods with hospital care increased consistently from 2.7 per cent for the under 15 to 7.4 per cent for the 65 and over age group. The proportion was almost twice as high for females as for males in the 15-44 year range and somewhat higher for males than for females in the other age groups.

During the survey year 122 hospital periods were reported for every 1,000 persons in the population; 100 for every 1,000 males and 145 for every 1,000 females (see column (g) of Table 9). The lowest average numbers of hospital periods were reported for males in the 15-24 year age group (55 per 1,000) and the 25-44 year age group (66 per 1,000). The highest average numbers were reported for females in the 25-44 year age group (185 per 1,000) and the 15-24 year age group (142 per 1,000).

The average number of days spent in hospital per person during the survey year increased generally with age, ranging from about one day per person (1,099 per 1,000 persons) for the under 15 age group to almost four days per person (3,737 per 1,000 persons) for the 65 and over age group (see column (c) of Table 10 and Chart 4). A comparatively high average number of hospital days was reported for the newborn, however, and if the newborn are excluded from the under 15 age group the figure for this age group becomes much lower yet, viz. 644 per 1,000 persons. Females, on the average, reported more days of hospitalization (1,849 per 1,000 persons) than males (1,609 per 1,000 persons).

The average number of days spent in hospital during the survey year by those persons who reported some hospital care was about 17 (16,916 per 1,000 persons—see column (d) of Table 10). While females in general reported more hospital care than males in general, those males who did report hospital care reported more per person (19,541 days per 1,000 persons) than females who did report care (15,123 days per 1,000 persons).

The average length of hospital periods for the under 15 age group was 8 days (see column (e) of Table 10). The average length of hospital periods increased successively with older age groups, reaching a high average of 28 days per period for the 65 and over age group. For males the average length (16 days) was somewhat higher than for females (13 days).

Home Nursing Services

Tables 12-15

Only 150,000 persons, just over one per cent of the population, reported home nursing care during the survey year (see Tables 13A and 13C). For most age-sex groups the number of persons reporting home nursing care was too small to permit of estimates with a sufficient degree of accuracy. Overall figures show, however, that a far greater number of females (110,000) reported home nursing care than males (40,000). The difference was especially great in the case of non-graduate nursing care which was reported by about 40,000 females, but hardly any males. In the case of graduate nursing care the difference was less pronounced with 70,000 females reporting care as against 40,000 males. Home nursing care was reported most frequently by females in the 25-44 year age group (the child-bearing age). Half of the females reporting any nursing care and and two-thirds of those reporting non-graduate nursing care were in this age group (see Table 13B). Nursing care was reported next most frequently by the older age groups.

While more than twice as many persons reported graduate nursing care than reported non-graduate nursing care, the number of days of non-graduate nursing care reported (1,040,000) was slightly in excess of the number of visits of graduate nursing care reported (990,000) (see Table 15A). Females received three-quarters (740,000) of graduate nursing visits and more than half of these (410,000) for persons in the 25-64 year age bracket. For both sexes combined persons over 45 received the largest share of graduate nursing care (600,000 visits out of the total of 990,000). Almost all the days of nongraduate nursing care were reported by females (980,000 days out of the total of 1,040,000). Females in the 25-44 year age group received 360,000 days of non-graduate nursing care.

The average number of visits reported by persons who received graduate nursing care was about nine (9,106 per 1,000 persons reporting graduate nursing care—see Table 15D). In the older age groups (45 years and over) an average of 17 visits per person reporting graduate nursing care (16,984 per 1,000 persons) was recorded. For the under 15 age group the average number of visits was less than four (3,586 per 1,000 persons) and for the 25-44 age group it was less than seven (6,713 per 1,000 persons). The average number of days reported by persons who received non-graduate nursing care was about 21 (21,298 per 1,000 persons). Females in the

25-44 year age group, who received most of the nongraduate nursing care, reported on the average 13 days per person (12,801 per 1,000 persons).

Operations

Table 16 and Chart 6

About 510,000 persons, or one in 27 of the population (3.7 per cent), reported operations during the survey year. Operations were reported by 3.5 per cent of males and by 4.0 per cent of females. The age group under 15 had the highest proportion reporting operations (4.3 per cent) and the 15-24 year age group the lowest (2.9 per cent).

The number of operations reported during the survey year was about 540,000, or 40 per 1,000 persons of the population. The 15-24 year age group reported proportionately the smallest number of operations (29 per 1,000 persons) and the age group under 15 the highest number (45 per 1,000 persons). Females reported proportionately more operations (43 per 1,000 persons) than males (37 per 1,000 persons). Operations were reported particularly frequently by females in the 25-44 year age group (48 per 1,000 persons), by females in the 45-64 year age group (47 per 1,000 persons), and by males in the age group under 15 (47 per 1,000 persons).

Few persons reported more than one operation: there were 1,055 operations for every 1,000 persons reporting operations. Females in the 45-64 year age group most frequently reported more than one operation (1,158 operations per 1,000 persons reporting).

Dental Care

Table 17 and Chart 7

About 2,000,000 persons, or one in every seven of the population (14.7 per cent) reported dental care during the survey year (excluding care given at school dental clinics). The 15-24 year age group had the highest proportion of persons reporting dental care (20.9 per cent). The proportion reporting dental care was successively lower for older age groups and amounted only to 4.0 per cent of persons 65 and over. Dental care was reported by 13.0 per cent of males and by 16.5 per cent of females. The proportion of females reporting dental care was higher than the proportion of male. In all age groups under 65.

The number of dental visits reported during the survey year was about 4,370,000, or 323 per 1,000 persons of the population. The 15-24 year age group reported proportionately the largest number of dental visits (530 per 1,000 persons) and the 65 and over age group the smallest number (77 per 1,000 persons). Females reported proportionately more dental visits than males in all age groups under 65. The total average for males was 264 dental visits per 1,000 persons and for females 383 dental visits per 1,000 persons.

A large number of persons reporting dental care reported more than one dental visit: there were 2,191 visits for every 1,000 persons reporting dental care. Multiple dental visits were reported most frequently in the 15-24 year age group (2,535 visits per 1,000 persons reporting). Multiple visits were reported more frequently by females (2,312 visits per 1,000 persons reporting) than by males (2,040 visits per 1,000 persons reporting).

Examinations for Glasses

Table 18 and Chart 8

About 450,000 persons, or one in every 30 of the population (3.3 per cent), reported examinations for glasses (other than by physicians) during the survey year. Examinations for glasses were reported by 2.6 per cent of males and by 4.0 per cent of females. The 45-64 year age group had the highest proportion reporting examinations for glasses (5.4 per cent) and the age group under 15 the lowest (2.0 per cent).

The number of examinations for glasses reported during the survey year was about 540,000, or 40 per 1,000 persons of the population. The 45-64 year age group reported proportionately the largest number of examinations for glasses (62 per 1,000 persons) and the age group under 15 the smallest number (25 per 1,000 persons). Females reported proportionately more examinations for glasses (47 per 1,000 persons) than males (32 per 1,000 persons).

Relatively few persons reported more than one examination for glasses: there were 1,188 examinations for every 1,000 persons reporting examinations for glasses. Multiple examinations for glasses were reported most frequently by the 15-24 year age group (1,268 examinations per 1,000 persons reporting) and by the age group under 15 (1,244 examinations per 1,000 persons reporting).

Miscellaneous Health Care

Table 19

About 130,000 persons, or one in every 106 of the population (0.9 per cent) reported chiropractic treatment during the survey year. Other miscellaneous health care was reported by about 90,000 persons, or one in every 156 persons of the population (0.6 per cent). In both cases the proportion of females reporting was slightly larger than the proportion of males reporting.

The number of chiropractic treatments reported during the year was about 960,000, or 71 per 1,000 persons of the population. Proportionately more chiropractic treatments were reported by females (84 per 1,000 persons of the population) than by males (58 per 1,000 persons of the population). The number of treatments of other categories of miscellaneous health care reported during the survey year was about 860,000, or 64 per 1,000 persons of the

population. Proportionately more of these treatments were reported by males (68 per 1,000 persons of the population) than by females (60 per 1,000 persons of the population).

While comparatively few persons reported chiropractic care, those who did report such care averaged seven to eight treatments per person (7,501 treatments per 1,000 persons reporting). Persons reporting other categories of miscellaneous health care averaged about 10 treatments per person (9,963 treatments per 1,000 persons reporting).

Total Health Care

Tables 1-2 and Chart 1

More than half of the Canadian population (53.1 per cent) reported some health care during the survey year (see Table 1C). The category of health care received by the largest number of persons was physicians' care, reported by 43.2 per cent of the population. Next came dental care, reported by 14.7 per cent, and in-patient hospital care, reported by 10.2 per cent of the population. Operations were reported by 3.7 per cent of the population and examinations for glasses by 3.3 per cent. Least frequently reported were miscellaneous health care (1.6 per cent of the population) and home nursing care (1.1 per cent of the population).

For each category of health care the proportion of females reporting care was appreciably higher than the proportion of males. For all categories taken together the proportions reporting care were 57.7 per cent for females and 48.6 per cent for males. In the case of most categories of health care (dental care being an exception) the proportion of males under 15 reporting care was fractionally higher than the proportion of females under 15 reporting care. But for the older age groups the proportion of females reporting care was generally much higher than the proportion of males. The difference was especially pronounced in the 25-44 year age group, and was also considerable in the 15-24 and 45-64 year age groups. The relatively small number of persons of 65 and over did not allow reliable estimates to be made for this age group in many cases. While for all categories of health care combined a larger proportion of females than of males in the 65 and over age group reported care, it is not at all certain that such was the case for each separate category of health care.

For all categories of health care combined the smallest proportion of males reporting health care was found in the 15-24 year age group (42.6 per cent). Successively larger proportions of the older male age groups reported health care, but the largest proportion of all was recorded for males in the age group under 15 (52.5 per cent). This general pattern did not hold for all the separate categories of health care. Dental care, for instance, showed a reverse pattern, with the 15-24 year male age group having the highest proportion of persons reporting care and the older male age groups having successively lower proportions reporting care. The under 15 age group. with a generally high proportion reporting care, had a very small proportion reporting examinations for glasses and miscellaneous health care.

The proportion of females under 15 reporting any health care (52.0 per cent) was only fractionally lower than the proportion of males under 15 reporting any health care (52.5 per cent). Yet, while the under 15 age group had a larger proportion reporting health care than any other male age group, it had a smaller proportion reporting health care than any other female age group. The proportion reporting any health care increased successively with the older female age groups, with the exception that the 25-44 year age group had the largest proportion of all reporting health care (65.2 per cent). There were a few significant exceptions to this general pattern. As with males, the highest proportion of persons reporting dental care was found in the 15-24 year age group. with older age groups having successively lower proportions reporting care. The female age groups with the lowest proportion reporting operations were the 45-64 year age group and the 65 and over age group. The highest proportions reporting examinations for glasses were in the female age groups of 45-64 years and 15-24 years.

For both sexes and all categories of health care combined the smallest proportion of persons reporting care (49.1 per cent) was recorded for the 15-24 year age group (particularly because of the very small proportion of males in this age group reporting care). The largest proportion of persons reporting care (56.3 per cent) was recorded for the 25-44 year age group (particularly because of the very large proportion of females in this age group reporting care). The proportion reporting care in the 65 and over age group (54.1 per cent) was a little higher than for the two remaining age groups of under 15 and of 45-64 years (both 52.3 per cent).

ESTIMATED NUMBER OF PERSONS REPORTING VARIOUS CLASSES OF HEALTH CARE

PERSONS

PHYSICIANS'

(SOURCE: TABLE IA)

REPORTING


IN - PATIENT HOSPITAL CARE



HOME NURSING




OPERATIONS



DENTAL CARE



EXAMINATIONS FOR GLASSES



MISCELLANEOUS HEALTH CARE



EACH FIGURE REPRESENTS 100,000 PERSONS REPORTING

TABLE 1. Persons Reporting Health Care during Survey Year A. Estimated number of persons reporting various classes of health care in each age-sex group

		•				Person	s reportin	g			, .
Age-sex group	Population	Any physicians' care	Any in-patient hospital -care	Any medical care	Any home nursing care	Any opera- tions	Any dental care	Any examina- tions for glasses	Any miscel- laneous health care	Any health care	No health care
	(a)	(b) ¹	(c) ²	(d)	(e) ³	(f)4	(g) ⁵	(h) ⁶	(i) ⁷	(i)	(k)
	<u> </u>	•		·——-		(Millions).				
Both sexes		ŧ	1	1 1		1 1	ı 1		1	l	I
All ages	13.54	5. 85	1.38	6.05	0.15	0.51	2.00	0.45	0. 21	7. 19	6, 35
Under 15	4.12 2.05 3.90 2.42 1.05	1.75 0.72 1.78 1.07 0.52	0.43 0.19 0.44 0.23 0.10	1.84 0.75 1.83 1.10 0.54		0.18 0.06 0.14 0.10	0.56 0.43 0.70 0.26 0.04	0.08 0.07 0.13 0.13	0.10 0.06	- 2.15 1.01 2.20 1.26 0.57	1.96 1.04 1.71 1.15 0.48
Male											
All ages	6.82	2.65	0.56	2.75	0.04	0.24	0.88	0.18	0.10	3.32	3.50
Under 15	2.10 1.01 1.93 1.25 0.53	0.90 0.29 0.71 0.51 0.24	0.23 0.05 0.12 0.11 0.05	0.94 0.31 0.73 0.52 0.25	0.02	0.09 0.05 0.05	0.28 0.18 0.28 0.12	0.04	0.05 0.04	1.10 0.43 0.91 0.60 0.27	1.00 0.58 1.02 0.65 0.26
Female											
All ages	6, 72	3.20	0.82	3, 30	0.11	0.27	1.11	0.27	0.11	3.87	
Under 15	2.02 1.04 1.97 1.17 0.52	0.85 0.43 1.08 0.56 0.28	0.20 0.14 0.32 0.11 0.05	0.89 0.44 1.10 0.58 0.29	0.02	•••	0. 29 0. 25 0. 43 0. 13	0.05 0.08 0.08		1.05 0.58 1.28 0.66 0.30	

TABLE 1. Persons Reporting Health Care during Survey Year B. Estimated number of persons in each age group reporting various classes of health care as percentage of total persons reporting these classes of health care, by sex1

						Person	s reportin	g	·		
Age-sex group	Population	Any physicians' care	Any in-patient hospital care	Any medical care	Any home nursing care	Any opera- tions	Any dental care	Any examina- tions for glasses	Any miscel- laneous health care	Any health care	No health care
	(a)	(b)	(c)	· (d)	(e).	(f)	(g)	. (h)	_ (i)	_ (J)	(k)
Both sexes	100.0	100.0	100.0	100.0	100.0	109.0	100.0	100.0	100.0	100.0	100.0
Under 15	30. 4 15. 1 28. 8 17. 8 7. 8	29.9 12.4 30.5 18.3 8.9	30.9 13.6 31.6 16.4 7.5	30.4 12.4 30.2 18.2 8.8	16.7 39.4 32.1	34.8 11.6 28.3 18.8	28.3 21.5 35.3 12.8 2.1	18.1 16.2 28.4 28.9	47.1 26.8	29.9 14.0 30.6 17.6 7.9	30.9 16.4 26.9 18.1 7.6
Male	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Under 15	30.8 14.8 28.3 18.3 7.8	34.0 11.1 26.7 19.2 9.0	40.4 8.9 20.9 20.5 9.3	34.3 11.3 26.6 18.8 9.0		39.5 21.8 20.3	31.4 20.5 31.3 13.8	24.4 30.2	46.4 40.1	33.2 13.0 27.5 18.1 8.2	28.5 16.5 29:1 18.4 7.5
Female All ages	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Under 15	30.0 15.5 29.3 17.4 7.7	26.5 13.4 33.7 17.6 8.9		27.1 13.4 33.3 17.6 8.7	23.4 50.1	30.8 33.9 17.6	25.8 22.3 38.5 12.1	19.0 30.7 28.0	47.8 39.2	27.1 14.9 33.2 17.1 7.7	34.0 16.3 24.1 17.8 7.8

^{1.} Derived from Table 1A: each age group as percentage of "all ages", by sex.

^{1.} See Table 4A, column (f).
2. See Table 8, column (b).
3. See Table 13A, column (d).
4. See Table 16, column (b).

See Table 17, column (b).
 See Table 10, column (b).
 See Table 19, column (b).

TABLE 1. Persons Reporting Realth Care during Survey Year C. Estimated number of persons reporting various classes ofhealth care as percentage of total persons in each age-sex group 1

•						Person	s reporti	ng 			
Age-sex group	Population	physicians' care	Any in-patient hospital care	Care	Any home nursing care	Any opera- tions	Any dental care	Any examina- tions for glasses	Any miscel- laneous health care	Any health care	No healti care
	(a)	(b)	(c)	(d)	(e)	(f)	(g)	(h)	(i)	(J)	(k)
•		l.									
Both sexes											
All ages	100.0	43. 2	10, 2	44. 7	1. 1	3. 7	14. 7	3.3	1. 6	53, 1	46, 9
Jnder 15	100, 0 100, 0 100, 0 100, 0 100, 0	42.5 35.3 45.7 44.3 49.6	10. 4 9. 2 11. 2 9. 4 9. 8	44. 6 36. 6 46. 8 45. 5 50. 8	0.6 1.5 1.3	4.3 2.9 3.7 4.0	13. 7 20. 9 18. 0 10. 6 4. 0	2. 0 3. 6 3. 3 5. 4	2. 5 2. 3	52. 3 49. 1 56. 3 52. 3 54. 1	47. 50. 9 43. 47. 45. 9
Male											
All ages	100. D	38, 9	8, 2	40.4	0. 6	3. 5	13, 0	2, 6	1.5	48.6	51.4
Inder 15	100.0 100.0 100.0 100.0 100.0	42.9 29.2 36.6 40.8 44.8	10. 8 5. 0 6. 1 9. 2 9. 7	45. 0 30. 8 37. 9 41. 7 46. 6	0.4	{ 4.4 2.7 3.8	13. 2 18. 0 14. 3 9. 8	2.1	2, 4	52.5 42.6 47.2 48.2 50.8	47. 5 57. 4 52. 7 51. 8 49. 2
Female											
All ages	100, 0	47, 6	12, 2	49. 0	1, 6	4.0	16. 5	4.0	1. 6	57, 7	42, 3
75-24	100.0 100.0 100.0 100.0 100.0	42. 1 41. 1 54. 7 48. 1 54. 7	10.0 13.3 16.2 9.6 9.8	44. 3 42. 3 55. 6 49. 5 55. 1	0.8 2.7	4.2 4.7 4.1	14. 2 23. 7 21. 7 11. 5	4.9 4.2 6.5	2. 7 2. 6	52.0 55.4 65.2 56.7	48. (44.) 34. (43.)

^{1.} Derived from Table 1A: columns (b)-(k) as percentage of column (a).

TABLE 2. Complaint Periods under Health Care during Survey Year A. Estimated number of complaint periods under various classes of health care for each age-sex group

	-			Period	s with		
Age-sex group	Total number of complaint periods	Any physicians' care	Any in-patient hospital care	Any home nursing care	Any physicians' care or in- patient hos- pital care or home nursing care	Any other health care	No health care
	(a)	(b) ¹	(¢) ²	(d) ³	(e)	(f)	(g)
				(Millions)			-
Both sexes]	1		1	1	
All ages	29, 47	7, 45	1, 39	0, 15	8, 61	0, 47	21, 13
Under 15	11.70 3.53 8.17 4.25 1.81	2. 32 0. 86 2. 22 1. 37 0. 68	0. 32 0. 20 0. 48 0. 26 0. 13		2. 44 0. 98 2. 46 1. 42 0. 71	0. 10 0. 07 0. 17 0. 09	9, 18 2, 50 5, 60 2, 77 1, 08
Male							
All ages	13, 76	3.42	0. 55	0. 04	3, 59	0, 23	10, 01
Under 15	5. 99 1. 53 3. 42 1. 97 0. 85	1. 23 0. 36 0. 90 0. 63 0. 30	0. 17 0. 05 0. 13 0. 13 0. 07	0.02	1. 29 0. 38 0. 94 0. 65 0. 32	0. 05 0. 08 0. 05	4.66 1.13 2.42 1.29 0.52
Female			į				
All ages	15, 71	4, 02	0. 84	0, 11	4, 43	0. 24	11, 12
Under 15	5. 71 2. 01 4. 76 2. 28 0. 96	1, 09 0, 50 1, 32 0, 73 0, 38	0. 15 0. 15 0. 36 0. 13 0. 06	0.02 0.05	{ 1. 15 0. 60 1. 52 0. 77 0. 39	0. 05 0. 09 0. 04	4. 52 1. 38 3. 18 1. 48 0. 56

See Table 5A, colv (1).
 See Table 9, column (b).
 See Table 14A, column (d).

TABLE 2. Complaint Periods under Health Care during Survey Year

B. Estimated number of complaint periods under various classes of health care for each age group as percentage of total complaint periods under these classes of health care, by sex 1

				Period	ls with		
Age-sex group	Total number of complaint periods	Any physicians' care	Any in-patient hospital care (c)	Any home nursing care	Any physicians' care or in- patient hos- pital care or home nursing care (e)	Any other health care	No health care
	(a.)	(0)	(6)	(u)	(6)	(1)	(6)
Both sexes						•	
All ages	100. 0	100.0	100, 0	100, 0	100,0	100, 0	100, 0
Under 15	39.7 12.0 27.7 14.4 6.2	31. 1 11. 5 29. 9 18. 3 9. 2	23. 0 14. 4 34. 6 18. 4 9. 6	15.4 35.6 37.4	30. 4 12. 2 30. 7 17. 7 8. 9	20, 9 15, 5 37, 5 19, 3	43. 4 11. 8 26. 5 13. 1 5. 1
Male		į					•
All ages	100, 0	100. 0	100. 0	100.0	100.0	100, 8	100. 0
Under 15	43.5 11.1 24.8 14.3 6.2	35. 8 10. 5 26. 4 18. 4 8. 8	30. 9 9. 9 22. 9 23. 5 12. 8	} 51.4	35. 9 10. 6 26. 3 18. 2 9. 0	21, 2 37, 3 20, 8	46.5 11.2 24.2 12.9 5.2
Female :							
All ages	100, 0	100. 0	100, 0	100, 0	100.0	100.0	100. 0
Under 15	36. 3 12. 8 30. 3 14. 5 6. 1	27. 1 12. 4 32. 8 18. 3 9. 4	17. 9 17. 3 42. 1 15. 1 7. 6	} 21.1 45.9	25, 9 13, 5 34, 3 17, 4 8, 9	20. 7 37. 8 . 18. 0	40.7 12.4 28.6 13.3 5.1

^{1.} Derived from Table 2A: each age group as percentage of "all ages", by sex.

TABLE 2. Complaint Periods under Health Care during Survey Year

C. Estimated number of complaint periods under various classes of health care as percentage of total complaint periods reported for each age-sex group $^{\rm 1}$

				Perio	is with		
- · Age-sex group	Total number of complaint periods	Any physicians' care	Any in-patient hospital care	Any home nursing care	Any physicians' care or in- patient hos- pital care or home nursing care	Any other health care	No health care
	(a)	(b)	(c)	(d)	(e) .	(n)	(g)
Both sexes							
All ages	100. 0	25. 3	4. 7	0.5	27, 2	1. 6	71. 7
Under 15	100. 0 100. 0 100. 0 100. 0 100. 0	19. 8 24. 3 27. 2 32. 1 37. 6	2. 7 5. 7 5. 9 6. 0 7. 4	0.2 0.7 1.0	20.8 27.7 30.1 33.4 39.4	0.8 2.0 2.1 2.1	78. 5 70. 9 68. 5 65. 1 59. 7
Male				·			
All ages	100, 0	24, 9	4.0	0.3	26.1	1, 6	72. 8
Under 15	100. 0 100. 0 100. 0 100. 0 100. 0	20, 5 23, 4 26, 5 32, 0 35, 5	2.8 3.6 3.7 6.5 8.2	0.2	21.5 25.0 27.6 33.1 37.8	0.8 2.5 2.4	77. 8 73. 8 70. 8 65. 3 60. 9
Female							
All ages	100. 0	25. 6	5.4	0. 7	28.2	1.5	70, 8
Under 15	100. 0 100. 0 100. 0 100. 0 100. 0	19. 1 24. 9 27. 7 32. 3 39. 5	2.6 7.3 7.5 5.6 6.6) 0.3 1.1 ::	20. 1 29. 8 32. 0 33. 7 40. 9	0.9 1.9 1.9	79. 2 68. 7 66. 8 65. 0 58. 6

^{1.} Derived from Table 2A: columns (b)-(g) as percentage of column (a).

TABLE 3. Physicians' Services during Survey Year Estimates of persons reporting physicians' care, complaint periods under physicians' care, and doctors' calls or clinic visits reported, by type of call or visit

		Persons reporting physicians' care			nt periods icians' care	Calls or visits reported			
Type of call or visit	Population (millions)	Number	Percentage	Number	Percentage of total	Number	Rate per 1,	000 persons	
	(a)	(millions) (b) ¹	of total persons (c) ²	(millions) (d) ³	complaint periods (e) ⁴	(millions) (f) ⁵	In population (g) ⁶	Reporting (h) ⁷	
Doctors' office calls]	4,42	32.7	4.78	16, 2	15,31	1,131	3,462	
Doctors' home calls		2.48	18.3	3, 10	10.5	6.93	512	2,793	
Doctors' office and/or home calls	13.54	5,67	41.9	7. 24	24.6	22.24	1,643	3,920	
Clinic visits	13.54	0.63	4.7	0.48	1,6	1.94	143	3,077	
Total (any) doctors' calls and/or clinic visits	j	5,85	43,2	7,45	25.3	24.18	1,786	4, 132	

TABLE 4. Persons Reporting Physicians' Care during Survey Year A. Estimated number of persons reporting doctors' calls or clinic visits in each age-sex group

•				Persons re	eporting		
Age-sex group	Population _	I	octors' calls			Any doctors' calls and/or clinic visits	No
		Office	Home	Office and/or home	Clinic visits		doctors' calls and/or clinic visits
	(a)	(b)	(c)	(d)	(e)	(f)	(g)
•	•			(Millions)			•
Both sexes				1	,	<i>.</i>	
All ages	13,54	4, 42	2,48	5.67	Q 63	5,85	7,69
Under 15	4,12	1,21	0,87	1,69	0,18	1.75	2, 37
15-24 ,	2,05	0.57	0, 26	0.70	0,08	0,72	1, 33
25-44	3,90	1.48	0.64	1,74	0,19	1.78	2.12
45-64	2, 42	0.83	0.43	1.04	0.13	1.07	1.34
65 and over	1,05	0,34	0.28	0.51	0,05	0,52	0,53
Male							
All ages	6, 82	2,00	1,04	2.56	Q 31	2,65	4.17
Under 15	2, 10	0,63	0, 45	0.87	0.10	0,90	1,20
15-24	1.01	0.23	0.08	0.28		0.29	0,71
25-44	1,93	0.57	0,21	0.68	0,08	0.71	1,23
45-64	1.25	0.40	0.19	0.49	0.07	0.51	0.74
65 and over	0,53	0.17	0,12	0, 23	••	0.24	0.30
Female							
All ages	6,72	2,42	1,44	3, 12	0,32	3,20	3,52
Under 15	2.02	0.58	0.42	0.82	0,08	0,85	1.17
15-24	1.04	0.34	0.18	0.42	.,	0.43	0, 61
25-44	1.97	0.90	0.42	1.06	0,11	1.08	0.89
45-64	1,17	0.43	0, 24	0.54	0.06	0,56	0,61
65 and over	0,52	0,17	0, 17	0.28	• •	0, 28	0, 24

Top line of Table 4A.
 Top line of Table 4C.
 Top line of Table 5A.
 Top line of Table 5C.

Top line of Table 6A,
 Top line of Table 6D,
 Top line of Table 6E,

TABLE 4. Persons Reporting Physicians' Care during Survey Year

B. Estimated number of persons in each age group reporting doctors' calls or clinic visits as percentage of total persons reporting doctors' calls or clinic visits, by sex 1

,				Persons r	eporting		
Age-sex group	Population		Doctors' calls			Any	No
	(a)	Office (b)	Home (c)	Office and/or home (d)	Clinic visits (e)	doctors' calls and/or clinic visits (f)	doctors' calls and/or clinic visits (g)
Both sexes							
All ages	100.0	100.0	100.0	100.0	100,0	100.0	100,0
Under 15 15-24 25-44 45-64 65 and over	30.4 15.1 28.8 17.8 7.8	27.3 12.9 33.4 18.7 7.7	35.1 10.4 25.6 17.4 11.5	29.7 12.4 30.6 18.3 9.0	28.6 12.1 30.4 20.9 8.0	29.9 12.4 30.5 18.3 8.9	30.8 17.3 27.6 17.5 6.9
Male				ļ		j	
All ages	100,0	100, 0	100.0	100,0	100,0	100,0	100.0
Under 15	30.8 14.8 28.3 18.3 7.8	31.4 11.6 28.6 20.0 8.3	43.0 7.6 20.4 17.9	33.9 11.0 26.7 19.2 9.1	34.2 25.5 22.6	34.0 11.1 26.7 19.2 9.0	28.7 17.1 29.4 17.7 7.1
Female		-					
All ages	100, 0	100,0	100.0	100, 0	100, 0	100, 0	100, 0
Under 15	30.0 15.5 29.3 17.4 7.7	23,8 14,0 37,3 17,7 7,2	29, 4 12, 4 29, 4 17, 0 11, 8	26.3 13.5 33.9 17.5 8.9	23,3 35,1 19,4	26.5 13.4 33.7 17.6 8.9	33.2 17.4 25.4 17.3 6.7

^{1.} Derived from Table 4A: each age group as percentage of "all ages", by sex.

TABLE 4. Persons Reporting Physicians' Care during Survey Year

C. Estimated number of persons reporting doctors' calls or clinic visits as percentage of total persons in each age-sex group 1

				Persons r	eporting		
Age-sex group	Population	Ε	octors' calls		·	Any doctors' calls and/or clinic visits	No doctors'
,	-	Office	Home	Office and/or home	Clinic visits		calls and/or clinic visits
	(a)	(b)	(c)	(d)	(e)	(f)	(g)
Both sexes					•		
All age groups	100,0	32.7	18, 3	41.9	4,7	43,2	56, 8
Under 15 15-24 25-44 45-64 65 and over	100.0 100.0 100.0 100.0 100.0	29, 3 27, 9 37, 8 34, 3 32, 2	21.2 12.6 16.3 17.8 27.0	41.0 34.3 44.5 42.9 48.4	4.4 3.7 4.9 5.5 4.8	42.5 35.3 45.7 44.3 49.6	57. 5 64. 7 54. 3 55. 7 50. 4
Male							
All age groups	180,0	29,4	15,3	37,5	4,5	38,9	61,1
Under 15	100.0 100.0 100.0 100.0 100.0	30.0 23.1 29.7 32.1 31.2	21.4 7.9 11.0 15.0 21.6	41.3 28.0 35.3 39.4 43.5	5.0 4.0 5.5	42, 9- 29, 2 36, 6 40, 8 44, 8	57. 1 70. 8 63. 4 59. 2 55. 2
Female					•		
All age groups	100,0	36, 0	21.4	46.4	4,8	47, 6	52,4
Under 15	100,0 100,0 100,0 100,0 100,0	28.6 32.6 45.8 36.6 33.3	21.0 17.2 21.4 20.9 32.5	40.6 40.3 53.5 46.5 53.4	3,8 5.8 5,4	42, 1 41, 1 54, 7 48, 1 54, 7	57, 9 58, 9 45, 3 51, 9 45, 3

^{1.} Derived from table 4A: columns (b) - (g) as percentage of column (a).

TABLE 5. Complaint Periods under Physicians' Care during Survey Year

A. Estimated number of complaint periods with doctors' calls or clinic visits for each age-sex group

·				Periods	with		
Age-sex group	Total number of		Octors' calls			Any doctors'	No doctors' calls and/or clinic visits
	complaint periods	Office	Home	Office and/or home	Clinic visits	calls and/or clinic visits	
	(a)	(b)	(c)	(d)	(e)	(1)	(g)
				(Millions)			
Both sexes		1		1 1	•	1	
All ages	29, 47	4, 73	3, 10	7. 24	0.48	7,45	22,02
Under 15	11. 70 3. 53 8. 17 4. 25 1. 81	1. 26 0. 59 1. 62 0. 95 0. 37	1.14 0.30 0.74 0.54 0.38	2, 25 0, 83 2, 17 1, 33 0, 67	0.13 0.06 0.14 0.11 0.47	2.32 0.86 2.22 1.37 0.68	9, 38 2, 68 5, 95 2, 89 1, 13
Male				,			
All ages	13, 76	2, 26	1. 30	3, 32	0, 25	3, 42	10. 34
Under 15	5. 99 1. 53 3. 42 1. 97 0. 85	0. 67 0. 27 0. 70 0. 46 0. 17	0.60 0.09 0.24 0.22 0.15	1. 19 0. 35 0. 88 0. 61 0. 30	0.79 0.07 0.06	1. 23 0. 36 0. 90 0. 63 0. 30	4. 76 1. 17 2. 51 1. 34 0. 55
Female			į				
All ages	15.71	2, 52	1, 79	3, 92	0,23	4.02	11, 69
Under 15	5.71 2.01 4.76 2.28 0.96	0.59 0.32 0.92 0.49 0.19	0.55 0.21 0.50 0.32 0.22	1.06 0.49 1.29 0.72 0.37	0.05 0.08 0.05	1.09 0.50 1.32 0.73 0.38	4. 62 1. 51 3. 44 1. 54 0. 58

TABLE 5. Complaint Periods under Physicians' Care during Survey Year

B. Estimated number of complaint periods with doctors' calls or clinic visits for each age group as percentage of total complaint periods with these doctors' calls or clinic visits, by sex'

				Periods	with		
Age-sex group	Total number of	Í	Doctors' calls			Any	No doctors' calls and/or clinic visits
	complaint periods	Office	Home (c)	Office and/or home (d)	Clinic visits (e)	doctors' calls and/or clinic visits (f)	
Both sexes	(4)	(0)	(0,		(6)	2	(8)
All ages	100, 0	, 100,0	100.0	100.0	100.0	100.0	100.0
Under 15	39. 7 12. 0 27. 7 14. 4 6. 2	26. 2 12. 4 33. 9 19. 8 7. 7	37. 0 9. 7 23. 8 17. 4 12. 2	31. 1 11. 5 29. 9 18. 3 9. 2	25. 9 11. 6 29. 7 23. 0 9. 8	31. 1 11. 5 29. 9 18. 3 9. 2	42. 6 12. 1 27. 0 13. 1 5. 1
Male							
All ages	100, 0	100,0	100, 0	100.0	100,0	100.0	100.0
Under 15	43. 5 11. 1 24. 8 14. 3 6. 2	29. 5 11. 8 30. 9 20. 2 7. 6	45. 8 7. 2 18. 4 16. 8 11. 8	35. 7 10. 4 26. 5 18. 5 8. 9	31. 4 26. 2 24. 4	35. 8 10. 5 26. 4 18. 4 8. 8	46. 1 11. 3 24. 3 13. 0 5. 3
Female				1			
All ages	100.0	. 100,0	100.0	100.0	100.0	100.0	100.0
Under 15	36. 3 12. 8 30. 3 14. 5 6. 1	23. 3 12. 9 36. 7 19. 4 7. 7	30. 5 11. 5 27. 7 17. 8 12. 5	27. 1 12. 4 32. 8 18. 2 9. 5	20. 1 33. 5 21. 6	27. 1 12. 4 32. 8 18. 3	39. 5 12. 9 29. 4 13. 2 5.0

^{1.} Derived from Table 5A: each age group as percentage of "all ages", by sex.

TABLE 5. Complaint Periods under Physicians' Care during Survey Year

C. Estimated number of complaint periods with doctors' calls or clinic visits as percentage of total complaint periods reported for each age-sex group'

			-	Periods	with		
Age-sex group	Total number of]	Doctors' calls	İ		Any	No doctors' calls and/or clinic visits (g)
nge sox group	complaint periods (a)	Office (b)	Home (c)	Office and/or home (d)	Clinic visits (e)	doctors' calls and/or clinic visits (f)	
Both sexes							
All ages	100,0	16, 2	10, 5	24. 6	1, 6	25, 3	74. 7
Under 15 15-24 25-44 45-64 65 and over	100.0 100.0 100.0 100.0 100.0	10.7 16.8 19.9 22.3 20.2	9, 8 8, 5 9, 0 12, 7 20, 8	19. 2 23. 5 26. 5 31. 2 36. 8	1. 1 1. 6 1. 8 2. 6 2. 6	19. 8 24. 3 27. 2 32. 1 37. 6	80. 2 75. 7 72. 8 67. 9 62. 4
Male			·				
All ages	100.0	16, 5	9, 5	24, 1	1.8	24.9	75, 1
Under 15 15-24 15-64 15-64 55 and over	100.0 100.0 100.0 100.0 100.0	11. 1 17. 5 20. 5 23. 2 20. 2	10. 0 6. 2 7. 0 11. 1 18. 0	19. 8 22. 7 25. 7 31. 0 34. 6	1. 3 1. 9 3. 1	20. 5 23. 4 26. 5 32. 0 35. 5	79. 5 76. 6 73. 6 68. 0 6 4. 5
Female							
All ages	100, 0	16, 0	11,4	25,0	1, 5	25, 6	74.4
Under 15 15-24 25-44 45-64 85 and over	100.0 100.0 100.0 100.0 100.0	10. 3 16. 2 19. 4 21. 5 20. 2	9. 6 10. 2 10. 4 14. 0 23. 3	18. 6 24. 2 27. 1 31. 4 38. 7	0.8 1.6 2.2	19. 1 24. 9 27. 7 32. 3 39. 5	80. 9 75. 1 72. 3 67. 7 60. 4

^{1.} Derived from Table 5A: columns (b)-(g) as percentage of column (a).

TABLE 6. Number of Doctors' Calls and Clinic visits during Survey Year

A. Estimated number of doctors' calls or clinic visits for each age-sex group

		Doctors' calls		Clinic	Total doctors' calls and
Age-sex group	Office (a)	Home (b)	Total (c)	visits (d)	cails and clinic visits (e)
			(Millions)		
Both sexes	,				
All ages	15, 31	6, 93	22, 24	1, 94	24. 18
Under 15. 15-24 25-44 45-64 65 and over	2, 86 1, 96 5, 65 3, 49 1, 35	1.72 0.62 1.70 1.57 1.33	4, 58 2, 57 7, 35 5, 06 2, 68	0.40 0.20 0.63 0.49 0.22	4.98 2.77 7.98 5.55 2.90
Male					
All ages	6, 14	2, 73	8, 87	1.03	9, 90
Under 15	1.52 0.63 1.81 1.59 0.60	0, 87 0, 18 0, 41 0, 66 0, 60	2, 39 0, 81 2, 22 2, 25 1, 20	0.26 0.29 0.32	2. 65 0. 90 2. 51 2. 57 1. 27
Female			•		
All ages	9, 16	4, 20	13.37	0.90	14. 27
Under 15	1. 35 1. 33 3. 84 1. 90 0. 75	0.85 0.44 1.28 0.91 0.72	2. 19 1. 77 5. 13 2. 81 1. 47	0. 14 0. 34 0. 16	2. 34 1. 87 5. 47 2. 97 1. 62

TABLE 6. Number of Doctors' Calls and Clinic Visits during Survey Year

B. Estimated number of doctors' calls or clinic visits reported for each age group as percentage of total doctors 'calls or clinic visits reported, by sex 1

	1	Doctors' calls			Total doctors'
. Age-sex group	Office (a)	Home (b)	Total (c)	Clinic visits (d)	calls and clinic visits (e)
Both sexes					
All ages	100.0	100.0	100.0	100, 0	100.0
Under 15 15-24 25-44 45-64 65 and over	18.7 12.8 36.9 22.8 8.8	24.8 8.9 24.5 22.6 19.1	20.6 11.6 33.0 22.8 12.0	20.6 10.2 32.5 25.1 11.5	20.6 11.5 33.0 22.9 12.0
Male					
All ages	100,0	100.0	100, 0	100.0	100, 0
Under 15	24.7 10.3 29.4 25.8 9.8	31.9 6.6 15.2 25.3 22.0	26.9 9.1 25.0 25.3 13.6	25. 1 27. 8 31. 4	26.7 9.1 25.3 26.0 12.9
Female					
All ages	100.0	100.0	100.0	100.0	100.0
Under 15	14.7 14.5 41.9 20.8 8.1	20. 2 10. 4 30. 5 21. 6 17. 2	16.4 13.2 38.3 21.0 11.0	15.6 37.9 17.9	16. 4 13. 1 38. 3 20. 8 11. 4

^{1.} Derived from Table 6A: each age group as percentage of "all ages", by -ex.

TABLE 6. Number of Doctors' Calls and Clinic Visits during Survey Year

C. Estimated number of doctors' call or clinic visits as percentage of total number of these doctors' calls and clinic visits for each age-sex group 1

Tor each	age-sex group	μ - 	<u> </u>			
	. 1	Doctors' calls			Total doctors'	
Age-sex group	Office (a.)	Home (b)	Total (c)	Clinic visits (d)	calls and clinic visits . (e)	
- Both sexes						
All ages	63,3	28. 7	92.0	8.0	100, 0	
Under 15 15 - 24 25 - 44 45 - 64 65 and over	57.5 70.6 70.8 62.9 46.6	34.5 22.3 21.3 28.3 45.7	92.0 92.9 92.1 91.2 92.3	8.0 7.1 7.9 8.8 7.7	100.0 100.0 100.0 100.0 100.0	
Male	ľ					
· All ages	62.0	27.5	89. 6	10.4	100.0	
Under 15	57.3 69.9 72.0 61.7 47.4	32.9 19.9 16.5 25.7 47.1	90.2 89.7 88.5 87.4 94.5	9.8 11.5 12.6	100.0 100.0 100.0 100.0 100.0	
Female						
All ages	64.2	29.5	93. 7	6.3	100.0	
Under 15	57.6 70.9 70.3 64.0 45.9	36.3 23.5 23.4 30.6 44.6	94.0 94.4 93.7 94.5 90.5	6.0 6.3 5.5	100.0 100.0 100.0 100.0 100.0	

^{1.} Derived from Table 6A: columns (a)-(d) as percentage of column (e).




TABLE 6. Number of Doctors' Calls and Clinic Visits during Survey Year D. Estimated average number of doctors' calls or clinic visits reported per 1,000 persons of each age-sex group 1

		Doctors' calls		Citata	Total doctors* calls and	
. Age-sex group	Office	Home (b)	Total (c)	Cliniç visits (d)	clinic visits (e)	
Both sexes						
All ages	1, 131	512	1, 643	143	1, 786	
Under 15	695 955 1, 448 1, 444 1, 281	418 302 434 650 1,258	1, 113 1, 256 1, 882 2, 094 2, 538	97 97 161 202 212	1, 210 1, 353 2, 044 2, 296 2, 751	
Male					`	
All ages	901	400	1, 301	. 152	1, 452	
Under 15	723 625 935 1, 273 1, 130	415 178 214 531 1, 125	1, 138 803 1, 149 1, 804 2, 255	124 149 261	1, 261 895 1, 298 2, 065 2, 386	
Female						
All ages	1, 364	626	1, 989	135	2, 124	
Under 15 15 - 24 25 - 44 45 - 64 65 and over	667 1, 273 1, 950 1, 626 1, 435	421 421 651 778 1, 394	1, 088 1, 694 2, 601 2, 404 2, 829	70 1 74 139	1, 158 1, 795 2, 775 2, 542 3, 125	

^{1.} Columns (a)-(e) of Table 6A divided by one thousandth of column (a) of Table 4A.

TABLE 6. Number of Doctors' Calls and Clinic Visits during Survey Year E. Estimated average number of doctors' calls or clinic visits reported per 1,000 persons reporting doctors' calls or clinic visits for each age-sex group

		Doctors' calls		Clinic	Total doctors' calls and	
Age-sex group	Office (a.) ¹	Home (b) ²	Total (c) ³	visits (d)4	clinic clinic visits (e) ⁵	
Both sexes						
All ages	3,462	2,794	3, 920	3, 077	4, 132	
Under 15	2, 374 3, 418 3, 828 4, 214 3, 972	1,973 2,392 2,668 3,650 4,655	2, 716 3, 667 4, 226 4, 884 5, 244	2, 223 2, 601 3, 285 3, 690 4, 454	2, 848 3, 836 4, 470 5, 180 5, 541	
Majes			•			
All ages	3, 069	2, 614	3,469	3, 389	3, 738	
Under 15	2, 411 2, 710 3, 151 3, 964 3, 625	1, 942 2, 248 1, 942 3, 550 5, 198	2, 752 2, 868 3, 251 4, 572 5, 180	2, 485 3, 697 4, 704	2, 937 3, 061 3, 550 5, 063 5, 330	
Female						
All ages	3, 787	2,924	4, 289	2, 784	4, 459	
Under 15 15-24	2, 334 3, 903 4, 258 4, 448 4, 306	2,006 2,456 3,033 3,726 4,284	2, 678 4, 204 4, 857 5, 165 5, 298	1, 863 3, 004 2, 578	2, 753 4, 369 5, 073 5, 286 5, 718	

Column (a) of Table 6A divided by one thousandth of column (b) of Table 4A.
 Column (b) of Table 6A divided by one thousandth of column (c) of Table 4A.
 Column (c) of Table 6A divided by one thousandth of column (d) of Table 4A.
 Column (d) of Table 6A divided by one thousandth of column (e) of Table 4A.
 Column (e) of Table 6A divided by one thousandth of column (f) of Table 4A.

TABLE 6. Number of Doctors' Calls and Clinic Visits during Survey Year

F. Estimated number of doctors' calls or clinic visits for complaint periods for each age-sex group

·		Doctors' calls	, .	6 11-1-	Total doctor's calls and clinic visits (e)	
Age-sex group	Office (a)	Home (b)	Total (c)	Clinic visits (d)		
			(Millions)			
Both sexes	1	1	1]	
All ages	11.68	6.76	18.44	1.53	19.97	
Under 15	2.03 1.32 4.06 3.10 1.17	1.69 0.60 1.61 1.56 1.30	3.72 1.92 5.67 4.66 2.47	0.28 0.13 0.50 0.43 0.19	4.00 2.05 6.17 5.09 2.66	
Malé					-	
All ages	5.18	2.69	7.87	0.85	8, 72	
Under 15. 15-24 25-44 45-64 65 and over	1.09 0.58 1.61 1.41 0.50	0.86 0.18 0.41 0.66 0.59	1.94 0.75 2.02 2.07 1.09	0.19 0.25 0.30	2.14 0.81 2.27 2.36 1.14	
Female	\$,	
All ages	6.51	4.97	10.57	0.68	11.25	
Under 15	0.95 0.74 2.45 1.70 0.67	0.83 0.42 1.20 0.90 0.71	1.78 1.16 3.66 2.59 1.38	0.08 0.25 0.13	1.86 1.24 3.91 2.73 1.52	

TABLE 6. Number of Doctors' Calls and Clinic Visits during Survey Year
G. Estimated average number of doctors' calls or clinic visits per complaint period for each age-sex group 1

		Doctors' calls			Total
Age-sex group	Office (a)	Home (b)	Total (c)	Clinic visits (d)	doctors' calls and clinic visits (e)
Both sexes					
All ages	0.396	0.229	0.626	0.052	0.678
Under 15	0.174 0.372 0.497 0.730 0.646	0.144 0.169 0.197 0.367 0.718	0.318 0.542 0.694 1.096 1.364	0.024 0.037 0.061 0.101 0.106	0.342 0.579 0.756 1.197 1.470
Male		,			•
All ages	0, 376	0.196	0.572	0.062	0. 634
Under 15	0.181 0.377 0.470 0.712 0.588	0.143 0.116 0.120 0.334 0.690	0.324 0.493 0.591 1.047 1.278	0.032 0.073 0.150	0.524 0.531 0.664 1.196 1.340
Female					
All ages	0.414	0.259	0.673	0.043	0.716
Under 15	0.166 0.369 0.516 0.744 0.697	0.145 0.210 0.253 0.394 0.743	0.311 0.579 0.769 1.139 1.440	0.015 0.053 0.058	- 0.326 0.616 0.821 1.197 1.585

^{1.} Columns (a)-(e) of Table 6F divided by column (a) of Table 5A.

TABLE 6. Number of Doctors' Calls and Clinic Visits during Survey Year H. Estimated average number of doctors' calls or clinic visits per complaint period with doctors' calls or clinic visits for each age-sex group

		Doctors' calls			Total doctors'	
Age-sex group	Office (a) ¹	Home (b) ²	Total (c) ³	Clinic visits (d) ⁴	calls and clinic visits (e) ⁵	
Both sexes						
All ages	2, 184	2. 442	2, 547	3. 153	2. 682	
Under 15	1.475 2.000 2.194 2.892 3.447	1. 620 2. 222 2. 501 3. 279 3. 195	1.655 2.302 2.620 3.509 3.710	2, 197 2, 351 3, 473 3, 837 4, 057	1. 726 2. 384 2. 777 3. 726 3. 904	
Male		<u>.</u>		,		
All ages	2, 066	2, 286	2, 373	3.385	2, 547	
Under 15	1.436 1.882 1.714 3.007 3.835	1.628 2.150 2.297 3.075 2.913	1.640 2.173 2.300 3.375 3.692	2. 445 3. 817 4. 825	1.741 2.266 2.508 3.743 3.774	
Female						
All ages	2, 270	2, 582	2, 694	2, 904	2. 797	
Under 15	1.518 2.055 2.426 2.814 3.182	1, 610 2, 282 2, 655 3, 469 3, 446	1. 670 2. 394 2. 838 3. 624 3. 725	1. 782 3. 185 2. 637	1. 710 2. 468 2: 961 3. 711 4. 007	

TABLE 7. Frequency of Doctors' Calls and Clinic Visits during Survey Year A. Estimated number of persons in each age-sex group with various numbers of doctors' calls and clinic visits

			Persons v	vith follow	ing total n	umber of de	octors' cal	ls and clin	nic visits:	
Age-sex group	Popula- tion (a)	0 (b)	1 (c)	2 (d)	3 (e)	4 (f)	5+9 (g)	10-14 (h)	15-19 (i)	20 or more (J)
Both sexes			ı	1	(Milli	ons)				
All ages	13, 54	7. 69	2. 04	1, 09	0, 72	0.48	1, 00	0, 31	0.09	0, 12
Under 15	4. 12 2. 05 3. 90 2. 42 1. 05	2. 37 1. 33 2. 12 1. 34 0. 53	0. 75 0. 25 0. 54 0. 35 0. 16	0.35 0.14 0.34 0.17 0.09	0. 22 0. 09 0. 21 0. 14 0. 05	0. 13 0.07 0. 15 0. 09 0. 05	0. 24 0. 11 0. 37 0. 18 0. 10	0.04 0.11 0.08	::	••
Male			į							1
All ages	6, 82	4. 17	1, 04	0,51	0, 32	0, 21	0, 37	0, 11	•• .	0, 05
Under 15	2. 10 1. 01 1. 93 1. 25 0. 53	1. 20 0. 71 1. 23 0. 74 0. 30	0.39 0.12 0.28 0.17 0.08	0.16 0.07 0.15 0.08 0.04	0.12 0.08 0.07	0.07 0.05 0.05	0. 13 0. 10 0. 08 0. 04	••	 	••
Female		,								
All ages	6, 72	3, 52	1.00	0.58	0.40	0, 27	0.62	0, 20	0.06	0.07
Under 15	2.02 1.04 1.97 1.17 0.52	1. 17 0. 61 0. 89 0. 61 0. 24	0.36 0.12 0.26 0.17 0.08	0.19 0.07 0.19 0.09 0.05	0.10 0.06 0.14 0.08	0.06 0.09 0.04	0.11 0.09 0.27 0.10 0.06	0. 08 0. 05	::	••

Column (a) of Table SF divided by column (b) of Table 5A.
 Column (b) of Table SF divided by column (c) of Table 5A.
 Column (c) of Table SF divided by column (d) of Table 5A.
 Column (d) of Table SF divided by column (e) of Table 5A.
 Column (e) of Table SF divided by column (f) of Table 5A.




TABLE 7. Frequency of Doctors' Calls and Clinic Visits during Survey Year

B. Estimated number of persons in each age-sex group with various numbers of doctors' calls and clinic visits as percentage of total persons with various numbers of doctors' calls and clinic visits, by sex1

			Persons w	ith followi	ng total nu	mber of do	ctors' cal	is and clin	ic visits:	
Age-sex group	Popula- tion (a)	(p)	1 (c)	2 (d)	3 (e)	4 (f)	5-9 (g)	10-14 (h)	15-19 (i) 100. 0	20 or more (j)
Both sexes					İ					
All ages	100.0	100. 0	100.0	100.0	100. 0	100. 0	100. 0	100.0	100. 0	100.0
Under 15	30. 4 15. 1 28. 8 17. 8 7. 8	30.8 17.2 27.6 17.5 7.0	36, 7 12, 2 26, 5 16, 9 7, 7	31.9 13.0 31.2 15.6 8.3	31.0 12.3 29.3 19.9 7.4	27. 9 14. 0 30. 4 18. 2 9. 4	23, 9 11, 3 36, 7 17, 7 10, 4	14. 4 34. 6 26. 4		••
Male					ļ					
All ages	100.0	100.0	100. 0	100.0	100.0	100.0	100.0	100. 0	100.0	100, 0
Under 15	30.8 14.8 28.3 18.3 7.8	28. 7 17. 1 29. 4 17. 7 7. 1	37. 4 11. 9 26. 5 16. 5 7. 7	31, 8 13, 8 30, 1 16, 2 8, 2	37. 8 23. 5 20. 9	32. 6 25. 1 21. 1	34.8 26.8 21.0 10.8	••	••	••
Female								ļ		
All ages	100. 0	100. 0	100.0	100. 0	100.0	100. 0	100.0	100. 0	100.0	100.0
Under 15	30. 0 15. 5 29. 3 17. 4 7. 7	33. 2 17. 4 25. 4 17. 3 6. 7	36. 0 12. 4 26. 5 17. 3 7. 7	32, 1 12, 3 32, 2 15, 1 8, 4	25. 4 14. 0 34. 1 19. 2	24. 2 34. 7 15. 9	17. 4 14. 0 42. 6 15. 8 10. 1	39. 3 25. 3	••	••

^{1.} Derived from Table 7A: each age group as percentage of "all ages", by sex.

TABLE 7. Frequency of Doctors' Calls and Clinic Visits during Survey Year

C. Estimated number of persons with various numbers of doctors' calls and clinic visits as percentage of total persons in each age-90x group!

	-	•	Persons w	ith followi	ng total nu	mper of do	ctors' cal	is and clin	ic visits:	
Age-sex group	Popula- tion (a)	0 (b)	1 (c)	2 (d)	3 (e)	4 (f)	5+9 (g)	10-14 (h)	15-19 (i)	or more (j)
Both sexes						ļ				
All ages	100.0	56.8	15. 1	8.1	5.3	3.5	7.4	2.3	0. 7	0.9
Under 15	100, 0 100, 0 100, 0 100, 0 100, 0	57, 5 64, 7 54, 3 55, 7 50, 4	18, 2 12, 1 13, 9 14, 3 14, 9	8. 5 6. 9 8. 7 7. 1 8. 6	.5, 4 4, 3 5, 4 5, 9 5, 1	3.3 3.7 3.6 4.3	- 5.8 5.5 9.4 7.3 9.8	1.1 2.7 3.4	• • • • • • • • • • • • • • • • • • •	**
Male							·			
All ages	100.0	61. 1	15.3	7. 5	4.7	3. 1	5. 5	1. 6	••	0.7
Under 15	100.0 100.0 100.0 100.0 100.0	57. 1 70. 8 63. 4 59. 2 55. 2	18.6 12.4 14.3 13.8 15.0	7. 7 7. 0 7. 9 6. 6 7. 8	5. 8 3. 9 5. 4	3, 3 2, 8 3, 6	6. 2 5. 1 6. 3 7. 5	••	••	••
Female		į			.					
All ages	100.0	52.4	14. 8	8. 7	5. 9	4, 0	9.3	3.0	0.9	1.1
Under 15	100.0 100.0 100.0 100.0 100.0	57. 9 58. 9 45. 3 51. 9 45. 3	17.8 11.9 13.4 14.8 14.8	9, 3 6, 9 9, 5 7, 6 9, 4	5. 0 5. 3 6. 9 6. 5	3. 2 4. 7 3. 6	5. 4 8. 4 13. 5 8. 4 12. 1	4.0 4.3	••	

^{1.} Derived from Table 7A: columns (b)-(j) as percentage of column (a).

TABLE 7. Frequency of Doctors' Calls and Clinic Visits during Survey Year

D. Estimated number of persons in each age-sex group with various numbers of doctors' calls and clinic visits (cumulative)

,		P	ersons with	following t	otal number	s of doctors	' calls and	clinic visit	s '
Age-sex group	Population	1	2	3	4	5	10	15	20
<u> </u>	(a)	or more (c)	or more (d)	or more (e)	or more (f)	or more (g)	or more (h)	or more (i)	or more (j)
,	•				(Millions)				
Both sexes		1						-	
All ages	13,54	5.85	3.81	2. 71	2, 90	1,51	0, 52	0, 21	0, 12
Under 15 15 - 24 25 - 44 45 - 64	4.12 2.05 3.90 2.42 1.05	1.75 0.72 1.78 1.10 0.52	1.00 0.47 1.24 0.73 0.37	0.65 0.33 0.90 0.55 - 0.28	0.43 0.24 0.70 0.41 0.22	0.29 0.18 0.54 0.32 0.18	0.06 0.06 0.18 0.15 0.07	0. 07 0. 07	**
Male									
All ages	6.82	2.65	1, 61	1,10	0.77	0, 56	0, 19	0.08	0, 05
Under 15	2.10 1.01 1.93 1.25 0.53	0.90 0.29 0.71 0.51 0.24	0.51 0.17 0.43 0.34 0.16	0, 35 0, 10 0, 28 0, 25 0, 12	0.23 0.07 0.20 0.19 0.09	0.16 0.04 0.15 0.14 0.07	0.05 0.06	• • •	••
Female							:	٠	
All ages	6.72	3, 20	2, 28	1, 62	1, 22	1, 00	0, 33	0, 13	0, 07
Under 15	2.02 1.04 1.97 1.17 0.52	0.85 0.43 1.08 0.56 0.28	0.49 0.30 0.81 0.39 0.21	0,30 0,23 0,63 0,30 0,16	0,20 0,18 0,50 0,23 0,13	0,14 0,13 0,40 0,18 0,10	0.13 0.08	o, ô5	••

TABLE 7. Frequency of Doctors' Calls and Clinic Visits during Survey Year

E. Estimated number of persons in each age-sex group with various numbers of doctors' calls and clinic visits as percentage of total persons with various numbers of doctors' calls and clinic visits, by sex (cumulative)

		Pe	ersons with	following t	otal number	s of doctors	' calls and	clinic visit	s:
Age-sex group	Population (a)	or more (c)	2 or more (d)	3 or more (e)	or more	or more (g)	or more (h)	or more (i)	20 or more (j)
Both sexes									
All ages	100, 0	100, 0	100.0	100. Q	100, 8	100, 0	100, 0	100.0	100, 0
Under 15 15-24 25-44 45-64 65 and over	30.4 15.1 28.8 17.8 7.8	29. 9 12. 4 30. 5 18. 3 8. 9	26. 2 12. 5 32. 6 19. 0 9. 6	24.0 12.2 33.2 20.4 10.2	21.4 12.2 34.6 20.6 11.1	19.4 11.6 36.0 21.4 11.7	10.7 12.3 34.5 28.3 14.2	34.4 31.1	••
Male									
All ages	100, 8	100, 0	100.0	100, 0	100, 0	100, 0	100, 0	100, 0	100, 0
Under 15	30.8 14.8 28.3 18.3 7.8	34.0 11.1 26.7 19.2 9.0	31.8 10.6 26.7 20.9 9.9	31.9 9.1 25.2 23.1 10.7	29, 4 8, 6 25, 9 24, 0 12, 0	28, 2 7, 4 26, 2 25, 2 13, 0	25. 1 33, 4	••	••
Female			•				İ		
All ages	100, 0	100,0	100, 0	100. 0	100, 0	100. 0	100. 0	100, 0	100, 0
Under 15	30.0 15.5 29.3 17.4 7.7	26.5 13.4 33.7 17.6 8.9	22.2 13.8 36.9 17.7 9.4	18.6 14.4 38.7 18.6 9.8	16, 4 14, 5 40, 1 18, 4 10, 6	14, 2 14, 1 41, 6 19, 2 10, 9	39.8 25.5	40.5	••

^{1.} Derived from Table 7D: each age group as percentage of "all ages", by sex.

TABLE 7. Frequency of Doctors' Calls and Clinic Visits during Survey Year F. Estimated number of persons with various numbers of doctors' calls and clinic visits as percentage of total persons in each age-sex group (cumulative).

		Persons with following total numbers of doctors' calls and clinic visits											
Age-sex group	Population	1	2	3	4	5	10	15	20				
	(a)	or more (c)	or more (d)	or more (e)	or more (f)	or more (g)	or more (h)	or more	or more (j)				
Both sexes						•							
All ages	100, 0	43, 2	28, 1	29, 0	14,7	11, 2	3, 8	1, 6	0, 9				
Under 15	100,0 100,0	42.5 35.3	24.3 23.1	15.8 16.2	10.4 11.9	7.1 8.6	1.3 3.1	::					
25 - 44	100.0	45.7	31.8	23, 1	17,7	13.9	4.6 6.1	1.9					
45-6465 and over	100.0 100,0	44.3 49.6	30.0 34.7	23.0 26.1	17.0 21.1	13.4 16.8	7.0	2, 7 3, 5	::				
· Male													
All ages	100.0	38.9	23.5	16.1	11.3	8, 2	2.7	1. 2	0.7				
Under 15	100.0	42.9 29.2	24.4 16.9	16.7 9.9	10.8 6.6	7.5 4.1	••	••	<i>i.</i>				
15 - 24	100.0 100.0	36,6	22.2	14.3	10,4	7.6	2.4	::	• • •				
45 - 64	100.0 100.0	40, 8 44, 8	26.9 29.8	20.3 22.0	14.9 17.4	11.3 13.6	5.0	::	• •				
OV BRU OTO MANAGEMENT CONTROL OF THE				,			- •		,,,				
Female	1												
All ages	100, 0	47. 6	32.8	24. 1	18. 2	14. 2	4.9	2, 0	1, 1				
Under 15	100.0 100.0	42.1 41.1	24.2 29.2	14.9 22.3	9.9 17.0	6,7 12.9	••	••	• •				
15 - 24 25 - 44	100.0	54,7	41.3	31,7	24.9	20.2	<u>6</u> . 7	2.7	• • • • • • • • • • • • • • • • • • • •				
45 - 64	100.0 100.0	48.1 54.7	33.3 39.8	25.7 30.4	19.2 24.8	15.6 20.0	7.2	::	••				
65 and over	100,0	54, Y	39.8	30,4	24,8	20.0	••	••	••				

^{1.} Derived from Table 7D: columns (c)-(j) as percentage of column (a).

TABLE 8. Persons Reporting In-Patient Hospital Care during Survey Year Estimates of persons reporting in-patient hospital care, by age and sex

		P ₀	ersons reporting ca	r.e
Age-sex group	Population (millions) (a)	Number (millions) (b)	As percentage of total persons reporting care (by sex)	As percentage of total persons in age-sex group (d)2
Both sexes	·			
All ages	13,54	1, 38	100.0	10, \$
Under 15 (except newborn) Under 15 15-24 25-44 45-64 65 and over	3.81 4.12 2.05 3.90 2.42 1.05	0, 26 0, 43 0, 19 0, 44 0, 23 0, 10	18.8 30.9 13.6 31.6 16.4 7.5	6, 8 10, 4 9, 2 11, 2 9, 4 9, 8
Maje				
All ages	6,82	0, 56	100. 0	8.2
Under 15 (except newborn) Under 15 15-24 25-44 45-64 65 and over	1, 94 2, 10 1, 01 1, 93 1, 25 0, 53	0, 14 0, 23 0, 05 0, 12 0, 11 0, 05	26, 1 40, 4 8, 9 20, 9 20, 5 9, 3	7, 2 10, 8 5, 0 6, 1 9, 2 9, 7
Female				
All ages	6.72	0, 82	100.0	12, 2
Under 15 (except newborn) Under 15 15-24 25-44 45-64 65 and over	1.87 2.02 1.04 1.97 1.17 0.52	0. 12 0. 20 0. 14 0. 32 0. 11 0. 05	14.5 24.4 16.8 38.9 13.6 6.2	6,4 10,0 13,3 16,2 9,6 9,8

Derived from column (b): each age group as percentage of "all ages", by sex.
 Column (b) as percentage of column (a).

TABLE 9. Periods of In-Patient Hospital Care during Survey Year Estimates of periods of in-patient hospital care, by age and sex

• • •		Cor	nplaint periods v	with care .		All hospital per	iods
Age-sex group	Total number of complaint periods (millions)	Number (millions)	As percentage of total complaint periods with care (by sex) (c)1	As percentage of total complaint periods for age-sex group (d) ²	Number (millions)	As percentage of total hospital periods (by sex)	Average number per 1,000 persons in population (g) ⁴
Both sexes							
All ages	29,47	1.39	100.0	4.7	1.65	100. 0	122
Under 15 (except newborn) Under 15. 15-24 25-44 45-64 65 and over	10.47 11.70 3.53 8.17 4.25 1.81	0.29 0.32 0.20 0.48 0.26 0.13	20.7 23.0 14.4 34.6 18.4 9.6	2.7 2.7 5.7 5.9 6.0 7.4	0.29 0.55 0.20 0.49 0.27 0.14	17.5 33.2 12.3 29.8 16.2 8.6	76 133 99 126 111
Male			·			·	
All ages	13.76	0.55	100.0	4.0	0.68	100.0	100
Under 15 (except newborn) Under 15	5.08 5.99 1.53 3.42 1.97 0.85	0.15 0.17 0.05 0.13 0.13	28. I 30. 9 9. 9 22. 9 23. 5 12. 8	3.0 2.8 3.6 3.7 6.5 8.2	0.16 0.29 0.06 0.13 0.13	22.8 42.3 8.1 18.8 19.8 11.0	80 137 55 66 108 140
Female		.					
All ages	15.71	0.84	100.0	5.4	0. 98	100.0	145
Under 15 (except newborn) Under 15	5, 39 5,71 2,01 4,76 2,28 0,96	0.13 0.15 0.15 0.36 0.13 0.06	15.9 17.9 17.3 42.1 15.1 7.6	2.5 2.6 7.3 7.5 5.6 6.6	0.13 0.26 0.15 0.37 0.13 0.07	13.8 26.8 15.2 37.5 13.7 6.9	72 130 142 185 114 129


Derived from column (c): each age group as percentage of "all ages", by sex.
 Column (b) as percentage of column (a).
 Derived from column (e): each age group as percentage of "all ages", by sex.
 Column (e) of Table 9 divided by one thousandth of column (a) of Table 8.

TABLE 10. Number of Person-Days of Hospitalization during Survey Year Estimates of person-days of hospitalization, by age and sex

	Number of person-days for all	As percentage of total		number of 000 persons	Average number	Number of person-days	Average number
Age-sex group	hospital periods (millions) (a)	person-days reported (by sex) (b)1	In population (c) ²	Reporting (d) ³	of days per hospital period (e) ⁴	for complaint periods (millions)	of days per complaint period (g) ⁵
Both sexes			-		"	,	
All ages	23.39	100, 0	1, 728	16, 916	14	21.68	0.736
Under 15 (except newborn)	2.45 4.52 2.40 6.70 5.84 3.94	10.5 19.3 10.2 28.6 25.0 16.8	644 1,099 1,169 1,716 2,417 3,737	9, 427 10, 581 12, 711 15, 333 25, 738 38, 134	8 8 12 14 22 28	2.48 2.86 2.39 6.64 5.84 3.94	0.237 0.245 0.678 0.813 1.373 2.173
Male					. •		
All ages	10.97	100.0	1, 609	19, 541	16	10.02	0. 728
Under 15 (except newborn)	1.53 2.60 0.86 2.90 2.82 1.80	13.9 23.7 7.8 26.4 25.7 16.4	786 1, 238 852 1, 499 2, 265 3, 362	10,870 11,453 17,119 24,652 24,566 34,483	10 9 16 23 21 24	1.55 1.70 0.86 2.84 2.82 1.80	0.305 0.284 0.562 0.832 1.429 2.107
Female							
All ages	12.42	100.0	1, 849	15, 123	13	11.66	0.742
Under 15 (except newborn) Under 15. 15-24 25-44 45-64 65 and over	0.92 1.93 1.54 3.80 3.02 2.14	7.4 15.5 12.4 30.6 24.3 17.2	495 955 1,475 1,929 2,578 4,122	7, 730 9, 597 11, 114 11, 905 26, 940 41, 849	7 7 10 10 23 32	0.93 1.16 1.54 3.80 3.02 2.14	0.172 0.204 0.766 0.799 1.325 2.231

Derived from column (a): each age group as percentage of "all ages", by sex.
 Column (a) of Table 10 divided by one thousandth of column (a) of Table 8.
 Column (a) of Table 10 divided by one thousandth of column (b) of Table 8.

CHART-4






TABLE 11. Duration of Hospitalization of Persons during Survey Year

A. Estimated number of persons in each age-sex group with various numbers of days in hospital

		Persons with following total number of days in hospital:												
Age-sex group	Popula- tion	0	1	2	3	4	5	6	7-13	14-20	21-27	28 or more		
	(a)	(b)	(c)	(d)	(e)	(f)	(g)	(h)	(i)	(j)	(k)	(1)		
,						(MI]]	lons)		-					
Both sexes	!		1	!		1			ا ا	· .		1		
All ages	13.54	12, 15	0, 12	0,08	0. 08	0. 08	0, 11	0. 11	0.47	0. 12	0.06	0. 14		
nder 15 (except newborn)	3. 81	3. 55	0.07	0.04	0.0	24			0.04		0. 03	1		
nder 15	4.12	3.69	0.07	0.04	0. (05	0.	08	0.14		0.05			
-44	2.05 3.90	1.86 3.47	0.0	\c		0.0		- 1	0.07		0.09			
-64	2.42	2. 19	0.0			0. 0.			0.18 0.06		0. 09 0. 10			
and over	1.05	0.95	::			•	-		. "		0.06	-		
Male														
All ages	6.82	· 6, 26	0.07	0. 63	0, (8	0. (07	0, 16	0. 05	0.	11		
ider 15 (except newborn)	1.94	1. 80	0.0	25		0.0	04		·	0.	05			
der 15	2. 10	1.87	Ŏ. C)6		0.0				ŏ.				
-24	11.01	0.96		'			• ·							
-44	1.93	1.81			0.0)5				0.	07			
-64and over	1. 25 0. 53	1. 13 0. 48			• •					0.				
200 0701 marinary	0, 33	0.40			••					0.	04			
Female											,			
All ages	6, 72	5, 90	0, 05	0. 05	0, 04	0, 05	0.07	0, 08	0, 32	0. 07	0,	09		
der 15 (except newborn)	1.87	1. 75	0,0	05	•	0.0	74		•	0.	33			
der 15	2.02	1.82	0.0)5 Î		0.0	6			0.0				
-24	1.04	0.90			0.0					0.1	08			
-64	1.97 1.17	1.65 1.06			0, 1			1		0.				
and over	0.52	0.47			••					0. 6				
					•••					0. (/T			

TABLE 11. Duration of Hospitalization of Persons during Survey Year

B. Estimated number of persons in each age-sex group with various numbers of days in hospital as percentage of total persons with various numbers of days in hospital, by sex ¹

,	D1-			Perso	ons with	following	total nu	nber of da	ays in hos	pital:	•	
Age-sex group	Popula- tion	0	1	2	3	4	5	6	7-13	14-20	21-27	28 or more
	(a)	(b)	(c)	(d)	(e)	(f)	(g)	(h)	(1)	(j)	(k)	(1)
Both sexes												
All ages	100. 6	100, 0	100.0	100. 0	100	. 0	100	0.0	100. 0		100. 0	
Under 15 (except newborn)	28. I 30. 4 15. 1 28. 8 17. 8 7. 8	29. 2 30. 3 15. 3 28. 5 18. 0 7. 8	53. 8 55. 6 23	0	26 31	. 3 17 32 12	7. 3 2. 4 2. 0	i. 5	9, 3 29, 2 15, 1 38, 1 13, 5		10, 5 16, 0 26, 5 30, 1 17, 4	
Male				!		٠.				• •		
All ages	100, 0	100. 0	100	. 0		. 100). 0			100	. 0	
Under 15 (except newborn)	28, 5 30, 8 14, 8	28. 8 29. 9 15. 3	<i>51.</i> 54.	7. 9	20	45	3. 4 5. 1			3	. 7 3. 4	
25-44 45-64 65 and over	28. 3 18. 3 7. 8	29. 0 18. 1 7. 7			20			ė.		21 25 11	.6 .7 .7	
Female				•								
All ages	100, 0	100.0	100.	.0 .		100), ()		•	100	. 0	
Under 15 (except newborn)	27. 8 30. 0 15. 5 29. 3 17. 4 7. 7	29. 6 30. 8 15. 3 28. 0 17. 9 7. 9	50. 53.	. 7 . 9	17. 35.	25 2 9	5. 2 5. 6			17 16 40 16	. 5 . 9	

^{1.} Derived from Table 11A: each age group as percentage of "all ages", by sex.

TABLE 11. Duration of Hospitalization of Persons during Survey Year

C. Estimated number of persons with various numbers of days in hospital as percentage of total persons in each age-sex group¹

				Perso	ns with f	ollowin	g total n	umber of da	ys in hos	pita <u>l</u>		
Age-sex group	Popula- tion	0	1	2	3	4	5	6	7-13	14-20	21-27	28 or more
	(a)	(b)	(c)	(b)	(e)	(f)	(g)	(h)	(i)	<u>(1)</u>	(k)	(1)
Both sexes												
		اممما			0. 6	0. 6	ا	8 0.8	3, 5	0.9	0.4	1. 1
All ages	100.0	89. 8	0. 9	0, 6	Ψ, ο	V, Q	' "•	ói mro	3.3	u, p	1 0.4	1 1.
Inder 15 (except newborn)	100.0	93. 2	1. 7	1.0	1	. 1		••	1. 2		0.9	
Inder 15	100.0	89, 6	1,6	1.1	1	, 2		1.8	3. 4		1. 3	
5-24	100.0	90.8		:			3. 2		3.5		2. 2	
5-44	100.0	88. 8	1.				3. 2 1. 9		4.6 2.6		2. 2	
5-64	100.0	90, 6		•					_, .		4.0 5.3	
5 and over	100.0	90, 2	•	•			• •		••		3, 3	
Male					•							
All ages	100, 0	91. 8	10	0. 5	1	. 1		1, 0	2. 3	0.7	1 1	1.6
Under 15 (except newborn)	100.0	92. 8	2	7			2. 2			2	2. 3	
Inder 15	100.0	89. 2	2.	7			2. 2 3. 2				. 9	
5-24	100.0	95.0									•	
5-44	100.0	93.9			2	. 6					. 5	
5-64	100.0	90.8				• •					i. 4	
55 and over	100.0	90.3				••				. 6	i. 8	
Female ·												
All ages	100.0	87. 8	0, 8	0.7	0.6	0. 7	1	0 1.1	4.7	1.1	;	1. 4
Under 15 (except newborn)	100.0	93. 6	9	Ω.			1 0			,	1. 7	
Inder 15 (except newborn)	100.0	90.0	2.	8			1. 9 3. 0		1		i. 3	
5-24	100.0	86. 7		'	5	. 5	0, 0		1		i. 7	
5 - 44	100.0	83.8				. ĭ			l). i	
5-64	100.0	90.4				• •			ļ		Š, 9	
is and over	100.0	90. 2				• •			1	7	1. 5	

^{1.} Derived from Table 11A: columns (b)-(l) as percentage of column (a).

TABLE 11. Duration of Hospitalization of Persons during Survey Year

D. Estimated number of persons in each age-sex group with various numbers of days in hospital (cumulative)

			1	Persons v	viti: follo	wing total	number o	of days in	hospital	:	
Age-sex group	Popula- tion	1 or more	2 or more	3 or more	4 or more	5 or more	6 or more	7 or more	14 or more	21 or more	28 or more
	(a)	(c)	(d)	(e)	(f)	(g)	(h)	(i)	(j)	(k)	(1)
						(Millions)					
Both sexes		ì	l .	1	I	I	1	1	ŀ	1	1
All ages	13. 54	1.38	1, 26	1, 18	1. 10	1. 02	0, 90	0.80	0, 32.	0. 20	0. 14
nder 15 (except newborn)	3. 81	0. 26	0. 19	0. 16	0. 13	0. 11	0.09	0.08	0.03	••	
nder 15	4. 12 2. 05	0.43	0.36 0.18	0.32	0. 29	0. 27 0. 14	0. 23 0. 12	0.19	0.05	::	::
5-44	3.90	0.44	0.41	0.39	0.37	0.34	0.31	0.27	0.09	0.05	
-64	2.42	0. 23	0. 21	0. 21	0. 19	0.18	0.17	0, 16	0.10	0.06	0.0
s and over	1.05	0.10	0. 10	0.10	0.09	0.08	0.08	0.08	0.00	••	,
Male	Ì								İ		
All ages	6. 82	0, 56	0, 49	0.46	0, 42	0. 38	0, 34	0.31	0. 16	0, 11	0.0
nder 15 (except newborn)	1.94	0. 14	0.11	0.09	0.07	0.06	0.05	0.05			
nder 15	2. 10	0. 23	0. 19	0.17	0.16	0.14 0.03	0.12	0.10	•••	• • •	
5 - 24	1.01	0.05	0.04	0.09	0.08	0.08	0.03	0.07	::	l ::	:
5-64	1. 25	0.11	0.11	0.10	0. 10	0.09	0.08	0.08	0.05		.
and over	0, 53	0.05	0. 05	0. 05	0.05	0.04	0.04	0.04		i	•
Female											
All ages	6, 72	0, 82	0. 77	0. 72	0, 68	0.63	0. 56	0.49	0.17	0. 09	0.0
nder 15 (except newborn)	1. 87	0. 12	0, 09	0.07	0.06		0.04	0.03			.
nder 15	2.02	0, 20	0. 17	0.15	0.13	0.12	0.11	0.09	•••	• • •	•
-24 -44	1.04 1.97	0.14	0. 13 0. 31	0.13	0. 12 0. 28	0.11 0.26	0.09 0.23	0.08	0.05	::	
} = 99	1.17	0. 11	0. 11	0. 10	0.10	0.09	0.09	0.08	0.05	::	
and over	0.52	0.05	0.05	0.05	0.04	0.04	0.04	0.04		••	

TABLE 11. Duration of Hospitalization of Persons during Survey Year

E. Estimated number of persons in each age-sex group with various numbers of days in hospital as percentage of total persons with various numbers of days in hospital, by sex (cumulative) 1

				Person	s with fo	lowing n	umber of	days in h	ospital:		
Age-sex group	Popula- tion (a)	or more	or more (d)	or more (e)	or more	5 or more (g)	6 or more (h)	or more	14 or more (j)	21 or more (k)	28 or more (1)
Both sexes											,
All ages	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100,0
Inder 15 (except newborn)	28. I 30, 4 15. 1 28. 8 17. 8 7. 8	18. 8 30. 9 13. 6 31. 6 16. 4 7. 5	15. 5 28. 5 14. 1 32. 6 17. 0 7. 8	13. 2 26. 9 14. 4 33. 1 17. 6 8. 1	11. 9 26. 3 14. 3 33. 4 17. 5 8. 4	11. 1 26. 1 13. 9 33. 8 17. 9 8. 3	10, 2 25, 2 13, 3 34, 0 18, 8 8, 6	9. 8 23. 8 13. 1 33. 4 20. 2 9. 5	10, 5 16, 0 26, 5 30, 1 17, 4	24. 8 32. 3	35.
Male		i			İ	ļ					
All ages	100.0	100.0	100.0	100.0	100.0	100.0	100,0	100.0	100.0	100.0	100,
inder 15 (except newborn)	28. 5 30. 8 14. 8 28. 3 18. 3 7. 8	25. 1 40. 4 8. 9 20. 9 20. 5 9. 3	21. 4 38. 7 8. 9 20. 8 21. 5 10. 0	19. 1 37. 2 9. 0 20. 3 22. 8 10. 7	17. 6 37. 2 8. 0 20. 3 23. 1 11. 4	16. 2 36. 9 7. 8 20. 6 23. 8 11. 0	15. 1 34. 9 7. 4 21. 7 24. 9 11. 0	14. 7 33. 4 7. 5 21. 6 25. 7 11. 7	30.4	••	
Female											
All ages	100.0	100.0	100.0	-100,0	100.0	100.0	100.0	100.0	100.0	100.0	100
Inder 15 (except newborn) S-24 5-44 5-64 5-64 5 and over	27. 8 30.0 15. 5 29. 3 17. 4 7. 7	14. 5 24. 4 16. 8 38. 9 13. 6 6. 2	11. 7 22.0 17. 4 40. 2 14. 1 6. 3	9. 4 20. 3 17. 9 41. 2 14. 2 6. 4	8. 4 19. 7 18. 2 41. 5 14. 1 6. 6	8.0 19.6 17.5 41.8 14.3 6.7	7. 2 19. 4 16. 9 41. 4 15. 1	6. 7 17. 7 16. 6 40. 9 16. 7 8. 1	29, 5 29, 7	••	

^{1.} Derived from Table 11D: each age group as percentage of "all ages", by sex.

TABLE 11. Duration of Hospitalization of Persons during Survey Year

F. Estimated number of persons with various numbers of days in hospital as percentage of total persons in each age-sex group (cumulative)

	1	<u> </u>		Damagna u	with follow	uina tatal		e dove do	. bassital		
Age-sex group	Popula- tion (a)	or more	. 2	or more (e)	or more	5 or more (g)	6 or more (h)	7 or more (i)	14 or more (j)	21 or more (k)	28 or more (1)
Both sexes	100.0	10.2	9.3	8.7	8. 1	7.5	6.7	5, 9	2. 4	1,5	1.1
Under 15 (except newborn)	100.0 100.0 100.0 100.0 100.0 100.0	6. 8 10. 4 9. 2 11. 2 9. 4 9. 8	5. 1 8. 8 8. 7 10. 5 8. 9 9. 3	4. 1 7. 7 8. 3 10. 0 8. 6 9. 1	3. 4 7. 0 7. 7 9. 4 8. 0 8. 8	2. 9 6. 4 6. 9 8. 8 7. 5 8. 0	2. 4 5. 5 5. 9 7. 9 7. 0 7. 4	2. 0 4. 6 5. 1 6. 8 6. 7 7. 2	0, 9 1, 3 2, 2 4, 0 5, 3	1.3 2.7	2. 1
Male	i00.0	8. 2	7.2	6.7	6.1	5.6	5.0	4, 6	2,3	1.6	1.2
Under 15 (except newborn)	100.0 100.0 100.0 100.0 100.0 100.0	7. 2 10. 8 5. 0 6. 1 9. 2 9. 7	5. 4 9. 1 4. 4 5. 3 8. 5 9. 3	4.5 8.1 4.1 4.8 8.4 9.2	3.8 7.4 3.3 4.4 7.7	3. 2 6. 7 2. 9 4. 1 7. 3 7. 9	2. 7 5. 7 2. 5 3. 8 6. 8 7. 0	2. 3 4. 9 2. 3 3. 5 6. 4 6. 8	3.8		
Female	100.0	19.0	11 4	10.7	10.1			70			
### All ages Under 15 (except newborn) Under 15	100.0 100.0 100.0 100.0 100.0 100.0	6. 4 10.0 13. 3 16. 2 9. 6 9. 8	11.4 4.8 8.4 12.8 15.7 9.2 9.3	3. 6 7. 2 12. 3 15. 0 8. 8 8. 9	3. I 6. 6 11. 9 14. 3 8. 2 8. 7	2. 7 6. 2 10. 7 13. 4 7. 8 8. 2	8.4 2.2 5.4 9.1 11.8 7.3 7.8	1. 7 4. 3 7. 7 10. 1 6. 9 7. 5	2.5 2.5 4.3	1.4	1.0

^{1.} Derived from Table 11D: columns (c) - (1) as percentage of column (a).

TABLE 12. Home Nursing Services during Survey Year Estimates of persons reporting home nursing care, complaint periods under home nursing care, and visits or days of home nursing care reported

Type of day or visit	Population (millions)	Persons visits	reporting or days	Complaint p	eriods under sing care	Visits or days reported			
			Percentage		Percentage		Rate per 1,000 persons		
		Number (millions) (b) ¹	of total persons	Number (millions) (d) ³	of total complaint periods (e)	Number (millions)	In population (g) ⁶	Reporting (h)	
Graduate nursing care		0.11	0.8	0. 11	0,4	0.99	73	9, 106	
Non-graduate nursing care	13. 54	0.05 0.15	- 0.4 1.1	0.05 0.15	0, 2 0, 5	1.04 n.a.	77 n.a.	21, 298 n.a.	

Top line of Table 13A.
Top line of Table 13C.
Top line of Table 14A.
Top line of Table 14A.
Top line of Table 15A.
Top line of Table 15C.
Top line of Table 15D.

TABLE 13. Persons Reporting Home Nursing Care during Survey Year A. Estimated number of persons reporting visits or days of home nursing care in each age-sex group

Age-sex group	Population (a)	Graduate nursing care (b)	Non-graduate nursing care (c)	Any nursing care (d)	. No nursing care (e)				
	(Millions)								
Both sexes	.]	-		· .]					
All ages	13,54	0, 11	0.05	0. 15	13, 39				
Jnder 15	4. 12	0.02		0.02	4. DS				
5-24	2.05				2.03				
35-44	3.90	0.04	0,03	0.06	3, 85				
5-64	2.42 1.05	0.04	{ ::	0.05	2, 39 1, 03				
Male									
All ages	6, 82	0.04	••	0.04	6, 7				
Jnder 15	2. 10	, ,	٠. ا	۱ . lr	2. 0!				
5-24	1.01	0.02	ŀ	} 0.02	1.00				
5-44	1.93			l I	1. 9				
5-64	1. 25	••	••		1. 24				
5 and over	0. 53	••	••		0.5				
·				• •					
. Female				,					
All ages	6, 72	0.07	0,04	, 0, 11	6. 61				
Under 15	2.02	••		ا ا	2.01				
5 - 24	1.04			0.02	1,03				
5-44	1.97	0,04	0.03	0.05	1.9				
5 - 64	. 1.17	. 0.04	·		- 1, 16				
5 and over	0.52			·	0,5				

n.a. - not applicable.

TABLE 13. Persons Reporting Home Nursing Care during Survey Year

B. Estimated number of persons in each age group reporting visits or days of home nursing care as percentage of total persons reporting visits or days of home nursing care, by sex¹

Age-sex group	Population (a)	Graduate nursing care (b)	Non-graduate nursing care (c)	Any nursing care (d)	No nursing care (e)
Both sexes					
All ages	100.0	100, 0	100, 0	100, 0	100, 0
Under 15	30. 4 15. 1 28. 8 17. 8 7. 8	20. 9 35. 0 33. 1	57.4	16. 7 39. 4 32. 1	30. 6 15. 2 28. 7 17. 9 7. 7
. Male				ı	
All ages	100. 0	100, 0	100. 0	100, 0	100, 0
Under 15	30.8 14.8 28.3 18.3 7.8	56.7	::	53.1	30. 8 14. 8 28. 4 18. 2 7. 7
Female					
All ages	100. 0	100, 0	100, 0	100, 0	100.0
Under 15	30, D 15, 5 29, 3 17, 4 7, 7	60.3	{ 64.7	33.4 50.1	30.3 15.6 29.0 17.5 7.6

^{1.} Derived from Table 13A: each age group as percentage of "all ages", by sex.

TABLE 13. Persons Reporting Home Nursing Care during Survey Year

C. Estimated number of persons reporting visits or days of home nursing care as percentage of total persons in each age-sex group!

Age-sex group	Population (a)	Graduate nursing care (b)	Non-graduate nursing care (c)	Any nursing care (d)	No nursing care (e)
Both sexes All ages	100, 0 100, 0 100, 0 100, 0 100, 0	0.8 0.5 1.0 1.0	0,4 0.7 	1.1 . 0.6 1.5 1.3	98. 9 99. 4 99. 2 98. 5 99. 1 97. 7
Male All ages Under 15 15-24 25-44 45-64 65 and over	100. 0 100. 0 100. 0 100. 0 100. 0	0.5 0.4	· · · · · · · · · · · · · · · · · · ·	0.6	99. 4 99. 6 99. 8 99. 8 99. 3
Female All ages	100. 0 100. 0 100. 0 100. 0 100. 0	1,1 1,4	0, 6 1.4	1.6 0.8 2.7	98. 4 99. 4 98. 7 97. 3 98. 8 97. 2

^{1.} Derived from Table 13A: columns (b) -(g) as percentage of column (a).

TABLE 14. Complaint Periods under Home Nursing Care during Survey Year

A. Estimated number of complaint periods with graduate or non-graduate home nursing care for each age-sex group

	Total	Periods with					
Age-sex group	number of complaint periods (a)	Graduate nursing care (b)	Non-graduate nursing care (c)	Any nursing care (d)	No nursing care (e)		
			(Millions)				
Both sexes			1	1	Ì		
All ages	29.47	0, 11	0, 05	0, 15	29, 32		
Under 15	11.70 3.53 8.17 4.25 1.81	0.02 0.04 0.04	0.03	0.02 0.06 0.06	11. 68 3. 52 8. 12 4. 22 1. 78		
Male							
All ages	13, 76	0, 04		0, 04	13. 72		
Under 15	5.99 1.53 3.42 1.97 0.85	0.02		0.02	\$ 5.98 1.52 3.41 1.96 0.84		
Female			II.				
All ages	15. 71	0.08	0, 05	0, 11	15, 60		
Under 15	5. 71 2. 01 4. 76 2. 28 0. 96) 0.04	{ 0.03	0.02 0.05	5.70 1.99 4.70 2.26 0.94		

TABLE 14. Complaint Periods under Home Nursing Care during Survey Year

B. Estimated number of complaint periods with graduate or non-graduate home nursing care for each age group as percentage of total complaint periods with graduate or non-graduate home nursing care, by sex 1

		Periods with					
Age-sex group	Total number of complaint periods (a)	Graduate nursing care (b)	Non-graduate nursing care (c)	Any nursing care (d)	No nursing care (0)		
Both sexes							
All ages	100.0	100, 0	100. 0	100, 0	100, 0		
Under 15	39. 7 12. 0 27. 7 14. 4 6. 2	19.6 31.4 38.0	53.4	15. 4 35. 6 37. 4	39. 8 12. 0 27. 7 14. 4 6. 1		
Male							
All ages	100, 0	100. 0	100. 0	100, 0	100. 0		
Under 15	43. 5 11. 1 24. 8 14. 3 6. 2	56. 6	 { ::	51.4	43.6 11.1 24.9 14.3 6.1		
Female							
All ages	100, 0	100, 0	100, 0	100. 0	100. 0		
Under 15	36. 3 12. 8 30. 3 14. 5 6. 1	59.6	{ 60.8	21. 1 45. 9	36. 5 12. 8 30. 2 14. 5 6. 0		

^{1.} Derived from Table 14A: each age group as percentage of "all ages", by sex.

TABLE 14. Complaint Periods under Home Nursing Care during Survey Year

C. Estimated number of complaint periods with graduate or non-graduate home nursing care as percentage of total complaint periods reported for each age-sex group

	1						
	Total	Periods with					
Age-sex group	number of complaint periods (a)	Graduate nursing care (b)	Non-graduate nursing care (c)	Any nursing care (d)	No nursing care (e)		
Both sexes							
All ages	100,0	0. 4	0.2	0.5	99, 5		
Under 15 15-24 25-44 45-64 65 and over	100.0 100.0 100.0 100.0 100.0	0.2	{ 0.3	0.2 0.7 1.0	99.8 99.5 99.3 99.3 99.3 98.4		
Male							
All ages	100, 0	0,3	••	0,3	99.7		
Under 15	100.0 100.0 100.0 100.0 100.0	0.2		0.2	99.8 99.7 99.9 99.4 98.8		
Female				·			
All ages	100,0	0, 5	0,3	` ' U. 7	99.3		
Under 15 15-24 25-44 45-64 65 and over	100, 0 100, 0 100, 0 100, 0 100, 0	}	0.6	0.3	99.8 99.3 98.9 99.2 98.0		

^{1.} Derived from Table 14A: columns (b)-(e) as percentage of column (a).

TABLE 15. Number of Visits or Days of Home Nursing Care during Survey Year

A. Estimated number of visits or days of home nursing care reported for each age-sex group

Age-sex group	Visits of graduate nursing care (a)	Days of non-graduate nursing care (b)
	(Mill	ions)
Both sexes .		
All ages	0,99	1,04
nder 15	0.08	
5 - 24	0, 26	0.36
- 64 and over	0,60	
and over ,	וי	
Male		
All ages	0, 25	••
nder 15	h !	ſ
+ 24	0.08	•
- 44	ľ ··· l	٠,
and over		•
Female]	
All ages	0, 74	0, 98
nder 15 ,		
- 24	ا ب	1 0.36
) - <u>44</u> - 64 <u></u>	} - 0.41	0.30
and over	[' · · ·	

TABLE 15. Number of Visits or Days of Home Nursing Care during Survey Year

B. Estimated number of visits or days of home nursing care reported for each age group as percentage of total visits or days of home nursing care reported, by sex 1

gradua	te	Days of non-graduate nursing care (b)
	100.0	100. D
	8.1	
		-24 5
}	60.7	
	100, 0	100.0
}	30.7	::
···· ´	::[`	••
	100,0	100, 0
1	55.3	36, 6
	gradua nursing	100,0 8.1 26.6 60.7 100,0 30.7 100,0

^{1.} Derived from Table 15A: each age group as percentage of "all ages", by sex.

TABLE 15. Number of Visits or Days of Home Nursing Care during Survey Year

C. Estimated average number of visits or days of home nursing care reported per 1,000 persons of each age-sex group 1

Age-sex group	Visits of graduate nursing care (a)	Days of non-graduate nursing care (b)
Both sexes		
All ages	73	.77
-	20	-
Under 15	l	::
25 - 44	. 68	92
65 and over	173	t ::
Male		
All ages	37	,
Under 15	15	
45 - 64	::	
OF SILL VICE AMERICAN AND AND AND AND AND AND AND AND AND A	••	••
Female		
All ages	110	145
Under 15		
15 - 24	h .::	(181
15-64	} 130	101
5 and over	••	••

^{1.} Columns (a)-(b) of Table 15A divided by one thousandth of column (a) of Table 13A.

TABLE 13. Number of Visits or Days of Home Nursing Care during Survey Year

D. Estimated average number of visits or days of home nursing care reported per 1,000 persons reporting visits or days of home nursing care for each age-sex group

Age-sex group	Visits of graduate nursing care (a) ¹	Days of non-graduate nursing care (b) ²
	- "	
Both sexes		
All ages	9, 106	21, 298
Inder 15.	3, 586	
5-24	6, 713	12, 808
15-64	16,984	{
is and over	,,	
Male		
All ages	ศ, 788	••
nder 15	l)	{
5-24	3, 540	•
5-64	ľl	••
5 and over	••	• •
Female		
All ages	10, 302	22, 631
nder 15	••	• •
- 24	··)	10.001
5-445-64	9. 457	12,801
5 and over	i′	

Column (a) of Table 15A divided by one thousandth of column (b) of Table 13A.
 Column (b) of Table 15A divided by one thousandth of column (c) of Table 13A.

TABLE 15. Number of Visits or Days of Home Nursing Care during Survey Year E. Estimated average number of visits or days of home nursing care per complaint period for each age-sex group 1

Age-sex group	Visits of graduate nursing care (a)	Days of non-graduate nursing care (b)
Both sexes		
All ages	0.034	0.035
Under 15	0.007	••
15-24	0.032	0.044
45-64 65 and over	0.099	{ ::
Male		
All ages	0.018	••
Under 15	0.007	{
45-64	::	` ::
Female		
All ages	0.047	0.062
Under 15	••	••
5-24	0.058	0.075
45-64) ""	::

^{1.} Columns (a)-(b) of Table 15A divided by column (a) of Table 14A.

TABLE'15. Number of Visits or Days of Home Nursing Care during Survey Year

F. Estimated average number of visits or days of home nursing care per complaint period with visits or days of home nursing care for each age-sex group

## Doth sexes All ages Under 15 15-24 25-44 45-64 65 and over Male All ages Under 15 15-24 25-44	8, 777 3, 628 7, 443 14, 014	20, 071 12, 966
Under 15 15-24 25-44 45-64 65 and over Male All ages Under 15 15-24 25-44	3, 628 7, 443	
15-24 25-44 45-64 65 and over Male All ages. Under 15 15-24 25-44	7, 443	12, 966
25-44. 45-64. 65 and over. Male All ages. Under 15. 15-24	· II	12, 966
## ## ## ## ## ## ## ## ## ## ## ## ##	· II	••
All ages		
Under 15		
Under 15	6, 629	••
	3, 597	•
5-64		•••
Female		
All ages	9, 864	21, 515
Jnder 15		
5-24	9, 154	12, 960

Column (a) of Table 15A divided by column (b) of Table 14A.
 Column (b) of Table 15A divided by column (c) of Table 14A.

TABLE 16. Operations during Survey Year

,		Persons	reporting ope	erations		Operations reported				
Age-sex group	Population (millions)	Number (millions)	As percentage of total persons	percentage of total persons	Number (millions)	As percentage of total number	Rate 1,000 p			
	(a)	reporting in age-sex (by sex) group	(e)	reported (by sex) (f) ³	In population (g) ⁴	Reporting (h) ⁵				
Both sexes										
All ages	13.54	0.51	100.0	3.7	0.54	100.0	40	1, 055		
Under 15	4.12 2.05 3.90 2.42 1.05	0.18 0.06 0.14 0.10	34.8 11.6 28.3 18.8	4.3 2.9 3.7 4.0	0.18 0.06 0.15 0.11	34.3 11.3 27.9 19.8	45 29 38 44	1, 040 1, 024 1, 043 1, 110		
Male						÷				
All ages	6, 82	0.24	100.0	3.5	0.25	100.0	37	1,060		
Under 15	2.10 1.01 1.93 1.25 0.53	0.09 0.05 0.05	39.5 21.8 20.3	4.4 2.7 3.8	0.10 0.05 0.05	39.5 21.8 20.3	47 28 41	1.060 1.062 1.063		
Female		}				•				
All ages	6. 72 2.02	0.28 0.08	100.0 30.8	4.0 4.2	0.29 0.09	100.0 29.8	43 42	1,051 1,019		
15-24 25-44 45-64 65 and o ver	1.04 1.97 1.17 0.52	0.09	33.9 17.6	4.7	0.10 0.06	33.3 19.4	48 47	1, 033 1, 158		

Derived from column (b): each age group as percentage of "all ages", by sex.
 Column (b) as percentage of column (a).
 Derived from column (e): each age group as percentage of "all ages", by sex.
 Column (e) divided by one thousandth of column (a).
 Column (e) divided by one thousandth of column (b).




TABLE 17. Dental Care during Survey Year Estimates of persons reporting dental care and of dental visits reported, by age and sex

and the second of the second		Persons reporting dental c			•	Dental visi	its reported		
Age-sex group	Population		As As percentage of total		Number	As percentage of total	Rate per 1,0	00 persons	
	(millions)	Number (millions)	persons reporting (by sex) (c)1	persons in age-sex group (d) ²	(millions)	number reported (by sex) (f) 3	In population (g) ⁴	Reporting (h)5	
			, -,		11		:		
Both sexes	,					:			
All ages	13, 54	2,00	100.0	14.7	4,37	100, 0	323	2, 191	
Under 15	4.12 2.05 3.90 2.42 1.05	0,56 0,43 0,70 0,26 0,04	28.3 21.5 35.3 12.8 2.1	13.7 20.9 18.0 10.6 4.0	1.10 1.09 1.55 0.56 0.08	25.2 24.9 35.4 12.7 1.9	268 530 396 230 77	1,952 2,535 2,196 2,174 1,936	
Male		,					4		
All ages	6. 82	0, 88	100,0	13, 0	1,80	100.0	264	2,040	
Under 15	2,10 1,01 1,93 1,25 0,53	0, 28 0, 18 0, 28 0, 12	31,4 20,5 31,3 13,8	13,2 18,0 14,3 9,8	0,52 0,43 0,56 0,24	29.1 23.9 31.1 13.2	250 427 290 191	1,890 2,373 2,027 1,953	
Female									
All ages	6.72	1, 11	100.0	16.5	2, 57	100,0	383	2,312	
Under 15	2.02 1.04 1.97 1.17 0.52	0, 29 0, 25 0, 43 0, 13	25.8 22.3 38.5 12.1	14.2 23.7 21.7 11.5	0.58 0.66 0.99 0.32	22.4 25.6 38.3 12.4	286 630 500 272	2,013 2,655 2,305 2,375	

Derived from column (b): each age group as percentage of "all ages", by sex.
 Column (b) as percentage of column (a).
 Derived from column (e): each age group as percentage of "all ages", by sex.
 Column (e) divided by one thousandth of column (a).
 Column (e) divided by one thousandth of column (b).




TABLE 18. Examinations for Glasses during Survey Year (other than by physicians) Estimates of persons reporting examinations for glasses and of examinations reported, by age and sex

Age-sex group	``	Persons	reporting exa for glasses	minations	Examinations reported				
	Population	Number	As percentage of total	\s percentage of total	Number	As percentage of total	Rate per 1,	000 persons	
	(millions)	•	(millions)	persons reporting (by sex) (c)	persons in age-sex group (d)2	(millions)	number reported (by sex) (f) ³	In population (g) ⁴	Reporting (h) ⁵
Both sexes	,								
All ages	13.54	0.45	100, 0	3.3	0, 54	100. 0	40	1,188	
Under 15	4.12 2.05 3.90 2.42 1.05	0.08 0.07 0.13 0.13	18.1 16.2 28.4 28.9	2.0 3.6 3.3 5.4	0.10 0.09 0.15 0.15	19.0 17.3 27.9 27.9	25 45 38 62	1,244 1,268 1,168 1,149	
Male									
All ages	. 6.82	0, 18	100.0	2, 6	0, 22	100,0	32	1,214	
Under 15	2,10 1,01	0.04	24.4	2.1	0.06	27, 5,	29	1,371	
25 - 44	1,93 1,25 0,53	0.05	30. 2	4.4	0 . 06	27, 2	48	1,094	
Female						,			
All ages	6, 72	0, 27	100.0	4, 0	0. 32	100.0	47	1, 171	
Under 15	2.02 1.04 1.97 1.17 0.52	0, 05 0, 08 0, 08	19.0 30.7 28.0	4.9 4.2 6.5	0.07 0.10 0.09	20.6 30.3 28.4	63 49 77	1,267 1,158 1,189	

Derived from column (b): each age group as percentage of "all ages", by sex.
 Column (b) as percentage of column (a).
 Derived from column (e): each age group as percentage of "all ages", by sex.
 Column (e) divided by one thousandth of column (a).
 Column (e) divided by one thousandth of column (b).

CHART - 8




TABLE 19. Miscellaneous Health Care during Survey Year Estimates of persons reporting miscellaneous health care and of treatments reported, by type of treatment and sex

	Population (millions)	Persons reporting miscellaneous health care		Treatments reported		
Type of treatment and sex		Number (millions)	As percentage of total persons (c) 1	Number (millions) (d)	Rate per 1,000 persons	
					In population (e) ²	Reporting (f) ³
•			·			
Chiropractic:						
Both sexes	13, 54	0,13	0,9	0.96	71	7,501
Male ,	6.82	0.06	0.9	0.40	58	6, 390
Female	6, 72	0.07	1.0	0.57	, 84	8, 543
Other*:			ľ		i	
Both sexes	13.54	0,09	0,6	0.86	64	9,963
Male	- 6.82	0.04	0,6	0.46	68	11,73
Female	6.72	0.05	0.7	0.40	60	8,50
Any miscellaneous health care:	,	,				
Both sexes	13, 54	. 0.21	1. 6	1. 83	135	8,653
Male	6.82	0, 10	1.5	0.86	126	8,495
Female	6.72	0, 11	1.6	0.97	144	8,797

Column (b) as percentage of column (a).
 Column (d) divided by one thousandth of column (a).
 Column (d) divided by one thousandth of column (b).

^{*} Includes treatment by chiropodists, herbalists, homoeopaths, naturopaths, osteopaths, physiotherapists, faith healers, "bone doctors", etc.

		•	
		,	
		•	
-			
	·		
			-
			•
,	•		
	•		
			•
	•		
		•	
	·		
			•
			•
	•		
	•		
	•		
•			

· ,					4	
•	-		•		•	
ı	-					
		٠.				
•		,				
	•	•		•		
	· ·					
	•					
			,			
	•					
					•	
			'			
			-			
		•	·			
					•	
	•					
				•		
	÷					
	,					
		•			•	
	•					
					٠	
		•				
		•				
	•					
		•				



DATE DUE DATE DE RETOUR

AUP 121	9804		
- M)	À 2 6 101)3	
FEB 0 1 2	716		
NLR 178			

CANADIAN SICKNESS SURVEY 1950-51 PREVIOUS PUBLICATIONS

No. 1	FAMILY EXPENDITURES FOR HEALTH SERVICES (NATIONAL ESTIMATES)	Price 25 cents
No. 2	FAMILY EXPENDITURES FOR HEALTH SERVICES BY INCOME GROUP (NATIONAL ESTIMATES)	25 cents
No. 3	FAMILY EXPENDITURES FOR HEALTH SERVICES BY EXPENDITURE GROUP (NATIONAL ESTIMATES)	25 cents
No. 4	FAMILY EXPENDITURES FOR HEALTH SERVICES (REGIONAL ESTIMATES)	25 cents
No. 5	VOLUME OF SICKNESS (NATIONAL ESTIMATES)	25 cents
No. 6	PERMANENT PHYSICAL DISABILITIES (NATIONAL ESTIMATES)	25 cents
No. 7	INCIDENCE AND PREVALENCE OF ILLNESS	25 cents