



Agriculture and
Agri-Food Canada

Agriculture et
Agroalimentaire Canada

Rural Municipality of Elton


Information Bulletin 96-11

Soils and Terrain

An introduction
to the land resource

Land Resource Unit
Brandon Research Centre



Canada 

Rural Municipality of Elton

Information Bulletin 96-11

Prepared by:

Manitoba Land Resource Unit,
Centre for Land and Biological Resources Research,
Agriculture and Agri-Food Canada.

Department of Soil Science, University of Manitoba.

Manitoba Soil Resource Section,
Soils and Crops Branch, Manitoba Agriculture.

PREFACE

This is one of a new series of information bulletins for individual rural municipalities of Manitoba. They serve to introduce the newly developed digital soil and terrain databases, and illustrate several typical derived map products for agricultural land use planning applications. The bulletins will also be available in diskette format for selected rural municipalities.

Information contained in this bulletin may be quoted and utilized with appropriate reference to the originating agencies. The authors and originating agencies assume no responsibility for the misuse, alteration, re-packaging, or re-interpretation of the information.

This information bulletin serves as an introduction to the land resource information available for the municipality. More detailed information, including copies of the primary soil and terrain maps at larger scales, may be obtained by contacting:

Manitoba Land Resource Unit
Room 360 Ellis Bldg, University of Manitoba,
Winnipeg, Manitoba R3T 2N2
Phone: 204-474-6118 FAX: 204-275-5817

CITATION

Manitoba Land Resource Unit, 1996. Soils and Terrain. An introduction to the land resource. Rural Municipality of Elton, Information Bulletin 96-11, Ellis Bldg, University of Manitoba, Winnipeg.

ACKNOWLEDGEMENTS

This project was supported under the Canada-Manitoba Agreement of Agricultural Sustainability.

The following individuals and agencies contributed significantly to the compilation, interpretation, and derivation of the information contained in this report.

Managerial and administrative support was provided by:

R.G. Eilers, Head, Manitoba Land Resource Unit, CLBRR, Agriculture and Agri-Food Canada.
G.J. Racz, Head, Dept. of Soil Science, University of Manitoba.
F. Wilson, Manager, Manitoba Land and Soil Programs, PFRA, Agriculture and Agri-Food Canada.
G.F. Mills, Manitoba Soil Survey, Soils and Crops Branch, Manitoba Agriculture.
K.S. McGill, Chief, Land Utilization and Soil Survey, Soils and Crops Branch, Manitoba Agriculture.

Technical support was provided by:

L. Fuller and G.W. Lelyk, Manitoba Land Resource Unit, CLBRR, Agriculture and Agri-Food Canada.
J. Fitzmaurice, N. Lindberg, K. Gehman, A. Waddell, and M. Fitzgerald, Dept. of Soil Science, University of Manitoba.
R. Lewis, PFRA, Agriculture and Agri-Food Canada.

Professional expertise for data conversion, correlation, and interpretation was provided by:

W.R. Fraser and W. Michalyna of the Manitoba Land Resource Unit, CLBRR, Agriculture and Agri-Food Canada.
P. Haluschak and G. Podolsky, Manitoba Soil Survey Unit, Soils and Crops Branch, Manitoba Agriculture.




Figure 3. Rural municipalities in southern Manitoba with digital soil and terrain map information.

LAND RESOURCE DATA

The soil and terrain (landscape) information were obtained as part of a larger project to provide a uniform level of land resource information for agricultural and regional planning purposes throughout Agro-Manitoba. This information was compiled and analyzed in digital form, using Geographic Information System (GIS) techniques. Three distinct layers of information were used, as shown in Figure 2.




Figure 2. Soil, Terrain, and Base Map data.

Base Layer

Digital base map information includes the municipality and Township boundaries, and major streams, roads and highways. The soil and terrain layers were added and aligned ("georeferenced") to the digital base map. Major rivers and lakes from the base layer were also used as common boundaries for the soil and terrain map layers. Water bodies larger than 25 ha in size were digitized as separate polygons.

Terrain Layer

A separate terrain layer was produced for municipalities for which only reconnaissance scale soil map coverage was available. This was compiled by aerial photo-interpretation techniques, using recent 1:50 000 scale stereo airphoto coverage. The terrain information was transferred from the photographs onto the standard RM base and digitized in the GIS. Where the soil and terrain boundaries coincided, such as along prominent escarpments and eroded stream channels, the new terrain line was used for both layers. The terrain line, delineated from modern airphoto interpretation, was considered more positionally accurate than the same boundary portrayed on the historical reconnaissance soil map. Each digital terrain polygon was assigned the following legend characteristics:

- Surface form
- Slope class
- Slope length class
- Percent wetlands
- Wetland size
- Erosional modifiers
- Extent of eroded knolls
- Polygon number

The first four legend fields are considered differentiating, that is, a change in any of these classes defines a new polygon.

Soil Layer

The most detailed soil information currently available was selected as the data source for the digital soil layer for each rural municipality.

Comprehensive detailed soil maps (1:20 000 to 1:50 000 scale) have been published for many rural municipalities. Where they were available, the individual soil map sheets were digitized and compiled as a single georeferenced layer to match the digital RM base. Map polygons have one or more soil series components, as well as slope and stoniness classes. Soil database information was produced for each polygon, to meet national standards (MacDonald and Valentine, 1992). Slope length classes were also added, based on photo-interpretation.

Older, reconnaissance scale soil maps (1:126 720 scale) represented the only available soil data source for many rural municipalities. These maps were compiled on a **soil association** basis, in which soil landscape patterns were identified with unique surficial geological deposits and textures. Each soil association consists of a range of different soils ("associates") each of which occurs in a repetitive position in the landscape. Each polygon digitized from the reconnaissance soil map was assigned the following legend characteristics:

Map symbol and modifier (overprinted symbol)
Soil Association or Complex name
Soil series and modifier codes
Polygon number

A modern soil series that best represents the soil association was identified for each soil polygon. The soil and modifier codes provide a link to additional databases of soil properties. In this way, both detailed and reconnaissance soil map polygons were related to soil drainage, surface texture, and other soil properties to produce various interpretive maps.

SOIL AND TERRAIN OVERVIEW

The Rural Municipality (RM) of Elton covers 6 Townships (approximately 58 000 ha) in south-western Manitoba. There are no major population centres in the municipality. Land use is predominantly agriculture.

Soils in the Rural Municipality of Elton have been previously mapped (at 1:126 720 scale) in the Reconnaissance Soil Survey of the Rossburn and Virden Map Sheet Areas (Ehrlich et al., 1956), and the Reconnaissance Soil Survey of the Carberry Map Sheet Area (Ehrlich et al., 1957). More detailed information for part of the RM is reported in the Soils of the Brandon Region Study Area (Michalyna et al., 1976).

Based on climatic data from Brandon Airport (Environment Canada, 1982) the mean annual temperature is 1.5°C; mean annual precipitation is 450 mm; frost-free period is 108 days; and degree days above 5°C is 1642. The seasonal moisture deficit for the period May to September is 250 to 300 mm and effective growing degree days (EGDD) above 5°C from seeding to first frost in fall is above 1500. This parameter provides an indication of heat energy available for crop growth (Agronomic Interpretations Working Group, 1995).

The municipality is located in portions of the Newdale Till Plain, the Brandon Lakes Plain and the Upper Assiniboine Delta. Elevations vary from 518 masl in the northwest part of the RM to 375 m in the southeast. Surface deposits are mainly strongly calcareous glacial till in the north half of the RM, and fluvial and lacustrine sediments ranging in texture from gravelly sands to clay in the southern portion. Relief is variable, generally less than 3 m with slopes up to 5% in the southern portion of the RM; areas of clayey and clay loam lacustrine sediments are dissected by occasional waterways with steeper slopes. The Newdale Plain in the northern portion is hummocky with local relief up to 3 m and slopes generally less than 5%. The till plain in the north eastern portion of the RM has areas of slightly higher relief (up to 5 m), some dissection and surface forms with slopes of 5 to 9%. Dissected

areas are characterized by steeper slopes and waterways. Surface forms are dominantly hummocky and undulating, with some level areas in the south central portions of the RM.

The soils in the RM are generally Orthic Black Chernozems in well drained positions, with Gleyed Black Chernozems in imperfectly drained lower slope positions. Areas of poorly drained soils with high seasonal water tables are typically classified as Rego Humic Gleysols.

Soils on the Newdale Till Plain are developed on strongly calcareous loam to clay loam glacial till derived from local bedrock (shale), limestone, and granitic sources. Most of this area is represented by the Newdale smooth phase. Newdale soils generally range in agriculture capability from class 2 to 3 with topography being the most common limiting factor. Localized areas of class 5 to 7 capability in this area are restricted to areas of severe soil erosion or wetness. The irrigation suitability of soils in the Newdale Plain varies from good to fair due to the uneven nature of the terrain and the incidence of poorly drained depressions throughout the landscape. The potential for environmental impact under irrigation is low to moderate. Risk of water erosion over the Newdale soils varies from low to severe depending upon slope gradients.

Soil associations in the Brandon Lakes Plain and the Upper Assiniboine Delta have developed on a range of fluvial and lacustrine materials described according to their dominant parent material and textural group. The Carroll soils developed on loam to clay loam lacustrine deposits are rated in agriculture capability class 1 to 2 on the better drained soils with level to gently undulating terrain, and class 3 or 4 on soils with steeper slopes or with appreciable erosion. Risk of water erosion is moderate to severe on the well drained sites depending upon topography. Imperfectly drained Carroll soils are considered class 2 to 3W (depending on the level of salinity), and poorly drained soils are placed in class 5W.

Beresford soils are similar to Carroll clay loam soils, differing mainly in the occurrence of stony glacial till within one meter of the surface. These soils are characterized by undulating to hummocky

topography with moderate to strong slopes. Agriculture capability of these soils is class 3 to 4T, depending on the slope and degree of erosion. Irrigation suitability is fair due to salinity and topography. The potential for adverse environmental impact when irrigated is low. Risk of water erosion is moderate to severe depending on the topography.

Sandy lacustrine soils occur in two areas of the RM. The Souris fine sandy loam soils in the south western portion of the RM are dominantly imperfectly drained whereas the Stockton fine sandy loam soils in the south eastern portion of the RM are mainly moderately well drained. Agriculture capability of these soils is mainly 2 to 3M due to periodic droughtiness; irrigation suitability is excellent to good and the potential for adverse environmental impact under irrigation is low to moderate. Land use is mainly cereal and forage production with some grazing. Poorly drained soils in the Stockton and Souris Associations have an agriculture capability of class 5W; land use is mainly hay production and grazing.

Harding clay soils occur extensively in the south central portion of the RM. These are mainly imperfectly drained soils developed on clayey lacustrine sediments. They have an agriculture capability of class 2W; irrigation suitability is poor. Risk of water erosion is low in level terrain but moderate to severe in steeply sloped areas adjacent to waterways. Local areas of soil salinity may occur; the Harding saline phase has an agriculture capability of class 3N. Areas of poorly drained soils are rated in class 5W for agriculture.

DERIVED AND INTERPRETIVE MAPS

A large variety of computer derived and interpretive maps can be generated, once the soil and terrain data are stored in digital format. These maps are based on selected combinations of database values and assumptions.

Derived maps show information that is given in one or more columns in the computer map legend (such as soil texture, drainage, stoniness, or slope class).

Interpretive maps portray a more complex evaluation of information presented in the legend which was combined in a unique way to arrive at an entirely new map.

Several examples of derived and interpretive maps are included in this information bulletin. The maps have all been reduced in size and generalized (simplified), in order to portray conditions for an entire rural municipality on one page. Only interpretations based on the dominant soil and terrain conditions in each polygon are shown at such reduced scales. These generalized maps provide a useful overview of conditions within a municipality, but are not intended to apply to site specific land parcels.

The digital databases may also contain more detailed information concerning significant inclusions of differing soil and slope conditions in each map polygon, particularly where they have been derived from modern detailed soil maps. This information can be portrayed at larger map scales.

Information concerning particular interpretive maps, and the primary soil and terrain map data, can be obtained by contacting the Manitoba Land Resource Unit.

Slope Map.

Slope describes the steepness of the landscape surface. The slope classes shown on this map are derived from the digital terrain layer database. Specific colours are used to indicate the most significant, limiting slope class for each terrain polygon in the RM. Additional slope classes can occur in each polygon area, but cannot be portrayed at this reduced map scale.


Table 1. Slope Classes¹

Slope Class	Area (ha)	Percent of RM
0 - 2 %	25581	44.0
2 - 5 %	19728	33.9
5 - 9 %	12115	20.8
9 - 15 %	19	0.0
15 - 30 %	650	1.1
> 30 %	66	0.1
Water	0	0.0
Unclassified	0	0.0
Total	58159	100.0

¹ Area has been assigned to the most significant limiting slope for each terrain polygon. Significant areas of lesser slope, and smaller areas of greater slope may occur in each terrain polygon.

Rural Municipality of Elton

Slope Map



Manitoba Land Resource Unit
Centre for Land and Biological Resources Research
March 1996

Surface Form Map.

Surface forms describe the overall shape of the earth's surface. The various surface forms may exhibit a regular (or irregular) pattern of convexities and concavities, and are commonly associated with characteristic ranges of slope gradients and slope lengths. They may also imply particular modes of origin. For example, scrolled and terraced surface forms are created by river and stream deposits, while undulating and hummocky surface forms are frequently associated with glacial moraines. A description of the various surface form classes are contained in a separate Soil and Terrain Classification System Manual (Manitoba Land Resource Unit, 1996).

Surface form and slope class are two key features of the digital terrain map layer. Both of these characteristics are important controlling and influencing factors to consider for sustainable land use planning and management.


Table 2. Surface Form and Slope Classes¹

Surface Form Slope Class	Area (ha)	Percent of RM
Hummocky	27634	47.5
C (2.0 to 5.0%)	16257	28.0
D (6.0 to 9.0%)	11358	19.5
E (10.0 to 15.0%)	19	0.0
Inclined	70	0.1
D (6.0 to 9.0%)	70	0.1
Level	4923	8.5
B (0.5 to 2.0%)	4923	8.5
Ridged	446	0.8
C (2.0 to 5.0%)	446	0.8
Steep	66	0.1
J (> 70.0%)	66	0.1
Undulating	22489	38.7
B (0.5 to 2.0%)	20657	35.5
C (2.0 to 5.0%)	1832	3.1
Runway	2531	4.4
C (2.0 to 5.0%)	1194	2.1
D (6.0 to 9.0%)	687	1.2
F (16.0 to 30.0%)	650	1.1
Total	58159	100.0

¹ Area has been assigned to the most significant limiting slope for each terrain polygon. Significant areas of lesser slope, and smaller areas of greater slope may occur in each terrain polygon.

Rural Municipality of Elton

Surface Form Map



Manitoba Land Resource Unit
Centre for Land and Biological Resources Research
March 1996

Generalized Soil Map.

All soil polygons on the original published reconnaissance maps were digitized to create the soil layer. In some cases, areas of overprinted symbols on the original maps were delineated as additional new soil polygons.

This generalized soil map has been reduced in size and simplified by grouping the original soil association polygons. The groups have been colour themed according to similar modes of origin, texture, and soil drainage. Soils derived from glacial till deposits (typically loam to clay loam in texture) have been assigned blue and green colours. Soils developed from glacial lake deposits are coloured yellow (sandy), orange (loam), or brown (clay). Sand and gravel deposits are coloured in pink.

The groups have been named after the dominant soil association, and the statistics for each the groups have been summarized (in bold). The original reconnaissance map symbol types and their areal extent in the municipality are shown within each group.


Table 3. Generalized Soil Association Groups

Association Group Associate	Area (ha)	Percent of RM
Carroll (CL - s)	853.5	
Bd (saline)	173	0.3
Ccl (saline)	680	1.2
Newdale (smooth - s)	350.6	
Nsp (saline)	350.6	
Newdale (smooth - m)	410.7	
Nsp (meadow)	410.7	
Harding	11241	19.3
Hc	4271	7.3
Hc (saline)	1748	3.0
Hc/T	5222	9.0

Association Group Associate	Area (ha)	Percent of RM
Carroll (CL)	7178	12.3
Bd	2065	3.5
Ccl	5031	8.7
Wd	82	0.1
Stockton (FSL)	2151	3.7
Sfsl	1542	2.7
Snfsl	609	1.0
Newdale (smooth)	33347	57.3
Nsp	33347	57.3
Marsh	526	0.9
Mh	526	0.9
Benchland Complex	287	0.5
Bcx	287	0.5
Stockton (LS)	60	0.1
Snls	60	0.1
Eroded Slopes	1402	2.4
Er	1402	2.4
Marringhurst	348	0.6
M	50	0.1
Ma	283	0.5
Ms	14	0.0
Total	58159	100.0

Rural Municipality of Elton

Generalized Soil Map



Manitoba Land Resource Unit,
Centre for Land and Biological Resources Research
March 1996

Agricultural Capability Map.

This evaluation utilizes the 7 class Canada Land Inventory system (CLI, 1965). Classes 1 to 3 represent the prime agricultural land, classes 4 and 5 represent marginal lands, and classes 6 and 7 are considered unsuitable for dryland agriculture.

This generalized interpretive map is based on the dominant modern soil type for the soil polygon, in combination with the dominant slope class identified from the terrain polygon layer. The nature of the CLI subclass limitations and the classification of subdominant components cannot be portrayed at this generalized map scale.


Table 4. Agricultural Capability¹

Class Subclass	Area (ha)	Percent of RM
1	63	0.1
2	33210	57.1
2T	15288	26.3
2TW	1863	3.2
2W	7800	13.4
2X	8260	14.2
3	21916	37.7
3	693	1.2
3I	11	0.0
3N	9951	17.1
3T	11260	19.4
4	1019	1.8
4	20	0.0
4M	999	1.7
5	1356	2.3
5	646	1.1
5M	293	0.5
5W	416	0.7
7	60	0.1
7	51	0.1
7T	9	0.0
Organic	518	0.9
Total	58140	100.0

¹ Based on **dominant** soil and slope of the respective soil and terrain maps.

Rural Municipality of Elton

Agriculture Capability Map



Manitoba Land Resource Unit
Winnipeg, Manitoba
June 2003

Irrigation Suitability Map.

Irrigation suitability is a four class rating system. Classes are **Excellent, Good, Fair, and Poor**. Irrigation ratings are based on an assessment of the most limiting combination of soil and landscape conditions. Soils in the same class have a similar relative suitability or degree of limitation for irrigation use, although the specific limiting factors may differ. These limiting factors are described by subclass symbols at detailed map scales. The irrigation rating system does not consider water availability, method of application, water quality, or economics of irrigated land use.

This generalized interpretive map is based on the dominant soil series for each soil polygon, in combination with the dominant slope class from the terrain layer database. The nature of the subclass limitations, and the classification of subdominant components is not shown at this generalized map scale.


Table 5. Irrigation Suitability¹

Class	Area (ha)	Percent of RM
Excellent	1410	2.4
Good	26144	45.0
Fair	16727	28.8
Poor	13878	23.9
Organic	0	0.0
Water	0	0.0
Unclassified	0	0.0
Total	58159	100.0

¹ Based on **dominant** soil and slope of the respective soil and terrain maps.

Rural Municipality of Elton

Irrigation Suitability Map



Manitoba Land Resource Unit
Centre for Land and Biological Resources Research
March 1996

Potential Environmental Impact Under Irrigation.

A major concern for land under irrigated crop production is the possibility that surface and/or groundwater may be impacted. The potential environmental impact assessment provides a relative rating of land into 4 classes (minimal, low, moderate and high) based on an evaluation of specific soil factors and landscape conditions that determine the impact potential.

Soil factors considered are those properties that determine water retention and movement through the soil; topographic features are those that affect runoff and redistribution of moisture in the landscape. Specifically considered are: soil texture, hydraulic conductivity, salinity, geological uniformity, depth to watertable and topography. The risk of altering surface and subsurface soil drainage regimes, soil salinity or the potential for runoff, erosion or flooding is determined by specific criteria for each property.


Use of this rating is intended to serve as a warning of potential environmental concern. It may be possible to design and/or give special consideration to soil-water-crop management practices that will mitigate any adverse impact.

This generalized interpretive map is based on the dominant soil series for each soil polygon, in combination with the dominant slope class from the terrain layer database. The nature of the subclass limitations, and the classification of subdominant components is not shown at this generalized map scale.

Table 6. Potential Environmental Impact Under Irrigation¹

Class	Area (ha)	Percent of RM
Minimal	6958	12.0
Low	36500	62.8
Moderate	12133	20.9
High	2568	4.4
Organic	0	0.0
Water	0	0.0
Unclassified	0	0.0
Total	58159	100.0

¹ Based on **dominant** soil, slope gradient, and slope length of the respective soil and terrain maps.

Rural Municipality of Elton**Potential Environmental Impact Under Irrigation**

Manitoba Land Resource Unit
Centre for Land and Biological Resources Research
March 1996

Water Erosion Risk Map.

The risk of water erosion was estimated using the universal soil loss equation (USLE) developed by Wischmeier and Smith (1965). The map shows 5 classes of soil erosion risk based on bare unprotected soil:

negligible
low
moderate
high
severe

Cropping and management practices will significantly reduce this risk depending on crop rotation program, soil type, and landscape features.


Table 7. Water Erosion Risk¹

Class	Area (ha)	Percent of RM
Negligible	1777	3.1
Low	24436	42.0
Moderate	19669	33.8
High	1836	3.2
Severe	10440	18.0
Water	0	0.0
Unclassified	0	0.0
Total	58159	100.0

¹ Based on **dominant** soil, slope gradient, and slope length of the respective soil and terrain maps.

Rural Municipality of Elton

Water Erosion Risk Map



Manitoba Land Resource Unit
Centre for Land and Biological Resources Research
March 1996

REFERENCES

Agronomic Interpretations Working Group. 1995. Land Suitability Rating System for Agricultural Crops: 1. Spring-seeded Small Grains. Edited by W.W. Pettapiece. Tech. Bull. 1995-6E. Centre for Land and Biological Resources Research, Agriculture and Agri-Food Canada, Ottawa. 90 pages, 2 maps.

Canada Land Inventory. 1965. Soil Capability Classification for Agriculture. Canada Land Inventory Report No. 2. ARDA, Dept. of Forestry, Canada, Ottawa.

Ehrlich, W.A., Pratt, L.E. and E.A. Poyser, E.A. 1956 Report of Reconnaissance Soil Survey of Rossburn and Virden Map Sheet Areas. Soils Report No. 6. Manitoba Soil Survey. Published by Manitoba Dept. of Agriculture. 121pp and 2 maps.

Ehrlich, W.A., Poyser, E.A. and Pratt, L.E. 1957. Report of the Reconnaissance Soil Survey of Carberry Map Sheet Area. Soils Report No.7. Manitoba Soil Survey. Published by Manitoba Dept. of Agriculture. 93 pp and 1 map.

Environment Canada. 1982. Canadian Climatic Normals 1951-1980. 1- Temperatures, Vol. 2; 2- Precipitation, Vol. 3; 3- Frost, Vol. 6; 4- Degree Days, Vol. 4. Atmospheric Environment, Downsview, Ontario.

Expert Committee on Soil Survey. 1987. The Canadian System of Soil Classification. Second Edition. Publ. No. 1646. Research Branch, Agriculture Canada.

MacDonald, K.B., and Valentine, K.W.G. 1992. CanSIS Manual 1 CanSIS/NSDB: A General Description. Land Resource Division, Centre for Land and Biological Resources Research, Research Branch, Agriculture Canada, Ottawa.

Manitoba Land Resource Unit. 1996. Soil and Terrain Classification System Manual. In preparation. Ellis Bldg. University of Manitoba. Winnipeg.

Michalyna, W., Podolsky, G.P. and Gardiner, Wm. 1976. Soils of the Brandon Region Study Area. Soils Report N0. 30. Manitoba Department of Municipal Affairs, Municipal Planning Branch, Province of Manitoba.

PFRA. 1964. Handbook for the Classification of Irrigated Land on the Prairie Provinces. PFRA, Regina, Saskatchewan.

Wishmeier, W.H. and Smith, D.D. 1965. Predicting Rainfall-erosion Loss from Cropland East of the Rocky Mountains. U.S. Department of Agriculture, Agriculture Handbook No. 282, U.S. Government Printing Office, Washington, D.C.

ADDENDUM

Land Use Map.

The land use classification of the RM has been interpreted from LANDSAT satellite imagery, using supervised computer classification techniques. Many individual spectral signatures were classified and grouped into the seven general land use classes shown here. Although land use changes over time, and some land use practices on individual parcels may occasionally result in similar spectral signatures, this map provides a general representation of the current land use in the RM.

The following is a brief description of the land use classes.

Annual Crop Land - land that is normally cultivated on an annual basis.

Forage - perennial forages, generally alfalfa or clover with blends of tame grasses.

Grasslands - areas of native or tame grasses, may contain scattered stands of shrubs.

Trees - lands that are primarily in tree cover.

Wetlands - areas that are wet, often with sedges, cattails, and rushes.

Water - open water - lakes, rivers streams, ponds, and lagoons.

Urban and Transportation - towns, roads, railways, quarries.

Table 10. Land Use¹

Class	Area (ha)	Percent of RM
Annual Crop Land	44229	75.5
Forage	1473	2.5
Grasslands	6942	11.9
Trees	1756	3.0
Wetlands	1457	2.5
Water	560	1.0
Urban and Transportation	2134	3.6
Total	58551	100.0

¹ Land use information (1995) and map supplied by Prairie Farm Rehabilitation Administration. Areas may vary from previous maps due to differences in analytical procedures.

