



Agriculture and  
Agri-Food Canada

Agriculture et  
Agroalimentaire Canada

**Rural Municipality of Shellmouth**


**Information Bulletin 97-27**

## **Soils and Terrain**

An introduction  
to the land resource

Land Resource Unit  
Brandon Research Centre



Canada 

## **Rural Municipality of Shellmouth**

### **Information Bulletin 97-27**

Prepared by:

Manitoba Land Resource Unit,  
Brandon Research Centre, Research Branch,  
Agriculture and Agri-Food Canada.

Department of Soil Science, University of Manitoba.

Manitoba Soil Resource Section,  
Soils and Crops Branch, Manitoba Agriculture.

Printed October, 1997

## PREFACE

This is one of a new series of information bulletins for individual rural municipalities of Manitoba. They serve to introduce the newly developed digital soil databases and illustrate several typical derived and interpretive map products for agricultural land use planning applications. The bulletins will also be available in diskette format for each rural municipality.

Information contained in this bulletin may be quoted and utilized with appropriate reference to the originating agencies. The authors and originating agencies assume no responsibility for the misuse, alteration, re-packaging, or re-interpretation of the information.

This information bulletin serves as an introduction to the land resource information available for the municipality. More detailed information, including copies of the primary soil and terrain maps at larger scales, may be obtained by contacting

Manitoba Land Resource Unit  
Room 360 Ellis Bldg, University of Manitoba,  
Winnipeg, Manitoba R3T 2N2  
Phone: 204-474-6118 FAX: 204-474-7633.

## CITATION

Manitoba Land Resource Unit, 1997. Soils and Terrain. An Introduction to the Land Resource. Rural Municipality of Shellmouth. Information Bulletin 97-27, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

## ACKNOWLEDGEMENTS

This project was financially supported in part by the Canada-Manitoba Agreement on Agricultural Sustainability, Prairie Farm Rehabilitation Administration (PFRA), and Agriculture and Agri-Food Canada.

The following individuals and agencies contributed significantly to the compilation, interpretation, and derivation of the information contained in this report.

Managerial and administrative support was provided by:


R.G. Eilers, Head, Manitoba Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.  
G.J. Racz, Head, Dept. of Soil Science, University of Manitoba.  
F. Wilson, Manager, Manitoba Land and Soil Programs, PFRA, Agriculture and Agri-Food Canada  
K.S. McGill, Chief, Land Utilization and Soil Survey, Soils and Crops Branch, Manitoba Agriculture.

Technical support was provided by:

G.W. Lelyk, Manitoba Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.  
J. Fitzmaurice, N. Lindberg, A. Waddell, M. Fitzgerald and S. Grift, Dept. of Soil Science, University of Manitoba.  
J. Griffiths, C. Aglugub, Manitoba Soil Resource Section, Soils and Crops Branch, Manitoba Agriculture.  
R. Lewis, PFRA, Agriculture and Agri-Food Canada.  
G.F. Mills, P.Ag, Winnipeg, Manitoba

Professional expertise for data conversion, correlation, and interpretation was provided by:

W.R. Fraser and W. Michalyna, Manitoba Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.  
P. Haluschak and G. Podolsky, Manitoba Soil Resource Section, Soils and Crops Branch, Manitoba Agriculture.



**Figure 1.** Rural municipalities in southern Manitoba.

## INTRODUCTION


The location of the Rural Municipality of Shellmouth is shown in Figure 1. A brief overview of the database information assembled, and general environmental conditions for the municipality are presented. A set of maps derived from the data for typical agricultural land use and planning applications are also included.

The soil map and database were compiled and registered using the computerized Geographic Information System (PAMAP GIS) facilities of the Manitoba Land Resource Unit. These databases were used in GIS to create the generalized, derived and interpretive maps and statistics contained in this report. The final maps were compiled and printed using Coreldraw.

This bulletin is available in printed or digital format. The digital bulletin is a Windows based executable file which offers addition display options, including the capability to print any portion of the bulletin.

## LAND RESOURCE DATA

The soil and terrain information presented in this bulletin was compiled as part of a larger project to provide a uniform level of land resource information for agricultural and regional planning purposes throughout Agro-Manitoba. This information was compiled and analysed in two distinct layers as shown in Figure 2.



**Figure 2.** Soil and Base Map data.

### Base Layer

Digital base map information includes the municipality and township boundaries, along with major streams, roads and highways. Major rivers and lakes from the base layer were also used as common boundaries for the soil map layer. Water bodies larger than 25 ha in size were digitized as separate polygons.

### Soil Layer

The most detailed soil information currently available was selected as the data source for the digital soil layer for each rural municipality.

Comprehensive detailed soil maps (1:20 000 to 1:50 000 scale) have been published for many rural municipalities. Where they were available, the individual soil map sheets were digitized and compiled as a single georeferenced layer to match the digital RM base. Map polygons have one or more soil series components, as well as slope and stoniness classes. Soil database information was produced for each polygon, to meet national standards (MacDonald and Valentine, 1992). Slope length classes were also added, based on photo-interpretation.

Older, reconnaissance scale soil maps (1:126 720 scale) represented the only available soil data source for many rural municipalities. These maps were compiled on a **soil association** basis, in which soil landscape patterns were identified with unique surficial geological deposits and textures. Each soil association consists of a range of different soils ("associates") each of which occurs in a repetitive position in the landscape. Modern soil series that best represent the soil association were identified for each soil polygon. The soil and modifier codes provide a link to additional databases of soil properties. In this way, both detailed and reconnaissance soil map polygons were related to soil drainage, surface texture, and other soil properties to produce various interpretive maps. Slope length classes were also added, based on photo-interpretation.

## SOIL AND TERRAIN OVERVIEW

The Rural Municipality (RM) of Shellmouth covers an area of 6 townships (approximately 58 126 hectares) of land in western Manitoba (page 3). There are no major population centres in the municipality and agriculture services generally are provided from larger towns in the surrounding region.

The climate in the municipality can be related to the weather data from Russell located just 8 km to the south. The mean annual temperature at Russell is 1.1°C and the mean annual precipitation is 484 mm (Environment Canada, 1993). The average frost-free period is 104 days and degree-days above 5°C average 1450 (Ash, 1991). The calculated seasonal moisture deficit for the period between May and September for the area is 200 to 250 mm. The estimated effective growing degree days (EGDD) above 5°C accumulated from date of seeding to the date of the first fall frost is 1200 to 1300 (Agronomic Interpretations Working Group, 1995). These parameters provide an indication of length of growing season and the moisture and heat energy available for crop growth.

Physiographically, the RM of Shellmouth is located mainly in the Newdale Plain and the St. Lazare Plain subsections of the Saskatchewan Plain (Canada-Manitoba Soil Survey, 1980). A small part of the Riding Mountain Upland occurs in the northeastern portion of the RM. Elevations range from 585 m asl in the southeast to 405 m where the Assiniboine River flows from the RM to the south. The land surface in the municipality is generally undulating to hummocky with local relief less than 3 m and slopes of 2 to 5 percent (page 9). Average relief of 3 to 8 m and slopes ranging from 5 to 9 percent occur along the eastern boundary of the RM. Greatest local relief is adjacent to two prominent glacial meltwater channels containing the Assiniboine and Shell Rivers. The depth of these valleys is 60 m with slopes commonly exceeding 30 percent along the valley walls and their tributaries. Near level terrain with local relief under 3 m and slopes less than 2 percent occurs in the southern portion of the RM and on the bottomlands within the Shell River valley.

The soil materials in this RM consist primarily of loamy textured glacial till deposits. Areas of sand and gravel, locally covered by overlays are common near the glacial meltwater channels. The deep ravines and river valleys are characterized by stream eroded glacial till and in places shale rock mantled with till, colluvium and slump debris (page 11).

Soils in the municipality have been mapped at a reconnaissance map scale of 1:126 720 and published in the soil survey reports for the Rossburn and Virden Map Sheet Areas (Ehrlich et al., 1956) and the Grandview Map Sheet Area (Ehrlich et al., 1959). According to the Canadian System of Soil Classification (Expert Committee on Soil Survey, 1987), soils in the municipality are classified as dominantly Black Chernozems (Newdale and Marringhurst Associations) and Dark Gray Chernozems (Erickson Association). Minor areas of Gray Luvisols (Waitville Association) occur along the northeast boundary of the RM. The Newdale and Erickson soils west of the Assiniboine Valley have been modified by glacial meltwaters leaving an erosion surface of coarse sediments, gravel and stones. Local areas of poorly drained soils (Gleysols) are common in depressional areas of the landscape. Regosolic soils occur on stratified stream deposits (alluvium) in the valleys and on steeply sloping areas of eroded slopes (page 11). A more detailed and complete description of the type, distribution and textural variability of soils in the municipality is provided in the published reconnaissance soil surveys (Ehrlich et al., 1956, 1959).

Surface drainage of hummocky terrain in the RM is largely local in nature. The majority of soils are well drained with minor areas of imperfect drainage on lower slopes. Surface runoff collects in poorly drained depressional areas, some of which contain shallow ponds and small lakes. Level areas of sand and gravel deposits are dominantly rapidly drained. Surface drainage of the RM flows through gullies and intermittent streams to enter the deeply cut valleys of the Shell and Assiniboine Rivers and their tributaries. The northern portion of the Assiniboine Valley contains the Lake of the Prairies formed behind the Shellmouth Dam (page 13).

Salinity is of minor importance in the RM, occurring as weakly saline areas in the south and usually in association with imperfectly drained soils on lower slopes in the landscape. Major management considerations are related to topography and coarse texture (page 15). Excessively wet soils are minor in extent and there are no significant bedrock outcrops. Although variably stony soils occur throughout the area, very stony conditions are of particular concern in the area west of the Assiniboine Valley and within the Shell Valley. Soils in these areas are modified by stream erosion and as a result, are coarse textured and in many places very stony.

Approximately one third of the land in the RM is rated as **Class 2** and one third is rated as **Class 3** for agriculture capability (page 17). About two thirds of the land area is rated **Good to Fair** for irrigation suitability (page 19). Topography and stoniness are the main limitations for agriculture capability. Well drained sandy and gravelly soils in level landscapes are rated as **Class 5** for agriculture and **Poor** for irrigation. Steeply sloping land is rated in **Class 6** for agriculture and **Poor** for irrigation. Poor drainage affects some 10.9 percent of the soils in the RM which are rated in **Class 5** for agriculture capability and **Poor** for irrigation.

A major issue currently receiving considerable attention is the sustainability of agricultural practices and their potential impact on the soil and groundwater environment. To assist in highlighting this concern to land planners and agricultural producers, an assessment of potential environmental impact (EI) under irrigation has been included in this bulletin (page 21). As shown, the majority of the RM is at **Low** risk of degradation. However, areas of deep sandy and gravelly soils and steeply sloping soils are rated as having a **High** potential for impact on the environment under irrigation. These conditions increase the risk for deep leaching of potential contaminants on the soil surface and the potential for rapid runoff from the soil surface into adjacent wetlands or water bodies. This EI map is intended to be used in association with the irrigation suitability map.

Another issue of concern to producers, soil conservationists and land use specialists is soil erosion caused by agricultural cropping

and tillage practices. To highlight areas with potential for water erosion, a risk map has been included to show where special practices should be adopted to mitigate this risk (page 23). About 27 percent of the land in the RM is at **Severe** risk of degradation and an additional 28 percent of the area has a **High** risk.. Slightly over 25 percent of the land in the RM is considered to have a **Moderate** risk of water erosion. Although agriculture is the dominant land use in the RM, many of the sandy and gravelly soils are in grass and steeply sloping soils remain in tree cover. Management practices for land in annual crop focus primarily on maintaining adequate crop residues to provide sufficient surface cover. However, adequate protection of the steeper sloping lands most at risk to water erosion may require a shift in land use away from annual cultivation to production of perennial forages and pasture or permanent tree cover.

An assessment of the status of land use in the RM of Shellmouth in 1994 was obtained through analysis of satellite imagery. It showed that nearly 42 percent of the land in the RM is in annual cropland and an additional 32 percent of the area is in grassland. Most of the wooded areas occur on steeper sloping lands. Production of perennial forages occurs on 2 percent of the area and most of the grasslands are used for hay and pasture. Natural wetlands are minor but together with the Lake of the Prairies constitute 5 percent of the RM. Various non-agricultural uses such as recreation and infrastructure for urban areas and transportation occupy 2.5 percent of the RM (page 25).

While the majority of the soils in the RM of Shellmouth have moderate to moderately severe limitations for arable agriculture, careful choice of crops and maintenance of adequate surface cover is essential for the management of sensitive lands with coarse texture or steeper slopes. This includes leaving adequate crop residues on the surface to provide sufficient trash cover during the early spring period. Implementation of minimum tillage practices and crop rotations including forage on a site by site basis will help to reduce the risk of soil degradation, maintain productivity and insure that agriculture land-use is sustainable over the long-term.

## DERIVED AND INTERPRETIVE MAPS

A large variety of computer derived maps can be generated from the digital and landscape datadases are stored in digital format. These maps are based on selected combinations of database values and assumptions.

**Derived maps** show information that is given in one or more columns in the computer map legend (such as soil drainage or slope class).

**Interpretive maps** portray more complex land evaluations based on a combination of soil and landscape information. Interpretative maps typically show land capabilities, suitabilities, or risks related to sustainability.

Several examples of derived and interpretive maps are included in this information bulletin:

### Derived Maps

Slope Classes

Generalized Soil

Drainage

Management Consideration

### Interpretative Maps

Agricultural Capabilities

Irrigation Suitability

Potential Environmental Impact

Water Erosion Risk

Land Use.

Digital databases derived from recent detailed soil inventories contain additional detailed information about significant inclusions of differing soil and slope conditions in each map polygon. This information can be portrayed at larger map scale than shown in this bulletin.

Information concerning particular interpretive maps, and the primary soil and terrain map data, can be obtained by contacting the Manitoba Soil Resource Section of Manitoba Agriculture, the local PFRA office, or the Manitoba Land Resource Unit.

The maps have all been reduced in size and generalized (simplified) in order to portray conditions for an entire rural municipality on one page. These generalized maps provide a useful overview of conditions within a municipality, but are not intended to apply to site specific land parcels. On-site evaluations are recommended for localized site specific land use suitability requirements.



**Slope Map.**


Slope describes the steepness of the landscape surface. The slope classes shown on this map are derived from the digital soil layer database. Specific colours are used to indicate the dominant slope class for each soil polygon in the RM. Additional slope classes may occur in each polygon area, but cannot be portrayed at this reduced map scale.

**Table 1. Slope Classes<sup>1</sup>**

<b>Slope Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>0 - 2 %</b>	<b>11911</b>	<b>20.5</b>
<b>2 - 5 %</b>	<b>29075</b>	<b>50.0</b>
<b>5 - 9 %</b>	<b>3384</b>	<b>5.8</b>
<b>9 - 15 %</b>	<b>1775</b>	<b>3.1</b>
<b>15 - 30 %</b>	<b>0</b>	<b>0.0</b>
<b>&gt; 30 %</b>	<b>8978</b>	<b>15.4</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3003</b>	<b>5.2</b>
<b>Total</b>	<b>58126</b>	<b>100.0</b>

<sup>1</sup> Area has been assigned to the most significant limiting slope for each terrain polygon. Significant areas of lesser slope, and smaller areas of greater slope may occur in each terrain polygon.

## Slope Map



### Generalized Soil Map.

The most recently available soil maps were digitized to produce the new digital soil map. For older reconnaissance soil maps, areas of overprinted symbols or significant differences in topography have been delineated as new polygons. All soil polygons have been digitized and translated into modern soil series equivalents.

The general soil groups provide a very simplified overview of the soil information contained in the digital soil map. The hundreds of individual soil polygons have been simplified into broad groups of soils with similar parent material origins, textures, and drainage classes. The dominant soil in each polygon determines the soil group, area, and colour for the generalized soil map. Gleysolic soils groups have poor to very poor drainage, while other mineral soil groups typically have a range of rapid, well, or imperfectly drained soils.


More detailed maps showing the dominant and subdominant soils in each polygon can also be produced at larger map scales.

**Table 2. Generalized Soil Groups<sup>1</sup>**

<b>Soil Group</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Sand and Gravel with overlays</b>	<b>1899</b>	<b>3.3</b>
<b>Variable Textured Alluvium (Gleysols)</b>	<b>812</b>	<b>1.4</b>
<b>Loamy Till (Luvisols)</b>	<b>367</b>	<b>0.6</b>
<b>Loamy Till (Dark Gray Chernozem)</b>	<b>14989</b>	<b>25.8</b>
<b>Clay over Shale Bedrock</b>	<b>23</b>	<b>0.0</b>
<b>Loamy Lacustrine</b>	<b>30</b>	<b>0.1</b>
<b>Loamy Till (Gleysols)</b>	<b>1626</b>	<b>2.8</b>
<b>Loamy Till (Black Chernozem)</b>	<b>19730</b>	<b>33.9</b>
<b>Loamy Till with water worked surfaces</b>	<b>762</b>	<b>1.3</b>
<b>Variable Textured Alluvium (Regosols)</b>	<b>1828</b>	<b>3.1</b>
<b>Sand and Gravel (Gleysols)</b>	<b>347</b>	<b>0.6</b>
<b>Eroded Slopes</b>	<b>8978</b>	<b>15.4</b>
<b>Sand and Gravel</b>	<b>3733</b>	<b>6.4</b>
<b>Water</b>	<b>3003</b>	<b>5.2</b>
<b>Total</b>	<b>58126</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil series for each soil polygon.

## Generalized Soil Map



**Soil Drainage Map.**

Drainage is described on the basis of actual moisture content in excess of field capacity, and the length of the saturation period within the plant root zone. Six drainage classes plus four land classes are shown on this map.

**Very Poor** - Water is removed from the soil so slowly that the water table remains at or on the soil surface for the greater part of the time the soil is not frozen. Excess water is present in the soil throughout most of the year.

**Poor** - Water is removed so slowly in relation to supply that the soil remains wet for a large part of the time the soil is not frozen. Excess water is available within the soil for a large part of the time.

**Imperfect** - Water is removed from the soil sufficiently slowly in relation to supply to keep the soil wet for a significant part of the growing season. Excess water moves slowly down the profile if precipitation is the major source.

**Moderately Well** - Water is removed from the soil somewhat slowly in relation to supply. Excess water is removed somewhat slowly due to low perviousness, shallow water table, lack of hydraulic gradient, or some combination of these.

**Well** - Water is removed from the soil readily but not rapidly. Excess water flows downward readily into underlying materials or laterally as subsurface flow.

**Rapid** - Water is removed from the soil rapidly in relation to supply. Excess water flows downward if underlying material is pervious. Subsurface flow may occur on steep slopes during heavy rainfall.


Drainage classification is based on the dominant soil series within each individual soil polygon.

**Table 3. Drainage Classes<sup>1</sup>**

<b>Drainage Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Very Poor</b>	<b>195</b>	<b>0.3</b>
<b>Poor</b>	<b>2590</b>	<b>4.5</b>
<b>Imperfect</b>	<b>1970</b>	<b>3.4</b>
<b>Moderately Well</b>	<b>0</b>	<b>0.0</b>
<b>Well</b>	<b>37658</b>	<b>64.8</b>
<b>Rapid</b>	<b>12710</b>	<b>21.9</b>
<b>Marsh</b>	<b>0</b>	<b>0.0</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3003</b>	<b>5.2</b>
<b>Total</b>	<b>58126</b>	<b>100.0</b>

<sup>1</sup> Area has been assigned to the dominant drainage class for each soil polygon.

## Soil Drainage Map



**Management Considerations Map.**

Management consideration maps are provided to focus on awareness of land resource characteristics important to land use. This map does not presume a specific land use. Rather it portrays the most common and wide spread attributes that apply to most soil landscapes in the province.

These maps **highlight attributes** of soil-landscapes that the land manager must consider for any intended land use.

- **Topography**
- **Wetness**
- **Coarse texture**
- **Medium texture**
- **Fine texture**
- **Organic**
- **Bedrock**

**F = Fine texture** - soil landscapes with **fine textured soils (clays and silty clays)**, and thus low infiltration and internal permeability rates. These require special considerations to mitigate surface ponding (water logging), runoff, and trafficability. Timing and type of tillage practices used may be restricted.

**C = Coarse texture** - soil landscapes with **coarse to very coarse textured soils (loamy sands, sands and gravels)** have a high permeability throughout the profile, and require special management practices related to application of agricultural chemicals, animal wastes, and municipal effluent to protect and sustain the long term quality of the soil and water resources. The risk of soil erosion can be minimized through the use of shelterbelts and maintenance of crop residues.

**M = Medium texture** - soil landscapes with medium to moderately fine textures (**loams to clay loams**), and good water and nutrient retention properties. Good management and cropping practices are required to minimize leaching and the risk of erosion.

**T = Topography** - soil landscapes with **slopes greater than 5 %** are steep enough to require special management practices to minimize the risk of erosion.

**W = Wetness** - soil landscapes that have **poorly drained soils and/or >50 % wetlands** (due to seasonal and annual flooding, surface ponding, permanent water bodies (sloughs), and/or high water tables), require special management practices to mitigate adverse impact on water quality, protect subsurface aquifers, and sustain crop production during periods of high risk of water logging.

**O = Organic** - soil landscapes with organic soils, requiring special management considerations of drainage, tillage, and cropping to sustain productivity and minimize subsidence and erosion.

**R = Bedrock** - soil landscapes that have **shallow depth to bedrock (< 50 cm) and/or exposed bedrock** which may prevent the use of some or all tillage practices as well as the range of potential crops. They require special cropping and management practices to sustain agricultural production.


**Table 5. Management Considerations<sup>1</sup>**

<b>Land Resource Characteristics</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Fine Texture</b>	<b>724</b>	<b>1.2</b>
<b>Fine Texture and Wetness</b>	<b>17</b>	<b>0.0</b>
<b>Fine Texture and Topography</b>	<b>0</b>	<b>0.0</b>
<b>Medium Texture</b>	<b>33793</b>	<b>58.1</b>
<b>Coarse Texture</b>	<b>3733</b>	<b>6.4</b>
<b>Coarse Texture and Wetness</b>	<b>347</b>	<b>0.6</b>
<b>Coarse Texture and Topography</b>	<b>0</b>	<b>0.0</b>
<b>Topography</b>	<b>14066</b>	<b>24.2</b>
<b>Topography and Bedrock</b>	<b>0</b>	<b>0.0</b>
<b>Wetness</b>	<b>2421</b>	<b>4.2</b>
<b>Wetness and Topography</b>	<b>0</b>	<b>0.0</b>
<b>Bedrock</b>	<b>0</b>	<b>0.0</b>
<b>Organic</b>	<b>0</b>	<b>0.0</b>
<b>Marsh</b>	<b>0</b>	<b>0.0</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3003</b>	<b>5.2</b>
<b>Total</b>	<b>58126</b>	<b>100.0</b>

<sup>1</sup> Based on **dominant** soil series for each soil polygon.



## Management Considerations Map



**Agricultural Capability Map.**

This evaluation utilizes the 7 class Canada Land Inventory system (CLI, 1965). Classes 1 to 3 represent the prime agricultural land, class 4 land is marginal for sustained cultivation, class 5 land is capable of perennial forages and improvement is feasible, class 6 land is capable of producing native forages and pasture but improvement is not feasible, and class 7 land is considered unsuitable for dryland agriculture. Subclass modifiers include structure and/or permeability (D), erosion (E), inundation (I), moisture limitation (M), salinity (N), stoniness (P), consolidated bedrock (R), topography (T), excess water (W) and cumulative minor adverse characteristics (X).


This generalized interpretive map is based on the dominant soil series and phases for each soil polygon. The CLI subclass limitations cannot be portrayed at this generalized map scale.

**Table 6. Agricultural Capability<sup>1</sup>**

<b>Class Subclass</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>2</b>	<b>19199</b>	<b>33.0</b>
2P	2247	3.9
2T	16356	28.1
2TP	304	0.5
2W	39	0.1
2X	253	0.4
<b>3</b>	<b>18604</b>	<b>32.0</b>
3I	1832	3.2
3M	1633	2.8
3MP	1048	1.8
3N	103	0.2
3P	10659	18.3
3T	3328	5.7
<b>4</b>	<b>1799</b>	<b>3.1</b>
4R	24	0.0
4T	1775	3.1
<b>5</b>	<b>6306</b>	<b>10.9</b>
5M	3732	6.4
5W	1938	3.3
5WI	635	1.1
<b>6</b>	<b>9194</b>	<b>15.8</b>
6T	9000	15.5
6W	28	0.0
6WI	166	0.3
<b>Water</b>	<b>3011</b>	<b>5.2</b>
<b>Total</b>	<b>58112</b>	<b>100.0</b>

<sup>1</sup> Based on **dominant** soil, slope gradient, and slope length of each soil polygon.

## Agriculture Capability Map



**Irrigation Suitability Map.**

Irrigation ratings are based on an assessment of the most limiting combination of soil and landscape conditions. Soils in the same class have a similar relative suitability or degree of limitation for irrigation use, although the specific limiting factors may differ. These limiting factors are described by subclass symbols at detailed map scales. The irrigation rating system does not consider water availability, method of application, water quality, or economics of irrigated land use.

Irrigation suitability is a four class rating system. Areas with no or slight soil and or landscape limitation are rated **Excellent** to **Good** and can be considered irrigable. Areas with moderate soil and/or landscape limitations are rated as **Fair** and considered marginal for irrigation providing adequate management exists so that the soil and adjacent areas are not adversely affected by water application. Soil and landscape areas rated as **Poor** have severe limitations for irrigation.


This generalized interpretive map is based on the dominant soil series for each soil polygon, in combination with the dominant slope class. The nature of the subclass limitations and the classification of subdominant components is not shown at this generalized map scale.

**Table 7. Irrigation Suitability<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Excellent</b>	<b>0</b>	<b>0.0</b>
<b>Good</b>	<b>32476</b>	<b>55.9</b>
<b>Fair</b>	<b>7129</b>	<b>12.3</b>
<b>Poor</b>	<b>15518</b>	<b>26.7</b>
<b>Organic</b>	<b>0</b>	<b>0.0</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3003</b>	<b>5.2</b>
<b>Total</b>	<b>58126</b>	<b>100.0</b>

<sup>1</sup> Based on **dominant** soil, slope gradient, and slope length of each soil polygon.

## Irrigation Suitability Map



**Potential Environmental Impact Under Irrigation Map.**

A major concern for land under irrigated crop production is the possibility that surface and/or ground water may be impacted. The potential environmental impact assessment provides a relative rating of land into 4 classes (minimal, low, moderate and high) based on an evaluation of specific soil factors and landscape conditions that determine the impact potential.

Soil factors considered are those properties that determine water retention and movement through the soil; topographic features are those that affect runoff and redistribution of moisture in the landscape. Several factors are specifically considered: soil texture, hydraulic conductivity, salinity, geological uniformity, depth to water table and topography. The risk of altering surface and subsurface soil drainage regimes, soil salinity, potential for runoff, erosion and flooding is determined by specific criteria for each property.

Use of this rating is intended to serve as a warning of potential environmental concern. It may be possible to design and/or give special consideration to soil-water-crop management practices that will mitigate any adverse impact.


This generalized interpretive map is based on the dominant soil series and slope class for each soil polygon. The nature of the subclass limitations, and the classification of subdominant components is not shown at this generalized map scale.

**Table 8. Potential Environmental Impact Under Irrigation<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Minimal</b>	<b>0</b>	<b>0.0</b>
<b>Low</b>	<b>33304</b>	<b>57.3</b>
<b>Moderate</b>	<b>6069</b>	<b>10.4</b>
<b>High</b>	<b>15750</b>	<b>27.1</b>
<b>Organic</b>	<b>0</b>	<b>0.0</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3003</b>	<b>5.2</b>
<b>Total</b>	<b>58126</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil, slope gradient, and slope length of each soil polygon.

## Potential Environmental Impact Under Irrigation





**Water Erosion Risk Map.**

The risk of water erosion was estimated using the universal soil loss equation (USLE) developed by Wischmeier and Smith (1965). The map shows 5 classes of soil erosion risk based on bare unprotected soil:

**negligible**  
**low**  
**moderate**  
**high**  
**severe.**

Cropping and residue management practices will significantly reduce this risk depending on crop rotation program, soil type, and landscape features.


**Table 9. Water Erosion Risk<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Negligible</b>	<b>6552</b>	<b>11.3</b>
<b>Low</b>	<b>1798</b>	<b>3.1</b>
<b>Moderate</b>	<b>14884</b>	<b>25.6</b>
<b>High</b>	<b>16161</b>	<b>27.8</b>
<b>Severe</b>	<b>15728</b>	<b>27.1</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3003</b>	<b>5.2</b>
<b>Total</b>	<b>58126</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil, slope gradient, and slope length of each soil polygon.



## Water Erosion Risk Map



**Land Use Map.**

The land use classification of the RM has been interpreted from LANDSAT satellite imagery, using supervised computer classification techniques. Many individual spectral signatures were classified and grouped into the seven general land use classes shown here. Although land use changes over time, and some land use practices on individual parcels may occasionally result in similar spectral signatures, this map provides a general representation of the current land use in the RM.

The following is a brief description of the land use classes:

**Annual Crop Land** - land that is normally cultivated on an annual basis.

**Forage** - perennial forages, generally alfalfa or clover with blends of tame grasses.

**Grasslands** - areas of native or tame grasses, may contain scattered stands of shrubs.

**Trees** - lands that are primarily in tree cover.

**Wetlands** - areas that are wet, often with sedges, cattails, and rushes.


**Water** - open water - lakes, rivers streams, ponds, and lagoons.

**Urban and Transportation** - towns, roads, railways, quarries.

**Table 10. Land Use<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Annual Crop Land</b>	<b>24522</b>	<b>41.8</b>
<b>Forage</b>	<b>1274</b>	<b>2.2</b>
<b>Grasslands</b>	<b>19014</b>	<b>32.4</b>
<b>Trees</b>	<b>9054</b>	<b>15.4</b>
<b>Wetland</b>	<b>331</b>	<b>0.6</b>
<b>Water</b>	<b>2952</b>	<b>5.0</b>
<b>Urban and Transportation</b>	<b>1488</b>	<b>2.5</b>
<b>Total</b>	<b>58635</b>	<b>100.0</b>

<sup>1</sup> Land use information (1994) and map supplied by Prairie Farm Rehabilitation Administration. Areas may vary from previous maps due to differences in analytical procedures.



**REFERENCES**

Agronomic Interpretations Working Group. 1995. Land Suitability Rating System for Agricultural Crops: 1. Spring-seeded Small Grains. Edited by W.W. Pettapeice. Tech Bull. 1995-6E. Centre for Land and Biological Resources Research, Agriculture and AgriFood Canada, Ottawa. 90 pages, 2 maps.

Canada Land Inventory. 1965. Soil Capability Classification for Agriculture. Canada Land Inventory Report No. 2. ARDA, Dept. of Forestry, Canada, Ottawa.

Canada-Manitoba Soil Survey. 1980. Physiographic Regions of Manitoba. Ellis Bldg., University of Manitoba, Winnipeg. Revised. Unpublished Report.

Canada-Manitoba Soil Survey. 1979. Ecological Regions and Subregions in Manitoba. Ellis Bldg., University of Manitoba, Winnipeg. Revised. Unpublished Report.

Ehrlich, W.A., Pratt, L.E. and Poyser, E.A. 1956 Report of Reconnaissance Soil Survey of Rossburn and Virden Map Sheet Areas. Soils Report No. 6. Manitoba Soil Survey. Published by Manitoba Dept. of Agriculture. 121pp and 2 maps.

Ehrlich, W.A., Pratt, L.E. and Leclaire, F.P. 1959 Report of Reconnaissance Soil Survey of Grandview Map Sheet Area. Soils Report No. 9. Manitoba Soil Survey. Published by Manitoba Dept. of Agriculture. 121pp and 2 maps.

Eilers R.G. 1983. Soils of the Roblin Area. Report No. D47. Canada-Manitoba Soil Survey. Winnipeg.

Environment Canada. 1982. Canadian Climatic Normals 1951-1980. Frost, Vol. 6. Atmospheric Environment, Downsview, Ontario.

Environment Canada. 1993. Canadian Climatic Normals 1961-1990. Prairie Provinces. Atmospheric Environment, Downsview, Ontario.

Expert Committee on Soil Survey. 1987. The Canadian System of Soil Classification. Second Edition. Publ. No. 1646. Research Branch, Agriculture Canada.

Irrigation Suitability Classification Working Group. 1987. An Irrigation Suitability Classification System for the Canadian Prairies. LRRC contribution no. 87-83, Land Resource Research Centre, Research Branch, Agriculture Canada, Ottawa

MacDonald, K.B., and Valentine, K.W.G. 1992. CanSIS Manual 1 CanSIS/NSDB: A General Description. Land Resource Division, Centre for Land and Biological Resources Research, Research Branch, Agriculture Canada, Ottawa.

Manitoba Land Resource Unit. 1997. Soil and Terrain Classification System Manual. In preparation. Ellis Bldg. University of Manitoba. Winnipeg.

Podolsky, G. 1985. Soils of the Binscarth and Russell Areas. Report No. D54. Canada-Manitoba Soil Survey. Winnipeg.

Wischmeier, W.H. and Smith, D.D. 1965. Predicting Rainfall-erosion Loss from Cropland East of the Rocky Mountains. U.S. Department of Agriculture, Agriculture Handbook No. 282, U.S. Government Printing Office, Washington, D.C.