



Agriculture and  
Agri-Food Canada

Agriculture et  
Agroalimentaire Canada

**Rural Municipality of Odanah**


**Information Bulletin 98-17**

## **Soils and Terrain**

An introduction  
to the land resource

Land Resource Unit  
Brandon Research Centre



Canada 

## **Rural Municipality of Odanah**

### **Information Bulletin 98-17**

Prepared by:

Land Resource Unit,  
Brandon Research Centre, Research Branch,  
Agriculture and Agri-Food Canada.

Department of Soil Science, University of Manitoba.

Manitoba Soil Resource Section,  
Soils and Crops Branch, Manitoba Agriculture.

Printed December, 1998

## PREFACE

This is one of a new series of information bulletins for individual rural municipalities of Manitoba. They serve to introduce the newly developed digital soil databases and illustrate several typical derived and interpretive map products for agricultural land use planning applications. The bulletins will also be available in diskette format for each rural municipality.

Information contained in this bulletin may be quoted and utilized with appropriate reference to the originating agencies. The authors and originating agencies assume no responsibility for the misuse, alteration, re-packaging, or re-interpretation of the information.

This information bulletin serves as an introduction to the land resource information available for the municipality. More detailed information, including copies of the primary soil and terrain maps at larger scales, may be obtained by contacting

Land Resource Unit  
Room 360 Ellis Bldg, University of Manitoba  
Winnipeg, Manitoba R3T 2N2  
Phone: 204-474-6118 FAX: 204-474-7633

## CITATION

Land Resource Unit, 1998. Soils and Terrain. An Introduction to the Land Resource. Rural Municipality of Odanah. Information Bulletin 98-17, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada

## ACKNOWLEDGMENTS

Continuing support for this project has been provided by Brandon Research Centre and PFRA Manitoba. The project was initiated by the Land Resource Unit under the Canada-Manitoba Agreement of Agricultural Sustainability.

The following individuals and agencies contributed significantly to the compilation, interpretation, and derivation of the information contained in this report.

Managerial and administrative support was provided by:

R.G. Eilers, Head, Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

G.J. Racz, Head, Department of Soil Science, University of Manitoba.

Technical support was provided by:

G.W. Lelyk, and P. Cyr, Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

J. Fitzmaurice, N. Lindberg, and A. Waddell, Department of Soil Science, University of Manitoba.

G.F. Mills, P. Ag, Winnipeg, Manitoba


J. Griffiths, and C. Aglugub, Soil Resource Section, Soils and Crops Branch, Manitoba Agriculture.

R. Lewis, PFRA, Agriculture and Agri-Food Canada.

Professional expertise for data conversion, correlation, and interpretation was provided by:

W.R. Fraser and R.G. Eilers, Land Resource Unit, Brandon Research Centre, Research Branch, Agriculture and Agri-Food Canada.

P. Haluschak and G. Podolsky, Soil Resource Section, Soils and Crops Branch, Manitoba Agriculture.



**Figure 1.** Rural municipalities of southern Manitoba.

The location of the Rural Municipality of Odanah is shown in Figure 1. A brief overview of the database information, and general environmental conditions for the municipality are presented. A set of maps derived from the data for typical agricultural land use and planning applications are also included.

The soil map and database were compiled and registered using the Geographic Information System (PAMAP GIS) facilities of the Land Resource Unit. These databases were used in the GIS to create the generalized, derived and interpretive maps and statistics in this report. The final maps were compiled and printed using Coreldraw.

This bulletin is available in printed or digital format. The digital bulletin is a Windows based executable file which offers additional display options, including the capability to print any portion of the bulletin.

## LAND RESOURCE DATA

The soil and terrain information presented in this bulletin was compiled as part of a larger project to provide a uniform level of land resource information for agricultural and regional planning purposes throughout Agro-Manitoba. This information was compiled and analysed in two distinct layers as shown in Figure 2.

### Base Layer

Digital base map information includes the municipality and township boundaries, along with major streams, roads and highways. Major rivers and lakes from the base layer were also used as common boundaries for the soil map layer. Water bodies larger than 25 ha in size were digitized as separate polygons.

### Terrain Layer

A separate terrain layer was produced for municipalities for which only reconnaissance scale soil map coverage was available. This was compiled by aerial photo-interpretation techniques, using recent 1:50 000 scale stereo airphoto coverage. The terrain information was transferred from the photographs onto the standard RM base and digitized in the GIS. Where the soil and terrain boundaries coincided, such as along prominent escarpments and eroded stream channels, the new terrain line was used for both layers. The terrain line, delineated from modern airphoto interpretation, was considered more positionally accurate than the same boundary portrayed on the historical reconnaissance soil map. Each digital terrain polygon was assigned the following legend characteristics:

Surface form	Wetland size
Slope	Erosional modifiers
Slope length	Extent of eroded knolls
Percent wetlands	

The four legend characteristics on the left are considered

differentiating, that is, a change in any of these classes defines a new polygon.

### Soil Layer

Older, reconnaissance scale soil maps (1:126 720 scale) represented the only available soil data source for many rural municipalities. These maps were compiled on a **soil association** basis, in which soil landscape patterns were identified with unique surficial geological deposits and textures. Each soil association consists of a range of different soils ("associates") each of which occurs in a repetitive position in the landscape. Modern soil series that best represent the soil association were identified for each soil polygon. The soil and modifier codes provide a link to additional databases of soil properties. In this way, reconnaissance soil map polygons were related to soil drainage, surface texture, and other soil properties to produce various interpretive maps.

## SOIL AND TERRAIN OVERVIEW

The Rural Municipality (RM) of Odanah covers an area of 4 townships (approximately 38 673 hectares) of land in western Manitoba (page 3). The population of the municipality is predominantly rural farm-based with small concentrations of people in the villages of Moorepark and Cordova. Most agricultural services are provided from larger centres outside the municipality such as Minnedosa or Neepawa.

The climate in the municipality can be related to weather data from Minnedosa located immediately north of the RM. The mean annual temperature at Minnedosa is 0.8°C and mean annual precipitation is 467 mm (Environment Canada, 1993). Average frost-free period is 91 days and degree-days above 5°C average 1441 (Ash, 1991). The calculated seasonal moisture deficit for the period between May and September for the area ranges from slightly less than 200 mm in the north to slightly in excess of 250 mm in the south. The estimated effective growing degree days (EGDD) above 5°C accumulated from date of seeding to the date of the first fall frost range from slightly less than 1300 in the north to about 1400 in the south (Agronomic Interpretations Working Group, 1995). These parameters provide an indication of length of growing season and moisture and energy available for crop growth. The growing conditions are generally adequate for production of a wide range of small grain and cool-season oilseed crops adapted to western Canada.

Physiographically, the RM of Odanah is located entirely in the Newdale Plain subsection of the Saskatchewan Plain (Canada-Manitoba Soil Survey, 1980). Elevations decrease gradually from a high of 549 metres above sea level (m asl) in the northwest corner of the municipality, to 413 m asl in the southeast corner, 502 m asl in the southwest corner and 480 m asl in the northeast corner. The land surface thus slopes gently in a southeasterly direction at a rate of about 5.5 metres per kilometer. A large glacial meltwater channel containing the Little Saskatchewan River cuts across the

northwest corner of the municipality. Topography is dominantly very gently sloping to irregular, gently undulating with local relief in the form of low knolls and slight depressions and slopes ranging from 2 to 9 percent. Greatest relief occurs along the glacial meltwater channel where it crosses the northwest corner of the area. This channel lies some 30 metres below the land surface and the slope of the valley walls and its tributary creeks and gullies commonly exceed 30 percent (page 9).

The soil materials in this municipality consist mainly of loamy textured glacial till (morainal) deposits. Stratified alluvial deposits occur within the Little Saskatchewan River Valley and local areas of water worked till and variably textured slump deposits are associated with the valley side walls (page 11).

Soils in the municipality have been mapped at a reconnaissance map scale of 1:126 720 and published in the soil survey report for the Carberry Map Sheet Area (Ehrlich et al, 1957). A detailed 1:20 000 scale soil map is published for a small area around Minnedosa in Soils of the Hallboro-Minnedosa Area (Podolsky, 1984). According to the Canadian System of Soil Classification (Expert Committee on Soil Survey, 1987), soils in the municipality are classified mainly as Black Chernozems of the Newdale Association. The smooth phase of the Newdale soils is dominant and is characterized by low local relief and gentle slopes. Local areas of poorly drained soils (Gleysols) are common in depressional areas of the landscape (page 11). A more detailed and complete description of the type, distribution and textural variability of soils in the municipality is provided in the published soil surveys.

Surface drainage of the municipality is provided by a sparse network of small creeks and intermittent streams flowing southeasterly across the municipality. The soils are dominantly well drained although the innumerable shallow depressions throughout the landscape are characterized by poorly drained soils usually surrounded by a narrow band of imperfectly drained soil. Surface runoff collects for short periods in these depressions which

may contain intermittent shallow ponds during spring snowmelt and periods of high rainfall (page 13).

Management considerations are related mainly to topography and the associated drainage pattern (page 15). The irregular topography of the smooth phase soils results in many fields being broken by numerous sloughs and shallow potholes. The percentage of arable land is lower in such landscapes as many of the poorly drained soils can not be cultivated because of inundation during the spring season and excessive salinity in some sites. However, salinity is a minor concern in the municipality, occurring mainly in poorly drained depressional areas and in lower slopes around the depressions. Crests and upper slopes of low knolls and ridges in this area are often calcareous and may be weakly saline, particularly in the subsoil. Although stony soils occur throughout the municipality, the condition is variable, being more pronounced on crests and upper slopes of low ridges and knolls.

Approximately 43 percent of the land in the RM is rated as **Class 2** for agriculture capability (page 17) and **Good** for irrigation suitability (page 19). About 56 percent of the land is rated as **Class 3** for agriculture capability and 57 percent of the land area is rated **Fair** for irrigation suitability. Topographic pattern is the main limitation for agriculture capability. Poorly drained soils are rated in **Class 5** for agriculture capability and **Poor** for irrigation. Steeply sloping land is rated in **Class 6** and **7** for agriculture and **Poor** for irrigation.

A major issue currently receiving considerable attention is the sustainability of agricultural practices and their potential impact on the soil and groundwater environment. To assist in highlighting this concern to land planners and agricultural producers, an assessment of potential environmental impact (EI) under irrigation has been included in this bulletin (page 21). As shown, nearly 43 percent of the land in the RM is at a **Low** risk of degradation. About 57 percent of the land area with slightly higher local relief and higher slopes is rated as having a **Moderate** risk of degradation due to

greater risk of runoff to streams and water bodies in the landscape. This EI map is intended to be used in association with the irrigation suitability map.

Another issue of concern to producers, soil conservationists and land use specialists is soil erosion caused by agricultural cropping and tillage practices. To highlight areas with potential for water erosion, a risk map has been included to show where special practices should be adopted to mitigate this risk (page 23). About 0.8 percent of the land in the RM is at **Severe** risk of degradation and an additional 36 percent of the area has a **High** risk.. Nearly 58 percent of the land in the RM is considered to have a **Moderate** risk of water erosion. Management practices for these soils most at risk from water erosion focus primarily on maintaining adequate crop residues to provide sufficient surface cover during the early spring period. These practices include minimum tillage and suitable crop rotations.

An assessment of the status of land use in the RM of Odanah in 1993 was obtained through analysis of satellite imagery. It showed that 62 percent of the land in the RM is in annual cropland and an additional 17.8 percent of the area is in grassland. Wooded areas occupy 3.9 percent of the area, occurring mainly around the periphery of wet depressions. Production of perennial forages occurs on only 0.1 percent of the area. Natural wetlands constitute 11.7 percent of the area and water bodies cover 1.1 percent of the RM. Various non-agricultural uses such as recreation and infrastructure for urban areas and transportation occupy 3.1 percent of the RM (page 25).

While the majority of the soils in the RM of Odanah have moderate to moderately severe limitations for arable agriculture, careful choice of crops and maintenance of adequate surface cover is essential for the management of sensitive lands with steeper slopes. Implementation of minimum tillage practices and crop rotations including forage on a site by site basis will help to reduce the risk of soil degradation, maintain productivity and insure that agriculture land-use is sustainable over the long-term.

## DERIVED AND INTERPRETIVE MAPS

A large variety of computer derived and interpretive maps can be generated from the digital soil and landscape databases. These maps are based on selected combinations of database values and assumptions.

**Derived maps** show information that is given in one or more columns in the computer map legend (such as soil drainage or slope class).

**Interpretive maps** portray more complex land evaluations based on a combination of soil and landscape information. Interpretations are based on soil and landscape conditions in each polygon. Interpretative maps typically show land capabilities, suitabilities, or risks related to sustainability.

Several examples of derived and interpretive maps are included in this information bulletin:

### Derived Maps

Slope

Generalized Soil

Drainage

Management Considerations

### Interpretative Maps

Agricultural Capability

Irrigation Suitability

Potential Environmental Impact

Water Erosion Risk

Land Use

Digital databases derived from recent detailed soil inventories contain additional detailed information about significant inclusions of differing soil and slope conditions in each map polygon. This information can be portrayed at larger map scale than shown in this bulletin.

Information concerning particular interpretive maps, and the primary soil and terrain map data, can be obtained by contacting the Manitoba Soil Resource Section of Manitoba Agriculture, the local PFRA office, or the Land Resource Unit.

The maps have all been reduced in size and generalized (simplified) in order to portray conditions for an entire rural municipality on one page. These generalized maps provide a useful overview of conditions within a municipality, but are not intended to apply to site specific land parcels. On-site evaluations are recommended for localized site specific land use suitability requirements.



**Slope Map.**


Slope describes the steepness of the landscape surface. The slope classes shown on this map are derived from the digital soil and terrain layer database. Specific colours are used to indicate the dominant slope class for each polygon in the RM. Additional slope classes may occur in each polygon area, but cannot be portrayed at this reduced map scale.

**Table 1. Slope Classes<sup>1</sup>**

<b>Slope Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>0 - 2 %</b>	<b>0</b>	<b>0.0</b>
<b>2 - 5 %</b>	<b>16478</b>	<b>42.6</b>
<b>5 - 9 %</b>	<b>21935</b>	<b>56.7</b>
<b>9 - 15 %</b>	<b>163</b>	<b>0.4</b>
<b>15 - 30 %</b>	<b>0</b>	<b>0.0</b>
<b>&gt; 30 %</b>	<b>93</b>	<b>0.2</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3</b>	<b>0.0</b>
<b>Total</b>	<b>38673</b>	<b>100.0</b>

<sup>1</sup> Area has been assigned to the dominant slope in each soil polygon.

## Slope Map



### Generalized Soil Map.

The most recently available soil maps were digitized to produce the new digital soil map. For older reconnaissance soil maps, areas of overprinted symbols or significant differences in topography have been delineated as new polygons. All soil polygons have been digitized and translated into modern soil series equivalents.

The general soil groups provide a very simplified overview of the soil information contained in the digital soil map. The hundreds of individual soil polygons have been simplified into broad groups of soils with similar parent material origins, textures, and drainage classes. The dominant soil in each polygon determines the soil group, area, and colour for the generalized soil map. Gleysolic soils groups have poor to very poor drainage, while other mineral soil groups typically have a range of rapid, well, or imperfectly drained soils.


More detailed maps showing the dominant and subdominant soils in each polygon can also be produced at larger map scales.

**Table 2. Generalized Soil Groups<sup>1</sup>**

<b>Soil Groups</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Loamy Till (Gleysols)</b>	<b>215</b>	<b>0.6</b>
<b>Loamy Till (Black Chernozem)</b>	<b>38328</b>	<b>99.1</b>
<b>Variable Textured Alluvium (Regosols)</b>	<b>34</b>	<b>0.1</b>
<b>Eroded Slopes</b>	<b>93</b>	<b>0.2</b>
<b>Water</b>	<b>3</b>	<b>0.0</b>
<b>Total</b>	<b>38673</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil series for each soil polygon.

## Generalized Soil Map



**Soil Drainage Map.**

Drainage is described on the basis of actual moisture content in excess of field capacity, and the length of the saturation period within the plant root zone. Five drainage classes plus three land classes are shown on this map.

**Very Poor** - Water is removed from the soil so slowly that the water table remains at or on the soil surface for the greater part of the time the soil is not frozen. Excess water is present in the soil throughout most of the year.

**Poor** - Water is removed so slowly in relation to supply that the soil remains wet for a large part of the time the soil is not frozen. Excess water is available within the soil for a large part of the time.

**Imperfect** - Water is removed from the soil sufficiently slowly in relation to supply to keep the soil wet for a significant part of the growing season. Excess water moves slowly down the profile if precipitation is the major source.

**Well** - Water is removed from the soil readily but not rapidly. Excess water flows downward readily into underlying materials or laterally as subsurface flow.

**Rapid** - Water is removed from the soil rapidly in relation to supply. Excess water flows downward if underlying material is pervious. Subsurface flow may occur on steep slopes during heavy rainfall.


Drainage classification is based on the dominant soil series within each individual soil polygon.

**Table 3. Drainage Classes<sup>1</sup>**

<b>Drainage Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Very Poor</b>	<b>0</b>	<b>0.0</b>
<b>Poor</b>	<b>215</b>	<b>0.6</b>
<b>Imperfect</b>	<b>706</b>	<b>1.8</b>
<b>Well</b>	<b>37656</b>	<b>97.4</b>
<b>Rapid</b>	<b>93</b>	<b>0.2</b>
<b>Marsh</b>	<b>0</b>	<b>0.0</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3</b>	<b>0.0</b>
<b>Total</b>	<b>38673</b>	<b>100.0</b>

<sup>1</sup> Area has been assigned to the dominant drainage class for each soil polygon.

## Soil Drainage Map



### Management Considerations Map.

Management consideration maps are provided to focus on awareness of land resource characteristics important to land use. This map does not presume a specific land use. Rather it portrays the most common and wide spread attributes that apply to most soil landscapes in the province.

These maps **highlight attributes** of soil-landscapes that the land manager must consider for any intended land use.

- Fine texture
- Medium texture
- Coarse texture
- Topography
- Wetness
- Organic
- Bedrock

**F = Fine texture** - soil landscapes with **fine textured soils (clays and silty clays)**, and thus low infiltration and internal permeability rates. These require special considerations to mitigate surface ponding (water logging), runoff, and trafficability. Timing and type of tillage practices used may be restricted.

**M = Medium texture** - soil landscapes with medium to moderately fine textures (**loams to clay loams**), and good water and nutrient retention properties. Good management and cropping practices are required to minimize leaching and the risk of erosion.

**C = Coarse texture** - soil landscapes with **coarse to very coarse textured soils (loamy sands, sands and gravels)**, have a high permeability throughout the profile, and require special management practices related to application of agricultural chemicals, animal wastes, and municipal effluent to protect and sustain the long term quality of the soil and water resources. The risk of soil erosion can be minimized through the use of shelterbelts and maintenance of crop residues.

**T = Topography** - soil landscapes with **slopes greater than 5 %** are steep enough to require special management practices to minimize the risk of erosion.

**W = Wetness** - soil landscapes that have **poorly drained soils and/or >50 % wetlands** (due to seasonal and annual flooding, surface ponding, permanent water bodies (sloughs), and/or high water tables), require special management practices to mitigate adverse impact on water quality, protect subsurface aquifers, and sustain crop production during periods of high risk of water logging.

**O = Organic** - soil landscapes with organic soils, requiring special management considerations of drainage, tillage, and cropping to sustain productivity and minimize subsidence and erosion.


**R = Bedrock** - soil landscapes that have **shallow depth to bedrock (< 50 cm) and/or exposed bedrock** which may prevent the use of some or all tillage practices as well as the range of potential crops. They require special cropping and management practices to sustain agricultural production.

**Table 4. Management Considerations<sup>1</sup>**

Land Resource Characteristics	Area (ha)	Percent of RM
Fine Texture	0	0.0
Fine Texture and Wetness	0	0.0
Fine Texture and Topography	0	0.0
Medium Texture	16459	42.6
Coarse Texture	0	0.0
Coarse Texture and Wetness	0	0.0
Coarse Texture and Topography	0	0.0
Topography	21996	56.9
Topography and Bedrock	0	0.0
Wetness	215	0.6
Bedrock	0	0.0
Organic	0	0.0
Marsh	0	0.0
Unclassified	0	0.0
Water	3	0.0
<b>Total</b>	<b>38673</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil series and slope gradient within each polygon.

## Management Considerations Map





**Agricultural Capability Map.**

This evaluation utilizes the 7 class Canada Land Inventory system (CLI, 1965). Classes 1 to 3 represent the prime agricultural land, class 4 land is marginal for sustained cultivation, class 5 land is capable of perennial forages and improvement is feasible, class 6 land is capable of producing native forages and pasture but improvement is not feasible, and class 7 land is considered unsuitable for dryland agriculture. Subclass modifiers include structure and/or permeability (D), erosion (E), inundation (I), moisture limitation (M), salinity (N), stoniness (P), consolidated bedrock (R), topography (T), excess water (W) and cumulative minor adverse characteristics (X).


This generalized interpretive map is based on the dominant soil series and phases for each soil polygon. The CLI subclass limitations cannot be portrayed at this generalized map scale.

**Table 5. Agricultural Capability<sup>1</sup>**

<b>Class Subclass</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>2</b>	<b>16430</b>	<b>42.5</b>
2T	16154	41.8
2TW	276	0.7
<b>3</b>	<b>21721</b>	<b>56.2</b>
3I	34	0.1
3T	21688	56.2
<b>4</b>	<b>163</b>	<b>0.4</b>
4T	163	0.4
<b>5</b>	<b>214</b>	<b>0.6</b>
5W	214	0.6
<b>6</b>	<b>92</b>	<b>0.2</b>
6T	92	0.2
<b>Water</b>	<b>3</b>	<b>0.0</b>
<b>Total</b>	<b>38624</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil series and slope gradient within each polygon.

## Agriculture Capability Map



**Irrigation Suitability Map.**

Irrigation ratings are based on an assessment of the most limiting combination of soil and landscape conditions. Soils in the same class have a similar relative suitability or degree of limitation for irrigation use, although the specific limiting factors may differ. These limiting factors are described by subclass symbols at detailed map scales. The irrigation rating system does not consider water availability, method of application, water quality, or economics of irrigated land use.

Irrigation suitability is a four class rating system. Areas with no or slight soil and/or landscape limitations are rated **Excellent** to **Good** and can be considered irrigable. Areas with moderate soil and/or landscape limitations are rated as **Fair** and considered marginal for irrigation providing adequate management exists so that the soil and adjacent areas are not adversely affected by water application. Soil and landscape areas rated as **Poor** have severe limitations for irrigation.


This generalized interpretive map is based on the dominant soil series for each soil polygon, in combination with the dominant slope class. The nature of the subclass limitations and the classification of subdominant components is not shown at this generalized map scale.

**Table 6. Irrigation Suitability<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Excellent</b>	<b>0</b>	<b>0.0</b>
<b>Good</b>	<b>16149</b>	<b>41.8</b>
<b>Fair</b>	<b>22212</b>	<b>57.4</b>
<b>Poor</b>	<b>308</b>	<b>0.8</b>
<b>Organic</b>	<b>0</b>	<b>0.0</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3</b>	<b>0.0</b>
<b>Total</b>	<b>38673</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil series and slope gradient within each polygon.

## Irrigation Suitability Map



### Potential Environmental Impact Under Irrigation Map.

A major environmental concern for land under irrigated crop production is the possibility that surface and/or ground water may be impacted. The potential environmental impact assessment provides a relative rating of land into 4 classes (minimal, low, moderate and high) based on an evaluation of specific soil factors and landscape conditions that determine the impact potential.

Soil factors considered are those properties that determine water retention and movement through the soil; topographic features are those that affect runoff and redistribution of moisture in the landscape. Several factors are specifically considered: soil texture, hydraulic conductivity, salinity, geological uniformity, depth to water table and topography. The risk of altering surface and subsurface soil drainage regimes, soil salinity, potential for runoff, erosion and flooding is determined by specific criteria for each property.

Use of this rating is intended to serve as a warning of potential environmental concern. It may be possible to design and/or give special consideration to soil-water-crop management practices that will mitigate any adverse impact.


This generalized interpretive map is based on the dominant soil series and slope class for each soil polygon. The nature of the subclass limitations, and the classification of subdominant components is not shown at this generalized map scale.

**Table 7. Potential Environmental Impact Under Irrigation<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Minimal</b>	<b>0</b>	<b>0.0</b>
<b>Low</b>	<b>16478</b>	<b>42.6</b>
<b>Moderate</b>	<b>21935</b>	<b>56.7</b>
<b>High</b>	<b>256</b>	<b>0.7</b>
<b>Organic</b>	<b>0</b>	<b>0.0</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3</b>	<b>0.0</b>
<b>Total</b>	<b>38673</b>	<b>100.0</b>

<sup>1</sup> Based on the **dominant** soil series and slope gradient within each polygon.

## Potential Environmental Impact Under Irrigation



**Water Erosion Risk Map.**

The risk of water erosion was estimated using the universal soil loss equation (USLE) developed by Wischmeier and Smith (1965). The USLE predicted soil loss (tons/hectare/year) is calculated for each soil component in each soil map polygon. Erosion risk classes are assigned based on the weighted average soil loss for each map polygon. Water erosion risk factors include mean annual rainfall, slope length, slope gradient, vegetation cover, management practices, and soil erodibility. The map shows 5 classes of soil erosion risk based on bare unprotected soil:

**negligible**  
**low**  
**moderate**  
**high**  
**severe.**


Cropping and residue management practices will significantly reduce this risk depending on crop rotation program, soil type, and landscape features.

**Table 8. Water Erosion Risk<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Negligible</b>	<b>920</b>	<b>2.4</b>
<b>Low</b>	<b>1147</b>	<b>3.0</b>
<b>Moderate</b>	<b>22386</b>	<b>57.9</b>
<b>High</b>	<b>13921</b>	<b>36.0</b>
<b>Severe</b>	<b>294</b>	<b>0.8</b>
<b>Unclassified</b>	<b>0</b>	<b>0.0</b>
<b>Water</b>	<b>3</b>	<b>0.0</b>
<b>Total</b>	<b>38673</b>	<b>100.0</b>

<sup>1</sup> Based on the **weighted average** USLE predicted soil loss within each polygon, assuming a bare unprotected soil.

## Water Erosion Risk Map





**Land Use Map.**

The land use classification of the RM has been interpreted from LANDSAT satellite imagery, using supervised computer classification techniques. Many individual spectral signatures were classified and grouped into the seven general land use classes shown here. Although land use changes over time, and some land use practices on individual parcels may occasionally result in similar spectral signatures, this map provides a general representation of the current land use in the RM.

The following is a brief description of the land use classes:

**Annual Crop Land** - land that is normally cultivated on an annual basis.

**Forage** - perennial forages, generally alfalfa or clover with blends of tame grasses.

**Grasslands** - areas of native or tame grasses, may contain scattered stands of shrubs.

**Trees** - lands that are primarily in tree cover.

**Wetlands** - areas that are wet, often with sedges, cattails, and rushes.


**Water** - open water - lakes, rivers streams, ponds, and lagoons.

**Urban and Transportation** - towns, roads, railways, quarries.

**Table 9. Land Use<sup>1</sup>**

<b>Class</b>	<b>Area (ha)</b>	<b>Percent of RM</b>
<b>Annual Crop Land</b>	<b>24230</b>	<b>62.2</b>
<b>Forage</b>	<b>25</b>	<b>0.1</b>
<b>Grasslands</b>	<b>6949</b>	<b>17.8</b>
<b>Trees</b>	<b>1525</b>	<b>3.9</b>
<b>Wetlands</b>	<b>4551</b>	<b>11.7</b>
<b>Water</b>	<b>439</b>	<b>1.1</b>
<b>Urban and Transportation</b>	<b>1222</b>	<b>3.1</b>
<b>Total</b>	<b>38941</b>	<b>100.0</b>

<sup>1</sup> Land use information (1995) and map supplied by Prairie Farm Rehabilitation Administration. Areas may vary from previous maps due to differences in analytical procedures.



**REFERENCES**

Agronomic Interpretations Working Group. 1995. Land Suitability Rating System for Agricultural Crops: 1. Spring-seeded Small Grains. Edited by W.W. Pettapiece. Tech. Bull. 1995-6E. Centre for Land and Biological Resources Research, Agriculture and Agri-Food Canada, Ottawa. 90 pages, 2 maps.

Ash, G.H.B. 1991. An Agroclimatic Risk Assessment of Southern Manitoba and Southeastern Saskatchewan. M.A. Thesis. Department of Geography, University of Manitoba, Winnipeg.

Canada Land Inventory. 1965. Soil Capability Classification for Agriculture. Canada Land Inventory Report No. 2. ARDA, Dept. of Forestry, Canada, Ottawa.

Canada-Manitoba Soil Survey. 1980. Physiographic Regions of Manitoba. Ellis Bldg., University of Manitoba, Winnipeg. Revised. Unpublished Report.

Canada-Manitoba Soil Survey. 1979. Ecological Regions and Subregions in Manitoba. Ellis Bldg., University of Manitoba, Winnipeg. Revised. Unpublished Report.

Ehrlich, W.A., Pratt, L.E. Poyser, E.A. and Leclaire, F.P. 1958 Report of Reconnaissance Soil Survey of West-Lake Map Sheet Area. Soils Report No. 8. Manitoba Soil Survey. Published by Manitoba Dept. of Agriculture. 100pp and 1 maps.

Environment Canada. 1982. Canadian Climatic Normals 1951-1980. Frost, Vol. 6; Atmospheric Environment, Downsview, Ontario.

Environment Canada. 1993. Canadian Climatic Normals 1961-1990. Prairie Provinces. Atmospheric Environment, Downsview, Ontario.

Expert Committee on Soil Survey. 1987. The Canadian System of Soil Classification. Second Edition. Publ. No. 1646. Research Branch, Agriculture Canada.

Irrigation Suitability Classification Working Group. 1987. An Irrigation Suitability Classification System for the Canadian Prairies. LRRC contribution no. 87-83, Land Resource Research Centre, Research Branch, Agriculture Canada, Ottawa

Land Resource Unit. 1998. Soil and Terrain Classification System Manual. In preparation. Ellis Bldg. University of Manitoba. Winnipeg.

MacDonald, K.B., and Valentine, K.W.G. 1992. CanSIS Manual 1 CanSIS/NSDB: A General Description. Land Resource Division, Centre for Land and Biological Resources Research, Research Branch, Agriculture Canada, Ottawa.

Podolsky, I.G. 1984. Soils of the Hallboro-Minnedosa Area. Report No. D58. Canada-Manitoba Soil Survey. Winnipeg.

Wischmeier, W.H. and Smith, D.D. 1965. Predicting Rainfall-erosion Loss from Cropland East of the Rocky Mountains. U.S. Department of Agriculture, Agriculture Handbook No. 282, U.S. Government Printing Office, Washington, D.C.