



Environment and
Climate Change Canada

Environnement et
Changement climatique Canada



ESTUARY ISLANDS NATIONAL WILDLIFE AREA MANAGEMENT PLAN 2018

Canada

Acknowledgements:

This management plan was developed by Benoît Roberge of the Canadian Wildlife Service of Environment and Climate Change Canada. Thanks are extended to Canadian Wildlife Service employees who were involved in the development or review of the document: Luc Bélanger, Marielou Verge, Olaf Jensen, Renée Langevin, Matthieu Allard, Martine Benoit, Benoît Jobin, Jean-François Rail, Stéphanie Gagnon, Josée Tardif, Christine Lepage, Diane Dauphin and Francine Rousseau. Special thanks are addressed to Christiane Foley for her input in this project and to Jean Bédard and Kim Marineau for their work on the initial drafts. The Canadian Wildlife Service would also like to thank Jean Bédard, Jean-François Giroux, Jean Huot, Cindy Garneau and Yvon Mercier from the Société Duvetor Ltée, Marc Lapointe from the Société protectrice des eiders de l'estuaire as well as Florence Parcoret and Jessie Moreau from the Essipit Innu First Nation Council who agreed to comment on this document.

Copies of this plan are available at the following addresses:

Environment and Climate Change Canada
Public Inquiries Centre
7th floor, Fontaine Building
200 Sacré-Coeur Boulevard
Gatineau QC K1A 0H3
Telephone: 819-938-3860
Toll free number: 1-800-668-6767 (in Canada only)
Email: ec.enviroinfo.ec@canada.ca

Environment and Climate Change Canada – Canadian Wildlife Service
Quebec Region
801-1550 D'Estimauville Avenue
Québec QC G1J 0C3

Environment and Climate Change Canada – Protected Areas website: www.ec.gc.ca/ap-pa

ISBN: 978-0-660-27384-6
Cat. No.: CW66-409/2018E-PDF

How to cite this document:

Environment and Climate Change Canada. 2016. Management Plan for the Estuary Islands National Wildlife Area. Environment and Climate Change Canada, Canadian Wildlife Service, Québec, 55 p.

Unless otherwise specified, you may not reproduce materials in this publication, in whole or in part, for the purposes of commercial redistribution without prior written permission from Environment and Climate Change Canada's copyright administrator. To obtain permission to reproduce Government of Canada materials for commercial purposes, apply for Crown Copyright Clearance by contacting:

Environment and Climate Change Canada
Public Inquiries Centre
7th floor, Fontaine Building
200 Sacré-Coeur Boulevard
Gatineau QC K1A 0H3
Telephone: 819-938-3860
Toll free number: 1-800-668-6767 (in Canada only)
Email: ec.enviroinfo.ec@canada.ca

Cover page photos: © Environment and Climate Change Canada – M. Dionne

© Her Majesty the Queen in Right of Canada, represented by the Minister of Environment and Climate Change, 2016

Aussi disponible en français

About Environment and Climate Change Canada's Protected Areas and Management Plans

What are Environment and Climate Change Canada Protected Areas?

Environment and Climate Change Canada establishes marine and terrestrial National Wildlife Areas for the purposes of conservation, research and interpretation. National Wildlife Areas are established to protect migratory birds, species at risk, and other wildlife and their habitats. National Wildlife Areas are established under the authority of the *Canada Wildlife Act* and are, first and foremost, places for wildlife. Migratory Bird Sanctuaries are established under the authority of the *Migratory Birds Convention Act, 1994* and provide a refuge for migratory birds in marine and terrestrial environments.

What is the Size of the Environment and Climate Change Canada Protected Areas Network?

The current Protected Areas Network consists of 54 National Wildlife Areas and 92 Migratory Bird Sanctuaries comprising more than 12 million hectares across Canada.

What is a Management Plan?

A management plan provides the framework in which management decisions are made. It is intended to be used by Environment and Climate Change Canada staff to guide decision making, notably with respect to permitting. Management is undertaken in order to maintain the ecological integrity of the protected area and to maintain the attributes for which the protected area was established. Environment and Climate Change Canada prepares a management plan for each protected area in consultation with First Nations and other stakeholders.

A management plan specifies activities that are allowed and identifies other activities that may be undertaken under the authority of a permit. It may also describe the necessary improvements needed in the habitat, and specify where and when these improvements should be made. A management plan identifies Aboriginal rights and allowable practices specified under land claims agreements. Further, measures carried out for the conservation of wildlife must not be inconsistent with any law respecting wildlife in the province in which the protected area is situated.

What is Protected Area Management?

Management includes monitoring wildlife, maintaining and improving wildlife habitat, periodic inspections, enforcement of regulations, as well as the maintenance of facilities and infrastructure. Research is also an important activity in protected areas; hence, Environment and Climate Change Canada staff carries out or coordinates research in some sites.

The Series

All of the National Wildlife Areas are to have a management plan. All of these management plans will be initially reviewed five years after the approval of the first plan, and every ten years thereafter.

To Learn More

To learn more about Environment and Climate Change Canada's protected areas, please visit the Department's website at www.ec.gc.ca/ap-pa or contact the Canadian Wildlife Service.

Estuary Islands National Wildlife Area

Estuary Islands National Wildlife Area (NWA) is made up of approximately a dozen islands or portions of islands. It encompasses a total area of 404 hectares and extends over 120 kilometres in the St. Lawrence Lower and Upper Estuaries between Kamouraska and Rimouski (Le Bic). This wildlife area was created in 1986 by the Canadian Wildlife Service of Environment and Climate Change Canada with the purpose of protecting important nesting sites for migratory birds, notably colonial seabirds, and particularly the Common Eider.

These rocky islands are surrounded by brackish or salt water and bordered by wide, muddy or rocky intertidal flats with communities of mixed algae. The islands themselves are covered in Balsam Fir-White Birch stands as well as White Spruce and herbaceous vegetation.

This protected area is home to approximately one hundred species of birds, a large proportion of which are nesters. Furthermore, five of these islands have been designated Important Bird Areas (IBA). The NWA accommodates several colonies of seabirds such as the Common Eider, the NWA's most abundant bird, as well as the Razorbill, Black Guillemot, Black-legged Kittiwake, Double-crested Cormorant, Great Black-backed Gull and Herring Gull. Approximately 11 500 pairs of Common Eiders nest in the NWA; this accounts for just over half of this species' nesting pairs in the St. Lawrence Estuary. The Île Bicquette colony alone numbers close to 7 000 pairs, making it one of the largest colonies of Common Eiders in North America. Waterfowl species other than the Common Eider are infrequent during the nesting season, but thousands of Brants, Snow Geese, American Black Ducks, Scoters and Goldeneyes, as well as numerous shorebirds use these islands, along with the intertidal flats and adjacent waters during migration. Three species at risk frequent the NWA and its surroundings: the Peregrine Falcon, Red Knot and Barrow's Goldeneye. The NWA is also home to about a dozen species of terrestrial mammals, the most common of which are the Red Fox, Snowshoe Hare and Muskrat. Moreover, the Grey Seal and the Harbour Seal use the shores of certain islands as haul-out sites.

Estuary Islands NWA is exposed to a range of threats and presents a number of management challenges, particularly wildlife diseases, habitat degradation, the impact of predators, the impact of human activities, invasive plant species, accidental spills, fragmentation, and facilities, infrastructure, and land maintenance as well as gaps in scientific knowledge.

Owing to the fragility of the lands and of the wildlife species that inhabit it, public access to the NWA is prohibited, except on Le Pot du Phare island. Access to this island is allowed for public education purposes, but only after the seabird nesting season, from mid-July to

mid-October, and is conditional on using the transportation service provided by the agency authorized by Environment and Climate Change Canada.

The goals for this management plan are: 1) to protect and improve habitats that are important for species at risk, priority bird species and other wildlife; 2) to reduce the impact of human activities on the NWA; 3) to consolidate the NWA's land holdings and promote the conservation of natural habitats on adjacent islands; 4) to ensure ecological monitoring of the NWA and improve knowledge about its wildlife and their habitats.

This document is the first approved management plan for Estuary Islands NWA. It will be implemented over 10 years based on priorities and available resources.

For greater certainty, nothing in this management plan shall be construed so as to abrogate or derogate from the protection provided for existing Aboriginal or treaty rights of the Aboriginal peoples of Canada by the recognition and affirmation of those rights in section 35 of the *Constitution Act, 1982*.

Table of Contents

1	DESCRIPTION OF THE PROTECTED AREA	1
1.1	Regional Context.....	5
1.2	Historical Background	7
1.2.1	<i>Prehistory</i>	7
1.2.2	<i>History</i>	7
1.3	Land Ownership.....	9
1.4	Facilities and Infrastructure	9
2	ECOLOGICAL RESOURCES.....	14
2.1	Terrestrial and Aquatic Habitats.....	14
2.2	Wildlife Species.....	17
2.2.1	<i>Invertebrates</i>	17
2.2.2	<i>Fish</i>	18
2.2.3	<i>Amphibians and Reptiles</i>	18
2.2.4	<i>Birds</i>	19
2.2.5	<i>Mammals</i>	25
2.3	Species at Risk	26
2.4	Invasive Species	27
3	MANAGEMENT CHALLENGES AND THREATS.....	29
3.1	Wildlife Diseases.....	29
3.2	Habitat Degradation	29
3.3	Impact of Predators.....	30
3.4	Human Impact.....	30
3.5	Invasive Plant Species.....	31
3.6	Accidental Spills.....	31
3.7	NWA Fragmentation.....	31
3.8	Facility, Infrastructure and Land Maintenance	31
3.9	Gaps in Scientific Knowledge.....	32
4	GOALS AND OBJECTIVES	33
4.1	Vision	33
4.2	Goals and Objectives.....	33
4.3	Evaluation	34
5	MANAGEMENT APPROACHES.....	38
5.1	Habitat Management.....	38
5.2	Wildlife Management.....	38
5.3	Monitoring	39
5.4	Research.....	39
5.5	Public Information and Outreach.....	40

6	AUTHORIZATIONS AND PROHIBITIONS	41
6.1	Prohibition of Entry.....	41
6.2	Authorized Activities.....	41
6.3	Authorizations	42
6.4	Exceptions.....	43
6.5	Other Federal and Provincial Authorizations.....	43
7	HEALTH AND SAFETY	44
8	ENFORCEMENT	45
9	PLAN IMPLEMENTATION	46
10	COLLABORATORS	47
11	REFERENCES	52
APPENDIX I: ENVIRONMENT AND CLIMATE CHANGE CANADA NOTICE		
	PUBLISHED IN 2016	55

1 DESCRIPTION OF THE PROTECTED AREA

Estuary Islands NWA was created in 1986 by the Canadian Wildlife Service of Environment and Climate Change Canada to protect important nesting sites for migratory birds, especially colonial seabirds and particularly the Common Eider (*Somateria mollissima*). It is home to this species' largest colony in North America. Table 1 summarizes general information about this NWA.

This protected area of 404 hectares is located in the brackish and salt waters of the St. Lawrence River Lower and Upper estuaries. It is made up of approximately a dozen islands or portions of islands, which are spread out over 120 kilometres between Kamouraska and Rimouski (Le Bic) (Figures 1 and 2 and Table 2) and are at a maximum elevation of approximately 30 metres. The distance separating the islands from the south shore varies between two and ten kilometres. The larger islands are mostly colonized by Balsam Fir-White Birch stands and White Spruce stands; while the smaller islands are generally covered in wet meadows composed of herbaceous vegetation.

Table 1: Information on Estuary Islands National Wildlife Area

Protected Area Designation	National Wildlife Area
Province or Territory	Québec – Municipalities of Kamouraska, Saint André and Rimouski (Le Bic); Regional County Municipality (RCM) of Kamouraska and RCM of Rimouski-Neigette
Latitude and Longitude	48°04'N and 69°29'W
Size	404 ha: 83.5 ha of lands and 320 ha of intertidal flats
Protected Area Selection Criteria (Protected Areas Manual¹)	Criteria 1a – The area's islands play a vital role in seabird nesting in the St. Lawrence Estuary, especially for the Common Eider, and they are an important resting and feeding place for waterfowl and shorebirds. Criteria 3a – The area provides colonial birds of the St. Lawrence Estuary with an island habitat that is relatively protected from the threats of predators and human disturbances.
Protected Area Classification System (Protected Areas Manual¹)	Category A – Conservation of species or critical habitat
International Union for Conservation of Nature (IUCN²) Classification	Category Ia – Strict nature reserve
Order-in-Council Number	SOR/DORS/2000-123

Directory of Federal Real Property (DFRP) Number	27013 (Île Bicquette only)
---	----------------------------

Table 1: Information on Estuary Islands National Wildlife Area (continued)

Gazetted	1986
Additional Designations	The following islands in the NWA are internationally designated as Important Bird Areas (IBA): Le Long Pèlerin (Les Pèlerins), Île aux Fraises, Le Pot du Phare (Brandypot Islands), Île Blanche and Île Bicquette.
Faunistic³ and Floristic⁴ Importance	More than half of the estuary's Common Eider pairs, approximately 11 500 couples, nest in the NWA. The NWA also supports several colonies of Razorbills and Black Guillemots. La Grande Île is home to Quebec's westernmost colony of Black-legged Kittiwakes.
Invasive Species⁴	On certain islands, exotic (or introduced) plant species make up 25 to 35% of all plant species in open areas. Some of them are considered invasive, including the Reed Canary Grass, the Wild Radish and the Smooth Bedstraw. Some animal species such as the Double-crested Cormorant can cause damage to habitat should they become abundant.
Species at Risk^{3, 4}	The area is home to a species that is at risk under Canada's <i>Species at Risk Act</i> (SARA) and under Quebec's <i>Act respecting threatened or vulnerable species</i> (ARTVS), namely the Peregrine Falcon, which nests in the NWA. The Red Knot probably uses the NWA's intertidal flats and the Barrow's Goldeneye uses the adjacent waters.
Management Agency	Environment and Climate Change Canada – Canadian Wildlife Service
Public Access and Use	Restricted public access only on Le Pot du Phare, one of the Brandypot Islands (Îles du Pot à l'Eau-de-Vie). Hiking, wildlife observation and photography are authorized only in designated areas (trails, lookouts) and at certain times of the year.

¹ Environment Canada, 2005

² IUCN, 2008

³ Names of vertebrate wildlife species used by MFFP, 2015.

⁴ Names of plant species used by Brouillet et al., 2010+ (VASCAN, accepted names).




Figure 1: Estuary Islands National Wildlife Area

Table 2: Islands, parts of islands and intertidal flats in Estuary Islands National Wildlife Area ¹

Island or part of island	Estimated area (ha) ²	
	NWA	Island (total area)
<i>Îles de Kamouraska</i>		
Île Brûlée	14.74	14.74
Les Rochers (southwest of Île Brûlée) ³	2.48	2.48
Les Rochers (northeast of Île Brûlée) ⁴	1.43	1.43
Île de la Providence	3.72	3.72
La Grande Île (part)	18.38	19.33
<i>Les Pèlerins</i>		
Le Long Pèlerin (part)	3.80	71.04
<i>Brandypot Islands (Îles du Pot à l'Eau-de-Vie)</i>		
Le Pot du Phare (part)	7.76	9.33
Le Pot du Phare intertidal flats	4.68	–
<i>Others</i>		
Île Blanche and reefs	5.95	5.95
Île Blanche intertidal flats	202.73	–
Île aux Fraises and reefs	8.50	8.50
Île aux Fraises intertidal flats	113.17	–
Île Bicquette (part)	16.73	17.38
TOTAL (terrestrial)	83.50	
TOTAL (flats)	320.58	
GRAND TOTAL	404.08	

¹ The NWA's islands, parts of islands and flats are the property of Environment and Climate Change Canada and are managed by the Department.

² Source: Environment Canada, 2013a

³ Sometimes called the western reefs.

⁴ Sometimes called the eastern reefs.



Figure 2: La Grande Île in Estuary Islands National Wildlife Area

Photo: Chantal Lepire © Environment and Climate Change Canada, Canadian Wildlife Service

1.1 REGIONAL CONTEXT

Estuary Islands NWA extends over 120 kilometres between Kamouraska and Rimouski (Le Bic). As a result of its geographic spread, it straddles several administrative entities. It is part of the municipalities of Kamouraska, Saint-André and Rimouski, and the regional county municipalities (RCM) of Kamouraska and Rimouski-Neigette, located in the Bas-Saint-Laurent administrative region. The territory of the RCM of Kamouraska, which is home to 22 000 people, is involved in agriculture, forestry, energy, and tourism, among others. The manufacturing sector plays an important role and benefits from the presence of major employers. The RCM of Rimouski-Neigette is home to more than 54 000 people, 85% of whom live in Rimouski. This city, which is an important regional economic driver, supports a commercial seaport, a high concentration of businesses, services, educational and health institutions, numerous corporate headquarters and major administrative centres. The RCM also includes a large area dedicated to agriculture, forestry and tourism. The Bas-Saint-Laurent region has various tourist attractions including parks, gardens, bike paths, hiking trails, museums, and historical sites. Further, discovery activities on the St. Lawrence, such as kayak excursions and whale watching cruises,

are offered at locations such as Saint-André, Rivière-du-Loup, Trois-Pistoles, and Le Bic (Rimouski).

None of the islands in the NWA are inhabited or open to the public, with the exception of Le Pot du Phare, in the Brandypot archipelago (Îles du Pot à l'Eau-de-Vie), where nature observation, interpretation activities and hiking are permitted during the summer. These activities are offered by Société Duvetnor Ltée, who also offers a lodging service in the lighthouse located on the north-eastern tip of the island, outside the NWA. Close to 1 000 tourists visit the island each year. Seasonal access to the island is allowed under an agreement between Environment and Climate Change Canada and Duvetnor, which owns the lighthouse as well as the other islands in the archipelago (Le Gros Pot and Le Petit Pot). The collection of eiderdown, which does not threaten the species in any way, is carried out in some of the NWA's islands by Société Duvetnor and the Société protectrice des eiders de l'estuaire (SPEE), two non-profit organizations. Interpretive activities and down collection are authorized by Environment and Climate Change Canada under commercial permits. These two organizations have been active in the preservation and development of the area for several years through service delivery contracts awarded by Environment and Climate Change Canada. In addition to the recreation and tourism activities it manages at Le Pot du Phare, Société Duvetnor monitors and protects the NWA's main islands. This organization has significantly contributed to the knowledge and protection of resources for more than 30 years. The SPEE has monitored and protected Île Bicquette for close to 25 years.

The NWA is part of a network of sites dedicated to preserving the natural heritage of the St. Lawrence Estuary and its coastline, including Parc national du Bic, a terrestrial provincial park located west of Rimouski, and Saguenay-St. Lawrence Marine Park. The latter encompasses a large portion of the St. Lawrence Estuary and almost all of the Saguenay River Fjord, and is jointly managed by Parks Canada and Parcs Québec. Several of the estuary's islands are protected by non-profit organizations, including the Société Duvetnor (which owns the île du Jardin and île du Milieu in Les Pèlerins archipelago and Le Gros Pot and Le Petit Pot islands in the Brandypot archipelago), the Nature Conservancy of Canada (which owns Le Petit Pèlerin island as well as a portion of Le Long Pèlerin and of Le Gros Pèlerin islands) and the Société Provencher (which owns the Razades islands and Île aux Basques). In addition, a large portion of Île aux Lièvres, formerly owned by Duvetnor, was acquired in 2012 by Québec's Ministère du Développement durable, de l'Environnement, de la Faune et des Parcs (MDDEFP), who created the Réserve de biodiversité projetée de l'Île-aux-Lièvres (the planned Île-aux-Lièvres Biodiversity Reserve). Duvetnor still owns a small portion of the island (59 ha).

1.2 HISTORICAL BACKGROUND

1.2.1 Prehistory

Approximately 10 000 years ago, the Bas-Saint-Laurent region lay under a post-glacial sea (Dionne 1977). As the waters receded, the area became habitable. Archeological sites discovered in Rimouski, notably in Le Bic, that date back to the oldest prehistoric period in north-eastern America, the Paleo-Indian (11 000 to 7 000 before present), suggest the existence of human settlement in the Bas-Saint-Laurent prior to 8 000 before present. There is little evidence left of these Paleo-Indian's way of life, but it appears that they lived off hunting, fishing and gathering (Fortin *et al.* 1993). Several sites from subsequent periods (Archaic and Woodland, 7 000 to 500 before present) were discovered in the Bas-Saint-Laurent. They indicate that for several centuries Aboriginals frequented the shores of the post-glacial sea and, later, of the estuary as well as inland areas, searching for land and water game (Fortin *et al.* 1993).

1.2.2 History

European Contact

When contact with Europeans occurred (historical period, around the 1500s), the Aboriginals living in the region that is now southern Quebec divided into two major groups or linguistic families: the Algonquians (e.g., the Montagnais or Innus and the Maliseets) and the Iroquoians (first the St. Lawrence Iroquoians, who disappeared from the St. Lawrence Valley after the passage of Jacques Cartier, and later the Mohawks and Hurons). Although the Iroquoians frequented the Bas-Saint-Laurent during long journeys from their wintering areas located further upstream, the region was mainly frequented or inhabited by Algonquians, including Innus, Mi'kmaq, Maliseets and possibly the Abenakis of the East (Fortin *et al.* 1993).

Between 1550 and 1652, the Innu hunting grounds covered the entire Bas-Saint-Laurent (between Rivière du Loup and Rivière Matane) and a large portion of the north shore (between La Malbaie, Sept-Îles and Lake Mistassini) (Fortin *et al.* 1993). They also occupied the islands next to the south shore (F. Parcoret, pers. comm. 2013). Around the same time, the Mi'kmaq traditional territory extended between Prince Edward Island and Gaspé. Therefore, present-day Bas-Saint-Laurent was located on the northwestern edge of their territory. The Maliseets inhabited a vast territory that included a large part of present-day Bas-Saint-Laurent, New Brunswick and Maine, the heart of which was the Saint John River valley (N.B.). These people's way of life was disrupted to a large extent by the arrival of the Europeans. The Innu population declined in the middle of the 17th century (starting in 1652). The Mi'kmaq and Maliseets were still very present in the Bas-Saint-Laurent area during the 17th and 18th centuries, but they

subsequently experienced periods of crisis and decline (Fortin et al. 1993). Today, the Bas-Saint-Laurent's only aboriginal community is the Maliseets of Viger First Nation. It owns an area in the Township of Whitworth, near Rivière-du-Loup, and a small plot of land in Cacouna, which is the smallest aboriginal reserve in Canada (Maliseets of Viger First Nation 2014).

Colonization and Seigneurial System

The earliest settlements of a sedentary population in the Bas-Saint-Laurent were created under the French seigneurial system (1653-1854). This system involved granting a portion of land to entrepreneurs (the “seigneurs”), where they would establish inhabitants (“censitaires” or settlers). Between 1653 and 1751, 19 seigneuries were granted in the Bas-Saint-Laurent (Fortin 2003; Fortin *et al.* 1993). The NWA's islands were allocated at that time, and were the subject of countless transactions, gifts, legacies and sales over the next three centuries. Bédard (2010) describes some of these transactions:

- Île aux Fraises, Île Blanche and Le Pot du Phare [as well as Île aux Lièvres] were granted to the Sieur de Saint-Lusson on November 7, 1672.
- The Îles de Kamouraska were granted to the Sieur de La Durantaye in 1674. They are all quite close to the St. Lawrence's south shore and several of them are accessible by foot at low tide. This proximity explains their marked human use and the existence of residual fishing rights. Several weir sites were set up on the shores of the archipelago's islands under rights often vested by deeds.
- Île Bicquette was granted to Monsieur de Vitré on May 6, 1675 (along with the Île du Bic).
- The Les Pèlerins archipelago was granted to François Jean Baptiste Deschamps, Sieur de la Bouteillerie, and to Étienne Landron and Louis de Niort on May 11, 1697, as a fiefdom and a seigneurie. The archipelago was used mostly as a commercial fishing station.

During the first half of the 18th century, the settlement of the Bas-Saint-Laurent progressed very slowly. During that time, the population was concentrated in four seigneuries: Rivière-du-Loup, L'Isle-Verte, Trois-Pistoles and Rimouski. Even after the British Conquest (1759), the Bas-Saint-Laurent was too far from the centre of the colony and had little appeal for young families who preferred the good farming lands of the Côte-du-Sud (south shore of the river between Beaumont and Kamouraska). The overpopulation of the Côte-du-Sud seigneuries eventually led to increased settlement in the Bas-Saint-Laurent region in the early 1800s (Fortin 2003).

National Wildlife Area

Environment and Climate Change Canada acquired the NWA's islands and portions of islands from private owners (by mutual agreement) and through interdepartmental transfers. On June 19, 1986, Environment and Climate Change Canada created Estuary Islands National Wildlife Area to protect colonial seabirds that nest in the estuary and their habitats, specifically a significant portion of the Common Eider population. When the NWA was created, the Îles de Kamouraska, Les Pèlerins, Le Pot du Phare, Île aux Fraises and Île Blanche had Migratory Bird Sanctuary status. Since this status was no longer necessary given the new NWA status, these five sanctuaries were abolished in 1996 by an Order in Council.

This document is the first approved management plan for Estuary Islands NWA. A conservation plan of this protected area was published in 2003 (CWS 2003).

1.3 LAND OWNERSHIP

The islands, portions of islands and intertidal flats (or foreshores) that make up the NWA are owned and managed by Environment and Climate Change Canada. The total area of the islands and portions of islands is 83.5 hectares and the total area of the intertidal flats is 320.58 hectares. Only the intertidal flats surrounding Île Blanche, Île aux Fraises and Le Pot du Phare are part of the NWA.

Some of the islands and portions of islands adjacent to the NWA remain private property, while several belong to organizations dedicated to preserving and enhancing the environment. In addition, Fisheries and Oceans Canada owns parcels of land adjacent to the NWA on Île Bicquette and La Grande Île.

1.4 FACILITIES AND INFRASTRUCTURE

The NWA's facilities and infrastructure are described below and in Table 3. The portion of Île Bicquette that is in the NWA (Figure 3) includes some infrastructure, i.e. the former lighthouse keeper's house, a foghorn building, a pump building, a boathouse and a cross (Figure 5). There is a shed on the boundary between the NWA and a parcel of land that belongs to Fisheries and Oceans Canada (its exact location remains to be confirmed).

Le Pot du Phare (Figure 4), in the Brandypot Islands, includes three trails with a total length of about one kilometre, short boardwalks, wooden observation platforms and stairs (Figure 6) that are mostly located inside the NWA. On Le Long Pèlerin, a skeleton tower (automated light) and a brick tower (lighthouse ruins) located in the NWA belong to Fisheries and Oceans Canada (Figure 7).

Fisheries and Oceans Canada owns some land adjacent to the NWA as well as navigational aids and other infrastructure located within. This includes three automated lights installed on skeleton towers and located on La Grande Île, Le Pot du Phare and Île Bicquette. The lighthouses on Île Bicquette and Le Pot du Phare are still present, but are no longer in operation. The lighthouse on Le Pot du Phare (Figure 8) is a “Designated Heritage Lighthouse” (Parks Canada, 2016) owned by Société Duvetnor, who uses it as its operation centre (in addition to other infrastructure on site, including a dock, the lighthouse keeper's house, a kitchen and an information pavilion). Fisheries and Oceans Canada also owns two heliports, one on Île Bicquette and the other on Le Pot du Phare. The lighthouse and the foghorn building (or fog alarm) on Île Bicquette are designated as “Recognized Federal Heritage Buildings” (Federal Heritage Buildings Review Office or FHBRO, 2016) owing to their historical associations and their architectural and environmental value.

Table 3: Facilities and infrastructure in the Estuary Islands NWA

Island	Type of facility or infrastructure	Approximate dimensions	Owner
Le Long Pèlerin	Brick tower (lighthouse ruins) Skeleton tower (automated light)	10 m 13.9 m	Fisheries and Oceans Canada
Le Pot du Phare	Three trails : La Chaloupe La Boucle du Gardien L'Eider Wooden stairs, observation platforms, small bridges and walkways (boardwalks)	1 km	To be confirmed
Île Bicquette	Old lighthouse keeper's house Foghorn building Pump building Boathouse Cross Shed	160 m ² 145 m ² 9 m ² 80 m ² — 9m ²	Environment and Climate Change Canada (the exact location and ownership of the shed remains to be determined)




Figure 3: Facilities and infrastructure on Île Bicquette




Figure 4: Facilities and infrastructure on Le Pot du Phare



Figure 5: Cross, automated light, former lighthouse and foghorn building on Île Bicquette

Photo: Patrick Labonté © Environment and Climate Change Canada, Canadian Wildlife Service



Figure 6: Stairs on La Chaloupe trail located on Le Pot du Phare

Photo: Stéphane Turgeon © Environment and Climate Change Canada, Canadian Wildlife Service



Figure 7: Automated light and old lighthouse on Le Long Pèlerin

Photo: Sylvain Giguère © Environment and Climate Change Canada, Canadian Wildlife Service



Figure 8: Lighthouse and buildings on Le Pot du Phare (outside the NWA), Société Duvetnor's operations centre

Photo: Benoît Roberge © Environment and Climate Change Canada, Canadian Wildlife Service

2 ECOLOGICAL RESOURCES

2.1 TERRESTRIAL AND AQUATIC HABITATS

The Estuary Islands NWA's underlying bedrock is made up of alternating layers of shale, sandstone blocks and conglomerates formed between the beginning of the Cambrian and the Ordovician, 570 to 440 million years ago.

This protected area is subject to daily semi-diurnal tides (two high tides and two low tides per lunar day). The majority of the NWA's islands are located in the St. Lawrence's Upper Estuary, which is characterized by brackish and turbid waters. The NWA's islands do not have any freshwater habitat Bédard, 2010).

The NWA is part of the Atlantic Maritime Ecoregion. It has a continental subpolar, subhumid climate. While most of the NWA's islands are located in the Sugar Maple – Yellow Birch bioclimatic zone, marine and weather conditions primarily support a Balsam Fir – White Birch vegetation type. Within the NWA, variation can be seen in the plant composition of terrestrial habitats from west to east. This variation depends on several factors, including substrate, type of soil, area and drainage. In general, the terrestrial portions of the NWA's islands are covered with Balsam Fir – White Birch stands and White Spruce stands (CWS, 2003) (Figure 9). Other common tree species include the American Mountain-Ash (*Sorbus americana*), the Pin Cherry (*Prunus pensylvanica*), the Trembling Aspen (*Populus tremuloides*) and the Balsam Poplar (*Populus balsamifera*).

The herbaceous vegetation that characterizes some of the non-forested islands and parts of islands was described by Reed (1975 in CWS, 2003) as being dominated by reedgrasses (*Calamagrostis*) and a mixture of annual plants. Muskrats (*Ondatra zibethicus*) may be abundant in these herbaceous areas, and their digging can contribute to the presence of plants typical of disturbed environments such as the Great Burdock (*Arctium lappa*) (Bédard and Guérin, 1991 in CWS, 2003).

More recently, plant surveys were carried out on three of the NWA's islands by Morisset (2010a, b and c in Bédard, 2010). Within the scope of the study 88 species of plants were identified on Île aux Fraises, 50 species on Île Blanche, and 151 species on Le Pot du Phare and the southern edge of Le Gros Pot, including several open country introduced species (Morisset, 2010a and b in Bédard, 2010). Morisset did not identify any rare plants on the three islands, but Asselin (1994 in Bédard, 2010) reports the presence of a very rare primrose, the Cowslip Primrose (*Primula veris*), on Le Gros Pot, located only a few dozen metres away from the NWA.

On all of the NWA's islands, the forest stands are highly disturbed as a result of multiple factors, such as intense grazing by the Snowshoe Hare (*Lepus americanus*), which has destroyed or severely reduced large numbers of forest perennials characteristic of similar stands on the mainland (Bédard et al., 1997 in Bédard, 2010). In addition, the Spruce Budworm (*Choristoneura fumiferana*) caused major damage on all of the islands during the 1970s, particularly to the fir stands on Îles de Kamouraska and on Le Pot du Phare. However, this insect's impact was not as severe in the archipelago of Les Pèlerins, the forests of which are dominated by the Black Spruce (*Picea mariana*). It spared a portion of the Balsam Fir (*Abies balsamea*) forest on Île Bicquette, which is otherwise in decline for unknown reasons. This forest is unique because of its old age (up to 136 years old), the absence of regeneration, the extreme density of its trees and the absence of herbaceous and moss flora on the ground (Bélanger and Bédard, 1997 in Bédard, 2010). The Double-crested Cormorant (*Phalacrocorax auritus*) has also had a significant impact on the forest habitat. Its excrement (or guano) kills trees and has already destroyed forests on several occasions on certain islands, especially Île Blanche, Île Brûlée, La Grande Île and Le Pot du Phare (CWS, 2003). Spatio-temporal changes in the NWA's riparian and terrestrial vegetation, which have been observed since the 1970s, are well documented in the Labrecque and Jobin study (2012).

The succession, community composition and biomass of macrophyte algae in nearshore marine environments has been described in several parts of the estuary (Himmelman *et al.* 1983 and Bourget et al., 1994 in CWS, 2003), but never within the boundaries of the NWA or on equivalent coastal points (St. Lawrence Centre, 1996 in CWS, 2003). The macrophyte algae communities differ widely from one substrate to another. Very large beds of *Laminaria* can be found surrounding Île du Bic, between Île du Bic and Île Bicquette, and surrounding the Brandypot Islands. In addition, various sized beds of *Ascophyllum* and *Fucus* can be found in all of the islands' nearshore environments (Figure 10) and in the top portion of the sublittoral zone. These algal beds seem better developed and more widespread downstream than upstream (CWS 2003).



Figure 9: Forest on La Grande Île

Photo: Benoît Roberge © Environment and Climate Change Canada, Canadian Wildlife Service



Figure 10: Île aux Fraises intertidal flats

Photo: Benoît Roberge © Environment and Climate Change Canada, Canadian Wildlife Service

2.2 WILDLIFE SPECIES

The NWA's wildlife is described in this section based on knowledge gathered over the past 30 years as a result of studies and surveys conducted by the Canadian Wildlife Service and various contributors such as Société Duvetnor. In addition, data on the organisms found in the waters adjacent to the NWA, including invertebrates, fish, and marine mammals, are presented as complementary information, given their ecological links with the NWA.

2.2.1 Invertebrates

Mollusks and Crustaceans

Several species of mollusks and crustaceans could be expected to be found in the sediments bordering the NWA. Softshell Clam (*Mya arenaria*) deposits are likely to be found in the Île Blanche and Île aux Fraises intertidal flats, while the Stimpson's Surfclam (*Mactromeris polynyma*) can be found around Île Bicquette and the Green Sea Urchin (*Strongylocentrotus droebachiensis*) is plentiful around Île Bicquette and Le Pot du Phare (DFO-SIGHAP 2002 in CWS, 2003). Green Sea Urchins are fished in the spring (April and May) and the fall (from the end of October until ice formation begins) almost exclusively near Île Blanche and Île Bicquette. Some of the urchins are taken at the lower boundary of the intertidal flats (Bédard 2010). Other species, such as the Atlantic Razor Clam (*Ensis directus*), Sea Cucumber (*Cucumaria frondosa*), Sand Shrimp (*Crangon septemspinosa*), Icelandic Scallop (*Chlamys islandic*), Rock Crab (*Cancer irroratus*) and Waved Whelk (*Buccinum undatum*) are found in the NWA (Brunel *et al.* 1998 in CWS, 2003). It appears that the American Lobster (*Homarus americanus*) and the Snow Crab (*Chionoecetes opilio*) also inhabit the waters surrounding Île Bicquette. It should be noted that these data (MPO-SIGHAP 2002 in CWS, 2003) were extracted for a one-kilometre zone around the NWA.

Insects and Spiders

In 1994 and 1995, sampling of insects and spiders was performed on approximately twenty islands in the estuary, including five islands that are part of the NWA (Nadeau *et al.*, 2009). Insects from at least four orders (Coleoptera, Diptera, Hymenoptera and Orthoptera) and Arachnids (the class of animals that includes spiders) were identified in the area. In 1995, this sampling dealt more specifically with beetles from the Carabidae family. Forty species of insects in this family were identified. The most harvested species (47% of captured individuals) found on the greatest number of islands (17 out of 20 islands) was the *Pterostichus adstrictus* beetle (Nadeau *et al.* 2009).

Benthos

No studies have been carried out on benthic fauna in the NWA's mud and rock intertidal flats.

2.2.2 Fish

The NWA's islands are too small to have any permanent freshwater river system that would support a community of fish. However, the surrounding waters of the St. Lawrence Estuary support several species of marine fish, including forage species consumed by various aquatic birds that use the area. The main species found are the Atlantic Herring (*Clupea harengus*), which has two seasonal spawning cycles around the western islands (Les Pèlerins, Île aux Lièvres) (Munro et al. 1998 in CWS, 2003), as well as the American Sand Lance (*Ammodytes americanus*) and the Capelin (*Mallotus villosus*). The latter may spawn around the Îles de Kamouraska (DFO-SIGHAP 2002 in CWS, 2003). The Atlantic Sturgeon (*Acipenser oxyrinchus*) has been subject to significant fishing, especially on the Saint-André bank (DFO-SIGHAP 2002 and Caron, 2002 in CWS, 2003), between the archipelagos of Kamouraska and Les Pèlerins. The Atlantic Herring, Blackspotted Stickleback (*Gasterosteus wheatlandi*), Threespine Stickleback (*Gasterosteus aculeatus*) and Winter Flounder (*Pseudopleuronectes americanus*) inhabit the waters surrounding the NWA's islands. The Atlantic Halibut (*Hippoglossus hippoglossus*) was commercially fished near the Brandypot Islands (until approximately 1999) and near the Îles de Kamouraska. The American Shad (*Alosa sapidissima*) is also found in the Kamouraska and Les Pèlerins archipelagos. Finally, the American Eel (*Anguilla rostrata*) is found in the Îles de Kamouraska area in the fall, and the Atlantic Cod (*Gadus morhua*) is present off the coast of Île Bicquette (DFO-SIGHAP 2002 in CWS, 2003).

2.2.3 Amphibians and Reptiles

The NWA provides very little habitats suitable for amphibians and reptiles owing to the absence of freshwater. No surveys of herpetofauna have been carried out in the NWA. To date, only one species of amphibian has been identified on the islands of the upper estuary: the Blue-spotted Salamander (*Ambystoma laterale*), which was observed on Île aux Lièvres (Société Duvetnor, unpublished data in CWS, 2003). Only one species of reptile, the Common Garter Snake (*Thamnophis sirtalis*), is common and widely distributed across the island. It is possible that both of these species were introduced while transporting forage and agricultural equipment during agroforestry activities practised on Île aux Lièvres from 1920 to 1923 and from 1950 to 1953.

2.2.4 Birds

The islands in the estuary are essential for bird conservation because of their location in a significant migratory bird flyway and their role in bird reproduction and feeding. They are also critical in maintaining populations of several species of colonial birds (Chapdelaine and Rail, 2002). Five of the islands in Estuary Islands NWA (île Bicquette, île Blanche, Le Pot du Phare, île aux Fraises and Le Long Pèlerin), as well as other surrounding islands, have been designated Important Bird Areas (IBA).

Colonial Seabirds

Significant proportions of certain populations of colonial seabirds in Quebec reproduce in the NWA (Table 4). The numbers and distribution of the most common species on these islands – including the Common Eider, the Herring Gull (*Larus argentatus*), the Great Black-backed Gull (*Larus marinus*), the Black-legged Kittiwake (*Rissa tridactyla*), the Common Murre (*Uria aalge*), the Black Guillemot (*Cepphus grylle*), and the Razorbill (*Alca torda*) as well as two aquatic colonial species (not necessarily marine), i.e. the Great Blue Heron (*Ardea herodias*) and the Black-crowned Night Heron (*Nycticorax nycticorax*) – have experienced continual fluctuations (Bédard 2010). Nonetheless, the main colonies have generally remained concentrated in the same areas.

Common Eider

The Common Eider population (*dresseri* subspecies) that nests in the estuary is thought to be one of the largest in North America. In 2009, it numbered 19 100 nesting pairs (Savard and Lepage 2013). Approximately 55% of these pairs nest in the NWA, mainly on Île Bicquette (6 716 nesting pairs or 13 432 individuals in 2014), Île Blanche (2 585 pairs or 5 170 individuals in 2015) and Île aux Fraises (1 427 pairs or 2 854 individuals in 2015) (BIOMQ 2015; Duvetnor 2015). This bird is, therefore, the most abundant in the NWA despite significant fluctuations in its numbers owing to epidemics of avian cholera, hunting and juvenile mortality (Joint Working Group on the Management of the Common Eider 2004). It appears, however, that the population in the estuary has decreased at a rate of approximately 2% per year over the last 25 years, which is worrisome (Bédard 2010).

Impressive synchronous fluctuations in the Common Eider population in the estuary in recent decades suggest that common factors are at work on all islands, including the ones within the NWA (Giroux 2008 in Bédard, 2010): three severe declines (1984 to 1985, 1992 to 1994, 2001 to 2002) coincided with three avian cholera outbreaks followed by a slow recovery. In 2002, approximately 10 000 birds (out of a total of close to 64 000) died from this bacterial

disease (Giroux *et al.* 2002 in CWS, 2003). That year, the foci of mortality were more extensive than ever (Île Bicquette, Île Blanche, Île aux Pommes and Île aux Fraises), but the colonies on the north shore of the estuary were not affected (CWS 2003).

Between 1985 and 1990, a joint effort by the Canadian Wildlife Service, Société Duvetnor and Ducks Unlimited Canada resulted in a major redevelopment of the Île Blanche habitat in order to eliminate the suspected cause of the infestation. These projects were followed by an intervention program on several islands in the estuary, which included planting shrubs and coniferous trees, controlled burning, and installing nest boxes for the Common Eider (Joint Working Group on the Management of the Common Eider 2004). In addition, drainage swales were dug to prevent the formation of stagnant water ponds, which is conducive to the development of avian cholera (J.-F. Giroux, pers. comm. 2012). Epidemiological factors of this disease among the Common Eider are still not well understood. There may be a carry-over effect (an epidemic at the end of the summer that could lead to another epidemic the following spring). The characteristics of the habitat appear to have fewer effects than what was believed in the 1980s (e.g. the Île Blanche redevelopment did not eliminate the problem even though it has possibly reduced its incidence and scope) (Joint Working Group on the Management of the Common Eider 2004).

Gulls and Black-Legged Kittiwake

The Herring Gull is an important species in the NWA because of its abundance; however, its numbers have significantly declined over the years (Bédard 1999 in CWS, 2003) for undetermined reasons (CWS 2003). A significant decline has been observed for several decades: the number of pairs identified on Île aux Fraises went from 2200 in 1967 to only 67 in 1999; however, it was at 146 in 2001 (BIOMQ 2015). Further, on other islands like Le Pot du Phare, the number of individuals in the species remained stable during the same period (Bédard 1999 in CWS, 2003). The Herring Gull competes with a larger species in the same genus, the Great Black-backed Gull. These two species of gulls exert predation pressure on the majority of aquatic birds. Predation on juvenile Common Eiders by Great Black-backed Gulls is a real concern throughout the NWA, but especially on Île Blanche and Île Bicquette (Bédard 2010). The Ring-billed Gull (*Larus delawarensis*) also nests in the NWA. The species nested on Île Blanche in 1986, when 265 couples were counted, but was not observed on the island in 1990 (BIOMQ 2015). This gull also nested on Île Brûlée in 2006, when 60 couples were recorded, but was not seen on the island in 2011 (BIOMQ 2015). It seems that the species had difficulties to become established at Île Blanche and also at Île aux Fraises at the end of the 1980s because

of predation by the other two species of gulls (CWS 2003). The species was however present on these islands in 2015 (J. Bédard, pers. comm. 2015).

The Black-legged Kittiwake nests on the Brandypot Islands and Les Pèlerins, but outside the NWA (CWS 2003). Inside the NWA, this species nests on Île Bicquette where varying numbers of pairs were identified between 1986 and 2007, including 130 pairs in 1986, 693 pairs in 2001 and 350 pairs in 2007. It also uses La Grande Île, in the Îles de Kamouraska, where 18 pairs were observed in 2011 (BIOMQ 2015). This is the westernmost colony of this species in Quebec.

Razorbill, Black Guillemot and Common Murre

Data collected between 1990 and 2001 indicate that there were only a few pairs of Razorbills in the NWA, on Le Long Pèlerin, Île Brûlée, La Grande Île, Le Pot du Phare (Bédard 2002 in CWS, 2003) and Île Bicquette (SPEE 1998). In 1999, the Razorbill colony in the archipelago of Les Pèlerins (the part outside the NWA) was the second largest in Canada with more than 1800 pairs identified (Bédard 1999 and Chapdelaine *et al.* 2001 in CWS, 2003). More recent data show that it is still the main colony in this sector with approximately 331 pairs identified in 2011 (BIOMQ 2015).

Some nesting pairs of Black Guillemots were observed on the Îles de Kamouraska (La Grande Île, Île Brûlée and Les Rochers), Le Long Pèlerin, Le Pot du Phare (63 pairs observed in 2011) and Île Bicquette (BIOMQ 2015). The Common Murre can be found in small numbers among groups of Razorbills on Le Pot du Phare and Le Long Pèlerin, but there is now a large concentration of this species on the other Brandypot islands (Le Petit Pot and Le Gros Pot), with more than 1000 individuals observed in each location in 2011 (BIOMQ 2015).

Double-Crested Cormorant

Occasional surveys of the Double-crested Cormorant show that several hundred pairs nest on the NWA's islands, mainly Île Bicquette, Île aux Fraises and La Grande Île (Table 4) (BIOMQ 2015). Significant increases in numbers for this species were observed between 1978 and 1990 in the St. Lawrence. These were due in part to the decrease in human disturbance and persecution of the species as well as the abundance of food near the colonies (Chapdelaine and Bédard 1995). The Double-crested Cormorant has an impact on the islands' forests, because its droppings can kill trees and alter the physical and chemical characteristics of the soil. As such, it devastated the NWA's forests in places where it had established colonies, including on Île Blanche between 1970 and 1985, Île Brûlée, La Grande Île, Le Pot du Phare and other islands located outside the NWA (CWS, 2003). In 1989, owing to the impact this species

was having on the ecosystems, Quebec's Ministère du Loisir, de la Chasse et de la Pêche (MLCP) undertook a control program aimed at reducing its numbers on Île Brûlée, La Grande Île, Île aux Fraises, Île Blanche and the reefs of Île Bicquette. This program included culling adults and spraying eggs with oil (Bédard *et al.* 1995). Despite the effectiveness of this control, the Double-crested Cormorant's population remains present and continues to impact the environment.

Great Blue Heron and Black-Crowned Night Heron

Aerial surveys conducted in 2001 identified 34 Great Blue Heron nests on Île Bicquette, 2 on Le Long Pèlerin, 31 on La Grande Île and 3 on Île Brûlée (Alain Desrosiers, pers. comm. in CWS 2003). Le Gros Pot (outside the NWA, Brandypot Islands) has long supported a large colony of this species. Surveys carried out on this island identified three nests in 1977, 112 in 1992 and 24 in 2010, but no nests were seen in 2011 (BIOMQ 2015).

The Black-crowned Night Heron's numbers have declined sharply on three of the NWA's islands, where the species had previously been abundant. On Le Pot du Phare, 443 nests were identified in 1991 and only 38 nests in 2002 (unpublished data, Duvetnor in CWS, 2003). On Île aux Fraises, close to 200 nests were observed in 1988, but the species was not seen there again. On Île Brûlée, the number of nests went from 537 in 1975 to only 6 in 1990 and the species was not seen there again (no nests between 2006 and 2008 and in 2011) (BIOMQ, 2015).

Waterfowl (other than the Common Eider)

Waterfowl species, other than the Common Eider, are uncommon in the NWA during nesting season, but significant numbers halt there during migration. Thousands of Brants (*Branta bernicla*) stage on Île Blanche during the spring, and hundreds of individuals were also observed on Île Bicquette in spring 1998 (CWS 2003). Several thousand Snow Geese (*Chen caerulescens*) stage near Île aux Fraises and Île Bicquette in the spring and close to the Îles de Kamouraska in the fall (G. Verreault and A. Bérubé, pers. comm., FAPAQ de Rivière-du-Loup in CWS 2003). The Black Scoter (*Melanitta americana*) and the Surf Scoter (*Melanitta perspicillata*) are also common in the islands' subtidal zone during migration. In addition, the Surf Scoter uses the Saint-André bank sector (between the Îles de Kamouraska and Les Pèlerins) during moulting (CWS 2003). Thousands of American Black Ducks (*Anas rubripes*) use the intertidal flats surrounding Île Blanche, Île aux Fraises, Île aux Lièvres, the Brandypot Islands and the Îles de Kamouraska from mid-July to the beginning of November (approximately

Table 4: Number¹ of colonial seabirds² in Estuary Islands NWA (Source: BIOMQ, 2015³, Duvetnor, 2015⁴)

Location	Common Eider	Great Black-backed Gull	Herring Gull	Black-legged Kittiwake	Black Guillemot	Common Murre	Razorbill	Double-crested Cormorant
Îles de Kamouraska								
Île Brûlée	92 (2012)	14 (2011)	370 (2011)		20 (2011)		8 (2011)	2 (2011)
Les Rochers	112 (2012)	34 (2011)	732 (2011)		2 (2011)		78 (2011)	0 (2011)
La Grande Île	380 (1996)	12 (2011)	472 (2011)	36 (2011)	30 (2011)		152 (2011)	1 332 (2011)
Les Pèlerins								
Le Long Pèlerin	0 (2008)	14 (2011)	252 (2011)	0 (2011)	16 (2011)	2 (2011)	663 (2011)	
Brandypot Islands (Îles du Pot à l'Eau-de-Vie)								
Le Pot du Phare	724 (2015)	22 (2011)	1 126 (2011)		126 (2011)	1 (2011)	8 (2015)	48 (2015)
Other islands								
Île Blanche	5 170 (2015)	336 (2001)	292 (2001)					
Île aux Fraises	2 854 (2015)	394 (2001)	292 (2001)					190 (2015)
Île Bicquette	13 432 (2014)	518 (2007)	964 (2007)	700 (2007)	90 (2007)	2(2012)	120 (2007)	654 (2007)

¹ The figures shown indicate the number of nesting individuals and the year of the last data entry. For the Common Eider, Great Black-backed Gull, Herring Gull, Black-legged Kittiwake and Double-crested Cormorant, the number of individuals was obtained by multiplying the number of nests identified (therefore, the number of nesting pairs) by two. For the Black Guillemot, Common Murre and Razorbill, the number of individuals and not the number of nests was identified.

² The Great Blue Heron and the Black-crowned Night Heron, colonial waterbirds (not necessarily marine), do not appear in the table, but they are briefly described above.

³ The *Banque informatisée des oiseaux marins du Québec* (BIOMQ, 2015; contact: Jean-François Rail, CWS) contains the survey results for 28 species of seabirds during reproduction in Quebec, mostly since 1976, but also data extracted from literature going back to 1833. These data mostly cover the Estuary and the Gulf of St. Lawrence. For the islands where only a portion is located in the NWA (e.g. Le Long Pèlerin), the survey data apply to the entire island.

⁴ The 2015 data are all from Duvetnor, 2015.

5000 birds around these islands) (Bédard, Ouellet and Giroux 1987 and Bédard, Ouellet, Giroux and Savard 1988 in CWS, 2003), and small numbers winter in the sector (Gauthier, Choinière and Savard 1992 in CWS, 2003).

In both spring and fall, tens of thousands of scoters, eiders and Long-tailed Ducks (*Clangula hyemalis*) gather in large numbers in the St. Lawrence Lower Estuary (Lepage and Savard 2013; Savard and Lepage 2013; Lepage and Cotter 2013). The estuary is also a major wintering site for several duck species, including the Long-tailed Duck (CWS 2003), the Red-breasted Merganser (*Mergus serrator*) and the American Black Duck. The latter is particularly abundant between Tadoussac and Cap de Bon-Désir, and is especially common in the NWA's Île Bicquette sector (Robert *et al.* 2003). CWS surveys (Robert *et al.* 2003) showed that the St. Lawrence Estuary is also home to a large number of Common Goldeneyes (*Bucephala clangula*) and Barrow's Goldeneyes (*Bucephala islandica*, Eastern population) in winter and serves as the latter's main wintering ground (see 2.3 Species at Risk).

Shorebirds

Many species of shorebirds are common along the St. Lawrence during fall migration (Aubry and Cotter 2007) and several of these species more than likely stopover in the NWA, especially the Red Knot (*Calidris canutus rufa*). No systematic survey of shorebirds has been carried out on the NWA's islands. However, one-time data collected by the CWS in fall 2008 (S. Giguère, pers. comm. 2012) helped to identify 11 species of shorebirds on three of the NWA's islands or nearby (Île aux Fraises, Île Blanche, Île Bicquette), including the Semipalmated Plover (*Charadrius semipalmatus*), the Black-bellied Plover (*Pluvialis squatarola*), the Semipalmated Sandpiper (*Calidris pusilla*), the Dunlin (*Calidris alpina*), the Sanderling (*Calidris alba*), and the Least Sandpiper (*Calidris minutilla*). In addition, large numbers of Purple Sandpipers (*Calidris maritima*) are common on some of the NWA's islands (Île Blanche, Île aux Fraises, Île Bicquette) during the winter (Robert *et al.* 2003; Aubry and Cotter 2007).

Landbirds

In 2008, the CWS surveyed landbirds on the NWA's eight main islands (S. Giguère, pers. comm. 2012). The initial analysis indicates the presence of 40 species of landbirds, even though the data collected are preliminary and were collected in a single season. Since this survey was conducted in June, during the nesting season, the species identified possibly breed there. The data indicate that the large islands are home to a more diverse range of species than the smaller islands. La Grande Île (Îles de Kamouraska) is where the largest number of species (25) was identified, followed by (in descending order): Île Bicquette (19 species), Le Long Pèlerin

(13 species), Île Brûlée (12 species), Le Pot du Phare (11 species), Île de la Providence (9 species) and, finally, Île aux Fraises (3 species) and Île Blanche (2 species). Only 8 of the 40 species identified were observed on four or more islands, with the most common being the American Crow (*Corvus brachyrhynchos*) (8 islands), the Song Sparrow (*Melospiza melodia*) (8 islands), the White-throated Sparrow (*Zonotrichia albicollis*) (6 islands), the Fox Sparrow (*Passerella iliaca*) (5 islands) and the Golden-crowned Kinglet (*Regulus satrapa*) (5 islands). Among birds of prey, only the Rough-legged Hawk (*Buteo lagopus*) was observed (on La Grande Île), but the only nesting bird of prey confirmed in the area is the Peregrine Falcon (*Falco peregrinus anatum/tundrius*) (see 2.3 Species at Risk).

2.2.5 Mammals

The quality of the NWA's islands as habitat for colonial seabirds is primarily determined by the relative lack of terrestrial predators. However, small numbers of such predators sometimes access the islands, where they have a considerable impact on the abundance of birds (Bédard 2010). The Red Fox (*Vulpes vulpes*), the Long-tailed Weasel (*Mustela frenata*) and the American Black Bear (*Ursus americanus*) were all reported in various parts of the NWA, but the Red Fox in particular is responsible for the largest decline in nesting seabirds. It's population persists on Île du Bic and Île aux Lièvres (outside the NWA), where it feeds on a variety of prey. When ice cover allows for movement between the islands, it also invades the NWA's other islands spreading out from the population "hubs" (Bédard 2010).

The Muskrat and the American Mink (*Neovison vison*) are present in the wildlife area, especially on Île aux Fraises and Île Blanche. The Snowshoe Hare is also present. Intense grazing by this species has decimated or eliminated an assemblage of the NWA's forest plants that characterize similar communities on the mainland (Bédard *et al.* 1997b in Bédard, 2010). The White-tailed Deer (*Odocoileus virginianus*) was spotted on several occasions swimming off the shore of the NWA. A juvenile moose (*Alces americanus*) wintered on Le Gros Pèlerin (outside the reserve) in 1986-1987 (CWS 2003) and another individual was observed on La Grande Île (within the area's boundaries) in 2012 (Jean Bédard, pers. comm. 2012).

During a survey of micromammals conducted in 2007 by the CWS on several of the NWA's islands, only the Meadow Vole (*Microtus pennsylvanicus*) was captured on Île de la Providence, Île Brûlée and La Grande Île in the Kamouraska archipelago (S. Giguère, pers. comm. 2012). No micromammals were captured on Île Bicquette, Île aux Fraises, or Île Blanche. During a survey conducted in 1995, the Meadow Vole was also identified on the portion of Le Long Pèlerin that is within the NWA (Nadeau *et al.* 2009). This same survey identified several

species of micromammals on some of the islands in the estuary that are not part of the NWA, including the Masked Shrew (*Sorex cinereus*), the Deer Mouse (*Peromyscus maniculatus*), the Southern Red-backed Vole (*Myodes gapperi*), and the Meadow Vole.

A significant number of Grey Seals (*Halichoerus grypus*) use the NWA as a haul-out site, particularly on the intertidal flats of Île aux Fraises and at the western point of the intertidal flats on Île Blanche. Harbour Seals (*Phoca vitulina vitulina*) also haul out on the Îles de Kamouraska and Les Pèlerins. These two pinniped species coexist in several locations, including the reefs and intertidal flats surrounding Île Bicquette and Île Blanche. Moreover, baby Harbour Seals are observed regularly in May on the shore of several islands (Bédard 2010). Finally, the brackish and salt waters that surround the NWA's islands are used by several species of cetaceans, including the Beluga Whale (*Delphinapterus leucas*), a permanent resident of the St. Lawrence Estuary, whose population is considered as threatened according to Quebec's *Act respecting threatened and vulnerable species* (ARTVS) and Canada's *Species at Risk Act* (SARA; the “threatened” SARA status is in the process of being replaced by the “endangered” status) and, since 2014, as endangered according to the Committee on the Status of Endangered Wildlife in Canada (COSEWIC).

2.3 SPECIES AT RISK

Table 5 presents species at risk that carry out part of their life cycle in the NWA (islands and intertidal flats) or in its surrounding waters.

Between 2002 and 2010, the Peregrine Falcon, a species of special concern in Canada, was observed repeatedly on a cliff on Le Long Pèlerin, in the archipelago of Les Pèlerins, where its nesting has been confirmed. In addition, in July and August 2002, the species was observed on a few occasions on Le Pot du Phare, in the Brandypot Islands (Bédard 2002 in CWS, 2003). One or several juvenile Bald Eagles (*Haliaeetus leucocephalus*) and Golden Eagles (*Aquila chrysaetos*) sometimes spend long periods on Le Pot du Phare in June and July (Bédard 2002 in Bédard, 2010), but no nesting by these species has been confirmed in the NWA. Both are species of special concern in Quebec.

The St. Lawrence Estuary is the main wintering ground for the eastern population of the Barrow's Goldeneye (Robert *et al.* 2003), a species of special concern in Canada. Its numbers are estimated at a maximum of 4500 individuals, of which at least half find refuge in the estuary during the colder months. The species is then observed in the waters adjacent to Île aux Fraises, Île Blanche, the Brandypot Islands (a portion of this archipelago is not part of the NWA) and Île aux Lièvres (outside the NWA).

The Red Knot of the *rufa* sub-species is found in the St. Lawrence Upper Estuary and Lower Estuary primarily during fall migration (Aubry and Cotter 2007). Although this shorebird most likely uses the intertidal flats on the NWA's islands during this period, its presence has not yet been documented. This sub-species is endangered in Canada and is likely to be designated as threatened or vulnerable in Quebec.

The *Centre de données sur le patrimoine naturel du Québec* (CDPNQ 2012) does not contain any entries of plants, amphibians, reptiles, or mammals at risk (in Canada or Quebec) within the NWA.

Table 5: Species at Risk in Estuary Islands NWA

Common and scientific names of the species	Status		
	Canada		Quebec
	SARA ¹	COSEWIC ²	ARTVS ³
Birds			
Peregrine Falcon <i>Falco peregrinus anatum/tundrius</i>	Special Concern*	Special Concern*	Vulnerable** or SLDTV****
Barrow's Goldeneye <i>Bucephala islandica</i> Eastern Population	Special Concern	Special Concern	Vulnerable
Red Knot <i>Calidris canutus rufa</i>	Endangered	Endangered	SLDTV ⁴

¹ Canadian *Species at Risk Act* (Species at Risk Public Registry, 2014)

² Committee on the Status of Endangered Wildlife in Canada (COSEWIC, 2014)

³ Quebec *Act respecting threatened or vulnerable species* (MFFP, 2016)

⁴ Species likely to be designated threatened or vulnerable in Quebec (MFFP, 2016)

* Status assigned to both *anatum/tundrius* sub-species (*F. peregrinus anatum/tundrius*)

** Status assigned to the *anatum* sub-species

*** Sub-species *tundrius*

2.4 INVASIVE SPECIES

No complete study on invasive species has been conducted in the NWA. However, Morisset's work (2010a, b, c in Bédard, 2010) highlighted the significance of exotic plants (referred to as introduced plants by the author) in open environments on three of the NWA's islands: Île Blanche, Île aux Fraises and Le Pot du Phare (one of the Brandypot Islands). On the first two islands, 35% of the flora is made up of exotic species while 25% of the species on the Brandypot Islands are exotic.

On Île Blanche, one of these plants, the Reed Canary Grass (*Phalaris arundinacea*) invaded the entire surface of the island to the extent that it makes up almost the entire herbaceous layer. This plant was sown in 1985 and 1986 on the entire island, except for the coastal belt, during work carried out after the avian cholera epidemic to encourage nesting of the Common Eider (J.-F. Giroux, pers. comm. in Morisset, 2010). The Canada Reedgrass

(*Calamagrostis canadensis*), a native grass species which would be expected to occur here, has not even been observed, possibly owing to the introduction and expansion of the Reed Canary Grass.

On Île aux Fraises, the Purple Loosestrife (*Lythrum salicaria*), an exotic plant considered to be invasive, was found in a single location (a single individual observed south of the western cove); other introduced species such as the Common Timothy (*Phleum pratense*) and the Canada Thistle (*Cirsium arvense*) are distributed throughout the island but are not considered invasive (Morisset 2010 in Bédard, 2010). On this same island, invasive exotic species like the Wild Radish (*Raphanus raphanistrum*) and the Smooth Bedstraw (*Galium mollugo*) often cover a considerable area. In places, the radish invades the entire area normally occupied by Sea Lyme grass (*Elymus arenarius*) or forms a band up to three metres wide in the zone of typical coastal shoreline plants, such as the Beach Pea (*Lathyrus japonicus*). It seems that this island was used during the early and mid-20th century by farmers from Saint-Siméon who came to graze their livestock. This could explain, in part, the presence of certain exotic species (Bédard 2010).

On Île aux Lièvres, close to the NWA, small colonies of the Purple Loosestrife and the European Reed (*Phragmites australis*, exotic) have been observed on some shorelines (CWS 2003).

3 MANAGEMENT CHALLENGES AND THREATS

Estuary Islands National Wildlife Area is exposed to numerous threats and presents several management challenges, including wildlife diseases, habitat degradation, the impact of predators, the impact of human activities, invasive plant species, accidental spills, area fragmentation, facility, infrastructure, and lands maintenance, as well as gaps in scientific knowledge. These are described below in relative order of importance since the scope of several of them is not well known.

3.1 WILDLIFE DISEASES

The NWA's colonial seabirds are susceptible to animal diseases, which can strike a species or a group of species. The Common Eider is occasionally affected by avian cholera, a bacterial disease that triggers potentially devastating epidemics within the species, such as one that decimated close to 10 000 individuals in 2002 (Giroux *et al.* 2002 in CWS, 2003). Factors that contribute to this disease and its spread among the Common Eider are still not well understood. An emergency plan in the event of mass bird mortality was developed by the CWS in order to react to epizootic diseases and to mitigate their impacts.

3.2 HABITAT DEGRADATION

Some of the NWA's animal populations become so abundant at times that they can cause habitat deterioration and affect other species. The increase in the Double-crested Cormorant's population between 1978 and 1990 in the St. Lawrence and the impact of its droppings on the ecosystems of the estuary islands prompted the Quebec Ministère du Loisir, de la Chasse et de la Pêche (MLCP) to launch a species control program in 1989 (Bédard *et al.*, 1995). The CWS is also concerned with the impacts of this species on the evolution of the NWA's habitats and the integrity of bird populations that nest there. Other animals can also disrupt vegetation on the NWA's islands, especially the Snowshoe Hare through its intense grazing (Bédard 2010). Cyclical insect epidemics periodically affect the NWA's forests, such as the Spruce Budworm epidemic that caused major damage on all of the NWA's islands during the 1970s (CWS 2003). The insect pest spread rapidly in 2014 and 2015 and could cause significant disturbances to the forested islands of the NWA, notably on Le Pot du Phare and Bicquette islands (J. Bédard, pers. comm. 2015).

In the last three years, large areas where trees have blown down have appeared on the south side of Île Bicquette as a result of strong winds (M. Lapointe, pers. comm. 2014).

In recent years, shore erosion, leading to the degradation of coastal habitats (receding by more than 5 m), has been observed on some of the NWA's low-lying islands, especially Île aux Fraises and Île Blanche, where the underlying rock is covered with unconsolidated materials (sand and gravel). This phenomenon has also been observed in the eastern part of Île Bicquette (M. Lapointe, pers. comm. 2014). This erosion could be due to storm frequency and intensity as well as high tides, which can be exacerbated by climate change.

3.3 IMPACT OF PREDATORS

Terrestrial predators can occasionally access the NWA's islands (e.g. by swimming or via ice bridges that form between the mainland and the islands). These predators, especially the Red Fox, and certain species of birds, such as gulls, can significantly reduce colonial seabird populations through egg or chick predation (Bédard 2010). Although the presence of these predators on the islands and their predation of birds is a natural phenomenon, controlling certain ones, particularly the fox, is occasionally necessary in order to avoid significant impacts on the affected populations of nesting seabirds. Predator control is subject to Environment and Climate Change Canada's "Predator Management Policy", which takes into account a series of factors to guide interventions.

3.4 HUMAN IMPACT

Despite regulations prohibiting public access (except on Le Pot du Phare), the presence of boaters and kayakers in the NWA can have harmful consequences for the integrity of ecosystems and successful reproduction among colonial seabirds. Trampling by these visitors can disturb wildlife and degrade riparian habitats and flora. These types of unauthorized activities take place primarily on La Grande Île and Le Long Pèlerin, within the NWA's boundaries, where visitors have left their mark (tent prints, fires and garbage). Visitors are also frequently seen on the Îles de Kamouraska (Bédard 2010), but no records are kept of these sightings.

Resource operations around the NWA (fishing, seaweed and sea urchin harvesting) could also have an impact on NWA's wildlife, such as the Common Eider or the Surf Scoter. A project to develop oil and gas reserves (exploration and production) in the sub-surface of the St. Lawrence, specifically in the basin of the Lower Estuary and the north-western Gulf of St. Lawrence, raises fears about the impact these activities could have on wildlife and the environment (AECOM Tecscult Inc. 2010).

Finally, the proposed construction of an oil pipeline and a transshipment terminal near the NWA could lead to increased maritime traffic and wildlife disturbances and an increased risk of accidental toxic substance spills.

3.5 INVASIVE PLANT SPECIES

No comprehensive studies have been done on the NWA's invasive species, but floral surveys conducted on three islands (Île Blanche, Île aux Fraises and Le Pot du Phare) show that exotic (introduced) plants make up 25 to 35% of the vegetation in these islands' open environments (Morisset, 2010 in Bédard, 2010). Some of these species are invasive or cover large areas, including the Reed Canary Grass, the Wild Radish and the Smooth Bedstraw, and can result in loss of biodiversity and natural habitats (Bédard 2010).

3.6 ACCIDENTAL SPILLS

A large number of commercial and passenger cruise vessels travel close to the NWA on the St. Lawrence Seaway every year. An accidental oil or chemical spill in the estuary could result in aquatic bird mortality and have serious consequences for the NWA's shores and ecosystems. Environment and Climate Change Canada and its collaborators have established an emergency response plan (ERP) to implement relevant bird protection measures if such a spill were to occur.

3.7 NWA FRAGMENTATION

The NWA's management and conservation challenges are, in large part, related to its geography. The NWA forms a discontinuous entity made up of islands and portions of islands interspersed with other islands or parts of islands that are not part of the NWA. Some of the islands without NWA status are protected, but others are inhabited or used as vacation spots and could be developed and exploited. For example, on Île Bicquette, the lighthouse owned by Fisheries and Oceans Canada was recently declared surplus and could be turned into a development project. In addition, the NWA's islands are spread out over a long distance (120 km) and are relatively far from the mainland (2 to 10 km from the coast). This limits the ecological connectivity of both habitats and species, which is already naturally weak in island environments. Finally, the NWA's discontinuity, sprawl and remoteness make posting its boundaries, monitoring the area and law enforcement challenging.

3.8 FACILITY, INFRASTRUCTURE AND LAND MAINTENANCE

The NWA has some facilities and infrastructure that require maintenance and restoration, including trails, boardwalks and stairs on Le Pot du Phare, and some buildings (former

lighthouse keeper's house, foghorn building, pump building and boathouse) and a cross on Île Bicquette. In addition, rehabilitation work are necessary on Île Bicquette, Le Pot du Phare, Le Long Pèlerin, La Grande Île and Île de la Providence due to the presence of garbage and small quantities of contaminants left behind before the NWA was created.

3.9 GAPS IN SCIENTIFIC KNOWLEDGE

Since its creation, the NWA has been the subject of several surveys by the CWS and Société Duvetnor. However, gaps remain in the scientific knowledge regarding habitats, flora, plant species at risk, certain animals (including amphibians and reptiles), and threats to the NWA's integrity. Ecological monitoring of the NWA and more knowledge is required to support habitat and species management as well as conservation decisions, such as those concerning the decreasing population of Common Eiders in the estuary.

4 GOALS AND OBJECTIVES

4.1 VISION

Estuary Islands National Wildlife Area protects habitats that are important for species at risk, priority bird species, particularly colonial seabirds, and other wildlife. Priority bird species are those mentioned in the *Bird Conservation Strategy for Bird Conservation Region 14, Quebec Region: Atlantic Northern Forest* (Environment Canada 2013b).

4.2 GOALS AND OBJECTIVES

The goals and objectives below elaborate on the vision for the management plan and take into account management threats and challenges. Achieving these goals and objectives will result in the implementation of the actions presented in Table 6 (Management Approaches for Estuary Islands NWA), which will be carried out according to the resources available.

Goal 1: Protect and improve habitats that are important to species at risk, priority bird species and other wildlife.

Objectives:

- 1.1 Protect colonial seabird populations by restricting natural pressures linked to wildlife diseases, abundant animal populations and terrestrial predators;
- 1.2 Conserve the NWA's seabird colonies, and waterfowl and shorebird staging areas by limiting the effects of human activity (disturbances, accidental spills, etc.);
- 1.3 Implement the management and recovery measures required to protect bird species at risk and priority bird species;
- 1.4 Prevent the proliferation of invasive plant species;
- 1.5 Evaluate erosion of the islands and limit its impact.

Goal 2: Reduce the impact of human activities on the NWA.

Objectives:

- 2.1 Adequately mark the NWA's boundaries in the field, so as to protect wildlife and plants from the impacts of human activity (disturbances from boaters, hunting, etc.);
- 2.2 Promote local and public awareness of the NWA's mission and applicable regulations to reduce instances of non-compliance;

- 2.3 Communicate concerns to managers of adjacent lands and waters regarding resource development near the NWA;
- 2.4 Ensure that Environment and Climate Change Canada's facilities, infrastructure and lands (especially on Île Bicquette and Le Pot du Phare) are in good and safe condition.

Goal 3: Consolidate the NWA's land holdings and promote the conservation of natural habitats on adjacent islands.

Objectives:

- 3.1 Incorporate into the NWA adjacent, federally-owned lands with significant conservation value;
- 3.2 Determine which of the islands located near the NWA have strong conservation potential and develop an acquisition strategy as required;
- 3.3 Encourage use of the islands and lands adjacent to the NWA in a way that is compatible with the NWA's conservation mandate.

Goal 4: Ensure ecological monitoring of the NWA and improve knowledge about its wildlife and their habitats.

Objectives:

- 4.1 Develop and implement an ecological monitoring plan;
- 4.2 Determine gaps in scientific knowledge and address those that are identified as priorities.

4.3 EVALUATION

An annual review of the steps taken and the results achieved will be conducted depending on the availability of financial and human resources. This review will help to identify future priorities for action and resource investment. The management plan itself will be re-evaluated five years after its initial approval and will be reviewed and updated every decade thereafter.

Table 6: Management approaches for Estuary Islands NWA

Goals	Objectives	Actions (Priority Level¹)
<p>Goal 1: Protect and improve habitats that are important to species at risk, priority bird species and other wildlife.</p> <p>Threats and challenges:</p> <ul style="list-style-type: none"> • Wildlife diseases • Habitat degradation • Impact of predators • Human impact • Accidental spills • Invasive plant species 	<p>Objective 1.1: Protect colonial seabird populations by limiting natural pressures linked to wildlife diseases, abundant animal populations and terrestrial predators.</p>	<ul style="list-style-type: none"> • Continue to monitor the estuary's seabird populations every five years and implement the protection measures recommended by ECCC CWS specialists and external scientists. (1) • Monitor if possible the Common Eider during down collection and attempt to understand the factors that have contributed to the decline in the population through specific studies. (1) • Monitor the health of bird populations, implement the emergency plan in the event of mass bird mortality and respond quickly with collaborators in the event of disease. (1) • Maintain an optimal forest habitat for nesting of the Common Eider on Île Bicquette. (1) • In collaboration with others, evaluate the status of abundant bird populations (cormorants, gulls) as needed and their impacts on habitats, and take appropriate management actions. (3) • Conduct annual monitoring of predators and control them if necessary (especially the fox) to avoid losses of birds in major seabird colonies. (1)
	<p>Objective 1.2: Conserve the NWA's seabird colonies, and waterfowl and shorebird staging areas by limiting the effects of human activities (e.g. disturbances, accidental spills).</p>	<ul style="list-style-type: none"> • Produce and post or distribute information posters and pamphlets intended for kayakers and boaters on the piers in Rivière-du-Loup, Kamouraska, Saint-Fabien and Saint-Siméon to limit disturbances to seabirds, waterfowl and shorebirds in the NWA. (2) • Map the NWA's sensitive wildlife habitats and participate in implementing the ECCC CWS emergency response plan and in disseminating it to collaborators in order to protect wildlife and sensitive habitats. (1)
	<p>Objective 1.3: Implement the management and recovery measures required to protect bird species at risk and priority bird species.</p>	<ul style="list-style-type: none"> • Protect the identified critical habitat for species at risk and habitats that are important for priority bird species. (2) • Apply the measures suggested in the planning documents for the recovery of bird species at risk that are common in the NWA or that nest there (e.g. Peregrine Falcon). (2)
	<p>Objective 1.4: Prevent the overgrowth of invasive plant species.</p>	<ul style="list-style-type: none"> • Assess the impact of invasive plant species. (2) • Implement measures to restrict the expansion of these species if needed. (2)
	<p>Objective 1.5: Evaluate island erosion and limit its impact.</p>	<ul style="list-style-type: none"> • Assess erosion on the shores of the NWA's low-lying islands. (2) • Assess the ecological relevance and feasibility of erosion control measures. (3)

Table 6: Management approaches for Estuary Islands NWA (continued)

Goals	Objectives	Actions (Priority Level ¹)
Goal 2: Reduce the impact of human activities on the NWA. Threats and challenges: <ul style="list-style-type: none"> • Human impact • Facility, infrastructure and land maintenance 	Objective 2.1: Adequately post the NWA's boundaries in the field to protect wildlife and plants from the impacts of human activities (disturbances from boaters, hunting, etc.).	<ul style="list-style-type: none"> • Improve signage on the NWA's boundaries by making signs more visible and by installing them in strategic locations throughout the NWA to prevent people from landing on the islands. (1)
	Objective 2.2: Promote local and public awareness of the NWA's mission and applicable regulations to reduce instances of non-compliance.	<ul style="list-style-type: none"> • Install regulatory signage inside and outside the NWA. (1) • Issue public notices of the NWA's regulations in newspapers and share general information about it (in newspapers, magazines, pamphlets, etc.). (1) • Share information on the NWA's importance with regional organizations, local communities and the public, in cooperation with different stakeholders. (3) • Target the necessity of monitoring, bird and wildlife protection, and law enforcement in the area with regards to prohibiting access and hunting during critical periods. (1) • Support area monitoring by Wildlife Enforcement Directorate officers, provincial conservation officers and private collaborators such as Duvetnor and the SPEE. (1)
	Objective 2.3: Communicate concerns to managers of adjacent lands and waters regarding resource development near the NWA.	<ul style="list-style-type: none"> • Participate in regional round tables dealing with resource development around the NWA, including fishing, seaweed harvesting and oil and gas exploration and production. (3)
	Objective 2.4: Ensure that Environment and Climate Change Canada's facilities, infrastructure and lands (especially on Île Bicquette and Le Pot du Phare) are in good and safe condition.	<ul style="list-style-type: none"> • Assess and perform required maintenance. (1) • Characterize and assess the level of risk on contaminated sites, including on Île Bicquette, La Grande Île, Le Long Pèlerin and Île de la Providence, and rehabilitate them, if required. (2)

Table 6: Management approaches for Estuary Islands NWA (continued)

Goals	Objectives	Actions (Level of priority ¹)
Goal 3: Consolidate the NWA's land holdings and promote the conservation of natural habitats on adjacent islands. Threats and challenges: <ul style="list-style-type: none"> • NWA fragmentation 	Objective 3.1: Incorporate into the NWA adjacent, federally-owned lands with significant conservation value.	<ul style="list-style-type: none"> • Evaluate whether federally-owned lands could be integrated into the NWA. (2) • Take the steps required to integrate these lands into the NWA. (3)
	Objective 3.2: Determine which of the islands located near the NWA have strong conservation potential and develop an acquisition strategy as required.	<ul style="list-style-type: none"> • Conduct an analysis of the ecological value and conservation potential of the islands as well as the portions of islands adjacent to the NWA. (2) • When possible, acquire lands with strong conservation potential. (3)
	Objective 3.3: Encourage use of the islands and lands adjacent to the NWA in a way that is compatible with the NWA's conservation mandate.	<ul style="list-style-type: none"> • Make adjacent land owners aware of the importance of protecting the habitat, especially in the Île Bicquette and Îles de Kamouraska sectors. (2) • Work in cooperation with other island owners and users to conserve the estuary's islands. (3)
Goal 4: Ensure ecological monitoring of the NWA and improve knowledge about its wildlife and their habitats. Threats and challenges: <ul style="list-style-type: none"> • Gaps in scientific knowledge 	Objective 4.1: Implement an ecological monitoring plan.	<ul style="list-style-type: none"> • Determine the indicators and follow-up methodologies for an ecological monitoring plan. (1) • Implement the ecological monitoring plan. (1)
	Objective 4.2: Determine gaps in scientific knowledge and address those that are identified as priorities.	<ul style="list-style-type: none"> • Determine the gaps in scientific knowledge. (2) • Support the acquisition of knowledge of prime importance on priority birds, species at risk, stresses, sensitive habitats and threats to the NWA based on the knowledge acquisition plan. (3) • Use various existing data sources (e.g. ÉPOQ, SOS-POP, NGOs, ornithology clubs, research) to improve scientific knowledge. (3)

¹ Level of Priority: 1 (from 0 to 3 years); 2 (from 4 to 6 years); 3 (from 7 to 10 years)

Note: The levels of priority assigned to the actions relate to the implementation schedule and not the significance for resource conservation. They can change depending on the context and the available resources.

5 MANAGEMENT APPROACHES

This section summarizes the approaches and actions that are presented in Table 6 and are likely to be used in managing Estuary Islands NWA. Management actions will be indicated in more detail during the annual planning process and will be implemented depending on available financial and human resources and according to the approaches described below.

5.1 HABITAT MANAGEMENT

Habitat management will be aimed first at protecting plant communities. In some cases, restoring disturbed habitats (e.g. by the Double-crested Cormorant, the Snowshoe Hare and insects) or rehabilitating contaminated lands can be carried out to preserve natural environments, to mitigate the impacts of disturbances on plant biomass and plant succession in terrestrial environments, or to re-establish seabird populations (e.g. reforestation on Île Bicquette for the Common Eider population) and species at risk.

Integrating federal lands without NWA status as well as other islands of conservation significance into the current NWA could also help preserve wildlife habitats.

5.2 WILDLIFE MANAGEMENT

Wildlife management will be based on the knowledge gained during surveys that have been completed to date. This knowledge will help in taking stock of many components of the NWA's biodiversity and in making informed decisions.

Direct predator control could continue on some islands to protect seabird colonies, while taking into consideration the guidelines in Environment and Climate Change Canada's "Predator Management Policy". If needed, specific management measures will be taken (e.g. installing nest boxes, protecting nesting areas) in order to protect seabird populations (including the Common Eider) whose numbers may be modified due to wildlife diseases, predation, accidental spills and other factors.

To protect species at risk, the presence and abundance of the Peregrine Falcon and the Red Knot will need to be better documented. The recovery strategies for these species at risk will guide conservation activities in the NWA. Collaboration with other specialists, departments and universities will also be favoured in order to promote learning and the protection of these species.

5.3 MONITORING

An ecological monitoring plan for the NWA will be developed in the next five years to evaluate the NWA's health and to gather information that will help in making management decisions. This plan will be based on biological monitoring carried out by Environment and Climate Change Canada's CWS and could include collaboration with regional and provincial stakeholders. Ecological monitoring efforts will deal mostly with habitats (e.g. forests stressed by cormorant droppings, habitat dynamics), species at risk, species representative of the NWA, and ecological and human stresses affecting the area. Efforts may also be made to standardize certain methods used to survey nesting seabirds in the NWA. In addition, some specific monitoring programs conducted by the CWS or external organizations, such as the CWS' seabird inventory program and the emergency plan in the event of mass bird mortality, will be drawn on in the management of seabirds in the NWA.

5.4 RESEARCH

Knowledge acquisition and research needs have been established for several groups of species and various management issues in the *Conservation Plan of the Îles de l'estuaire National Wildlife Area* (CWS, 2003). Since then, various surveys have been performed in the NWA and have helped to address certain gaps. These priorities include better documenting the presence of rare or at-risk plant species, certain management problems (e.g. diseases affecting the Common Eider, the progressive reduction of its population over the past 25 years and the status of the forest on Île Bicquette) as well as ecological stresses and the impacts of human activities (e.g. waterfowl hunting, the extent of marine organism harvesting around the NWA and the presence of invasive species). In addition, it would be helpful to improve knowledge about certain elements of the NWA's biological diversity (e.g. insects, algae, wetland plants, amphibians, reptiles, plant life, habitats, shorebirds as well as the nesting and breeding of waterfowl).

Permits can be issued for research activities that are aligned with the priorities identified in the management plan and for scientific activities such as surveys, habitat restoration and enhancement works.

For permission to conduct research in Estuary Islands NWA and for instructions on the guidelines for a research proposal, please contact:

National Wildlife Area – Research Request
Environment and Climate Change Canada
Canadian Wildlife Service
Québec Region
801-1550 D'Estimauville Avenue
Québec QC G1J 0C3
Email: ec.permisscfquebec-cwsquebecpermit.ec@canada.ca

5.5 PUBLIC INFORMATION AND OUTREACH

Estuary Islands NWA is not open to the public, except Le Pot du Phare, where Environment and Climate Change Canada allows public outreach activities about conserving natural environments. These activities are currently carried out by Société Duvetnor Ltée, a local conservation organization, under a commercial permit issued by the CWS. Various public education and outreach measures are planned to prevent boaters and kayakers from landing in the NWA and to limit disturbances to wildlife. These measures include improving signage on the NWA's boundaries as well as distributing pamphlets, installing posters and signs, or publishing notices that present general information about the NWA and its applicable regulations.

6 AUTHORIZATIONS AND PROHIBITIONS

In the interest of wildlife and their environment, human activities are minimized and controlled in NWAs through the implementation of the *Wildlife Area Regulations*. These regulations set out the activities that are prohibited (subsection 3(1)) in NWAs and provide mechanisms for the Minister of Environment and Climate Change to authorize certain activities that are otherwise considered prohibited. The regulations also provide the authority for the Minister to prohibit entry into NWAs.

Activities within an NWA are authorized where notices have been posted at the entrance to or along the boundaries of the NWA or when notices have been published in local newspapers. All activities in an NWA are prohibited unless a notice has been posted or published authorizing the activity to take place (see example in Appendix I). However, in addition to notices, certain activities may be authorized by obtaining a permit from the Minister of Environment and Climate Change.

6.1 PROHIBITION OF ENTRY

Under the *Wildlife Area Regulations*, the Minister may publish a notice in a local newspaper or post notices at the entrance of any NWA or on the boundary of any part thereof prohibiting entry to any NWA or its boundaries. These notices can be issued when the Minister is of the opinion that entry is a public health and safety concern or may disturb wildlife and their habitats.

Owing to the fragility of the area and the species that are found there, **access to Estuary Islands NWA is prohibited at all times, except on Le Pot du Phare, unless authorized by a permit issued by the Minister.**

Notices prohibiting entry appear on signage in various locations in the area.

6.2 AUTHORIZED ACTIVITIES

Public access to Estuary Islands NWA is permitted only on Le Pot du Phare, but only after the seabird nesting period, which ranges from mid-July to mid-October, and is conditional on using the transportation service provided by the agency authorized by Environment and Climate Change Canada. Visitors must comply with all other restrictions unless they hold a permit issued by the Minister.

The following activities are authorized only on Le Pot du Phare: accommodating visitors, nature interpretation and observation, hiking, photography, and picnicking in designated locations. These activities are permitted only on Le Pot du Phare trails and at the facilities and

infrastructure provided for that purpose, such as look-outs and service areas, and only during authorized periods. In addition, eiderdown collection by certain conservation organizations is authorized under a commercial permit issued by Environment and Climate Change Canada.

Other Activities

Any activity, other than those permitted on Le Pot du Phare, is prohibited in the NWA, including hunting, camping, hiking, photography and picnicking.

A notice specifying the types of activities permitted in Estuary Islands NWA was published in local newspapers in 2016 (see Appendix I).

Note: If there is a discrepancy between the information presented in this document and the notice, the notice prevails as it is the legal instrument authorizing the activity.

6.3 AUTHORIZATIONS

Permits and notices authorizing an activity may be issued if the Minister is of the opinion that the activity is scientific research relating to wildlife or habitat conservation; or the activity benefits wildlife and their habitats or will contribute to wildlife conservation; or the activity is not inconsistent with the purpose for which the NWA was established and is consistent with this management plan.

The Minister may also add terms and conditions to permits in order to minimize the impact of an activity on wildlife and wildlife habitats.

All requests for permits or authorizations must be made (in writing or online) to the following address:

National Wildlife Area – Permit Request
Environment and Climate Change Canada
Canadian Wildlife Service
Quebec Region
801-1550 D'Estimauville Avenue
Québec QC G1J 0C3
E-mail: ec.permisscfquebec-cwsquebecpermit.ec@canada.ca

For further information, please consult the *Policy when Considering Permitting or Authorizing Prohibited Activities in Protected Areas Designated under the Canada Wildlife Act and Migratory Bird Convention Act, 1994* (December 2011). This Environment and Climate Change Canada policy document is available on the Protected Areas website at www.ec.gc.ca/ap-pa.

6.4 EXCEPTIONS

The following activities do not require a permit or authorization:

- Activities related to public safety, public health or national security, that are authorized under another act of Parliament or activities authorized under the *Health of Animals Act* and the *Plant Protection Act* to protect animal or plant health;
- Activities related to routine maintenance of NWAs, the implementation of management plans and enforcement activities conducted by an Environment and Climate Change Canada officer or employee.

6.5 OTHER FEDERAL AND PROVINCIAL AUTHORIZATIONS

Depending on the type of activity, other federal or provincial permits or authorizations may be required to undertake an activity in this NWA. For more information, please contact the regional office of the appropriate federal or provincial authority.

7 HEALTH AND SAFETY

All reasonable efforts will be made to protect the health and safety of the public including adequately informing visitors of any known or anticipated hazards or risks. Environment and Climate Change Canada staff will also take all reasonable and necessary precautions to protect their own health and safety as well as that of their co-workers. However, visitors (including researchers and contractors) must make all reasonable efforts to inform themselves of risks and hazards, and must be prepared and self-sufficient. Natural areas contain some inherent dangers and visitors must take proper precautions to insure their own security, recognizing that Environment and Climate Change Canada staff neither regularly patrol nor offer visitor safety services in this NWA.

Environment and Climate Change Canada intends to develop a public safety plan in order to limit the risks of accident and insure public safety in this NWA.

In the case of environmental emergencies, please contact the National Environmental Emergencies Centre at the following telephone numbers:

514-283-2333 or 1-866-283-2333

Incidents or emergency situations can be reported to:

- Environment and Climate Change Canada: 1-800-668-6767 or ec.enviroinfo.ec@info.ec.gc.ca
- Marine salvage (Canadian Coast Guard): 1-800-463-4393/cellular: *16
- Sûreté du Québec (Police): 310-4141/cellular: *4141
- Sécurité civile (Civil Security): 1-866-776-8345/cellular: 911
- SOS-Braconnage (Poaching): 1-800-463-2191
- Environmental emergencies: Environment and Climate Change Canada : 1-866-283-2333 or Ministère du Développement durable, de l'Environnement et de la Lutte contre les changements climatiques: 1-866-694-5454
- Forest fire (SOPFEU): 1-800-463-FEUX (3389)
- Local authorities (Police or Fire Department): 911

8 ENFORCEMENT

The management of National Wildlife Areas is based on three acts and the regulations thereunder:

- the *Migratory Birds Convention Act, 1994* and the *Migratory Birds Regulations*;
- the *Canadian Wildlife Act* and the *Wildlife Area Regulations*;
- the *Species at Risk Act*.

Environment and Climate Change Canada's wildlife enforcement officers are mandated to monitor compliance with the acts and regulations on an ongoing basis and to conduct investigations as required.

Here are some examples of activities that, if undertaken without authorization in NWAs, may constitute offences:

- Accessing the site;
- Destroying or disturbing migratory birds, their nests or eggs;
- Possessing a weapon or other instrument that could be used for hunting;
- Picnicking, camping or engaging in any other recreational activity;
- Lighting a fire;
- Removing or damaging any natural artefact, building, fence, poster, sign or other structure;
- Dumping or depositing waste or substances likely to reduce the quality of the natural environment;
- Letting a pet run free.

9 PLAN IMPLEMENTATION

The management plan will be implemented over a 10-year period. Annual work plans will be developed in accordance with priorities and budgets. Depending on available resources and opportunities, some actions may be accelerated, postponed or cancelled. Environment and Climate Change Canada will follow an adaptive management approach. The implementation of the plan will be evaluated five years after its publication, on the basis of the actions identified in Table 6.

10 COLLABORATORS

Collaborations with local agencies and organizations will be favoured for promoting the protection and conservation of wildlife and their habitats in Estuary Islands NWA. Cooperating with local organizations involved in conservation of the area, such as Société Duvetnor Ltée and the Société protectrice des eiders de l'estuaire (SPEE), is essential for the preservation of this isolated protected area in the middle of the St. Lawrence and must continue. Collaborations may also be developed or pursued with universities and research centres to fill scientific knowledge gaps, with the province for the implementation of species-at-risk recovery measures, especially those under provincial jurisdiction, and with non-governmental organizations and municipal authorities to educate the public about the NWA's conservation objectives.

The following is a list of the main organizations that may collaborate in the mission and activities of Estuary Islands NWA.

CEGEP de Rimouski
60 De l'Évêché Street West
Rimouski QC G5L 4H6
Telephone: 418-723-1880
Toll free number: 1-800-463-0617
Fax: 418-724-4961

City of Rimouski
205 De la Cathédrale Avenue
P.O. Box 710
Rimouski QC G5L 7C7
Telephone: 418-724-3171
Fax: 418-724-3183
Email: direction.generale@ville.rimouski.qc.ca

Club des ornithologues du Bas-Saint-Laurent (COBSL)
P.O. Box 66
Rimouski QC G5L 7B7
Email: cobsl@globetrotter.net

Comité ZIP du Sud-de-l'Estuaire (ZIP Committee of Sud-de-l'Estuaire)

88 Saint-Germain Street West, Suite 101

Rimouski QC G5L 4B5

Telephone: 418-722-8833

Fax: 418-722-8831

Email: zipse@globetrotter.net

Essipit Innu First Nation Council (Conseil de la Première Nation des Innus Essipit)

32, De la Réserve Street

Les Escoumins QC G0T 1K0

Telephone: 418-233-2509 poste 281

Fax: 418-233-2888

Email : aimun@essipit.com

Fisheries and Oceans Canada

104 Dalhousie Street

Québec QC G1K 7Y7

Telephone: 418-648-2239

Fax: 418-648-4758

Email: info@dfo-mpo.qc.ca (Ottawa)

Ministère du Développement durable, de l'Environnement et de la Lutte contre les changements climatiques (MDDELCC) (Quebec Ministry of Sustainable Development, Environment and Fight against Climate Change)

Direction générale de l'analyse et de l'expertise régionales et Direction régionale (Analysis and Regional Expertise Branch and Regional Branch)

212 Belzile Avenue

Rimouski QC G5L 3C3

Telephone: 418-727-3511

Fax: 418-727-3849

Email: bas-saint-laurent@mddelcc.gouv.qc.ca

Ministère des Forêts, de la Faune et des Parcs (MFFP)
Direction de la protection de la faune du Bas-Saint-Laurent et de la Gaspésie–Îles-de-la-Madeleine (Bas-Saint-Laurent and Îles-de-la-Madeleine Wildlife Protection Branch)
92 2nd Street West, Suite 207
Rimouski QC G5L 8B3
Telephone: 418-727-3710
Fax: 418-727-3735

Pointe-au-Père Local Office
365 Sainte-Anne Boulevard, Suite 1
Rimouski QC G5M 1E8
Telephone: 418-727-3516
Fax: 418-727-3773

S.O.S. Braconnage (Poaching)
Telephone : 1 800 463-2191
Email: centralesos@mffp.gouv.qc.ca

Ministère de l'Énergie et des Ressources naturelles (MERN) (Quebec Ministry of Energy and Natural Resources)
92 2nd Street West, Suite 207
Rimouski QC G5L 8B3
Telephone: 1-844-282-8277
Email: droit.terre.publique@mern.gouv.qc.ca

Municipality of Kamouraska
67 Morel Avenue
Kamouraska QC GOL 1MO
Telephone: 418-492-6523
Fax: 418-492-9789
Email: mychelle.levesque@kamouraska.ca (city manager)

Municipality of Saint-André
122A Principale Street
Saint-André QC G0L 2H0
Telephone: 418-493-2085
Email: munand@bellnet.ca

Nature Conservancy of Canada (NCC)
55 Mont-Royal Avenue West, Suite 1000
Montreal QC H2T 2S6
Telephone: 514-876-1606
Toll free number: 1-877-876-5444
Fax: 514-876-7901
Email: quebec@conservationdelanature.ca

Organisme des bassins versants du Nord-Est du Bas-Saint-Laurent (OBVNEBSL)
23 De l'Évêché Street West, Suite 200
Rimouski QC G5L 4H4
Telephone: 418-724-5154 ext. 219
Fax: 418-725-4567
Email: direction@obv.nordestbsl.org

Regional County Municipality of Kamouraska (all islands except Bicquette)
425 Patry Avenue
Saint-Pascal QC G0L 3Y0
Telephone: 418-492-1660
Fax: 418-492-2220
Email: info@mrckamouraska.com

Regional County Municipality of Rimouski-Neigette (Bicquette Island)
23 De l'Évêché Street West, Suite 200
Rimouski QC G5L 4H4
Telephone: 418-724-5154
Fax: 418-725-4567
Email: administration@mrkrimouskineigette.qc.ca

Société Duvetnor Ltée
200 Hayward Street
P.O. Box 305
Rivière-du-Loup QC G5R 3Y9
Telephone: 418-867-1660
Fax: 418-867-3639
Email: info@duvetnor.com

Société protectrice des eiders de l'estuaire (SPEE)
181 Saint-Paul Street
Le Bic QC G0L 1B0
Telephone: 418-736-4975
Email: spee@globetrotter.net

Université du Québec à Montréal – Département des sciences biologiques
141 Président-Kennedy
P. O. Box 8888, succ. Centre-ville
Montréal QC H3C 3P8
Telephone 514-987-4118
Fax 514-987-4647
Email: dept-biologie@uqam.ca

Université du Québec à Rimouski
300 Allée des Ursulines
P. O Box 3300, succ. A
Rimouski QC G5L 3A1
Telephone: 418-723-1986
Toll free number: 1-800-511-3382
Fax: 418-724-1525
Email: uqar@uqar.ca

11 REFERENCES

- AECOM Tecsalt Inc. 2010. Évaluation environnementale stratégique de la mise en valeur des hydrocarbures dans le bassin de l'estuaire maritime et du nord-ouest du golfe du Saint-Laurent. 05-19255 – Preliminary report supporting the consultations, July 2010, [www.ees.gouv.qc.ca/documents/ees_preliminaire_pour_consultation.pdf] (consulted February 28, 2012).
- Aubry, Y. and R. Cotter. 2007. Québec Shorebird Conservation Plan. Environment Canada, Canadian Wildlife Service, Quebec Region, Sainte-Foy, xvi + 203 pp.
- Bédard, J. 2010. Plan de gestion de la Réserve nationale de faune des îles de l'estuaire. Proposal. Environment Canada, Canadian Wildlife Service, Quebec Region, 93 pp. plus appendices.
- Bédard, J., A. Nadeau and M. Lepage. 1995. "Double-Crested Cormorant Culling In the St. Lawrence River Estuary". Colonial Waterbirds, vol. 18 (Special Publication 1), pp. 78-85.
- BIOMQ (Banque informatisée des oiseaux marins du Québec). 2015. Database, Canadian Wildlife Service, Environment Canada, Quebec Region (updated April 22, 2015).
- Brouillet, L., F. Coursol, S.J. Meades, M. Favreau, M. Anions, P. Bélisle and P. Desmet. 2010+. VASCAN, Database of Vascular Plants of Canada. [<http://data.canadensys.net/vscan/>] (consulted March 15, 2013 for plant names).
- CDPNQ (Centre de données sur le patrimoine naturel du Québec). 2012. Excerpts from the data system for the Îles de l'Estuaire National Wildlife Area. Ministère du Développement durable, de l'Environnement, de la Faune et des Parcs and Ministère des Ressources naturelles, Quebec.
- Chapdelaine, G. and J. Bédard. 1995. "Recent Changes in the Abundance and Distribution of the Double-crested Cormorant in the St. Lawrence River, Estuary and Gulf, Quebec, 1978-1990". Colonial Waterbirds, vol. 18 (Special Publication 1), pp. 70-77.
- Chapdelaine, G. and J.-F. Rail. 2002. Québec's Waterbird Conservation Plan. 1st draft. Environment Canada, Canadian Wildlife Service, Quebec Region, unpublished document, 80 pp. plus appendices.
- COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2014. [www.cosewic.gc.ca; research on wildlife species – status] (Last update on April 1st, 2014 for the species checked; consulted October 2, 2014).
- CWS (Canadian Wildlife Service). 2003. Conservation Plan of the Îles de l'estuaire National Wildlife Area. Environment Canada, Canadian Wildlife Service, Quebec Region, 76 pp. plus appendices.

- Dionne, J.-C. 1977. "La mer de Goldthwait au Québec". *Géographie physique et Quaternaire*, vol. 31, nos 1-2, pp. 61-80.
- Duvelnor (Société Duvelnor Ltée). 2015. Plan de protection – Réserve nationale de faune des Îles-de-l'Estuaire, Rapport final (période du 1^{er} mai au 20 octobre 2015). Unpublished report presented to the Canadian Wildlife Service, 19 p.
- Environment Canada. 2013a. Geographic Information System for the National Wildlife Areas of Quebec (GIS-NWA) [Access 14.0 and ArcGIS 10.0 data, unpublished], prepared by the Canadian Wildlife Service, Quebec Region (updated October 2013).
- Environment Canada. 2013b. Bird Conservation Strategy for Bird Conservation Region 14 in Quebec Region – Atlantic Northern Forest. Canadian Wildlife Service, Environment Canada, Quebec, 159 pages plus appendices.
- Environment Canada. 2005. Protected Areas Manual (Draft). Appendix 5: Selection Criteria for Candidate Areas – National Wildlife Area and Appendix 8: Policy on Establishing and Managing EC's Protected Areas (protected area classification system).
- Federal Heritage Buildings Review Office (FHBRO). 2016. http://www.pc.gc.ca/apps/dfhd/page_fhbros_eng.aspx?id=3093; http://www.pc.gc.ca/apps/dfhd/page_fhbros_eng.aspx?id=7484 (Accessed December 2016.)
- Fortin, J.-C. (INRS-Urbanisation, Culture et Société). 2003. "En marge de l'aire seigneuriale". In www.encyclobec.ca/main.php?docid=290, September 25, 2003 article (consulted March 6, 2013).
- Fortin, J.C., A. Lechasseur, Y. Morin, F. Harvey, J. Lemay and Y. Tremblay. 1993. Histoire du Bas-Saint-Laurent. Institut québécois de recherche sur la culture, Collection Les régions du Québec (5), Quebec, 860 pp.
- IUCN (International Union for the Conservation of Nature; Dudley, N., Publisher). 2008. Guidelines for Applying Protected Area Management Categories. Gland, Switzerland: IUCN. x +96 pp.
- Joint Working Group on the Management of the Common Eider. 2004. Quebec Management Plan for the Common Eider (*Somateria mollissima dresseri*). Special publication of the Joint Working Group on the Management of the Common Eider, Quebec, 44 pp.
- Labrecque, S. and B. Jobin. 2012. Dynamique des habitats et des pressions périphériques dans les RNF du Québec – volet 2. Environment Canada, Canadian Wildlife Service, Quebec Region, 88 pp.
- Lepage, C. and R. Cotter. 2013. Long-tailed duck, pp. 175-180 in Lepage, C. and D. Bordage eds. Status of Quebec Waterfowl Populations, 2009. Technical Report Series No. 525, Canadian Wildlife Service, Environment Canada, Quebec Region, Quebec City, 243 pp.

- Lepage, C. and J.-P.L. Savard. 2013. Surf Scoter, pp 160-167 in Lepage, C. and D. Bordage, eds. Status of Quebec Waterfowl Populations, 2009. Technical Report Series No. 525, Canadian Wildlife Service, Environment Canada, Quebec Region, Quebec City, 243 pp.
- MVFN (Maliseets of Viger First Nation). 2014. [www.malecites.ca] ("Histoire" section, accessed March 3, 2014).
- MFFP (Ministère des Forêts, de la Faune et des Parcs, gouvernement du Québec). 2016. Liste des espèces de la faune désignées menacées ou vulnérables au Québec; Liste des espèces de la faune susceptibles d'être désignées menacées ou vulnérables au Québec. [www3.mffp.gouv.qc.ca/faune/especes/menacees/liste.asp] (Accessed September 30, 2016).
- MFFP (Ministère des Forêts, de la Faune et des Parcs, gouvernement du Québec). 2015. Liste de la faune vertébrée du Québec. [www3.mffp.gouv.qc.ca/faune/verteebree/recherche/index.asp] (Accessed November 2015).
- Morisset, P. 2010. La flore vasculaire de l'île Blanche. Environment Canada. Contract no. K2C55-09-2540, 12 pp.
- Nadeau, A., Bédard, J. and Y. Turcotte. 2009. La biodiversité dans la réserve nationale de faune des îles de l'estuaire. Société Duvetnor Ltée for the Canadian Wildlife Service, Quebec Region. Contract no. K4C25-08-7808, 22 pp. plus appendices.
- Parks Canada. 2016. [<http://www.pc.gc.ca/eng/progs/lhn-nhs/pp-hl/page01.aspx>] (Accessed December 2016.)
- Robert, M., R. Benoit, C. Marcotte, J.-P.L. Savard, D. Bordage and D. Bourget. 2003. Le Garrot d'Islande dans l'estuaire du Saint-Laurent : calendrier de présence annuelle, répartition, abondance, âge-ratio et sex-ratio. Series of technical reports no. 398, Canadian Wildlife Service, Quebec Region, Environment Canada, Sainte-Foy, Quebec, 129 pp.
- Savard, J.-P.L. and C. Lepage. 2013. Common Eider, subspecies *dresseri*, pp 146-150 in Lepage, C. and D. Bordage, eds. Status of Quebec Waterfowl Populations, 2009. Technical Report Series No. 525, Canadian Wildlife Service, Environment Canada, Quebec Region, Quebec City, 243 pp.
- Species at Risk Public Registry (Government of Canada). 2014. [www.registrelep-sararegistry.gc.ca/; species research] (Modified August 30, 2012 [Goldeneye and Knot] and June 5, 2014 [Falcon], consulted October 2, 2014).
- SPEE (Société protectrice des eiders de l'estuaire). 1998. Rapport annuel des activités de conservation - île Bicquette (period of May 1, 1997 to April 30, 1998). Unpublished report submitted to the Canadian Wildlife Service, 2 pp.

APPENDIX I: ENVIRONMENT AND CLIMATE CHANGE CANADA NOTICE PUBLISHED IN 2016

Environment and Climate Change Canada would like to inform the public that the Estuary Islands National Wildlife Area (NWA) is a protected area as described in Schedule I of the *Wildlife Area Regulations*. The NWA is a cluster of about a dozen islands or parts of islands, including the mudflats and reefs that fringe some of them. The islands are scattered in the St. Lawrence Estuary in the area encompassing the municipalities of Rimouski, Saint-André and Kamouraska. Since its creation in 1986, the NWA has protected nesting sites for migratory birds, particularly seabird colonies.

The Department would also like to inform the public that they must follow certain rules dictated by the *Canada Wildlife Act* and its regulations. Any person who fails to follow these rules or comply with the applicable laws may be fined or prosecuted.

Under subsection 3(1) of the *Wildlife Area Regulations*, no person may do any of the following things in a wildlife area, unless the person has a permit issued under section 4 of the regulations:

- swim, picnic, camp or carry on any other unauthorized recreational activity or light a fire;
- operate a conveyance;
- damage, destroy or remove a plant;
- disturb or remove any soil, sand, gravel or other material;
- remove, damage or destroy any natural object, poster, sign or other structure;
- dump or deposit any waste material;
- hunt or fish;
- allow any domestic animal to run at large; or
- carry on any commercial activity.

Access to the Estuary Islands National Wildlife Area is prohibited, except under the authority of a permit issued under section 4 of the *Wildlife Area Regulations*. For information on all the applicable regulations, please refer to the *Canada Wildlife Act*, the *Wildlife Area Regulations*, the *Migratory Birds Convention Act, 1994* and the *Species at Risk Act* on the www.ec.gc.ca Web site (Acts and Regulations section).

To file a complaint or report a possible contravention of the regulations, please contact Environment and Climate Change Canada by phone at 1-800-668-6767 or by email at ec.enviroinfo.ec@canada.ca.

Nothing in this notice shall affect aboriginal or treaty rights.