



Natural Resources
Canada

Ressources naturelles
Canada

CANADIAN GEOSCIENCE MAP 311

BEDROCK GEOLOGY

LAC MAUNOIR

Northwest Territories



Map Information
Document



Geological Survey of Canada
Canadian Geoscience Maps

2018

Canada



MAP NUMBER

Natural Resources Canada, Geological Survey of Canada
Canadian Geoscience Map 311

TITLE

Bedrock geology, Lac Maunoir, Northwest Territories

SCALE

1:250 000

CATALOGUE INFORMATION

Catalogue No. M183-1/311-2017E-PDF

ISBN 978-0-660-07652-2

<https://doi.org/10.4095/306205>

COPYRIGHT

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources, 2018

Information contained in this publication or product may be reproduced, in part or in whole, and by any means, for personal or public non-commercial purposes, without charge or further permission, unless otherwise specified.

You are asked to:

- exercise due diligence in ensuring the accuracy of the materials reproduced;
- indicate the complete title of the materials reproduced, and the name of the author organization; and
- indicate that the reproduction is a copy of an official work that is published by Natural Resources Canada (NRCan) and that the reproduction has not been produced in affiliation with, or with the endorsement of, NRCan.

Commercial reproduction and distribution is prohibited except with written permission from NRCan. For more information, contact NRCan at nrcan.copyrightdroitdauteur.nrcan@canada.ca.

RECOMMENDED CITATION

Fallas, K.M., 2018. Bedrock geology, Lac Maunoir, Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 311, scale 1:250 000.

<https://doi.org/10.4095/306205>

ABSTRACT

The Lac Maunoir map area (NTS 96-N) lies within the Colville Hills region of the Northwest Territories. Sparse bedrock exposures in the area include carbonate and siliciclastic strata ranging from Cambrian to Paleogene. These strata were deformed in

the Cretaceous to Eocene by folding and contractional faulting associated with Cordilleran deformation. A pre-Cordilleran set of approximately north-trending extensional faults are preserved within subsurface Proterozoic and Cambrian strata, and were locally reactivated by Cordilleran deformation. A major unconformity between Devonian and Cretaceous strata is marked by tilted Paleozoic strata beneath the Cretaceous, and the absence of Devonian strata in the eastern part of the map area. Oil has been reported from petroleum exploration wells drilled into Mount Clark Formation (Cambrian) sandstone.

RÉSUMÉ

La région cartographique de Lac Maunoir (SNRC 96-N) se situe dans la région des collines Colville des Territoires du Nord-Ouest. Dans la région, des affleurements clairsemés du socle rocheux renferment des strates carbonatées et des strates silicoclastiques rapportées à l'intervalle du Cambrien au Paléogène. Ces strates ont été déformées dans l'intervalle du Crétacé à l'Éocène par des plis et des failles de compression associés à la déformation cordillérienne. Des failles de distension de direction à peu près nord appartenant à un ensemble pré-cordillérien sont conservées en profondeur dans les strates du Protérozoïque et du Cambrien et ont été réactivées localement par la déformation cordillérienne. Une discordance majeure entre les strates du Dévonien et celles du Crétacé est révélée par l'inclinaison des strates du Paléozoïque sous celles du Crétacé ainsi que par l'absence de strates du Dévonien dans la partie est de la région cartographique. On a signalé la présence de pétrole dans des puits d'exploration pétrolière forés dans le grès de la Formation de Mount Clark (Cambrien).

LICENCE AGREEMENT

View the license agreement at

<http://open.canada.ca/en/open-government-licence-canada>

ACCORD DE LICENCE

Voir l'accord de licence à

<http://ouvert.canada.ca/fr/licence-du-gouvernement-ouvert-canada>

SHEET 1 OF 1, BEDROCK GEOLOGY

GENERAL INFORMATION

Author: K.M. Fallas

Geological compilation by K.M. Fallas, 2015–2016

Geology conforms to Bedrock Data Model v. 4.0

Geological field observations by K.M. Fallas, R.B. MacNaughton, and M.J. Sommers, 2015; D.G. Cook and M. Ayling, 1968

Stratigraphic sections measured by E.C. Turner (Laurentian University), 2015; R.W.

Macqueen, 1968

Reflection-seismic data interpreted by B.C. MacLean and K.M. Fallas, 2015. Petroleum exploration well-picks selected by J. Dixon, 2016

Geomatics by K.M. Fallas and D.A. Lemay

Cartography by D.A. Lemay

Initiative of the Geological Survey of Canada, conducted under the auspices of the Mackenzie Project as part of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) program

Logistical support provided by the Polar Continental Shelf Program as part of its mandate to promote scientific research in the Canadian North. PCSP 05415

Map projection Universal Transverse Mercator, zone 10.
North American Datum 1983

Base map at the scale of 1:250 000 from Natural Resources Canada, with modifications.

Elevations in metres above mean sea level

Mean magnetic declination 2018, 21°11'E, decreasing 32.5' annually. Readings vary from 21°25'E in the NW corner to 20°54'E in the SE corner of the map.

This map is not to be used for navigational purposes.

Title photograph: View of Paleogene conglomerate outcrop on west flank of Maunoir Ridge in the Colville Hills, Northwest Territories. Photograph by K.M. Fallas. 2017-044

The Geological Survey of Canada welcomes corrections or additional information from users.

Data may include additional observations not portrayed on this map. See map info document accompanying the downloaded data for more information about this publication.

This publication is available for free download through GEOSCAN (<http://geoscan.nrcan.gc.ca/>).

MAP VIEWING FILES

The published map is distributed as a Portable Document File (PDF), and may contain a subset of the overall geological data for legibility reasons at the publication scale.

DESCRIPTIVE NOTES

Initial bedrock mapping and stratigraphic studies by the Geological Survey of Canada in the Lac Maunoir map area (NTS 96-N) were conducted in 1968 as part of Operation

Norman. This operation led to the release of a report and preliminary map of the area (Aitken and Cook, 1970; Cook and Aitken, 1971). Observations from the 1968 field activities have been incorporated into this compilation along with observations collected in 2015 as part of the Geo-mapping for Energy and Minerals (GEM) Program. Petroleum exploration wells and reflection-seismic data drilled or collected since 1970 have also helped constrain the map interpretation and geological relationships in the subsurface (Fig. 1, 2). Despite significant areas of unconsolidated Quaternary sediment cover, the bedrock units have been interpreted beneath that cover in an attempt to create a seamless bedrock interpretation.

Starting with the oldest bedrock units, changes to the stratigraphic map units from Aitken and Cook (1970) include the subdivision of the obsolete Ronning Group into the Franklin Mountain and Mount Kindle formations (Norford and Macqueen, 1975), separated by an unconformity spanning the Middle Ordovician. Erosion on the sub-Devonian unconformity has removed the Mount Kindle Formation along Maunoir Ridge ('Maunoir Arch' of Fig. 2). Morrow (1991) documents the lateral relationship between Devonian bedded carbonates of the Arnica and Landry formations and brecciated carbonate of the Bear Rock Formation. Limited outcrop exposure combined with irregular brecciation of this interval encouraged the adoption of 'Bear Rock assemblage' in this area to encompass lithologies found in each of these units as well as postulated occurrences of Delorme Group strata (Gouwy et al., 2017). Although common to the south in the Lac des Bois map area (NTS 96-K; Fallas, 2018), karst features within carbonate strata of the above mentioned units are not noticeable at surface.

Low-angle tilting and erosion of the Paleozoic units are evident beneath the sub-Cretaceous unconformity, with Mount Kindle and Bear Rock formations missing beneath the Cretaceous on the east flank of Maunoir Dome and east of Maunoir Ridge ('Maunoir Arch' in Fig. 2). Limited exposure of the Cretaceous units makes it difficult to map formation boundaries; however, some sandstone strata are assigned to Martin House Formation with some confidence (equivalent to Langton Bay Formation of the Anderson Plain; Dixon, 1999). An isolated outcrop of iron-stained conglomerate on the west flank of Maunoir Ridge has been dated as Paleogene (see accompanying digital data for more details). Areas shown as Quaternary sediment on the map include substantial areas of unconsolidated glacial (e.g. hummocky terrain, eskers) and fluvial deposits left behind by the Laurentide ice sheet (Hughes, 1987), but rare bedrock exposures may be present.

Proterozoic deformation of sedimentary strata in this map area is documented from reflection-seismic data (Cook and MacLean, 2004). No dominant structural trend is interpreted from the Proterozoic contractional features. Subsequent to Proterozoic deformation, extensional faults developed in the Cambrian (MacLean, 2011) with a dominant north to northeast trend. During Cordilleran deformation in Cretaceous to Eocene time, the pre-existing structures in the subsurface influenced the location and trend of Cordilleran structures, in some cases through reactivation of older structures. Reactivated structures typically have steeper dips on the fault plane cutting into Proterozoic strata and are therefore shown as reverse faults rather than thrust faults. In contrast, thrust faults show evidence of detachment in evaporite of the Cambrian Saline River Formation on reflection-seismic data.

Petroleum exploration wells in the Lac Maunoir area have targeted Cambrian sandstone of the Mount Clark Formation (Dixon and Stasiuk, 1998). An oil showing has been reported from the Lac Maunoir C-34 well. In agreement with the statement in Cook

and Aitken (1971), 2015 field activities did not observe the presence of any metallic minerals of economic significance in the map area. Deposits of sand and gravel within the Quaternary unit (Smith and Lesk-Winfield, 2010) may be useful for infrastructure development, but more detailed study of the surficial materials found in this area is required to identify suitable deposits.

ACKNOWLEDGMENTS

This work was carried out on lands within the Sahtu Settlement Area as identified in the Sahtu Dene and Métis Comprehensive Land Claim Agreement. The author gratefully acknowledges the gracious welcome and logistical support of the Behdzi Ahda First Nation (Colville Lake, Northwest Territories). Helicopter and fixed-wing transportation was provided by local service providers based in Norman Wells. Additional field assistance was cheerfully provided by B. Manuel, and the geology team was kept safe by the wildlife monitoring of F.J. Barnaby and C. Oudzi. C. Yakeleya reliably ensured that logistics were co-ordinated. The author also thanks L.D. Currie and L.S. Lane for critical reviews of the map.

REFERENCES

- Aitken, J.D. and Cook, D.G., 1970. Geology, Colville Lake and Coppermine, District of Mackenzie; Geological Survey of Canada, Preliminary Map 12-1970, scale 1:500 000. <https://doi.org/10.4095/220386>
- Cook, D.G. and Aitken, J.D., 1971. Geology, Colville Lake map-area and part of Coppermine map-area, Northwest Territories; Geological Survey of Canada, Paper 70-12, 42 p. <https://doi.org/10.4095/102357>
- Cook, D.G. and MacLean, B.C., 2004. Subsurface Proterozoic stratigraphy and tectonics of the western plains of the Northwest Territories; Geological Survey of Canada, Bulletin 575, 91 p. <https://doi.org/10.4095/215739>
- Dixon, J., 1999. Mesozoic-Cenozoic stratigraphy of the northern Interior Plains and plateaux, Northwest Territories; Geological Survey of Canada, Bulletin 536, 56 p. <https://doi.org/10.4095/210800>
- Dixon, J. and Stasiuk, L.D., 1998. Stratigraphy and hydrocarbon potential of Cambrian strata, northern Interior Plains, Northwest Territories; Bulletin of Canadian Petroleum Geology, v. 46, p. 445–470.
- Fallas, K.M., 2018. Bedrock geology, Lac des Bois, Northwest Territories; Geological Survey of Canada, Canadian Geoscience Map 308, scale 1:250 000. <https://doi.org/10.4095/306201>
- Gouwy, S.A., MacNaughton, R.B., and Fallas, K.M., 2017. New conodont data constraining the age of the 'Bear Rock assemblage' in the Colville Hills, Northwest Territories; Geological Survey of Canada, Current Research 2017-3, 11 p. <https://doi.org/10.4095/306171>

Hughes, O.L., 1987. Late Wisconsinan Laurentide glacial limits of northwestern Canada: the Tutsieta Lake and Kelly Lake phases; Geological Survey of Canada, Paper 85-25, 19 p. <https://doi.org/10.4095/122385>

MacLean, B.C., 2011. Tectonic and stratigraphic evolution of the Cambrian basin of northern Northwest Territories; Bulletin of Canadian Petroleum Geology, v 59, p. 172-194. <https://doi.org/10.2113/gscpgbull.59.2.172>

Morrow, D.W., 1991. The Silurian-Devonian sequence in the northern part of the Mackenzie Shelf, Northwest Territories; Geological Survey of Canada, Bulletin 413, 121 p. <https://doi.org/10.4095/132170>

Norford, B.S. and Macqueen, R.W., 1975. Lower Paleozoic Franklin Mountain and Mount Kindle formations, District of Mackenzie: their type sections and regional development; Geological Survey of Canada, Paper 74-34, 37 p. <https://doi.org/10.4095/102525>

Smith, I.R. and Lesk-Winfield, K., 2010. Seismic shothole drillers' log-derived thematic GIS, Northwest Territories and northern Yukon: bedrock outcrop and subcrop, geohazards, and muskeg thickness; Geological Survey of Canada, Open File 6410. <https://doi.org/10.4095/261784>

Geological Survey of Canada Paleontological Reports (available from GSC Calgary):

Bolton, T.E., 1959. Report on fossils from north of Great Bear Lake submitted by J.C. Sproule 1959; Geological Survey of Canada, Paleontological Report O2-B-59-60, 1 p.

Galloway, J.M., 2017. Palynological report on samples submitted by Robert MacNaughton and Karen Fallas, Geological Survey of Canada, Calgary, in support of the GEM Mackenzie Project (various NTS map sheets); Geological Survey of Canada, Paleontological Report JMG-2017-03, 7 p.

Gouwy, S.A., 2016. Report on conodonts and tentaculites from 12 Devonian samples from the Arnica, Landry, Bear Rock and Hare Indian formations, Colville Hills, NWT, NTS 96E, 96K, 96L, 96M, 96N & 97B collected and submitted by Karen Fallas and Robert MacNaughton (GEM-2 Shield to Selwyn) CON. NO. 1807; Geological Survey of Canada, Paleontological Report 2-SAG-2016, 13 p.

Hopkins, W.S., 1969. Palynology report on miscellaneous outcrop samples from the Northwest Territories, collected by D.G. Cook, 1969 (NTS 96E, F, K, N, 106O, P); Geological Survey of Canada, Paleontological Report K-12 WSH 1969, 2 p.

McCracken, A.D., 2016. Report on thirty-two Ordovician and Silurian conodont samples from the Franklin Mountain Formation and Mount Kindle Formation, Colville Hills, Northwest Territories, collected in 2015 by Robert MacNaughton, Karen Fallas (GSC Calgary) and Elizabeth Turner (Laurentian University) as part of GEM-2 Shield to Selwyn, NTS 096K/01, 096K/02, 096K/03, 096K/08, 096K/12, 096N/06, 096N/15, Con.

No. 1807-13 to 1807-44 (LSA 16-3-006); Geological Survey of Canada, Paleontological Report 5-ADM-2016, 15 p.

McIntyre, D.J., 1992. Report on palynological examination of four samples from Maunoir Ridge, Northwest Territories (NTS 96 N/3), Samples submitted by D.G. Cook; Geological Survey of Canada, Paleontological Report 7-DJM-1992, 1 p.

Norford, B.S., 1969. Report on nineteen lots of fossils from northwest District of Mackenzie, collected by J.D. Aitken and D.G. Cook, Operation Norman, 1968, (NTS 96E, F, K, M, N; 97A; 106G, H); Geological Survey of Canada, Paleontological Report C-O 2 BSN 1969, 3 p.

Norford, B.S., 1970. Report on seven lots of fossils from the northern District of Mackenzie submitted by Dr. D.G. Cooke, 1970 (NTS 96E, 96N, 97B); Geological Survey of Canada, Paleontological Report O-4 BSN 1970, 2 p.

ADDITIONAL INFORMATION

The Additional Information folder of this product's digital download contains figures and tables that appear in the map surround as well as additional geological information not depicted on the map, nor this document, nor the geodatabase.

- PDF of each figure/table that appears in the CGM surround.
- Excel file of the Master Legend Table (legend symbols, descriptions, headings, etc.).

AUTHOR CONTACT

Questions, suggestions, and comments regarding the geological information contained in the data sets should be addressed to:

K.M. Fallas
Geological Survey of Canada
3303 33 Street NW
Calgary AB
T2L 2A7
Karen.Fallas@canada.ca

COORDINATE SYSTEM

Projection: Universal Transverse Mercator
Units: metres
Zone: 10
Horizontal Datum: NAD83
Vertical Datum: mean sea level

BOUNDING COORDINATES

Western longitude: 126°00'00"W
Eastern longitude: 124°00'00"W
Northern latitude: 68°00'00"N
Southern latitude: 67°00'00"N

SOFTWARE VERSION

Data has been originally compiled and formatted for use with ArcGIS™ desktop version 10.2.2 developed by ESRI®.

DATA MODEL INFORMATION

Bedrock (Calgary) [remove whole section if not applicable]

Surface bedrock data are organized into feature classes and themes consistent with logical groupings of geological features. All field observation point data are related through the Station_ID property of the Station theme. These feature attribute names and definitions are identical in the shapefiles and the XML files.

Consult PDFs in Data folder for complete description of the feature classes, feature attributes, and attribute domains.

The Bedrock Data Model and the Bedrock Domains documents are intended to describe all bedrock features which may be compiled at the 1:50 000 scale. Therefore, some of the feature classes and feature attributes described in these documents may not be present.