



Natural Resources
Canada

Ressources naturelles
Canada

CANADIAN GEOSCIENCE MAP 323

SURFICIAL GEOLOGY

ABERDEEN LAKE

Nunavut

NTS 66-B

Map Information Document

Preliminary



Geological Survey of Canada
Canadian Geoscience Maps

2017

Canada



MAP NUMBER

Natural Resources Canada, Geological Survey of Canada
Canadian Geoscience Map 323 (Preliminary)

TITLE

Surficial geology, Aberdeen Lake, Nunavut, NTS 66-B

SCALE

1:125 000

CATALOGUE INFORMATION

Catalogue No. M183-1/323-2017E-PDF
ISBN 978-0-660-09001-6
<https://doi.org/10.4095/304279>

COPYRIGHT

© Her Majesty the Queen in Right of Canada, as represented by the Minister of Natural Resources, 2017

Information contained in this publication or product may be reproduced, in part or in whole, and by any means, for personal or public non-commercial purposes, without charge or further permission, unless otherwise specified.

You are asked to:

- exercise due diligence in ensuring the accuracy of the materials reproduced;
- indicate the complete title of the materials reproduced, and the name of the author organization; and
- indicate that the reproduction is a copy of an official work that is published by Natural Resources Canada (NRCan) and that the reproduction has not been produced in affiliation with, or with the endorsement of, NRCan.

Commercial reproduction and distribution is prohibited except with written permission from NRCan. For more information, contact NRCan at nrcan.copyrightdroitdauteur.rncan@canada.ca.

RECOMMENDED CITATION

Geological Survey of Canada, 2017. Surficial geology, Aberdeen Lake, Nunavut, NTS 66-B; Geological Survey of Canada, Canadian Geoscience Map 323 (preliminary, Surficial Data Model v. 2.3 conversion of Map 44-1989), scale 1:125 000.
<https://doi.org/10.4095/304279>

ABSTRACT

This new surficial geology map product represents the conversion of Map 44-1989 and its legend, using the Geological Survey of Canada's Surficial Data Model (SDM version 2.3) (Open File 8236). All geoscience knowledge and information from Map 44-1989 that conformed to the current SDM were maintained during the conversion process. Supplementary, limited legacy information was added to complement the converted geoscience data. This consists of glacial striations from Wright (1955), as well as striations and field data from McMMartin et al. (2017); these are identified in the accompanying geodatabase. The purpose of converting legacy map data to a common science language and a common legend is to enable and facilitate the efficient digital compilation, interpretation, management, and dissemination of geological map information in a structured and consistent manner. This provides an effective knowledge management tool designed around a geodatabase that can expand, following the type of information to appear on new surficial geology maps.

RÉSUMÉ

Ce nouveau produit cartographique de la géologie des formations superficielles correspond à la conversion de la Carte 44-1989 et de sa légende, en se servant du Modèle de données pour les formations superficielles (MDFS version 2.3) de la Commission géologique du Canada (Dossier public 8236). Toutes les connaissances et l'information de nature géoscientifique de la Carte 44-1989 qui sont en conformité avec le modèle de données ont été conservées pendant le processus de conversion. Une faible quantité d'information ancienne a été ajoutée en complément aux données géoscientifiques converties. Il s'agit de stries glaciaires tirées de Wright (1955), et des stries et données de terrain de McMMartin et al. (2017). Ces entités sont identifiées dans la géodatabase du présent produit cartographique. Le but de la conversion de cartes publiées antérieurement suivant un langage scientifique commun et une légende commune est de permettre et de faciliter la compilation, l'interprétation, la gestion et la diffusion efficaces de l'information géologique cartographique en mode numérique de façon structurée et cohérente. Cette façon de faire offre un outil efficace de gestion des connaissances élaboré à l'aide d'une géodatabase qui pourra évoluer suivant le type d'information à paraître sur les nouvelles cartes des formations superficielles.

LICENCE AGREEMENT

View the license agreement at

<http://open.canada.ca/en/open-government-licence-canada>

ACCORD DE LICENCE

Voir l'accord de licence à

<http://ouvert.canada.ca/fr/licence-du-gouvernement-ouvert-canada>

SHEET 1 OF 1, SURFICIAL GEOLOGY

GENERAL INFORMATION

Author: Geological Survey of Canada

Geology based on airphoto interpretation with ground observations by J.M. Aylsworth, 1980, 1983. Striation measurements J.G. Fyles, 1954.

Geology conforms to Surficial Data Model v. 2.3

Data conversion by D.E. Kerr, 2015, 2016

Geomatics by QSP Geographics Inc. and A. Noad

Cartography by E. Everett

Initiative of the Geological Survey of Canada, conducted under the auspices of Natural Resources Canada's Geo-mapping for Energy and Minerals (GEM) Program.

Map projection Universal Transverse Mercator, zone 14.
North American Datum 1983

Base map at the scale of 1:250 000 from Natural Resources Canada, with modifications.

Elevations in metres above mean sea level

Mean magnetic declination 2017, 1°08'E, decreasing 6' annually. Readings vary from 2°44'E in the SW corner to 0°35'W in the NE corner of the map.

This map is not to be used for navigational purposes.

The Geological Survey of Canada welcomes corrections or additional information from users.

Data may include additional observations not portrayed on this map. See map info document accompanying the downloaded data for more information about this publication.

This publication is available for free download through GEOSCAN (<http://geoscan.nrcan.gc.ca/>).

This publication has been scientifically reviewed, but it has not undergone a formal edit.

MAP VIEWING FILES

The published map is distributed as a Portable Document File (PDF), and may contain a subset of the overall geological data for legibility reasons at the publication scale.

REFERENCES

Aylsworth, J.M., 1990. Surficial geology, Aberdeen Lake, District of Keewatin, Northwest Territories; Geological Survey of Canada, Map 44-1989, scale 1:125 000.
<https://doi.org/10.4095/130944>

Deblonde, C., Cocking, R.B., Kerr, D.E., Campbell, J.E., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Parent, M., Plouffe, A., Robertson, L., Smith, I.R., and Weatherston, A., 2017. Surficial Data Model, version 2.3.0: revisions to the science language of the integrated Geological Survey of Canada data model for surficial geology maps; Geological Survey of Canada, Open File 8236, 1 .zip file.
<https://doi.org/10.4095/302717>

McMartin, I., Tremblay, T., and Godbout, P.-M., 2017. Report of 2017 field activities for the GEM-2 Rae glacial history activity in the Kivalliq region, Nunavut; Geological Survey of Canada, Open File 8320, 14 p. <https://doi.org/10.4095/306006>

Wright, G.M., 1955. Geological notes on central District of Keewatin, Northwest Territories; Geological Survey of Canada, Paper 55-17, 1 .zip file.
<https://doi.org/10.4095/101294>

AUTHOR CONTACT

Questions, suggestions, and comments regarding the geological information contained in the data sets should be addressed to:

D.E. Kerr
Geological Survey of Canada
601 Booth Street
Ottawa ON
K1A 0E8
Daniel.Kerr@canada.ca

COORDINATE SYSTEM

Projection: Universal Transverse Mercator

Units: metres

Zone: 14

Horizontal Datum: NAD83

Vertical Datum: mean sea level

BOUNDING COORDINATES

Western longitude: 100°00'00"W

Eastern longitude: 98°00'00"W

Northern latitude: 65°00'00"N

Southern latitude: 64°00'00"N

SOFTWARE VERSION

Data has been originally compiled and formatted for use with ArcGIS™ desktop version 10.2.2 developed by ESRI®.

DATA MODEL INFORMATION

Surficial

The Geological Survey of Canada (GSC) through the Geo-mapping for Energy and Minerals Program (GEM) has undertaken the Geological Map Flow to develop protocols for the collection, management (compilation, interpretation), and dissemination of

surficial and bedrock geology data and map information. To this end, a data model has been created.

The Surficial Data Model (SDM) was designed using ESRI geodatabase architecture. The XML workspace document provided can be imported into a geodatabase, and the geodatabase will then be populated with the feature datasets, feature classes, tables, relationship classes, subtypes, and domains.

Shapefile and table (.dbf) versions of the data are included within the data. Column names have been simplified and the text values have been maintained within the shapefile attributes. The direction columns are numerical, to display rotation for points, and the symbol fields will hold the correct values to be matched to the appropriate style file.

For a more in depth description of the data model please refer to the official publication:

Deblonde, C., Cocking, R.B., Kerr, D.E., Campbell, J.E., Eagles, S., Everett, D., Huntley, D.H., Inglis, E., Parent, M., Plouffe, A., Robertson, L., Smith, I.R., and Weatherston, A., 2017. Surficial Data Model, version 2.3.0: revisions to the science language of the integrated Geological Survey of Canada data model for surficial geology maps; Geological Survey of Canada, Open File 8236, 1 .zip file.
<https://doi.org/10.4095/302717>