

CATALOGUE No.

83-209 C.2

ANNUAL




CANADA

## TUBERCULOSIS STATISTICS

Supplement: PATIENTS IN INSTITUTIONS

1960



DOMINION BUREAU OF STATISTICS

Health and Welfare Division

Public Health Section



DOMINION BUREAU OF STATISTICS

Health and Welfare Division

Public Health Section

TUBERCULOSIS STATISTICS

Supplement: PATIENTS IN INSTITUTIONS

1960

*Published by Authority of*  
The Honourable George Hees, Minister of Trade and Commerce

April, 1962  
9005-542

Price 50 cents



## PREFACE

This is the third annual report dealing with the characteristics of patients in Canadian sanatoria and tuberculosis units of hospitals as at the year end, published by the Dominion Bureau of Statistics.

It complements the annual publication *Tuberculosis Statistics* and is based upon the admission data for individual patients supplied by the institutions in Canada.

The Dominion Bureau of Statistics wishes to thank the medical superintendents and other personnel of the reporting institutions and the provincial and federal authorities for their prompt and accurate reporting. Without their co-operation this report would not be possible.

The report was prepared in the Public Health Section of the Health and Welfare Division.

WALTER E. DUFFETT,

*Dominion Statistician.*

## **SYMBOLS**

The interpretation of the symbols used in the tables throughout this publication is as follows:

.. figures not available.

... used to replace percentages when the divisor is less than 10.

- nil or zero.

## TABLE OF CONTENTS

	Page
Introduction .....	7
Descriptive Text and Charts .....	7
Table	
1. All patients at December 31, 1936-60, by Province and Sex .....	14
2. Tuberculous Patients, by Age, Province, and Sex, 1960 .....	16
3. Tuberculous Patients, by Age, Diagnosis, and Sex, 1960 .....	18
4. Tuberculous Patients, 15 Years and Over, by Sex, Age, Diagnosis, and Marital Status, 1960 .....	20
5. All Patients, Time Since Admission, by Province, 1960 .....	21
6. All Patients, by Sex, Diagnosis, and Time of Admission, 1934-1960..	22
7. Tuberculous Patients, by Sex, Type of Admission, and Time of Admission, 1934-60 .....	24
8. Tuberculous Patients, by Age, Origin, and Sex, 1960 .....	26
9. All Indian and Eskimo Patients, by Sex and Diagnosis, 1960.....	27
10. Non-Tuberculous Patients, by Sex and Diagnosis, 1960 .....	27
11. All Patients, by Sex, Diagnosis, Province, and Institution, 1960 .....	28
Appendix — Patients Under Treatment for Tuberculosis in Mental Institutions, 1960 .....	32



## INTRODUCTION

This report has been produced to give a detailed statistical description of patients in tuberculosis institutions at December 31, 1960. The Dominion Bureau of Statistics has collected and published data on patients admitted to and separated from tuberculosis institutions during each calendar year since 1937. The patient population at any time has characteristics different from those of the patients' movement for a year. A description of patient movement within tuberculosis institutions can be found in the annual parent publication "Tuberculosis Statistics".

The following tables constitute a census of patients in residence in tuberculosis sanatoria and tuberculosis units of hospitals in Canada. The

data are based on records of admission documents and are subject to the following limitations: All patients who, on admission, had been reported as tuberculosis suspects are excluded from the tables on tuberculous patients, even when tuberculosis was confirmed during their hospital stay and reported only when separation data became available. This census also excludes 1,611 patients in mental institutions who were under treatment for tuberculosis, but had not been reported as an admission to a tuberculosis unit. Their distribution is included in the appendix on page 32.

The diagnostic classifications used in this report are taken from the "Manual of the International Statistical Classification of Diseases, Injuries, and Causes of Death, Geneva, 1957".

## DESCRIPTIVE TEXT AND CHARTS


### Historical

The number of patients in tuberculosis institutions had increased from 7,764 in 1936 to 16,051 in 1953. Since then, the number of patients has dropped rapidly to 7,757 at December 31, 1960, equalling the total in 1936 in spite of a 64 per cent increase in the general population over the 24 year period.

The patient-population ratio which stood at 72.0 per 100,000 population in 1937 reached a

peak in 1952 with 109.3 but was down to its lowest point with 43.0 per 100,000 population at December 31, 1960. Wide variations occur when comparisons of patient-population ratios are studied on a provincial basis. At the end of 1960, Newfoundland had the highest patient-population ratio with 84.1 per 100,000 population. New Brunswick and Prince Edward Island followed with a ratio of 75.6 and 63.8 respectively. The lowest ratios were in Ontario with 27.0 and British Columbia with 30.3.

GRAPH - I



### Sex and age

There were 7,043 tuberculous patients in residence at December 31, 1960, and 62.6 per cent were males. In 1959, 61.4 per cent of residents were males. The median age of the male patients was 46 years, and that of the female patients 32 years, both unchanged from 1959.


In order to evaluate to what degree the disease affects persons of different age and sex, ratios of patients per 100,000 of the general population are shown in Table 2 and Graph 2 for each five-year group by age and sex. The male ratio of 48.2 per 100,000 population was 13.9 per cent lower than in 1959, while the female ratio declined by 18.2 per cent to 29.6. The distribution of tuber-

culous patients by sex and five year age groups shows a different pattern for the two sexes. Between 5 and 25 years, ratios for females are higher, whereas for all other ages the male ratios are higher.

Provincial male ratios were lower in Ontario (32.0), British Columbia (40.0) and Saskatchewan (47.9) than the national ratio (48.2). Female ratios were lower than the national ratio (29.6) only in Ontario (16.9) and British Columbia (19.7).

The male/female ratio in 1960 was 1.67 (i.e., there were 167 males for every 100 females). In 1959 this ratio was 1.59. Provincial male/female ratio showed considerable variations ranging from 1.31 in Saskatchewan to 2.15 in British Columbia.

GRAPH - 2



### Diagnosis


Pulmonary tuberculosis constituted the largest proportion (80.8 per cent) of all tuberculosis diagnoses in 1960. The proportion was slightly smaller than in 1959 (82.7 per cent). The percentage of tuberculosis of other respiratory sites increased to 12.1 per cent in 1960 from 10.5 per cent in 1959. Tuberculosis other forms, in spite of a decrease in number were proportionately higher in 1960 than in 1959. They represented 7.1 per cent of all tuberculous diagnoses in 1960 compared with 6.7 per cent in 1959. Tuberculosis of the genito-urinary system accounted for 34.9 per cent of the tuberculosis other forms group in 1960 as compared with 30.3 per cent in 1959. The proportion

of males with pulmonary tuberculosis was 82.4 per cent (84.8 per cent in 1959) while females with this diagnosis made up 78.1 per cent (79.6 per cent in 1959) of all tuberculous females in hospital. The median age for patients with primary infection was 7 years in 1960 against 6 years in 1959, for minimal pulmonary tuberculosis it was 32 years, unchanged from 1959, moderately advanced 45 years (43 in 1959, and far advanced 50 years unchanged from 1959). The male/female ratio for primary infection was 1.17, minimal pulmonary tuberculosis was 1.10, moderately advanced 1.77, and far advanced 2.32. In all diagnoses, with the exception of minimal pulmonary tuberculosis, the ratio was higher than in 1959.

GRAPH - 3


## TUBERCULOUS IN-PATIENTS, BY AGE, AND BY PROVINCE, 1960

(RATIO PER 100,000 POPULATION)




GRAPH - 4

TUBERCULOUS IN-PATIENTS, 15 YEARS OF AGE OR OVER  
BY SEX, AGE, AND MARITAL STATUS, 1960  
(RATIO PER 100,000 POPULATION)



GRAPH - 5

TIME SINCE ADMISSION, TUBERCULOUS IN-PATIENTS, 1959, 1960  
(CUMULATIVE PERCENTAGE DISTRIBUTION)



### Marital status

Ratios of patients per 100,000 population, specific for marital status, by broad age groups, are shown in Table 4 and Graph 4. In all four age groups the ratios of married patients are much lower than single patients or widowed and divorced patients. This applies to both males and females. The proportion of single males in age group 45-64 is more than three times those of the married in the same age group. The widowed and divorced population comprise relatively more tuberculous hospital patients than the married population, especially among men. The figures for female widowed and divorced are only slightly higher than the corresponding figures for the married females.

### Time since admission

Time in hospital since admission for both tuberculous and non-tuberculous patients is shown in Table 5 and Table 6. Non-tuberculous patients formed 9.2 per cent of all patients in hospital at the end of 1950 compared with 8.8 per cent in 1959 and 7.5 per cent in 1958.

The proportion of non-tuberculars in hospital at December 31, 1960 was much less than the proportion of non-tuberculous admissions which in 1959 was 27.6 per cent of all admissions and in 1960 was 28.3 per cent. Of the 714 non-tuberculous patients in hospital 431 or 60.4 per cent were in Quebec. Quebec non-tuberculous patients were 15.3 per cent of the total number of patients in tuberculous hospitals in the Province. Ontario had 153 non-tuberculous patients or 21.4 per cent of all non-tuberculous. This was 9.2 per cent of the provincial total for Ontario.

At the end of 1960, one-fourth of the tuberculous patients had been in hospital for 11.2 months or more and another fourth had stayed 2.5 months or less. The median was 5.5 months, i.e., half of the tuberculous patients had been in hospital 5.5 months or less (Graph 5).

Time since admission for half the patients ranged from 4.0 months in Newfoundland to 8.0 in Manitoba. The median length of time since admission for patients in Quebec and Ontario was almost the same as the median for Canada.

Time since admission for half of the non-tuberculous patients (median) was 3.2 months in 1960 compared to 3.4 months in 1959.

Median length of time since admission for males with pulmonary tuberculosis was 5.7 months, for females it was 5.2 months. The corresponding figures for 1959 were 6.1 months for males and 5.7 months for females. At the end of 1960, 7.4

per cent of the patients with pulmonary tuberculosis had been in hospital for three years or more. Of this group, 71.7 per cent were males and 28.3 per cent were females. The male/female ratio for this particular group, was 2.53 (i.e., there were 253 males for 100 females) whereas in total for all periods, the male/female ratio was 1.76 (i.e., 176 males for 100 females).

### Type of admission

In Table 7 tuberculous patients are shown by type of latest admission. Of all tuberculous patients at the end of the year, 4,309 or 61.2 per cent were first admissions who had never had previous treatment for tuberculosis in hospital, while the remainder had received some previous treatment. These include patients (23.5 per cent) who had previously been discharged from a tuberculosis institution (readmissions), patients who had been discharged previously with the understanding that they would continue treatment at some future date (2.8 per cent), patients who had transferred directly from one tuberculosis institution to another (12.1 per cent) and a very few who had entered hospital late in the year for review and who were subsequently discharged inactive and thus not requiring treatment.

### Origin


The origin of patients is shown for Indians and Eskimos only. All other origins are combined. Registered Indians (those who live on Indian reserves), and unregistered Indians (where the Indian origin is traced through the father), are included under the category "Indian". Population ratios could not be calculated, as a distribution of the general population by origin was not available.

Tuberculous Indians in hospital at the end of 1960 were 15.7 per cent of all tuberculous patients while Eskimos were 3.8 per cent. Both these proportions were slightly higher than in 1959. For both Indians and Eskimos, females constituted 51.9 per cent of the patients whereas for all other origins they made up only 34 per cent. Median age for female Indians was 23 years, for Eskimos 24 years and for all others 36 years. Median age for male Indians was 26 years, for Eskimos 22 and for all other 49 years. Tuberculous Indian and Eskimo patients show higher proportions of "other respiratory tuberculosis" and of "tuberculosis, other forms" (Graph 6).

### Patients without tuberculosis

Of the 714 non-tuberculous patients in hospital, 307 or 43 per cent had no disease, were undiagnosed, or were tuberculosis suspects only. Of the remainder, a majority suffered from respiratory diseases like pneumonia and bronchitis.

GRAPH - 6



**TABLE 1. All Patients at December 31, 1936-60, by Province and Sex**

TABLE 1. All Patients at December 31, 1936-60, by Province and Sex

1945	1946	1947	1948	1949	1950	1951	1952	1953	1954	1955	1956	1957	1958	1959	1960	No.
10,721	12,006	12,407	12,950	13,598	14,930	15,090	16,047	16,051	15,220	14,157	13,072	11,670	10,433	8,966	7,757	1
5,883	6,756	6,668	6,784	7,086	7,806	7,992	8,723	8,899	8,706	8,248	7,676	6,998	6,266	5,461	4,836	2
4,838	5,250	5,741	6,186	6,512	7,124	7,098	7,324	7,152	6,514	5,909	5,396	4,672	4,167	3,505	2,921	3
..	..	..	..	..	368	409	429	568	772	803	663	564	483	455	392	4
..	..	..	..	..	170	215	213	290	405	425	329	299	250	250	231	5
..	..	..	..	..	198	194	216	278	367	378	334	265	233	205	161	6
88	135	129	135	146	145	146	142	133	149	95	92	85	80	69	67	7
40	58	55	58	65	70	62	72	70	85	50	58	54	49	44	43	8
48	77	74	77	81	75	84	70	63	64	45	34	31	31	25	24	9
595	861	857	658	626	611	650	841	781	751	664	620	499	477	430	359	10
286	523	508	316	307	306	326	454	419	439	421	367	294	288	281	239	11
309	338	349	342	319	305	324	387	362	312	243	253	205	189	149	120	12
515	659	816	815	846	805	839	858	793	699	701	632	582	602	464	458	13
237	304	393	394	424	406	407	438	403	370	382	358	325	320	272	279	14
278	355	423	421	422	399	432	420	390	329	319	274	257	282	192	179	15
3,343	3,528	3,527	3,654	4,111	4,789	4,951	5,111	4,961	4,556	4,278	4,082	3,645	3,274	3,014	2,758	16
1,706	1,856	1,814	1,866	2,075	2,405	2,495	2,618	2,658	2,528	2,463	2,377	2,175	2,019	1,846	1,724	17
1,637	1,672	1,713	1,788	2,036	2,384	2,456	2,493	2,303	2,028	1,815	1,705	1,470	1,255	1,168	1,034	18
3,379	3,818	3,559	3,965	3,918	4,123	4,083	4,171	4,242	3,987	3,823	3,568	3,100	2,736	2,151	1,666	19
2,049	2,349	2,024	2,273	2,237	2,388	2,440	2,483	2,571	2,504	2,372	2,253	2,010	1,779	1,424	1,096	20
1,330	1,469	1,535	1,692	1,681	1,735	1,643	1,688	1,671	1,483	1,451	1,315	1,090	957	727	570	21
823	825	1,007	1,100	1,161	1,159	1,137	1,111	1,116	1,064	1,014	999	940	799	614	457	22
456	430	527	527	577	578	577	581	563	566	550	557	547	448	337	270	23
367	395	480	573	584	581	560	530	553	498	464	442	393	351	277	187	24
775	775	763	739	759	795	735	790	805	833	738	658	589	507	417	409	25
436	424	400	373	378	400	358	389	413	436	410	362	312	291	236	233	26
339	351	363	366	381	395	377	401	392	397	328	296	277	216	181	176	27
386	588	645	693	800	860	893	978	940	887	785	770	724	661	626	603	28
198	320	320	333	393	456	457	495	496	459	427	399	391	342	313	347	29
188	268	325	360	407	404	436	483	444	428	358	371	333	319	313	256	30
817	817	1,104	1,191	1,231	1,275	1,247	1,414	1,364	1,245	1,038	843	777	661	579	494	31
475	492	625	624	630	627	655	877	844	789	658	549	510	411	395	338	32
342	325	479	567	601	648	592	537	520	456	380	294	267	250	184	156	33
..	..	..	..	..	..	..	202	348	277	218	145	165	153	147	94	34
..	..	..	..	..	..	..	103	172	125	90	67	81	69	63	36	35
..	..	..	..	..	..	..	99	176	152	128	78	84	84	84	58	36

TABLE 2. Tuberculous Patients, by Age, Province, and Sex, 1960

Note: This table and all succeeding tables refer to patients in institutions at December 31, 1960

Province and sex	All ages	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
Number											
Canada .....	T. 7,043	373	397	293	391	482	495	507	537	482	500
M. 4,409	200	190	140	197	240	257	269	312	301	354	
F. 2,634	173	207	153	194	242	238	238	225	181	146	
Newfoundland .....	T. 375	24	42	19	39	39	27	31	33	30	33
M. 219	11	22	8	20	19	14	19	18	19	28	
F. 156	13	20	11	19	20	13	12	15	11	5	
Prince Edward Island .....	T. 65	8	7	5	3	2	2	1	1	1	5
M. 42	3	4	2	3	1	—	1	—	—	—	5
F. 23	5	3	3	—	1	2	—	—	—	1	—
Nova Scotia .....	T. 343	10	2	9	17	22	20	25	31	25	39
M. 230	3	2	5	9	9	9	14	18	18	30	
F. 113	7	—	4	8	13	11	11	13	7	9	
New Brunswick .....	T. 432	22	35	33	30	21	26	24	27	34	30
M. 268	14	16	15	20	14	15	15	14	23	23	
F. 164	8	19	18	10	7	11	9	13	11	7	
Quebec .....	T. 2,327	54	87	50	125	177	192	177	205	199	179
M. 1,473	27	39	28	66	88	104	96	123	131	114	
F. 854	27	48	22	59	89	88	81	82	68	65	
Ontario .....	T. 1,513	72	56	47	63	87	96	99	125	103	97
M. 996	43	28	19	29	51	49	53	78	61	69	
F. 517	29	28	28	34	36	47	46	47	42	28	
Manitoba .....	T. 434	32	23	23	33	34	29	37	24	27	30
M. 253	20	8	11	18	14	9	17	12	14	22	
F. 181	12	15	12	15	20	20	20	12	13	8	
Saskatchewan .....	T. 399	35	63	33	16	22	21	24	25	13	20
M. 226	15	33	12	7	7	13	8	11	7	15	
F. 173	20	30	21	9	15	8	16	14	6	5	
Alberta .....	T. 584	68	41	33	26	44	52	45	40	30	31
M. 334	34	22	17	7	17	27	23	20	15	20	
F. 250	34	19	16	19	27	25	22	20	15	11	
British Columbia .....	T. 491	36	30	28	31	31	26	36	25	15	35
M. 335	24	12	15	18	19	16	20	17	11	28	
F. 156	12	18	13	13	12	10	16	8	4	7	
Northwest Territories .....	T. 80	12	11	13	8	3	4	8	1	5	1
M. 33	6	4	8	—	1	1	3	—	2	—	
F. 47	6	7	5	8	2	3	5	1	3	1	
Number											
Canada .....	T. 488	489	461	381	341	218	124	44	15	2	23
M. 372	363	360	296	267	160	74	31	10	1	1	15
F. 116	126	101	85	74	58	50	13	5	1	8	
Newfoundland .....	T. 15	14	5	8	9	4	1	1	—	—	1
M. 12	6	3	7	6	4	1	1	—	—	—	1
F. 3	8	2	1	3	—	—	—	—	—	—	—
Prince Edward Island .....	T. 5	10	2	3	3	1	2	2	1	—	1
M. 5	9	2	1	2	1	—	—	—	—	—	—
F. —	1	—	2	1	—	2	2	—	—	—	—
Nova Scotia .....	T. 33	27	24	24	18	8	7	2	—	—	—
M. 26	23	21	17	14	7	4	1	—	—	—	—
F. 7	4	3	7	4	1	3	1	—	—	—	—
New Brunswick .....	T. 26	26	25	19	22	18	8	3	3	—	—
M. 16	18	18	11	16	11	6	2	1	2	—	—
F. 10	8	7	8	6	7	2	2	—	—	—	—
Quebec .....	T. 208	171	152	128	97	74	32	5	2	—	13
M. 165	132	113	92	74	47	19	5	2	—	—	8
F. 43	39	39	36	23	27	13	—	—	—	—	5
Ontario .....	T. 103	122	132	102	87	50	43	19	6	1	3
M. 77	94	112	83	69	40	22	12	4	1	2	
F. 26	28	20	19	18	10	21	7	2	—	—	
Manitoba .....	T. 17	26	26	27	23	11	7	3	1	—	1
M. 14	13	19	25	18	10	5	2	1	—	—	
F. 3	13	7	2	5	1	2	1	—	—	—	
Saskatchewan .....	T. 20	19	24	18	23	13	6	2	1	—	1
M. 18	14	17	14	18	11	4	2	—	—	—	
F. 2	5	7	4	5	2	2	—	1	—	—	
Alberta .....	T. 29	43	35	21	22	16	4	3	—	—	1
M. 19	30	26	20	20	11	3	2	—	—	—	
F. 10	13	9	1	2	5	1	1	—	—	—	
British Columbia .....	T. 32	30	29	29	35	22	14	4	1	1	1
M. 20	24	24	25	29	18	10	4	1	—	—	
F. 12	6	5	4	6	4	4	—	—	—	—	
Northwest Territories .....	T. —	1	7	2	2	1	—	—	—	—	1
M. —	—	5	1	1	1	—	—	—	—	—	
F. —	1	2	1	1	1	—	—	—	—	—	

TABLE 2. Tuberculous Patients, by Age, Province, and Sex, 1960 — Concluded

Province and sex	All ages	0-4	5-9	10-14	15-19	20-24	25-29	30-34	35-39	40-44	45-49
Ratio per 100,000 population											
Canada .....	M. F.	48.2 29.6	17.3 15.7	18.2 20.5	15.5 17.7	28.0 28.3	38.0 39.5	40.7 40.0	41.4 37.7	50.9 36.7	53.8 32.4
Newfoundland .....	M. F.	90.5 69.6	29.5 36.2	68.3 63.1	... 40.0	93.5 86.8	110.5 122.7	88.6 100.0	127.5 99.2	128.6 124.0	140.7 98.2
Prince Edward Island .....	M. F.	78.9 44.4	...	...	...	...	...	—	...	...	—
Nova Scotia .....	M. F.	61.9 31.7	...	...	...	...	54.2	52.1	64.2 52.1	82.6 57.5	82.6 ...
New Brunswick .....	M. F.	87.8 54.6	34.5 ...	41.2 50.5	42.7 54.1	71.7 36.6	66.0 ...	85.2 60.4	88.2 ...	79.5 71.4	135.3 64.7
Quebec .....	M. F.	56.7 33.1	7.9 8.3	12.6 16.0	10.0 8.2	29.6 27.1	46.0 46.5	55.8 47.2	51.5 43.5	72.0 46.6	84.4 43.8
Ontario .....	M. F.	32.0 16.9	11.4 8.1	8.3 8.6	6.7 10.3	13.4 16.2	25.1 18.2	22.1 22.4	22.6 19.6	35.1 20.6	30.0 20.6
Manitoba .....	M. F.	54.8 40.5	36.7 22.8	...	24.7 28.7	50.8 44.6	45.3 66.7	...	56.7 71.2	40.0 66.7	49.8 38.8
Saskatchewan .....	M. F.	47.9 39.3	25.7 36.0	62.4 58.7	25.8 47.9	...	51.4	45.9 ...	...	37.7 56.5	47.9 ...
Alberta .....	M. F.	49.2 39.7	37.1 38.8	28.0 25.5	26.5 26.0	40.3	37.1 60.7	55.8 56.2	47.5 48.0	43.7 44.9	36.9 38.2
British Columbia .....	M. F.	40.0 19.7	25.0 12.9	14.2 21.7	20.5 18.6	31.6 24.2	35.4 24.6	27.3 19.8	33.2 28.9	29.8 ...	19.9 ...
Northwest Territories .....	M. F.	264.0 494.7	...	...	...	—	...	...	...	—	...
		50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99
Ratio per 100,000 population											
Canada .....	M. F.	85.9 28.0	100.6 36.8	124.7 35.0	126.2 36.2	134.5 37.3	126.7 40.2	102.5 69.3	172.2 36.0	138.9 ...	...
Newfoundland .....	M. F.	129.0 ...	...	...	...	...	...	...	...	—	—
Prince Edward Island .....	M. F.	...	...	—	...	...	—	...	...	—	—
Nova Scotia .....	M. F.	148.6 ...	166.7 ...	181.0 ...	166.7 ...	160.9 ...	...	...	...	—	—
New Brunswick .....	M. F.	120.3 82.6	165.1 ...	197.8 ...	139.2 ...	238.8 ...	261.9 ...	...	...	...	—
Quebec .....	M. F.	144.9 37.8	141.6 41.8	155.9 53.8	161.7 63.3	178.7 49.4	181.5 86.8	182.7 83.9	...	...	—
Ontario .....	M. F.	49.9 17.6	72.6 21.6	106.9 18.0	103.5 20.5	101.8 24.3	92.6 18.0	118.9 68.2	193.5 ...	...	...
Manitoba .....	M. F.	61.7 ...	68.1 71.4	116.6 ...	172.4 ...	141.7 ...	109.9 ...	...	...	...	...
Saskatchewan .....	M. F.	78.9 ...	69.7 ...	103.7 ...	90.3 ...	131.4 ...	110.0 ...	...	...	—	—
Alberta .....	M. F.	60.5 36.4	114.5 58.3	124.4 ...	117.6 ...	127.4 ...	104.8 ...	...	...	—	—
British Columbia .....	M. F.	45.5 30.7	67.0 ...	86.6 ...	96.2 ...	111.5 ...	92.3 ...	102.0 ...	...	...	...
Northwest Territories .....	M. F.	— —	— ...	...	...	...	— —	— —	— —	— —	— —

<sup>1</sup> Less than 50 persons.

Note: Canada totals include Yukon.

TABLE 3. Tuberculous Patients, by Age, Diagnosis, and Sex, 1960

No.	Diagnosis and sex	All ages	0-4	5-9	10-14	15-19	20-24	25-29	30-34	
			Number							
1	Totals .....	T.	7,043	373	397	293	391	482	495	507
2		M.	4,409	200	190	140	197	240	257	269
3		F.	2,634	173	207	153	194	242	238	238
4	Pulmonary (002.1-002.5) .....	T.	5,689	76	117	124	314	415	421	444
5		M.	3,631	42	52	45	155	207	215	226
6		F.	2,058	34	65	79	159	208	206	218
7	Minimal pulmonary (002.1) .....	T.	1,047	34	74	66	108	107	101	83
8		M.	548	20	32	24	46	45	52	33
9		F.	499	14	42	42	62	62	49	50
10	Moderately advanced pulmonary (002.2) .....	T.	2,429	16	24	37	129	192	176	199
11		M.	1,552	12	9	10	67	105	84	102
12		F.	877	4	15	27	62	87	92	97
13	Far advanced pulmonary (002.3) .....	T.	2,122	4	4	13	69	112	142	159
14		M.	1,483	1	3	6	40	56	78	89
15		F.	639	3	1	7	29	56	64	70
16	Other and unspecified pulmonary (002.4, 002.5) .....	T.	91	22	15	8	8	4	2	3
17		M.	48	9	8	5	2	1	1	2
18		F.	43	13	7	3	6	3	1	1
19	Other respiratory (001, 003-008) .....	T.	855	255	244	134	47	29	18	8
20		M.	500	137	122	80	28	16	10	4
21		F.	355	118	122	54	19	13	8	4
22	With occupational diseases of lung (001) .....	T.	67	—	—	—	—	—	—	—
23		M.	67	—	—	—	—	—	—	—
24		F.	—	—	—	—	—	—	—	—
25	Pleurisy (003) .....	T.	165	—	10	26	30	26	17	7
26		M.	101	—	7	11	19	16	10	4
27		F.	64	—	3	15	11	10	7	3
28	Primary infection (004) .....	T.	584	244	222	98	15	3	1	—
29		M.	315	135	111	61	8	—	—	—
30		F.	269	109	111	37	7	3	1	—
31	Tracheobronchial (005) .....	T.	37	10	12	10	2	—	—	—
32		M.	17	2	4	8	1	—	—	—
33		F.	20	8	8	2	1	—	—	—
34	Other respiratory organs (007, 008) .....	T.	2	1	—	—	—	—	—	1
35		M.	—	—	—	—	—	—	—	—
36		F.	2	1	—	—	—	—	—	1
37	Other forms (010-019) .....	T.	499	42	36	35	30	38	56	55
38		M.	278	21	16	15	14	17	32	39
39		F.	221	21	20	20	16	21	24	16
40	Meninges and central nervous system (010) .....	T.	55	23	14	5	6	3	2	—
41		M.	25	9	7	3	4	—	2	—
42		F.	30	14	7	2	2	3	—	—
43	Intestines, peritoneum, and mesenteric glands (011) .....	T.	11	—	1	—	3	1	—	1
44		M.	3	—	1	—	—	—	—	1
45		F.	8	—	—	—	3	1	—	—
46	Bones and joints (012, 013) .....	T.	169	6	10	14	11	8	16	14
47		M.	105	2	4	5	8	6	10	12
48		F.	64	4	6	9	3	2	6	2
49	Skin and subcutaneous cellular tissue (014) .....	T.	4	1	1	—	—	—	—	1
50		M.	1	1	—	—	—	—	—	—
51		F.	3	—	1	—	—	—	—	1
52	Lymphatic system (015) .....	T.	57	8	9	7	6	9	8	5
53		M.	27	6	3	3	4	4	6	1
54		F.	30	2	6	4	5	5	2	4
55	Genito-urinary system (016) .....	T.	174	1	—	3	2	15	28	29
56		M.	106	1	—	1	1	6	14	24
57		F.	68	—	—	2	1	9	14	5
58	Other organs (017-019) .....	T.	29	3	1	6	2	2	2	5
59		M.	11	2	1	3	—	1	—	1
60		F.	18	1	—	3	2	1	2	4
Ratio per 100,000 population										
61	Totals .....	M.	48.2	17.3	18.2	15.5	28.0	38.0	40.7	41.4
62		F.	29.6	15.7	20.5	17.7	28.3	39.5	40.0	37.7
63	Pulmonary .....	M.	39.7	3.6	5.0	5.0	22.0	32.8	34.1	34.8
64		F.	23.1	3.1	6.4	9.1	23.2	33.9	34.6	34.5
65	Other respiratory .....	M.	5.5	11.9	11.7	8.9	4.0	2.5	1.6	...
66		F.	4.0	10.7	12.1	6.2	2.8	2.1	—	...
67	Other forms .....	M.	3.0	1.8	1.5	1.7	2.0	2.7	5.1	6.0
68		F.	2.5	1.9	2.0	2.3	2.3	3.4	4.0	2.5

TABLE 3. Tuberculous Patients, by Age, Diagnosis, and Sex, 1960

35-39	40-44	45-49	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99	Not stated	No.
Number														
537	482	500	488	489	461	381	341	218	124	44	15	2	23	1
312	301	354	372	363	360	296	267	160	74	31	10	1	15	2
225	181	146	116	126	101	85	74	58	50	13	5	1	8	3
485	442	451	460	448	421	350	318	206	120	39	15	2	21	4
285	275	323	352	332	328	273	247	151	72	26	10	1	14	5
200	167	128	108	116	93	77	71	55	48	13	5	1	7	6
86	58	56	63	58	43	30	29	26	15	4	2	—	4	7
42	29	32	39	42	32	23	24	17	9	3	2	—	2	8
44	29	24	24	16	11	7	5	9	6	1	—	—	2	9
218	199	200	175	191	189	153	149	89	63	19	6	2	3	10
138	122	139	136	136	142	117	111	68	37	10	3	1	3	11
80	77	61	39	55	47	36	38	21	26	9	3	1	—	12
178	180	195	218	197	184	161	138	90	41	16	7	—	14	13
104	123	152	174	152	149	127	111	66	25	13	5	—	9	14
74	57	43	44	45	35	34	27	24	16	3	2	—	5	15
3	5	—	4	2	5	6	2	1	1	—	—	—	—	16
1	1	—	3	2	5	6	1	—	1	—	—	—	—	17
2	4	—	1	—	—	—	1	1	—	—	—	—	—	18
9	9	12	8	19	25	16	11	5	2	3	—	—	1	19
4	8	6	8	17	23	16	11	5	2	3	—	—	—	20
5	1	6	—	2	2	—	—	—	—	—	—	—	1	21
1	1	3	4	14	18	11	10	3	1	1	—	—	—	22
1	1	3	4	14	18	11	10	3	1	1	—	—	—	23
—	—	—	—	—	—	—	—	—	—	—	—	—	—	24
7	8	9	4	5	5	5	1	2	1	2	—	—	—	25
3	7	3	4	3	3	5	1	2	1	2	—	—	—	26
4	1	6	—	2	2	—	—	—	—	—	—	—	—	27
—	—	—	—	—	—	—	—	—	—	—	—	—	1	28
—	—	—	—	—	—	—	—	—	—	—	—	—	—	29
—	—	—	—	—	—	—	—	—	—	—	—	—	1	30
1	—	—	—	—	—	2	—	—	—	—	—	—	—	31
—	—	—	—	—	—	—	—	—	—	—	—	—	—	32
1	—	—	—	—	—	—	—	—	—	—	—	—	—	33
—	—	—	—	—	—	—	—	—	—	—	—	—	—	34
—	—	—	—	—	—	—	—	—	—	—	—	—	—	35
—	—	—	—	—	—	—	—	—	—	—	—	—	—	36
43	31	37	20	22	15	15	12	7	2	2	—	—	1	37
23	18	25	12	14	9	7	9	4	—	—	—	—	1	38
20	13	12	8	8	6	8	3	3	2	—	—	—	—	39
—	1	—	1	—	—	—	—	—	—	—	—	—	—	40
—	1	—	1	—	—	—	—	—	—	—	—	—	—	41
—	1	—	1	—	—	—	—	—	—	—	—	—	—	42
—	1	3	—	1	—	—	—	—	—	—	—	—	—	43
—	1	2	—	1	—	—	—	—	—	—	—	—	—	44
16	12	14	9	10	9	8	5	4	—	—	2	—	1	45
11	8	11	5	7	5	3	3	2	—	—	—	—	1	46
5	4	3	4	3	4	5	2	2	—	—	—	—	1	47
—	—	—	—	—	—	—	—	1	—	—	—	—	—	48
—	—	—	—	—	—	—	—	1	—	—	—	—	—	49
—	—	—	—	—	—	—	—	1	—	—	—	—	—	50
—	—	—	—	—	—	—	—	1	—	—	—	—	—	51
2	—	1	1	1	—	—	—	—	—	—	—	—	—	52
1	—	1	1	—	—	—	—	—	—	—	—	—	—	53
1	—	—	—	—	—	—	—	—	—	—	—	—	—	54
21	17	19	9	11	6	6	4	3	2	1	—	—	—	55
11	10	12	6	7	4	2	2	1	—	—	—	—	—	56
10	7	7	3	4	2	—	—	—	1	—	—	—	—	57
4	—	—	—	—	—	—	—	3	—	—	—	—	—	58
4	—	—	—	—	—	—	—	3	—	—	—	—	—	59
—	—	—	—	—	—	—	—	—	1	—	—	—	—	60
Ratio per 100,000 population														
50.9	53.8	70.1	85.9	100.0	124.7	126.2	134.5	126.7	102.5	172.2	138.9	...	...	61
36.7	32.4	30.0	28.0	36.8	35.0	36.2	37.3	40.2	69.3	36.0	...	...	...	62
46.5	49.2	63.9	81.3	92.0	113.6	116.4	124.4	119.6	99.7	144.4	138.9	...	...	63
32.6	29.9	26.3	26.0	33.8	32.2	32.8	35.8	38.1	66.5	36.0	...	...	...	64
...	...	...	...	...	4.7	8.0	6.8	5.5	...	...	—	—	—	65
...	...	...	—	...	...	—	—	—	...	...	—	—	—	66
3.7	3.2	4.9	2.8	3.9	...	...	...	...	—	...	—	—	—	67
3.3	2.3	2.5	...	...	...	...	...	...	...	...	—	—	—	68

TABLE 4. Tuberculous Patients, 15 Years and Over, by Sex, Age, Diagnosis, and Marital Status, 1960

Diagnosis and marital status	Male						Female					
	Total	15-24	25-44	45-64	65 or over	Not stated	Total	15-24	25-44	45-64	65 or over	Not stated
Number												
Totals.....	3,879	437	1,139	1,449	839	15	2,101	436	882	489	286	8
Single .....	1,469	392	468	409	196	4	661	311	205	94	48	3
Married .....	2,029	44	651	898	428	8	1,178	123	645	314	91	5
Widowed .....	325	1	6	109	206	3	143	1	21	77	144	—
Divorced.....	48	—	11	30	7	—	14	1	7	4	2	—
Not stated .....	8	—	3	3	2	—	5	—	4	—	1	—
Pulmonary .....	3,492	362	1,001	1,335	780	14	1,880	367	791	445	270	7
Single .....	1,297	321	412	373	187	4	575	262	183	83	45	2
Married .....	1,848	40	572	833	396	7	1,057	103	577	289	83	5
Widowed .....	297	1	6	99	188	3	229	1	20	69	139	—
Divorced.....	43	—	9	27	7	—	14	1	7	4	2	—
Not stated .....	7	—	2	3	2	—	5	—	4	—	1	—
Other respiratory .....	161	44	26	54	37	—	61	32	18	10	—	1
Single .....	67	41	8	12	6	—	33	23	6	3	—	1
Married .....	70	3	18	31	18	—	27	9	12	6	—	—
Widowed .....	21	—	—	8	13	—	1	—	—	1	—	—
Divorced.....	3	—	—	3	—	—	—	—	—	—	—	—
Not stated .....	—	—	—	—	—	—	—	—	—	—	—	—
Other forms .....	226	31	112	60	22	1	160	37	73	34	16	—
Single .....	105	30	48	24	3	—	53	26	16	8	3	—
Married .....	111	1	61	34	14	1	94	11	56	19	8	—
Widowed .....	7	—	—	2	5	—	13	—	1	7	5	—
Divorced.....	2	—	2	—	—	—	—	—	—	—	—	—
Not stated .....	1	—	1	—	—	—	—	—	—	—	—	—
Ratio per 100,000 population												
Totals.....	66.2	34.3	47.3	94.4	134.4		36.6	35.2	37.3	33.3	42.7	
Single .....	81.8	36.7	97.7	231.7	275.3		49.6	38.2	67.7	62.3	73.5	
Married .....	52.7	21.6	34.0	68.9	100.0		30.9	29.1	31.9	28.8	32.7	
Widowed and divorced .....	175.4	...	133.9	252.3	147.4		43.3	...	70.9	35.6	44.8	
Pulmonary .....	59.6	28.4	41.6	87.0	121.2		32.7	29.7	33.4	30.3	40.3	
Single .....	72.2	30.0	86.0	211.3	262.6		43.2	32.2	60.5	55.0	68.9	
Married .....	48.0	19.6	29.9	63.9	92.5		27.7	24.3	28.5	26.5	29.8	
Widowed and divorced .....	159.8	...	118.1	228.7	134.9		40.9	...	68.4	32.1	43.3	

TABLE 5. All Patients, Time Since Admission, by Province, 1960

Time since admission	Canada	Nfld.	P.E.I.	N.S.	N.B.	Que.	Ont.	Man.	Sask.	Alta.	B.C.	N.W.T.
Numerical distribution												
Totals .....	7,757	392	67	359	458	2,758	1,666	457	409	603	494	94
4 years or more .....	334	2	7	10	27	134	92	19	13	17	12	1
3-4 years .....	140	1	2	9	7	53	29	20	6	5	7	1
2-3 years .....	279	1	6	12	11	109	49	27	13	20	30	1
1-2 years .....	903	19	8	48	40	288	184	94	53	84	72	13
6-12 months .....	1,801	81	16	93	84	612	377	93	107	195	115	28
4-6 months .....	1,190	84	7	55	72	381	256	56	83	94	83	19
3-4 months .....	652	49	5	21	36	254	123	33	44	45	30	12
2-3 months .....	790	44	6	50	41	298	196	31	28	47	43	6
1-2 months .....	842	44	4	31	71	313	185	42	34	53	59	6
Under 1 month .....	826	67	6	30	69	316	175	42	28	43	43	7
Tuberculous .....	7,043	375	63	343	432	2,327	1,513	434	399	584	491	80
4 years or more .....	310	2	7	10	25	118	86	19	13	17	12	1
3-4 years .....	130	1	2	9	6	45	29	19	6	5	7	1
2-3 years .....	250	1	6	11	11	89	42	27	12	20	30	1
1-2 years .....	838	19	8	48	37	247	172	91	53	82	72	9
6-12 months .....	1,704	80	15	92	81	552	357	89	106	193	115	24
4-6 months .....	1,105	84	7	52	67	336	228	55	83	92	83	18
3-4 months .....	590	48	5	21	35	214	112	31	40	43	30	11
2-3 months .....	702	43	6	47	40	239	176	30	27	46	43	5
1-2 months .....	741	37	3	26	68	257	166	39	33	49	58	5
Under 1 month .....	673	60	6	27	62	230	145	34	26	37	41	5
Non-tuberculous .....	714	17	2	16	26	431	153	23	10	19	3	14
4 years or more .....	24	—	—	—	2	16	6	—	—	—	—	—
3-4 years .....	10	—	—	—	1	8	—	1	—	—	—	—
2-3 years .....	29	—	—	1	—	20	7	—	1	—	—	—
1-2 years .....	65	—	—	—	3	41	12	3	—	2	—	4
6-12 months .....	97	1	1	1	3	60	20	4	1	2	—	4
4-6 months .....	85	—	—	3	5	45	28	1	—	2	—	1
3-4 months .....	62	1	—	—	1	40	11	2	4	2	—	1
2-3 months .....	88	1	—	3	1	59	20	1	1	1	—	1
1-2 months .....	101	7	1	5	3	56	19	3	1	4	1	1
Under 1 month .....	153	7	—	3	7	86	30	8	2	6	2	2
Percentage distribution												
Totals .....	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Under 4 years .....	95.7	99.5	89.6	97.2	94.1	95.1	94.5	95.8	96.8	97.2	97.6	98.9
Under 3 years .....	93.9	99.2	86.6	94.7	92.6	93.2	92.7	91.5	95.4	96.4	96.2	97.9
Under 2 years .....	90.3	99.0	77.6	91.4	90.2	89.3	89.8	85.6	92.2	93.0	90.1	96.8
Under 1 year .....	78.7	94.1	65.7	78.0	81.4	78.8	78.8	65.0	79.2	79.1	75.5	83.0
Under 6 months .....	55.4	73.5	41.8	52.1	63.1	56.6	56.1	44.6	53.1	46.8	52.2	53.2
Under 4 months .....	40.1	52.0	31.3	36.8	47.4	42.8	40.8	32.4	32.8	31.2	35.4	33.0
Under 3 months .....	31.7	39.5	23.9	30.9	39.5	33.6	33.4	25.2	22.0	23.7	29.4	20.2
Under 2 months .....	21.5	28.3	14.9	17.0	30.6	22.8	21.6	18.4	15.2	15.9	20.6	13.8
Under 1 month .....	10.6	17.1	9.0	8.4	15.1	11.5	10.5	9.2	6.8	7.1	8.7	7.4
Tuberculous .....	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Under 4 years .....	95.6	99.5	89.2	97.1	94.2	94.9	94.3	95.6	96.7	97.1	97.6	98.8
Under 3 years .....	93.8	99.2	86.2	94.5	92.8	93.0	92.4	91.2	95.2	96.2	96.1	97.5
Under 2 years .....	90.2	98.9	76.9	91.3	90.3	89.2	89.6	85.0	92.2	92.8	90.0	96.3
Under 1 year .....	78.3	93.9	64.6	77.3	81.7	78.6	78.3	64.1	78.9	78.8	75.4	85.0
Under 6 months .....	54.1	72.5	41.5	50.4	63.0	54.8	54.7	43.5	52.4	45.7	51.9	55.0
Under 4 months .....	38.4	50.1	30.8	35.3	47.5	40.4	39.6	30.9	31.6	30.0	35.0	32.5
Under 3 months .....	30.0	37.3	23.1	29.2	39.4	31.2	32.2	23.7	21.6	22.6	28.9	18.8
Under 2 months .....	20.1	25.9	13.8	15.5	30.1	20.9	20.6	16.8	14.8	14.7	20.2	12.5
Under 1 month .....	9.6	16.0	9.2	7.9	14.4	9.9	9.6	7.8	6.5	6.3	8.4	6.3
Non-tuberculous .....	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Under 4 years .....	96.6	100.0	100.0	100.0	92.3	96.3	96.1	100.0	100.0	100.0	100.0	100.0
Under 3 years .....	95.2	100.0	100.0	100.0	88.5	94.4	96.1	95.7	100.0	100.0	100.0	100.0
Under 2 years .....	91.2	100.0	100.0	93.8	88.5	89.8	91.5	95.7	90.0	100.0	100.0	100.0
Under 1 year .....	82.1	100.0	100.0	93.8	76.9	80.3	83.7	82.6	90.0	89.5	100.0	71.4
Under 6 months .....	68.5	94.1	50.0	87.5	65.4	66.4	70.6	65.2	80.0	78.9	100.0	42.9
Under 4 months .....	56.6	94.1	50.0	68.8	46.2	55.9	52.3	60.9	80.0	68.4	100.0	35.7
Under 3 months .....	47.9	88.2	50.0	68.8	42.3	46.6	45.1	52.2	40.0	57.9	100.0	28.6
Under 2 months .....	35.6	82.4	50.0	50.0	38.5	32.9	32.0	47.8	30.0	52.6	100.0	21.4
Under 1 month .....	21.4	41.2	—	18.8	26.9	20.0	19.6	34.8	20.0	31.6	66.7	14.3

TABLE 6. All Patients, by Sex, Diagnosis, and Time of Admission, 1934-60

No.	Time of admission	Both sexes				
		All diagnoses	Pulmonary tuberculosis	Other respiratory tuberculosis	Other forms of tuberculosis	Non-tuberculous
1	Totals .....	7,757	5,689	855	499	714
2	1934 .....	1	1	—	—	—
3	1936 .....	2	2	—	—	—
4	1937 .....	1	1	—	—	—
5	1938 .....	2	2	—	—	—
6	1939 .....	1	1	—	—	—
7	1940 .....	2	2	—	—	—
8	1941 .....	2	2	—	—	—
9	1942 .....	1	1	—	—	—
10	1943 .....	1	1	—	—	—
11	1944 .....	5	5	—	—	—
12	1945 .....	9	9	—	—	—
13	1946 .....	6	5	—	1	—
14	1947 .....	2	1	1	—	—
15	1948 .....	12	9	—	2	1
16	1949 .....	12	12	—	—	—
17	1950 .....	17	17	—	—	—
18	1951 .....	18	18	—	—	—
19	1952 .....	21	19	—	—	2
20	1953 .....	26	24	1	1	—
21	1954 .....	42	37	—	2	3
22	1955 .....	55	46	2	1	6
23	1956 .....	96	80	1	3	12
24	1957 .....	140	126	3	1	10
25	1958-Totals .....	279	210	19	21	29
26	January-March .....	51	43	1	2	5
27	April-June .....	61	46	5	4	6
28	July-September .....	85	62	7	5	11
29	October-December .....	82	59	6	10	7
30	1959-Totals .....	903	682	96	60	65
31	January-March .....	123	98	11	5	9
32	April-June .....	180	135	14	20	11
33	July-September .....	276	204	37	14	21
34	October-December .....	324	245	34	21	24
35	1960-Totals .....	6,101	4,376	732	407	586
36	January .....	198	161	24	6	7
37	February .....	218	161	27	21	9
38	March .....	259	194	35	22	8
39	April .....	298	223	32	21	22
40	May .....	378	279	46	31	22
41	June .....	450	336	61	24	29
42	July .....	511	363	74	36	38
43	August .....	679	503	88	41	47
44	September .....	652	467	83	40	62
45	October .....	790	562	94	46	88
46	November .....	842	580	95	66	101
47	December .....	826	547	73	53	153

TABLE 6. All Patients, by Sex, Diagnosis, and Time of Admission, 1934-60

All diagnoses	Male				Female					No.
	Pulmonary tuberculosis	Other respiratory tuberculosis	Other forms of tuberculosis	Non-tuberculous	All diagnoses	Pulmonary tuberculosis	Other respiratory tuberculosis	Other forms of tuberculosis	Non-tuberculous	
4,836	3,631	500	278	427	2,921	2,058	355	221	287	1
1	1	—	—	—	—	—	—	—	—	2
2	2	—	—	—	—	—	—	—	—	3
1	1	—	—	—	—	—	—	—	—	4
1	1	—	—	—	1	1	—	—	—	5
1	1	—	—	—	—	—	—	—	—	6
2	2	—	—	—	—	—	—	—	—	7
1	1	—	—	—	1	1	—	—	—	8
1	1	—	—	—	—	—	—	—	—	9
—	—	—	—	—	1	1	—	—	—	10
3	3	—	—	—	2	2	—	—	—	11
3	3	—	—	—	6	6	—	—	—	12
5	4	—	1	—	1	1	—	—	—	13
2	1	1	—	—	—	—	—	—	—	14
7	6	—	1	—	5	3	—	1	1	15
8	8	—	—	—	4	4	—	—	—	16
12	12	—	—	—	5	5	—	—	—	17
13	13	—	—	—	5	5	—	—	—	18
15	14	—	—	1	6	5	—	—	—	19
16	16	—	—	—	10	8	1	1	—	20
26	25	—	—	1	16	12	—	2	2	21
38	31	2	1	4	17	15	—	—	—	22
70	60	1	3	6	26	20	—	—	6	23
102	96	—	1	5	38	30	3	—	5	24
181	143	8	13	17	98	67	11	8	12	25
32	27	—	2	3	19	16	1	—	2	26
45	36	3	3	3	16	10	2	1	3	27
52	41	2	2	7	33	21	5	3	4	28
52	39	3	6	4	30	20	3	4	3	29
571	446	53	35	37	332	236	43	25	28	30
78	59	10	4	5	45	39	1	1	4	31
113	86	8	12	7	67	49	6	8	4	32
164	125	20	7	12	112	79	17	7	9	33
216	176	15	12	13	108	69	19	9	11	34
3,754	2,740	435	223	356	2,347	1,636	297	184	230	35
124	109	11	2	2	74	52	13	4	5	36
134	106	13	11	4	84	55	14	10	5	37
164	124	22	14	4	95	70	13	8	4	38
197	147	20	16	14	101	76	12	5	8	39
234	173	33	17	11	144	106	13	14	11	40
268	205	30	15	18	182	131	31	9	11	41
297	213	46	20	18	214	150	28	16	20	42
401	302	50	22	27	278	201	38	19	20	43
390	283	47	19	41	262	184	36	21	21	44
486	346	58	29	53	304	216	36	17	35	45
520	364	57	34	65	322	216	38	32	36	46
539	368	48	24	99	287	179	25	29	54	47

TABLE 7. Tuberculous Patients, by Sex, Type of Admission, and Time of Admission, 1934-60

No.	Time of admission	Both sexes					
		Total	First admissions	Re-admissions	To continue treatment	Reviews	Transfers
1	Totals .....	7,043	4,309	1,655	200	28	851
2	1934 .....	1	—	1	—	—	—
3	1936 .....	2	1	1	—	—	—
4	1937 .....	1	—	1	—	—	—
5	1938 .....	2	1	1	—	—	—
6	1939 .....	1	1	—	—	—	—
7	1940 .....	2	2	—	—	—	—
8	1941 .....	2	1	1	—	—	—
9	1942 .....	1	1	—	—	—	—
10	1943 .....	1	—	1	—	—	—
11	1944 .....	5	2	3	—	—	—
12	1945 .....	9	5	4	—	—	—
13	1946 .....	6	2	4	—	—	—
14	1947 .....	2	2	—	—	—	—
15	1948 .....	11	6	3	—	—	2
16	1949 .....	12	7	4	—	—	1
17	1950 .....	17	8	7	1	—	1
18	1951 .....	18	6	10	—	—	2
19	1952 .....	19	7	10	—	—	2
20	1953 .....	26	10	12	1	—	3
21	1954 .....	39	6	21	3	—	9
22	1955 .....	49	24	17	—	—	8
23	1956 .....	84	34	27	2	—	21
24	1957 .....	130	47	48	7	—	28
25	1958-Totals .....	250	127	68	8	—	47
26	January-March .....	46	19	11	—	—	16
27	April-June .....	55	27	13	1	—	14
28	July-September .....	74	40	21	3	—	10
29	October-December .....	75	41	23	4	—	7
30	1959-Totals .....	838	448	205	27	—	158
31	January-March .....	114	43	32	3	—	36
32	April-June .....	169	102	34	7	—	26
33	July-September .....	255	141	64	10	—	40
34	October-December .....	300	162	75	7	—	56
35	1960-Totals .....	5,515	3,561	1,206	151	28	569
36	January .....	191	128	37	8	—	18
37	February .....	209	144	43	3	—	19
38	March .....	251	173	50	5	—	23
39	April .....	276	191	57	4	—	24
40	May .....	356	259	58	5	—	34
41	June .....	421	295	59	6	—	61
42	July .....	473	348	76	10	1	38
43	August .....	632	422	135	18	1	56
44	September .....	590	371	142	22	2	53
45	October .....	702	404	171	22	2	103
46	November .....	741	439	203	29	3	67
47	December .....	673	387	175	19	19	73

TABLE 7. Tuberculous Patients, by Sex, Type of Admission, and Time of Admission, 1934-60

Male						Female						No.
Total	First admissions	Re-admissions	To continue treatment	Reviews	Transfers	Total	First admissions	Re-admissions	To continue treatment	Reviews	Transfers	
4,409	2,613	1,072	137	20	567	2,634	1,696	583	63	8	284	1
1	—	1	—	—	—	—	—	—	—	—	—	2
2	1	1	—	—	—	—	—	—	—	—	—	3
1	—	1	—	—	—	—	—	—	—	—	—	4
1	1	—	—	—	—	1	—	1	—	—	—	5
1	1	—	—	—	—	—	—	—	—	—	—	6
2	2	—	—	—	—	—	—	—	—	—	—	7
1	—	1	—	—	—	1	1	—	—	—	—	8
1	1	—	—	—	—	—	—	—	—	—	—	9
—	—	—	—	—	—	1	—	1	—	—	—	10
3	1	2	—	—	—	2	1	1	—	—	—	11
3	1	2	—	—	—	6	4	2	—	—	—	12
5	2	3	—	—	—	1	—	1	—	—	—	13
2	2	—	—	—	—	—	—	—	—	—	—	14
7	5	1	—	—	—	1	4	1	2	—	—	15
8	5	2	—	—	—	1	4	2	2	—	—	16
12	6	4	1	—	—	1	5	2	3	—	—	17
13	4	7	—	—	—	2	5	2	3	—	—	18
14	5	7	—	—	—	2	5	2	3	—	—	19
16	5	10	—	—	—	1	10	5	2	1	—	20
25	3	15	1	—	—	6	14	3	6	2	—	21
34	15	13	—	—	—	6	15	9	4	—	—	22
64	29	17	2	—	—	16	20	5	10	—	—	23
97	29	40	5	—	23	33	18	8	2	—	—	24
164	81	43	4	—	36	86	46	25	4	—	—	25
29	12	6	—	—	11	17	7	5	—	—	5	26
42	20	9	—	—	13	13	7	4	1	—	1	27
45	23	15	2	—	5	29	17	6	1	—	5	28
48	26	13	2	—	7	27	15	10	2	—	—	29
534	287	133	18	—	96	304	161	72	9	—	62	30
73	28	23	2	—	20	41	15	9	1	—	16	31
106	68	18	5	—	15	63	34	16	2	—	11	32
152	84	38	8	—	22	103	57	26	2	—	18	33
203	107	54	3	—	39	97	55	21	4	—	17	34
3,398	2,127	769	106	20	376	2,117	1,434	437	45	8	193	35
122	82	23	6	—	11	69	46	14	2	—	7	36
130	91	27	2	—	10	79	53	16	1	—	9	37
160	105	35	4	—	16	91	68	15	1	—	7	38
183	131	33	3	—	16	93	60	24	1	—	8	39
223	158	40	3	—	22	133	101	18	2	—	12	40
250	170	34	2	—	44	171	125	25	4	—	17	41
279	202	45	6	1	25	194	146	31	4	—	13	42
374	246	73	12	1	42	258	176	62	6	—	14	43
349	204	95	14	1	35	241	167	47	8	1	18	44
433	239	119	18	1	56	269	165	52	4	1	47	45
455	253	126	23	3	50	286	186	77	6	—	17	46
440	246	119	13	13	49	233	141	56	6	6	24	47

## TUBERCULOSIS STATISTICS

TABLE 8. Tuberculous Patients, by Age, Origin, and Sex, 1960

Origin and sex	All ages	0 - 4	5 - 9	10 - 14	15 - 19	20 - 24	25 - 29	30 - 34	35 - 39	40 - 44	45 - 49
	T.	373	397	293	391	482	495	507	537	482	500
Totals .....	T.	7,043	373	397	293	391	482	495	507	537	482
	M.	4,409	200	190	140	197	240	257	269	312	301
	F.	2,634	173	207	153	194	242	238	238	225	181
Indian .....	T.	1,104	155	130	97	96	88	97	101	49	41
	M.	549	88	64	41	44	33	34	42	25	19
	F.	555	67	66	56	52	55	63	59	24	22
Eskimo .....	T.	265	34	30	31	24	22	23	18	15	22
	M.	110	16	8	20	6	12	7	8	4	10
	F.	155	18	22	11	18	10	16	10	11	12
Other .....	T.	5,662	184	236	165	271	372	375	388	473	419
	M.	3,741	96	117	79	147	195	216	219	283	272
	F.	1,921	88	119	86	124	177	159	169	190	147
Not stated .....	T.	12	—	1	—	—	—	—	—	—	3
	M.	9	—	1	—	—	—	—	—	—	3
	F.	3	—	—	—	—	—	—	—	—	—
	50 - 54	55 - 59	60 - 64	65 - 69	70 - 74	75 - 79	80 - 84	85 - 89	90 - 94	95 - 99	Not stated
Totals .....	T.	488	489	461	381	341	218	124	44	15	2
	M.	372	363	360	296	267	160	74	31	10	1
	F.	116	126	101	85	74	58	50	13	5	1
Indian .....	T.	36	38	44	34	28	12	10	3	1	5
	M.	22	25	27	24	21	8	5	1	1	—
	F.	14	13	17	10	7	4	5	2	—	4
Eskimo .....	T.	7	8	9	2	4	—	—	—	—	2
	M.	3	—	7	—	3	—	—	—	—	2
	F.	4	8	2	2	1	—	—	—	—	—
Other .....	T.	444	441	408	344	308	204	113	41	14	1
	M.	346	337	326	272	242	151	68	30	9	1
	F.	98	104	82	72	66	53	45	11	5	4
Not stated .....	T.	1	2	—	1	1	2	1	—	—	—
	M.	1	1	—	—	1	1	1	—	—	—
	F.	—	1	—	1	—	1	—	—	—	—

TABLE 9. All Indian and Eskimo Patients, by Sex and Diagnosis, 1960

Diagnosis	Indian			Eskimo		
	Both sexes	Male	Female	Both sexes	Male	Female
Totals .....	1,176	580	596	294	123	171
Pulmonary tuberculosis .....	714	355	359	191	73	118
Other respiratory tuberculosis .....	256	128	128	55	27	28
Other forms of tuberculosis .....	134	66	68	19	10	9
Non-tuberculous .....	72	31	41	29	13	16

TABLE 10. Non-tuberculous Patients, by Sex and Diagnosis, 1960

Diagnosis	Both sexes	Male	Female	Diagnosis	Both sexes	Male	Female
Totals .....	714	427	287	Lobar pneumonia .....	6	4	2
Septicaemia and pyaemia .....	2	2	—	Bronchopneumonia .....	11	7	4
Acute poliomyelitis .....	1	1	—	Primary atypical, other, and unspecified pneumonia .....	17	11	6
All other diseases classified as infective and parasitic .....	11	5	6	Acute bronchitis .....	6	2	4
Malignant neoplasm of intestine, except rectum .....	1	1	—	Bronchitis, chronic and unqualified .....	39	21	18
Malignant neoplasm of rectum .....	1	1	—	Empyema and abscess of lung .....	11	11	—
Malignant neoplasm of larynx .....	1	1	—	Pleurisy .....	13	11	2
Malignant neoplasm of trachea, bronchus, and lung, not specified as secondary .....	18	13	5	All other respiratory diseases .....	77	55	22
Malignant neoplasm of breast .....	1	—	1	Spontaneous pneumothorax .....	3	3	—
Malignant neoplasm of cervix uteri .....	1	—	1	Pulmonary congestion and hypostasis .....	14	7	7
Malignant neoplasm of all other and unspecified sites .....	4	3	1	Pneumoconiosis due to silica and silicates....	4	4	—
Leukaemia and aleukaemia .....	1	—	1	Other specified pneumoconiosis and pulmonary fibrosis of occupational origin .....	1	1	—
Lymphosarcoma and other neoplasms of lymphatic and haematopoietic system .....	1	1	—	Other chronic interstitial pneumonia .....	13	9	4
Benign neoplasms and neoplasms of unspecified nature .....	5	5	—	Bronchiectasis .....	24	19	5
Diabetes mellitus .....	8	6	2	Other diseases of lung and pleural cavity .....	18	12	6
Anaemias .....	3	—	3	Ulcer of stomach .....	1	1	—
Allergic disorders; all other endocrine, metabolic, and blood diseases .....	23	17	6	Gastro-enteritis and colitis, except diarrhoea of the newborn .....	2	—	2
Psychoses .....	3	1	2	Cholelithiasis and cholecystitis .....	3	1	2
Psychoneuroses and disorders of personality .....	10	4	6	Other diseases of digestive system .....	1	1	—
Vascular lesions affecting central nervous system .....	8	4	4	Chronic, other, and unspecified nephritis .....	1	—	1
Multiple sclerosis .....	2	—	2	Infections of kidney .....	1	1	—
Otitis media and mastoiditis .....	2	1	1	Other diseases of genito-urinary system .....	2	2	—
All other diseases of the nervous system and sense organs .....	19	8	11	Infections of skin and subcutaneous tissue .....	1	1	—
Rheumatic fever .....	1	—	1	Arthritis and spondylitis .....	8	4	4
Chronic rheumatic heart disease .....	2	1	1	Muscular rheumatism and rheumatism unspecified .....	1	—	1
Arteriosclerotic and degenerative heart disease ..	24	14	10	All other diseases of skin and musculoskeletal system .....	4	4	—
Other diseases of heart .....	12	7	5	Spina bifida and meningocele .....	1	—	1
Hypertension with heart disease .....	3	1	2	All other congenital malformations .....	3	—	3
Hypertension without mention of heart .....	8	4	4	Senility without mention of psychosis .....	4	—	4
Diseases of arteries .....	8	5	3	Ill-defined and unknown causes of morbidity and mortality .....	8	3	5
Other diseases of circulatory system .....	2	—	2	No disease, no diagnosis, suspected tuberculosis only, and positive reactor .....	307	181	126

TABLE 11. All Patients, by Sex, Diagnosis, Province, and Institution, 1960

Province and institution	Both sexes					Male					Female				
	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous
Number															
Canada .....	8,246	6,099	862	519	766	5,143	3,887	504	288	464	3,103	2,212	358	231	302
Newfoundland .....	395	255	73	50	17	231	145	40	34	12	164	110	33	16	5
Corner Brook .....	143	90	28	13	12	92	59	18	7	8	51	31	10	6	4
North West River .....	7	—	7	—	—	3	—	3	—	—	4	—	4	—	—
St. Anthony .....	30	20	2	6	2	17	12	1	2	2	13	8	1	4	—
St. John's .....	206	136	36	31	3	116	71	18	25	2	90	65	18	6	1
St. John's (Hosp. for M. and N. Diseases) .....	3	3	—	—	—	—	—	—	—	—	3	3	—	—	—
Twillingate .....	6	6	—	—	—	3	3	—	—	—	3	3	—	—	—
Prince Edward Island .....	67	41	17	7	2	43	30	9	3	1	24	11	8	4	1
Charlottetown .....	67	41	17	7	2	43	30	9	3	1	24	11	8	4	1
Nova Scotia .....	407	349	24	16	18	266	229	19	7	11	141	120	5	9	7
Dartmouth .....	11	11	—	—	—	7	7	—	—	—	4	4	—	—	—
Cole Harbour .....	23	21	—	1	1	14	13	—	—	1	9	8	—	1	—
Halifax .....	14	11	—	2	1	10	9	—	1	—	4	2	—	1	1
Kentville .....	213	192	12	5	4	143	129	10	1	3	70	63	2	4	1
Sydney (Cape Breton Hosp.) .....	14	13	—	—	1	6	5	—	—	1	8	8	—	—	—
Sydney (Point Edward Hosp.) .....	132	101	12	8	11	86	66	9	5	6	46	35	3	3	5
New Brunswick .....	458	336	74	22	26	279	213	42	13	11	179	123	32	9	15
St. Basile .....	95	65	20	2	8	61	46	12	1	2	34	19	8	1	6
St. John .....	137	102	19	11	5	99	75	14	6	4	38	27	5	5	1
The Glades .....	77	46	18	6	7	41	27	7	3	4	36	19	11	3	3
Vallée-Lourdes .....	149	123	17	3	6	78	65	9	3	1	71	58	8	—	5
Québec .....	2,758	2,107	163	57	431	1,724	1,348	95	30	251	1,034	759	68	27	180
Gaspé .....	196	62	16	2	116	126	40	8	1	77	70	22	8	1	39
Hull .....	52	39	10	2	1	33	28	5	—	—	19	11	5	2	1
Lac-Édouard .....	73	59	1	1	12	57	49	1	1	6	16	10	—	—	6
Lac-Etchemin .....	51	45	1	—	5	37	35	—	—	2	14	10	1	—	3
Macamic .....	73	49	2	4	18	36	23	1	2	10	37	26	1	2	8
Mont-Joli .....	338	245	31	4	58	207	147	21	3	36	131	98	10	1	22
Montreal (Alexandra Hosp.) .....	26	24	—	2	—	10	9	—	1	—	16	15	—	1	—
Montreal (Grace Dart Hosp.) .....	90	78	1	2	9	71	63	1	1	6	19	15	—	1	3
Montréal (Hôpital du Sacré-Coeur) .....	135	116	4	—	15	78	67	4	—	7	57	49	—	—	8
Montréal (Hôpital St-Joseph) .....	393	368	8	2	15	244	223	7	—	14	149	145	1	2	1
Montreal (Montreal Children's Hospital) .....	1	—	—	—	1	—	—	—	—	—	1	—	—	—	1
Montreal (Queen Mary Veterans' Hosp.) .....	9	4	—	—	5	9	4	—	—	5	—	—	—	—	—
Montreal (Royal Edward Laurentian Hosp.) .....	88	65	—	5	18	60	43	—	4	13	28	22	—	1	5
Roberval .....	309	235	51	5	18	176	138	24	3	11	133	97	27	2	7
Ste-Agathe-des-Monts (Mount Sinai San.) .....	106	93	3	2	8	87	77	2	2	6	19	16	1	—	2
Ste-Agathe-des-Monts (Royal Edward Laurentian Hosp.) .....	125	121	3	1	—	101	98	3	—	—	24	23	—	1	—
Ste-Anne-de-Bellevue .....	38	36	2	—	—	38	36	2	—	—	—	—	—	—	—
Ste-Foy .....	359	275	26	23	35	192	151	15	11	15	167	124	11	12	20
Sherbrooke .....	75	71	2	1	1	37	36	1	—	—	38	35	1	1	1
Trois-Rivières .....	221	122	2	1	96	125	81	—	1	43	96	41	2	—	53

TABLE 11. All Patients, by Sex, Diagnosis, Province, and Institution, 1960 - Continued

Province and institution	Both sexes					Male					Female				
	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous
	Number														
Ontario .....	1,792	1,312	175	129	176	1,179	875	115	74	115	613	437	60	55	61
Brantford .....	46	37	2	-	7	34	25	2	-	7	12	12	-	-	-
Comwall .....	53	44	3	2	4	32	28	1	-	3	21	16	2	2	1
Fort William .....	198	120	41	16	21	117	76	22	8	11	81	44	19	8	10
Hamilton .....	186	124	26	16	20	108	68	18	10	12	78	56	8	6	8
Kingston .....	68	50	1	2	15	52	39	1	1	11	16	11	-	1	4
Kitchener .....	40	28	8	1	3	31	21	7	1	2	9	7	1	-	1
London .....	182	142	11	10	19	131	102	10	7	12	51	40	1	3	7
Moose Factory .....	61	39	6	3	13	20	15	-	1	4	41	24	6	2	9
Ottawa .....	144	105	13	9	17	88	61	8	5	14	56	44	5	4	3
St. Catharines .....	97	77	9	8	3	64	49	9	4	2	33	28	-	4	1
Sioux Lookout .....	7	3	1	2	1	5	2	1	1	1	2	1	-	1	-
Sudbury .....	105	66	14	9	16	74	45	11	7	11	31	21	3	2	5
Weston .....	427	339	35	43	10	307	250	22	28	7	120	89	13	15	3
Windsor .....	52	40	2	6	4	33	28	2	-	3	19	12	-	6	1
Woodstock .....	126	98	3	2	23	83	66	1	1	15	43	32	2	1	8
Manitoba .....	484	382	30	48	24	288	220	20	30	18	196	162	10	18	6
Brandon .....	25	10	6	9	-	10	5	1	4	-	15	5	5	5	-
Brandon (Mental Hospital) .....	8	7	-	1	-	5	4	-	1	-	3	3	-	-	-
Ninette .....	165	137	6	12	10	109	85	4	11	9	56	52	2	1	1
Portage la Prairie .....	5	4	-	1	-	4	3	-	1	-	1	1	-	-	-
St. Vital .....	179	152	10	16	1	108	91	7	9	1	71	61	3	7	-
Selkirk .....	14	12	1	-	1	9	7	1	-	1	5	5	-	-	-
The Pas .....	73	51	7	5	10	36	20	7	3	6	37	31	-	2	4
Winnipeg .....	15	9	-	4	2	7	5	-	1	1	8	4	-	3	1
Saskatchewan .....	488	363	67	47	11	275	216	33	18	8	213	147	34	29	3
Fort Qu'Appelle .....	19	13	3	3	-	8	7	1	-	-	11	6	2	3	-
Fort San .....	108	84	11	7	6	68	55	5	3	5	40	29	6	4	1
Prince Albert .....	159	99	42	17	1	83	53	23	6	1	76	46	19	11	-
Saskatoon .....	123	93	10	17	3	74	61	4	8	1	49	32	6	9	2
Weyburn .....	79	74	1	3	1	42	40	-	1	1	37	34	1	2	-
Alberta .....	603	387	117	80	19	347	229	59	46	13	256	158	58	34	6
Calgary .....	169	113	38	15	3	117	79	23	12	3	52	34	15	3	-
Edmonton (Aberhart San.) .....	204	128	30	32	14	123	86	14	14	9	81	42	18	18	5
Edmonton (Charles Camsell Hosp.) .....	230	146	49	33	2	107	64	22	20	1	123	82	27	13	1
British Columbia .....	700	519	106	47	28	475	362	64	28	21	225	157	42	19	7
Essondale .....	206	167	2	12	25	137	111	2	6	18	69	56	-	6	7
Sardis (Indian Health Services) .....	225	114	90	21	-	127	65	48	14	-	98	49	42	7	-
Vancouver (Division of T.B. Control) .....	269	238	14	14	3	211	186	14	8	3	58	52	-	6	-
Northwest Territories .....	94	48	16	16	14	36	20	8	5	3	58	28	8	11	11
Aklavik (All Saints' Hosp.) .....	13	7	1	2	3	3	2	-	-	1	10	5	1	2	2
Aklavik (Immaculate Conception Hosp.) .....	17	7	2	-	8	6	3	1	-	2	11	4	1	-	6
Fort Rae .....	21	16	5	-	-	10	8	2	-	-	11	8	3	-	-
Fort Simpson .....	12	5	1	3	3	1	1	-	-	-	11	4	1	3	3
Fort Smith .....	24	8	5	11	-	10	2	3	5	-	14	6	2	6	-
Pangnirtung .....	7	5	2	-	-	6	4	2	-	-	1	1	-	-	-

TABLE 11. All Patients, by Sex, Diagnosis, Province, and Institution, 1960 — Continued

Province and institution	Both sexes					Male					Female				
	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous
Percentage distribution															
<b>Canada</b>	<b>100.00</b>	<b>74.0</b>	<b>10.5</b>	<b>6.3</b>	<b>9.3</b>	<b>100.0</b>	<b>75.6</b>	<b>9.8</b>	<b>5.6</b>	<b>9.0</b>	<b>100.0</b>	<b>71.3</b>	<b>11.5</b>	<b>7.4</b>	<b>9.7</b>
<b>Newfoundland</b>	<b>100.0</b>	<b>64.6</b>	<b>18.5</b>	<b>12.7</b>	<b>4.3</b>	<b>100.0</b>	<b>62.8</b>	<b>17.3</b>	<b>14.7</b>	<b>5.2</b>	<b>100.0</b>	<b>67.1</b>	<b>20.1</b>	<b>9.8</b>	<b>3.0</b>
Corner Brook	100.0	62.9	19.6	9.1	8.4	100.0	64.1	19.6	7.6	8.7	100.0	60.8	19.6	11.8	7.8
North West River	100.0	—	100.0	—	—	100.0	—	100.0	—	—	100.0	—	100.0	—	—
St. Anthony	100.0	66.7	6.7	20.0	6.7	100.0	70.6	5.9	11.8	11.8	100.0	61.5	7.7	30.8	—
St. John's	100.0	66.0	17.5	15.0	1.5	100.0	61.2	15.5	21.6	1.7	100.0	72.2	20.0	6.7	1.1
St. John's (Hosp. for M. and N. Diseases)	100.0	100.0	—	—	—	—	—	—	—	—	100.0	100.0	—	—	—
Twillingate	100.0	100.0	—	—	—	100.0	100.0	—	—	—	100.0	100.0	—	—	—
<b>Prince Edward Island</b>	<b>100.0</b>	<b>61.2</b>	<b>25.4</b>	<b>10.4</b>	<b>3.0</b>	<b>100.0</b>	<b>69.8</b>	<b>20.9</b>	<b>7.0</b>	<b>2.3</b>	<b>100.0</b>	<b>45.8</b>	<b>33.3</b>	<b>16.7</b>	<b>4.2</b>
Charlottetown	100.0	61.2	25.4	10.4	3.0	100.0	69.8	20.9	7.0	2.3	100.0	45.8	33.3	16.7	4.2
<b>Nova Scotia</b>	<b>100.0</b>	<b>85.7</b>	<b>5.9</b>	<b>3.9</b>	<b>4.4</b>	<b>100.0</b>	<b>86.1</b>	<b>7.1</b>	<b>2.6</b>	<b>4.1</b>	<b>100.0</b>	<b>85.1</b>	<b>3.5</b>	<b>6.4</b>	<b>5.0</b>
Dartmouth	100.0	100.0	—	—	—	100.0	100.0	—	—	—	100.0	100.0	—	—	—
Cole Harbour	100.0	91.3	—	4.3	4.3	100.0	92.9	—	—	7.1	100.0	88.9	—	11.1	—
Halifax	100.0	78.6	—	14.3	7.1	100.0	90.0	—	10.0	—	100.0	50.0	—	25.0	25.0
Kentville	100.0	90.1	5.6	2.3	1.9	100.0	90.2	7.0	0.7	2.1	100.0	90.0	2.9	5.7	1.4
Sydney (Cape Breton Hosp.)	100.0	92.9	—	—	7.1	100.0	83.3	—	—	16.7	100.0	100.0	—	—	—
Sydney (Point Edward Hosp.)	100.0	76.5	9.1	6.1	8.3	100.0	76.7	10.5	5.8	7.0	100.0	76.1	6.5	6.5	10.9
<b>New Brunswick</b>	<b>100.0</b>	<b>73.4</b>	<b>16.2</b>	<b>4.8</b>	<b>5.7</b>	<b>100.0</b>	<b>76.3</b>	<b>15.1</b>	<b>4.7</b>	<b>3.9</b>	<b>100.0</b>	<b>68.7</b>	<b>17.9</b>	<b>5.0</b>	<b>8.4</b>
St. Basile	100.0	68.4	21.1	2.1	8.4	100.0	75.4	19.7	1.6	3.3	100.0	55.9	23.5	2.9	17.6
St. John	100.0	74.5	13.9	8.0	3.6	100.0	75.8	14.1	6.1	4.0	100.0	71.1	13.2	13.2	2.6
The Glades	100.0	59.7	23.4	7.8	9.1	100.0	65.9	17.1	7.3	9.8	100.0	52.8	30.6	8.3	8.3
Vallée-Lourdes	100.0	82.6	11.4	2.0	4.0	100.0	83.3	11.5	3.8	1.3	100.0	81.7	11.3	—	7.0
<b>Québec</b>	<b>100.0</b>	<b>76.4</b>	<b>5.9</b>	<b>2.1</b>	<b>15.6</b>	<b>100.0</b>	<b>78.2</b>	<b>5.5</b>	<b>1.7</b>	<b>14.6</b>	<b>100.0</b>	<b>73.4</b>	<b>6.6</b>	<b>2.6</b>	<b>17.4</b>
Gaspé	100.0	31.6	8.2	1.0	59.2	100.0	31.7	6.3	0.8	61.1	100.0	31.4	11.4	1.4	55.7
Hull	100.0	75.0	19.2	3.8	1.9	100.0	84.8	15.2	—	—	100.0	57.9	26.3	10.5	5.3
Lac-Édouard	100.0	80.8	1.4	1.4	16.4	100.0	86.0	1.8	1.8	10.5	100.0	62.5	—	—	37.5
Lac-Etchemin	100.0	88.2	2.0	—	9.8	100.0	94.6	—	—	5.4	100.0	71.4	7.1	—	21.4
Macamic	100.0	67.1	2.7	5.5	24.7	100.0	63.9	2.8	5.6	27.8	100.0	70.3	2.7	5.4	21.6
Mont-Joli	100.0	72.5	9.2	1.2	17.2	100.0	71.0	10.1	1.4	17.4	100.0	74.8	7.6	0.8	16.8
Montreal (Alexandra Hosp.)	100.0	92.3	—	7.7	—	100.0	90.0	—	10.0	—	100.0	93.8	—	6.3	—
Montreal (Grace Dart Hosp.)	100.0	86.7	1.1	2.2	10.0	100.0	88.7	1.4	1.4	8.5	100.0	78.9	—	5.3	15.8
Montréal (Hôpital du Sacré-Coeur)	100.0	85.9	3.0	—	11.1	100.0	85.9	5.1	—	9.0	100.0	86.0	—	—	14.0
Montréal (Hôpital St-Joseph)	100.0	93.6	2.0	0.5	3.8	100.0	91.4	2.9	—	5.7	100.0	97.3	0.7	1.3	0.7
Montreal (Montreal Children's Hospital)	100.0	—	—	—	100.0	—	—	—	—	—	100.0	—	—	—	100.0
Montreal (Queen Mary Veterans' Hosp.)	100.0	44.4	—	—	55.6	100.0	44.4	—	—	55.6	—	—	—	—	—
Montreal (Royal Edward Laurentian Hosp.)	100.0	73.9	—	5.7	20.5	100.0	71.7	—	6.7	21.7	100.0	78.6	—	3.6	17.9
Roberval	100.0	76.1	16.5	1.6	5.8	100.0	78.4	13.6	1.7	6.2	100.0	72.9	20.3	1.5	5.3
Ste-Agathe-des-Monts (Mount Sinai San.)	100.0	87.7	2.8	1.9	7.5	100.0	88.5	2.3	2.3	6.9	100.0	84.2	5.3	—	10.5
Ste-Agathe-des-Monts (Royal Edward Laurentian Hosp.)	100.0	96.8	2.4	0.8	—	100.0	97.0	3.0	—	—	100.0	95.8	—	4.2	—
Ste-Anne-de-Bellevue	100.0	94.7	5.3	—	—	100.0	94.7	5.3	—	—	—	—	—	—	—
Ste-Foy	100.0	76.6	7.2	6.4	9.7	100.0	78.6	7.8	5.7	7.8	100.0	74.3	6.6	7.2	12.0
Sherbrooke	100.0	94.7	2.7	1.3	1.3	100.0	97.3	2.7	—	—	100.0	92.1	2.6	2.6	2.6
Trois-Rivières	100.0	55.2	0.9	0.5	43.4	100.0	64.8	—	0.8	34.4	100.0	42.7	2.1	—	55.2

TABLE 11. All Patients, by Sex, Diagnosis, Province, and Institution, 1960 — Concluded

Province and institution	Both sexes					Male					Female				
	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous	All diagnoses	Pulmonary	Other respiratory	Other forms	Non-tuberculous
Percentage distribution — Concluded															
Ontario .....	100.0	73.2	9.8	7.2	9.8	100.0	74.2	9.8	6.3	9.8	100.0	71.3	9.8	9.0	10.0
Brantford .....	100.0	80.4	4.3	—	15.2	100.0	73.5	5.9	—	20.6	100.0	100.0	—	—	—
Cornwall .....	100.0	83.0	5.7	3.8	7.5	100.0	87.5	3.1	—	9.4	100.0	76.2	9.5	9.5	4.8
Fort William .....	100.0	60.6	20.7	8.1	10.6	100.0	65.0	18.8	8.8	9.4	100.0	54.3	23.5	9.9	12.3
Hamilton .....	100.0	66.7	14.0	8.6	10.8	100.0	63.0	16.7	9.3	11.1	100.0	71.8	10.3	7.7	10.3
Kingston .....	100.0	73.5	1.5	2.9	22.1	100.0	75.0	1.9	1.9	21.2	100.0	68.8	—	6.3	25.0
Kitchener .....	100.0	70.0	20.0	2.5	7.5	100.0	67.7	22.6	3.2	6.5	100.0	77.8	11.1	—	11.1
London .....	100.0	78.0	6.0	5.5	10.4	100.0	77.9	7.6	5.3	9.2	100.0	78.4	2.0	5.9	13.7
Moose Factory .....	100.0	63.9	9.8	4.9	21.3	100.0	75.0	—	5.0	20.0	100.0	58.5	14.6	4.9	22.0
Ottawa .....	100.0	72.9	9.0	6.3	11.8	100.0	69.3	9.1	5.7	15.9	100.0	78.6	8.9	7.1	5.4
St. Catharines .....	100.0	79.4	9.3	8.2	3.1	100.0	76.6	14.1	6.3	3.1	100.0	84.8	—	12.1	3.0
Sioux Lookout .....	100.0	42.9	14.3	28.6	14.3	100.0	40.0	20.0	20.0	20.0	100.0	50.0	—	50.0	—
Sudbury .....	100.0	62.9	13.3	8.6	15.2	100.0	60.8	14.9	9.5	14.9	100.0	67.7	9.7	6.5	16.1
Weston .....	100.0	79.4	8.2	10.1	2.3	100.0	81.4	7.2	9.1	2.3	100.0	74.2	10.8	12.5	2.5
Windsor .....	100.0	76.9	3.8	11.5	7.7	100.0	84.8	6.1	—	9.1	100.0	63.2	—	31.6	5.3
Woodstock .....	100.0	77.8	2.4	1.6	18.3	100.0	79.5	1.2	1.2	18.1	100.0	74.4	4.7	2.3	18.6
Manitoba .....	100.0	78.9	6.2	9.9	5.0	100.0	76.4	6.9	10.4	6.3	100.0	82.7	5.1	9.2	3.1
Brandon .....	100.0	40.0	24.0	36.0	—	100.0	50.0	10.0	40.0	—	100.0	33.3	33.3	33.3	—
Brandon (Mental Hospital) .....	100.0	87.5	—	12.5	—	100.0	80.0	—	20.0	—	100.0	100.0	—	—	—
Ninette .....	100.0	83.0	3.6	7.3	6.1	100.0	78.0	3.7	10.1	8.3	100.0	92.9	3.6	1.8	1.8
Portage la Prairie .....	100.0	80.0	—	20.0	—	100.0	75.0	—	25.0	—	100.0	100.0	—	—	—
St. Vital .....	100.0	84.9	5.6	8.9	0.6	100.0	84.3	6.5	8.3	0.9	100.0	85.9	4.2	9.9	—
Selkirk .....	100.0	85.7	7.1	—	7.1	100.0	77.8	11.1	—	11.1	100.0	100.0	—	—	—
The Pas .....	100.0	69.9	9.6	6.8	13.7	100.0	55.6	19.4	8.3	16.7	100.0	83.8	—	5.4	10.8
Winnipeg .....	100.0	60.0	—	26.7	13.3	100.0	71.4	—	14.3	14.3	100.0	50.0	—	38.5	12.5
Saskatchewan .....	100.0	74.4	13.7	9.6	2.3	100.0	78.5	12.0	6.5	2.9	100.0	69.0	16.0	13.6	1.4
Fort Qu'Appelle .....	100.0	68.4	15.8	15.8	—	100.0	87.5	12.5	—	—	100.0	54.5	18.2	27.3	—
Fort San .....	100.0	77.8	10.2	6.5	5.6	100.0	80.9	7.4	4.4	7.4	100.0	72.5	15.0	10.0	2.5
Prince Albert .....	100.0	62.3	26.4	10.7	0.6	100.0	63.9	27.7	7.2	1.2	100.0	60.5	25.0	14.5	—
Saskatoon .....	100.0	75.6	8.1	13.8	2.4	100.0	82.4	5.4	10.8	1.4	100.0	65.3	12.2	18.4	4.1
Weyburn .....	100.0	93.7	1.3	3.8	1.3	100.0	95.2	—	2.4	2.4	100.0	91.9	2.7	5.4	—
Alberta .....	100.0	64.2	19.4	13.3	3.2	100.0	66.0	17.0	13.3	3.7	100.0	61.7	22.7	13.3	2.3
Calgary .....	100.0	66.9	22.5	8.9	1.8	100.0	67.5	19.7	10.3	2.6	100.0	65.4	28.8	5.8	—
Edmonton (Aberhart San.) .....	100.0	62.7	14.7	15.7	6.9	100.0	69.9	11.4	11.4	7.3	100.0	51.9	19.8	22.2	6.2
Edmonton (Charles Camell Hosp.) .....	100.0	63.5	21.3	14.3	0.9	100.0	59.8	20.6	18.7	0.9	100.0	66.7	22.0	10.6	0.8
British Columbia .....	100.0	74.1	15.1	6.7	4.0	100.0	76.2	13.5	5.9	4.4	100.0	69.8	18.7	8.4	3.1
Esondale .....	100.0	81.1	1.0	5.8	12.1	100.0	81.0	1.5	4.4	13.1	100.0	81.2	—	8.7	10.1
Sardis (Indian Health Services) .....	100.0	50.7	40.0	9.3	—	100.0	51.2	37.8	11.0	—	100.0	50.0	42.9	7.1	—
Vancouver (Division of T.B. Control) .....	100.0	88.5	5.2	5.2	1.1	100.0	88.2	6.6	3.8	1.4	100.0	89.7	—	10.3	—
Northwest Territories .....	100.0	51.1	17.0	17.0	14.9	100.0	55.6	22.2	13.9	8.3	100.0	48.3	13.8	19.0	19.0
Aklavik (All Saints' Hosp.) .....	100.0	53.8	7.7	15.4	23.1	100.0	66.7	—	—	33.3	100.0	50.0	10.0	20.0	20.0
Aklavik (Immaculate Conception Hosp.) .....	100.0	41.2	11.8	—	47.1	100.0	50.0	16.7	—	33.3	100.0	36.4	9.1	—	54.5
Fort Rae .....	100.0	76.2	23.8	—	—	100.0	80.0	20.0	—	—	100.0	72.7	27.3	—	—
Fort Simpson .....	100.0	41.7	8.3	25.0	25.0	100.0	100.0	—	—	—	100.0	36.4	9.1	27.3	27.3
Fort Smith .....	100.0	33.3	20.8	45.8	—	100.0	20.0	30.0	50.0	—	100.0	42.9	14.3	42.9	—
Pangnirtung .....	100.0	71.4	28.6	—	—	100.0	66.7	33.3	—	—	100.0	100.0	—	—	—

**APPENDIX****Patients Under Treatment for Tuberculosis in Mental Institutions, 1960<sup>1</sup>**

Province	Total	Active	Inactive
<b>Canada</b> .....	<b>1,611</b>	<b>276</b>	<b>1,335</b>
Newfoundland .....	—	—	—
Prince Edward Island .....	1	1	—
Nova Scotia .....	—	—	—
New Brunswick .....	23	8	15
Quebec .....	1,428	231	1,197
Ontario .....	18	7	11
Manitoba .....	—	—	—
Saskatchewan .....	15	3	12
Alberta .....	126	26	100
British Columbia .....	—	—	—

<sup>1</sup> Comprises only those patients for whom morbidity cards had not been submitted, and who consequently are excluded from Table 11.





1010484633

## SELECTED PUBLICATIONS ON HEALTH STATISTICS AND POPULATION

Obtainable from Information Services Division  
Dominion Bureau of Statistics or Queen's Printer, Ottawa, Canada

Catalogue number	VITAL STATISTICS	Price
84-001 - Monthly Report - Bil.	.....	per year \$1.00
84-201 - Preliminary Annual Report - Bil.	.....	.25
84-202 - Annual Report - Bil.	.....	2.00

## Analytical Reports:

84-503 - Leading Causes of Death by Major Age Groupings, 1921-55 - E.	.....	.25
84-504 - Accidental Drowning, 1956 - E.	.....	.25
84-505 - Mortality from Accidental Falls, 1956 - E.	.....	.25
84-506 - Mortality from Accidental Poisoning, 1956 - E.	.....	.25
84-507 - Mortality from Home Accidents, 1956 - E.	.....	.25
84-511 - Mortality from Suicide, 1921-58 - E.	.....	.50
84-508 - Causes of Stillbirths, 1943-55 - E.	.....	.25
84-509 - Canadian Life Tables, 1951 - Bil.	.....	.25
84-510 - Canadian Life Tables, 1950-52, 1955-57 - Bil.	.....	.50
84-512 - Canadian Life Tables, Provincial and Regional, 1950-52, 1955-57 - Bil.	.....	.75
84-513 - Fatal Accidents, Mortality Summary, 1959	.....	.75

## INSTITUTIONAL STATISTICS

83-201 - List of Canadian Hospitals and Related Health Facilities - A.E.	.....	.75
83-202 - Hospital Statistics, Vol. I, General Information, 1958 - A. Bil.	.....	1.50
83-203 - Hospital Statistics, Vol. II, Financial Information, 1958 - A. Bil.	.....	2.00
83-502 - The Large General Hospital, 1957 - E.	.....	.50
83-503 - The Small General Hospital, 1957 - E.	.....	.50
83-205 - Mental Health Statistics, Financial Supplement - A. Bil.	.....	.25
83-207 - Tuberculosis Statistics, Financial Supplement - A. Bil.	.....	.25
83-210 - Hospital Statistics, Vol. I, Hospital Beds, 1959 - A. Bil.	.....	1.50

## MORBIDITY STATISTICS

82-201 - Notifiable Diseases - A. Bil.	.....	.75
82-202 - Home Nursing Services - A.E.	.....	1.00
82-203 - Illness in the Civil Service - A. Bil.	.....	.75
82-204 - Poliomyelitis Trends - A.E.	.....	.25
82-512 - Summary of Notifiable Diseases, 1924-52 (annual supplements to 1960) - Bil.	.....	.75
82-516 - Influenza in Canada, Some Statistics on its Characteristics and Trends, 1957 - E.	.....	.50
83-204 - Mental Health Statistics - A. Bil.	.....	2.00
83-208 - Mental Health Statistics, Supplement: Patients in Institutions - A.E.	.....	.50
83-504 - Rates and Probabilities of Separation from Mental Institutions, 1956-58 - E.	.....	.25
83-206 - Tuberculosis Statistics - A. Bil.	.....	2.00
83-209 - Tuberculosis Statistics Supplement: Patients in Institutions - A.E.	.....	.50
83-001 - Incidence of Tuberculosis - Bil. (monthly)	.....	per year 1.00
82-205 - Tuberculosis Trends, 1960 - A.E.	.....	.25

## Canadian Sickness Survey (National Estimates), 1950-51 - E. and F.

82-502 - Family Expenditures for Health Services by Income Groups	.....	.25
82-505 - Volume of Sickness	.....	.25
82-506 - Permanent Physical Disabilities	.....	.25
82-507 - Incidence and Prevalence of Illness	.....	.25
82-508 - Volume of Health Care	.....	.25
82-509 - Volume of Health Care for Selected Income Groups	.....	.25
82-510 - Illness Frequency by Diagnostic Classification	.....	.25
82-511 - Illness Frequency by Class of Illness, Age and Sex	.....	.25
82-518 - Illness and Health Care in Canada, 1950-51 - E.	.....	2.00
82-519 - Injuries: Frequency, Severity, Health Care - E.	.....	.75
82-520 - Volume of Health Care for Insured and Non-Insured Persons - E.	.....	.75
82-521 - Disability Among the Gainfully Employed - E.	.....	.50

## POPULATION STATISTICS - Bil.

91-201 - Canada, by Provinces, Estimated for Intercensal Years, 1921-60 - A.	.....	.25
91-503 - Estimates (Age and Sex), 1921-52 - R. No. 40	.....	.25
91-504 - Estimates (Age and Sex), 1952-56 (Supp. to 91-503)	.....	.25
91-202 - Estimates, by Age and Sex Groups, Canada and Provinces, 1957-60 (A. Supp.)	.....	.25
91-203 - Estimates, by Marital Status, Age and Sex, Canada and Provinces - A.	.....	.25
91-204 - Estimates of Families in Canada - A.	.....	.50

A. - Annual

E. - English

F. - French

Bil. - Bilingual

R. - Reference

Supp. - Supplement

In addition to the selected publications listed above, the Dominion Bureau of Statistics publishes a wide range of statistical reports on Canadian economic and social affairs including detailed reports on the 1956 and 1951 Censuses. A comprehensive catalogue of all current publications is available free on request from the Dominion Bureau of Statistics, Ottawa 3.