


Analytical Paper

Life Tables, Canada, Provinces and Territories, 1980/1982 to 2014/2016

by Demography Division

Release date: June 28, 2018



Statistics
Canada

Statistique
Canada

Canada

How to obtain more information

For information about this product or the wide range of services and data available from Statistics Canada, visit our website, www.statcan.gc.ca.

You can also contact us by

email at STATCAN.infostats-infostats.STATCAN@canada.ca

telephone, from Monday to Friday, 8:30 a.m. to 4:30 p.m., at the following numbers:

- | | |
|---|----------------|
| • Statistical Information Service | 1-800-263-1136 |
| • National telecommunications device for the hearing impaired | 1-800-363-7629 |
| • Fax line | 1-514-283-9350 |

Depository Services Program

- | | |
|------------------|----------------|
| • Inquiries line | 1-800-635-7943 |
| • Fax line | 1-800-565-7757 |

Standards of service to the public

Statistics Canada is committed to serving its clients in a prompt, reliable and courteous manner. To this end, Statistics Canada has developed standards of service that its employees observe. To obtain a copy of these service standards, please contact Statistics Canada toll-free at 1-800-263-1136. The service standards are also published on www.statcan.gc.ca under “Contact us” > “Standards of service to the public.”

Note of appreciation

Canada owes the success of its statistical system to a long-standing partnership between Statistics Canada, the citizens of Canada, its businesses, governments and other institutions. Accurate and timely statistical information could not be produced without their continued co-operation and goodwill.

Standard table symbols

The following symbols are used in Statistics Canada publications:

- | | |
|----------------|--|
| . | not available for any reference period |
| .. | not available for a specific reference period |
| ... | not applicable |
| 0 | true zero or a value rounded to zero |
| 0 ^s | value rounded to 0 (zero) where there is a meaningful distinction between true zero and the value that was rounded |
| ^p | preliminary |
| ^r | revised |
| x | suppressed to meet the confidentiality requirements of the <i>Statistics Act</i> |
| ^E | use with caution |
| F | too unreliable to be published |
| * | significantly different from reference category ($p < 0.05$) |

Published by authority of the Minister responsible for Statistics Canada

© Her Majesty the Queen in Right of Canada as represented by the Minister of Industry, 2018

All rights reserved. Use of this publication is governed by the Statistics Canada [Open Licence Agreement](#).

An HTML version is also available.

Cette publication est aussi disponible en français.

LIFE TABLES, CANADA, PROVINCES AND TERRITORIES 1980/1982 TO 2014/2016

For the periods from 1980/1982 to 2014/2016, complete life tables for Canada, Newfoundland and Labrador, Nova Scotia, New Brunswick, Quebec, Ontario, Manitoba, Saskatchewan, Alberta and British Columbia are available in Table 13-10-0114-01.

Abridged life tables for Prince Edward Island, Yukon, Northwest Territories and Nunavut are available in Table 13-10-0140-01.

**A full set of life tables is also available in Excel on the publication's website:
<https://www150.statcan.gc.ca/n1/pub/84-537-x/84-537-x2018002-eng.htm>**

This document includes life tables for Canada, provinces and territories from the period 1980 to 1982 to the period 2014 to 2016. In addition to producing life expectancy, life tables also provide other indicators describing the mortality within a population during a given period of time: death probabilities, survival probabilities, survivors at specific ages and years of life lived.

Complete life tables, by single years of age and sex, are available for Canada as a whole and for nine provinces: Newfoundland and Labrador, Nova Scotia, New Brunswick, Quebec, Ontario, Manitoba, Saskatchewan, Alberta and British Columbia.

For Prince Edward Island, Yukon, the Northwest Territories and Nunavut, abridged life tables were computed, that is by five-year age groups and sex. The population size of this province and the three territories were too small to allow the calculation of complete life tables with accuracy. Northwest Territories and Nunavut are treated separately in the construction of 1998/2000 and 1999/2001 mortality tables, even though Nunavut was officially created on April 1st of 1999.

Methods

The methods used for the life tables are described in the document *Methods for Constructing Life Tables for Canada, Provinces and Territories*, catalogue no. 84-538. All of the Statistics Canada life tables computed in this series are based on this revised methodology which takes into account recent progress in the field of mortality studies.

Definitions of the elements included in the life tables

The following elements are available in all life tables included in this document.

Age (age interval):

The major visual difference between the complete and abridged life tables lies in the age groupings for which the estimates have been produced.

In complete life tables, there is only one age value per row, which indicates the exact age for the number of survivors, the cumulative number of life years lived and the life expectancy. For the number of deaths, death and survival probabilities, as well as the number of life years lived, the interval in the life table represent the interval between two exact ages. For example, death at age 30 means that the death occurred on or after the 30th birthday but before reaching its 31st birthday.

The presentation is the same in abridged life tables, but the age intervals are of the form $(x, x+(n-1))$; that is, both ages x and $x+(n-1)$ are included in the interval. For example, the age interval 40 to 44 comprises deaths occurring among 40 to 44 year-olds. Most age intervals in abridged life tables span five years. The exceptions occur in the first two rows of these tables and for the last row: the first row (age 0) represents a one-year interval and the second row, a four-year interval (ages 1 to 4). The last row is an open age interval, 90 years and over.

l_x (number of survivors at age x):

Number of persons in an initial cohort of 100,000 live births who are still alive at the beginning of each subsequent age interval. The number of survivors decreases as age increases, under the effect of mortality.

It is possible to compute, from the number of survivors, probabilities of survival between two ages. For example, if the number of survivors is 99,297 at age 10 and 98,935 at age 20, the probability of surviving from age 10 to age 20 is $98,935 / 99,297$, that is, 0.99635.

d_x (number of deaths between age x and x+n):

Number of deaths which occur in each age interval among the initial cohort of 100,000 live births at age 0.

q_x (death probability between age x and x+n):

Probability that a person of age x dies before reaching age x+n.

m.e. (q_x) (margin of error associated with the death probability):

Margin of error associated with the death probability. For example, a margin of error of 0.00020 for a death probability at age 0 of 0.00556 enables the construction of a 95% confidence interval with lower and upper limits of 0.00536 and 0.00576. In other words, the death probability is precise within a range of 0.00020, 19 times out of 20.

p_x (probability of survival between age x and x+n):

Probability that a person of age x survives up to year x+n.

L_x (number of life years lived between age x and x+n):

Number of life years lived by persons between age x and x+n. Life years lived are also considered as the stationary population of the life table. Except for ages from 0 to 4 where a separation factor is computed (see the document [*Methods for Constructing Life Tables for Canada, Provinces and Territories*](#), catalogue no. 84-538), the assumption made is that the deaths are distributed evenly over time within the age interval.

T_x (cumulative number of life years lived beyond age x):

Total number of life years lived by persons of age x and all those included in subsequent age intervals.

e_x (life expectancy at age x):

Average number of years remaining to be lived by persons at age x if these persons would experience, during their life, the mortality observed over the reference period.

m.e. (e_x) (margin of error associated with the life expectancy):

Margin of error associated with the life expectancy at age x. For example, a margin of error of 0.2 on a life expectancy at birth of 81.9 years enables the construction of a 95% confidence interval with lower and upper limits of 81.7 years and 82.1 years.