


06-215



Environment Canada  
Environnement Canada

Canada



NATIONAL WATER  
RESEARCH INSTITUTE

INSTITUT NATIONAL DE  
RECHERCHE SUR LES EAUX

TD  
226  
N87  
No.  
06-215

Food Chain effects of sublethal  
Ultraviolet radiation on subarctic  
*Daphnia pulex* - a field and laboratory  
study  
By:

I.D. Zellmer, M.T. Arts, V. Sustr

NWRI Contribution No: 06-215

## Food chain effects of sublethal ultraviolet radiation on subarctic *Daphnia pulex* – a field and laboratory study

I. D. Zellmer<sup>1</sup>, M. T. Arts<sup>2</sup> and Vladimír Šustr<sup>3</sup>

With 5 figures

**Abstract:** In shallow subarctic ponds *Daphnia* and their algal food are often exposed to solar ultraviolet radiation (UVR). It was previously demonstrated that direct exposure of *Daphnia* to UVR may result in sublethal damage to *Daphnia*'s intestinal system. This led to the hypothesis that, as consequences of such exposures, *Daphnia* may be experiencing a situation similar to starvation. We examined, in controlled experiments, the indirect effect of UVR-treated food algae on *Daphnia pulex* which, themselves, were also exposed to UVR. We specifically tested whether exposure of *D. pulex* to solar UVR affects food transport and enzymatic digestion. As expected, the UVR-exposed food and UVR-treated *Daphnia* combination produced the strongest effects on intestinal damage, and mortality. Some of these effects, as well as the grazer-related effects, were similar to those observed during starvation. The total activities of digestive enzymes (amylase and cellulase) were somewhat reduced in UVR treated *D. pulex*, but the function of enzymes is not seriously damaged as is clear from the increasing trend of protein-specific activities. We conclude that, in some ways, *Daphnia* undergoing sublethal UVR exposures may be experiencing a condition similar to what they would experience under very low food or even fasting conditions.

**Key words:** UV radiation, *Daphnia*, enzymes, sub-Arctic ponds, chronic stress.

### Introduction

Ponds reflect changes in environmental conditions very quickly due to their shallowness and small volume. Thus, phyto- and zooplankton inhabiting such

<sup>1</sup> **Authors' addresses:** Martin-Luther University, Institute of Plant Physiology, Weinbergweg 10, D-06120 Halle, Germany. Corresponding author.

<sup>2</sup> National Water Research Institute, Environment Canada, 867 Lakeshore Road, P. O. Box 5050, Burlington, Ontario, L7R 4A6, Canada.

<sup>3</sup> Academy of Sciences of the Czech Republic, B. C., Institute of Soil Biology, Na Sádkách 7, CZ-37005 České Budějovice, Czech Republic.

ecosystems are exposed to rapid changes in their physical (i.e. solar radiation, temperature, DOC dissolved organic carbon, pH) as well as biochemical environment. In high latitude ponds physical and chemical forcing of certain variables are even more extreme than for ponds at lower latitudes compelling biota to cope with sudden and drastic changes in their environment. Two of the most variable factors in high latitudes ponds are temperature and solar radiation. As a consequence of decreases in stratospheric ozone concentrations increases in solar ultraviolet radiation (UVR, 290–400 nm) have been recorded in some regions (PERIN & LEAN 2004). UVR is the most energetic and most harmful part of the solar spectrum and shows a maximum in early spring in northern latitudes. The attenuation of UVR in the water column depends mainly on the concentration of the chromophoric portion of DOC (ARTS et al. 2000). The shallowness of many subarctic ponds and their low DOC concentrations allows UVR to penetrate to the bottom of most ponds. Thus, the biota living in these ponds must tolerate at least some exposure to solar UVR especially in spring and early summer when many pond animals are reproducing and solar UVR levels are high.

Species in the genus *Daphnia* are herbivores that play a key role in subarctic pond systems by providing a critical link between primary producers and secondary consumers (invertebrate predators, fish and humans). Several investigators have examined the direct effect of UVR on *Daphnia* demonstrating for example, that melanization of *Daphnia*'s carapace increases UVR-tolerance (HEBERT & EMERY 1990, ZELLMER 1998) and that *Daphnia* employs avoidance strategies such as migration to the bottom of the pond to avoid high radiation conditions (RAUTIO et al. 2003, LEECH & WILLIAMSON 2001). Many experimental and field studies use mortality as the endpoint to assess the effects of UVR on biota, however, sublethal damage may affect the fitness and reproductive success of *Daphnia* populations long before outright mortality occurs and, in more subtle, but equally important, ways. For example, in high latitudes sublethal solar UVR may damage *Daphnia*'s intestine before mortality occurs (ZELLMER et al. 2004). Evidence of this form of UVR-induced damage led to the hypothesis that gut-damaged *Daphnia* may suffer from malnutrition comparable to starvation (ZELLMER et al. 2004). Hence, in the current study we addressed the following questions: a) which parts of the digestive system are affected by UVR?, b) is food transport through the gut affected?, c) is the food digestion (i.e. enzyme activity) process affected? In order to answer these questions we compared UVR-treated with control *D. pulex* as well as fasting versus fed *D. pulex*.

The second part of our study focussed on the indirect effect of UVR on the nutritional status of *D. pulex* due to UVR-induced alterations in their food (algae). Since most algae cannot precisely control their depth within the water column and are more subject to wind-induced mixing they are likely to be

even more exposed to the harmful effects of UVR than the more mobile grazers are. Several authors reported changes in the morphology and/or biochemistry of algae due to UVR-treatment (VINCENT & NEALE 2000). UVR-protecting substances (e.g. mycosporine-like amino acids) which are produced by some phytoplankton in the presence of UVR (LAURION et al. 2002) have been shown to be retained in copepods (TARTAROTTI et al. 2001) but such adaptations have not been reported in *Daphnia*. Changes in polysaccharide, protein and/or lipid content (ARTS & RAI 1997) as well as fatty acid (FA) composition (ZELLMER et al. 2004) following exposure to UVR may affect the nutritional status of algae.

Thus, we tested whether UVR-treated food algae modify the sublethal parameters examined in UVR-treated *D. pulex* compared to animals fed control food. Our goal was to gain insights into the sublethal effects that solar UVR may have on the planktonic food chain in subarctic ponds.

## Material and methods

### Site description

All field experiments were conducted with *Daphnia pulex* collected from Tibors Pond in June 2003 and June 2004 in the immediate vicinity of the biological research station of the University of Helsinki in Kilpisjärvi, Finland (69° 03' N, 20° 50' E). This pond is situated at the edge of the tree line and is, in general, extremely shallow (~20 cm depth) with a single small (2 m<sup>2</sup>), deep ( $Z_{\max}$  = 1 m) depression; the only region in this




Fig. 1. Radiation transmission (%) through 1 cm of water from Lake Kilpisjärvi and Tibors Pond. Scanning performed with a Bio 50 Spectrophotometer by Cary. Insert: Transmission through acetate and WIPA foils.

pond where *D. pulex* was found. Water transparency was high (Fig. 1) and there were very few aquatic macrophytes.

### Algal cultures

The green algae *Ankistrodesmus gracilis* (Chlorophyta) was used as food source for the *daphnids* and was grown in sterile medium (as in ZELLMER 1995) in 500 ml flasks. For the field experiments the cultures were placed inside a lab window and were thus exposed to long hours (midnight sun) of natural sunlight without UVR. Flasks were shaken several times per day. Two algae cultures (50 ml of media in 250 ml beakers) were started every day. Both beakers were covered with filters (UVR-absorbing WIPA foil, WIPA Technik, Tettmang, Germany, or UVR-transmitting acetate foil, Clarifoil, England, Fig. 1) and were placed in direct sunlight on a bench outside the lab. Both treatments (with or without UVR) therefore experienced different radiation regimes but were exposed to identical temperatures during the experiments. Cultures were grown for 2 d and thereafter fed to the *daphnids* according to the experimental treatments described below (for details see ZELLMER et al. 2004). For the laboratory experiments aliquots were taken from the stock cultures of *A. gracilis* and *Scenedesmus obliquus* growing in exponential phase, centrifuged and resuspended in boiled and filtered (folded paper filters, Schleich and Schuell, Germany) lake water (Lake Wallwitz, 51° 35' N, 11° 55' E) before feeding of surplus food quantities to stock cultures of *Daphnia*. For the life table experiments, an aliquot was taken from the stock culture three times a week, centrifuged and resuspended to an optical density of 1.1 at 750 nm (equivalent to 1.5 mg C L<sup>-1</sup>). For both treatment cultures (with or without UVR), 2 ml (1.4 µg DW) of the suspension were added to 50 ml of medium each into 200 ml plastic bags (Whirlpak, Neolab, Heidelberg, Germany). Both Whirlpaks were aerated and placed in a light bank with white light only. Three times a week, at 2 d intervals, the Whirlpak bag of the UVR-treatment was placed horizontally for 20 min under two Q-Panel bulbs (Pausch Messtechnik, Haan, Germany; intensity: UVB = 295 µW<sup>-1</sup> cm<sup>-2</sup> s<sup>-1</sup>, UVA = 52.1 µW<sup>-1</sup> cm<sup>-2</sup> s<sup>-1</sup>) covered with acetate foil to remove any UVC radiation. Thereafter the bag was returned to the light bank (recovery). Thus after 7 d the algae experienced three 20 min UVR treatments in total followed by 24 h of recovery each time. Following this 7 d period algae were centrifuged and resuspended in purified lake water to an optical density of 1.1 at 750 nm (equal to 1.6 µg DW ml<sup>-1</sup>).

### *Daphnia* cultures

For the field experiments *D. pulex* were caught in the morning prior to the experiment. 50 vigorous adult *D. pulex* were added to each experimental plastic container (150 ml lake Kilpisjärvi water, pre-filtered through 160-µm sieve, 4 cm deep water column). Four replicate containers were added for each treatment resulting in a total of 200 *Daphnia* per treatment. *D. pulex* were fed equal quantities of *A. gracilis* (equivalent to 1.5 mg C L<sup>-1</sup>) and exposed to natural sunlight in a 10 °C water bath for 5 or 8 d. Ungrazed algae settled to the bottom of the containers and thus were considered to represent *ad libitum* feeding conditions. The containers were either covered with acetate foil (plus UVR) or WIPA (minus UVR) foil.

In the laboratory, stock cultures of *D. pulex* were maintained in glass jars (15 °C, dim white light) in lake water (Lake Wallwitz), which had been boiled, filtered (folded paper filters, Schleich and Schüell, Germany) and stored in a cold room (7 °C) until use. To start the life table experiment gravid *D. pulex* were separated from the stock cultures in glass jars filled with purified lake water and fed *A. gracilis* or *S. obliquus* depending on the experiment. Released neonates of *D. pulex* were removed within 48 h and evenly distributed in 4 plastic jars (200 ml) with 150 ml of purified lake water each to a density of 50 neonates per beaker. Since we used neonates from a general population of gravid females, biases as a result of genetic differences between the treatment groups are unlikely. No attempt was made to isolate a specific genotype (clone) of *D. pulex* in these experiments in order to provide a more realistic simulation of a natural pond situation, i. e. where multiple clones of daphnids co-occur. Food quantity provided was 0.27 µg DW per *Daphnia* every other day (for details see ZELLMER et al. 2004). From the age of 3 d, UVR-daphnids were irradiated 3 times per week for 10 min (or 15 min or 20 min in the food transportation experiments) under the same two Q-panel bulbs as above. The beakers were covered with acetate foil to remove any UVC radiation. It was previously determined that this dose rate did not cause acute mortality and was thus considered sublethal.

#### Treatments and parameters measured

The following treatments were tested: 1) *Daphnia* not exposed to UVR fed algae not exposed to UVR 2) *Daphnia* not exposed to UVR fed UVR-exposed algae 3) *Daphnia* exposed to UVR fed algae not exposed to UVR and 4) *Daphnia* exposed to UVR fed algae exposed to UVR.

In the field experiments every evening and every 2 d in the laboratory experiments all *Daphnia* individuals were scored for the following parameters under a stereo microscope (25–25× magnification): survival (%), intestinal damage (any changes in form or structure of the intestinal tube, i. e. bends, breaks or changes in diameter were scored as damage) and an estimate of the length of the green-colored portion of the intestine. In *Daphnia*'s intestine roughly 75 % (foregut and midgut) usually appears bright green in actively feeding animals due to the chlorophyll still present in the incompletely digested algae. The hindgut (~25 % of the intestine) appears more yellowish or brownish due to pigment degradation as the algae are digested. We used a conservative scoring system to quantify changes in the digestive competency of *Daphnia* in the various treatments wherein a score of 1 or 0 was assigned to a *Daphnia* whose gut appeared entirely green or entirely brown, respectively. Intermediate stages were ranked 0.25, 0.5 or 0.75 depending on the extent of the green part of the gut. We assumed that the green part of the intestine represented the largely undigested mass of algae, thus providing a rough measure of the digestive competency of *D. pulex* (ZELLMER et al. 2004). This scoring method was fast and non-invasive. Animals appeared not to be stressed heavily by it since survival in the controls (no UV-treatments) was very high. Dead animals and neonates were removed.

After scoring, the *Daphnia* were returned to the containers and fed providing the algae enough time to settle during the night so as not to unduly alter the UVR-transmittance the next day due to suspended algae.

### Enzyme analysis

Enzyme analysis on *D. pulex* was conducted immediately after the termination of the experiment at the field station. Enzyme activities were measured in whole body homogenates of groups of *Daphnia* individuals ( $n = 11$  and  $15$  animals for the  $\alpha$ -amylase and cellulase assay, respectively). *D. pulex* were isolated from original pond water or the experimental containers by pipette. Water was filtered out using a plastic mesh and the mesh with animals was dried out carefully to remove residual source water. The animals were then placed into either  $3.5$  ml ( $\alpha$ -amylase assay) or  $2.5$  ml (cellulase assay) of cold sterile phosphate (Britton-Robinson) buffer  $\text{pH} = 8$  ( $\text{pH} = 7$  for cellulase assay) and homogenized in a glass homogenizer cooled by a surrounding ice bath. The homogenate was centrifuged at  $6000$  g for  $7$  min and the supernatant was taken for activity measurements at  $30^\circ\text{C}$  after  $1$  h ( $\alpha$ -amylase assay) or  $24$  h (cellulase assay). Total soluble protein content was measured in  $100\ \mu\text{l}$  sub-sample of homogenate using the Bradford reagent (SIGMA, product No. B6916).

**$\alpha$ -Amylase assay:** enzyme activity was measured using the tablet S-test (STU, Bratislava, Slovakia) with specific chromolytic substrate (URBÁŠEK & STARÝ 1994). One ml of supernatant was added to  $50$  mg of chromolytic substrate and a drop of toluene was added as a bactericide. This mixture was incubated at  $30^\circ\text{C}$  for  $1$  h. Incubation was finished by adding  $2$  ml of acetone solution ( $900$  ml of  $\text{H}_2\text{O}$ ,  $100$  ml acetone,  $10$  g  $\text{Na}_2\text{CO}_3$ ). After centrifugation ( $6000$  g,  $7$  min) the light absorbance of the final colored supernatant was measured at  $620$  nm. The results are expressed in mg of decomposed substratum per hour per individual (per mg of soluble protein).

**Cellulase assay:** enzyme activity was measured using carboxymethyl cellulose (CMC) as a substrate and assaying for the final concentration of glucose as the reaction product (ŠUSTR & CHALUPSKÝ 1996). A  $0.5$  ml aliquot of supernatant was incubated with a  $0.5$  ml suspension of CMC ( $0.1$  g of CMC in  $5$  ml of phosphate buffer). A drop of toluene was added as a bactericide. This mixture was incubated at  $30^\circ\text{C}$  for  $24$  h. Incubation was finished by adding  $0.3$  ml of the incubation mixture to  $1$  ml of trichloroacetic acid ( $50$  g  $\text{L}^{-1}$ ). After centrifugation  $0.2$  ml of the supernatant was added to  $1$  ml of glucose-oxidase reagent for enzymatic determination of glucose (GLUCOSE GOD-250, Lachema, Czech Republic). Absorbance at  $492$  nm was measured after  $30$  min incubation at room temperature (about  $22^\circ\text{C}$ ,  $30$  min at  $15$ – $25^\circ\text{C}$  is sufficient safely for oxidation of all glucose in the sample according to the manufacturer).

### Determination of the food transportation rate through *Daphnia*'s intestine

To test whether the rate of food transportation through the gut is affected by UVR-treatment of *D. pulex* and/or its food we measured the time until control or UVR-treated *S. obliquus* passed through the entire intestine of UVR-treated or control *Daphnia*. In this experiment we chose three different UVR-treatments for the *Daphnia*:  $10$  min,  $15$  min and  $20$  min of UVR-irradiation for the last  $5$  d before the start of the grazing experiments. At an age of  $13$  d, *Daphnia* were transferred into a yeast suspension for  $2$  d. In this way the entire gut of these animals was filled with cream-colored yeast cells. The *Daphnia* were then rinsed in fresh culture water and transferred into a

*S. obliquus* solution ( $OD_{750\text{nm}} = 0.68$ ). At 5 min intervals *Daphnia* were checked under the microscope and the length of gut occupied by algae (as shown by the advancing green color) was determined. *Daphnia* were allowed to graze for 30 min and were then returned to the beakers with the yeast solution. After 2 h (following which their intestines were again completely filled with cream colored yeast cells) the grazing experiment was repeated with UVR-treated *S. obliquus*. Each treatment had between 2 and 3 replicates with 20 animals each. Due to the high mortality in *Daphnia* from the 20 min UVR treatment this portion of the experiment had only 1 replicate.

UVR-radiation was measured with a calibrated, cosine-corrected UVR-meter (X9, Giga-Hertz Optik, Munich, Germany) with separate sensors for UVA and UVB. During the field experiments we continuously measured integrated UVA and UVB surface intensities from 7 a.m. to 7 p.m. On sunny days, the UVB maximum was  $0.22 \text{ mW cm}^{-2} \text{ s}^{-1}$  and the UVA maximum was  $0.04 \text{ mW cm}^{-2} \text{ s}^{-1}$ .

We used one-way ANOVAs (Origin ver. 5.0) to test for main effects between the three different treatments on each day of each of the experiments. Enzyme activity data were tested using a non-parametric Kruskal-Wallis ANOVA (STATISTICA ver. 6.0).

## Results

### Direct effect of UVR on *Daphnia pulex*

The effects of UVR-exposure on *Daphnia pulex* were similar between the field and the laboratory experiments (Figs 2 and 3). UVR treatment of *D. pulex* resulted in increased mortality, an extension in the length of the green-colored section of the intestine and an increase in gut damage compared to control *D. pulex*.

The transportation rate of algae through the intestine of the *D. pulex* increased with increasing UVR-exposure (Fig. 4). In control animals (no UVR) 80 % ( $\pm 28.3$  %; SD) of the animals had completely green intestines after 10 min., which was recorded already after 5 min in *D. pulex* with the longest (20 min, no SD available) UVR-exposure. In addition, 12.5 % (no SD available) of the UVR-treated animals did not ingest any algae during the first 10 min of grazing in comparison of 3.3 % ( $\pm 4.7$ ; SD) of the control animals.

Digestive enzymes activity of *D. pulex* exposed to solar UVR for 8 d in the field experiment is shown in Fig. 5 (left panel). Three different parameters were tested: the total activity of  $\alpha$ -amylase ( $\text{mg ind}^{-1} \text{ h}^{-1}$ ) and cellulase ( $\mu\text{M ind}^{-1} \text{ h}^{-1}$ ), the body protein content as a measure of biomass ( $\mu\text{g ind}^{-1}$ ) and the protein specific activity ( $\text{mg mg}^{-1} \text{ h}^{-1}$ ) or ( $\mu\text{M ind}^{-1} \text{ h}^{-1}$ ) as a measure of the biomass related  $\alpha$ -amylase and cellulase activities respectively. The total enzyme activity in UVR-exposed compared to control animals showed a decreasing trend. The body protein content tended to be lower in UVR-treated *D. pulex* compared to the controls, too. In contrast to this, the protein-specific ac-






Fig. 2. Clockwise from top left; survival (%), mean index (range 0 to 1) representing the length of the green-colored portion of the intestine (see Methods) and % *Daphnia pulex* with damaged guts. *D. pulex* from Tibors Pond exposed to solar radiation with/without UVR, fed *Ankistrodesmus gracilis* cultured with/without solar UVR in Finland (July 2003). Means and S.E. of 4 replicates with 50–70 *D. pulex*.

tivity of digestive enzymes showed a tendency to increase, however due to the small number of replicates the statistical analysis failed to find significances (Kruskal-Wallis test,  $p > 0.2$ ).

Change of enzyme activity in fasting control animals was not identical to that of UVR-exposed animals (Fig. 5 right panel). Although starvation of *D. pulex* (no UVR-treatment) resulted in a decreasing trend of body proteins as was the case in UVR-treated animals, total enzyme activities of fasting *D. pulex* showed a non-significant increase (Kruskal-Wallis,  $p > 0.2$ ) accompanied with non-significant increase of protein specific activity (Kruskal-Wallis,  $p > 0.24$  or  $0.12$  in  $\alpha$ -amylase and cellulase respectively) after 5 d of starvation.

#### Indirect effect of UVR-treated food on *Daphnia pulex*

While UVR-irradiation of *D. pulex* caused direct effects on the animal's physiology, indirect effects may also be caused due to modifications in the digestibility and/or quality of UV-exposed food. The life-table experiments in the laboratory and in the field (Figs 2 and 3) showed no effect of UVR-treated algae on the survival of *D. pulex*. However, if *D. pulex* were repeatedly exposed to UVR, UVR-treated food resulted in higher mortality compared to control




Fig. 3. Clockwise from top left; survival (%), mean index (range 0 to 1) representing the length of the green-colored portion of the intestine (see Methods), % *Daphnia pulex* with damaged guts, and % *D. pulex* with partly empty guts. *D. pulex* were exposed to standardized radiation treatments in the laboratory with/without UVR and were fed *Scenedesmus obliquus* cultured under standardized radiation with/without UVR. Means and S.E. of 50 *D. pulex*.

food in both experimental sets. In contrast, the parameters characterizing the intestine of UV-treated *Daphnids* were not additionally affected by UVR-treated food.

The gut passage rate of UVR-treated algae through the intestine of control *D. pulex* and *D. pulex* which had received increasing doses of UVR-irradiation is shown in Fig. 4b. There are clear differences in the gut passage rate of control algae (Fig. 4a). In control animals, the non-UVR-algae were transported very quickly through the entire gut. After 15 min of grazing, in 90% ( $\pm 14.1\%$  SD) of the control *D. pulex* the gut was completely filled (100%) with green algae and in 6.7% ( $\pm 9.4\%$  SD) it was filled to 75%. At the same time, only 15.6% ( $\pm 13.3\%$  SD) of the control animals showed a completely filled gut if UVR-treated algae were fed, yet in 78% ( $\pm 13.3\%$  SD) of the animals 75% of the intestine was green (Fig. 4b). Even 30 min of grazing only increased the percentage of animals with a completely green gut to 13.3% (no SD available) if UVR-treated food was provided 90% ( $\pm 14.1\%$  SD) in the control algae group after 15 min. This situation was modified if the *Daphnia* themselves had been UVR-exposed. As was recorded for the control-algae experiment, direct UVR-treatment of *D. pulex* accelerated the transport of UVR-treated algae




Fig. 4. Food transport through the intestine of *Daphnia pulex*. *D. pulex* had been exposed to standardized radiation treatments with/without UVR prior to the experiment. a) *Scenedesmus obliquus* with no UV-pre-treatment was provided as food source. b) *S. obliquus* with UV-pretreatment was provided as food source.

through the intestinal tube. With increasing UVR-exposure time of the *Daphnia*, the percentage of *D. pulex* with totally green guts increased and the time in which this stage was reached was shortened. In addition, the rate at which




Fig. 4b.

food was transported from the midgut (75 % of the gut green) to the hind gut (100 % of the gut green) became more rapid with increasing UVR-treatment of the *Daphnia*. This was true for all UVR-treatments of the daphnids.

The percentage of *D. pulex* which did not ingest any algae was much higher in *D. pulex* exposed to high UV doses (20 min) (80 % (no SD available) after 5 min of grazing, Fig. 4b) compared to *D. pulex* exposed to a lower dose of UVR (0 % in all shorter UV-treatments after 5 min of grazing).




Fig. 5. Total and protein-specific activity of digestive enzymes and body protein content of *Daphnia pulex* exposed to UVR (left) and fasting (right). Left y-axis refers to total activity and to body protein content in cellulase, right one refers to protein-specific activity and to body protein content in amylase. DkAuv = control *Daphnia* fed on UVR treated algae, DuvAk = UVR treated *Daphnia* fed on control algae, DuvAuv = UVR treated *Daphnia* fed on UVR treated algae. \* = significant difference ( $p \leq 0.05$ ). Means, S.E.

Feeding of UVR-treated algae resulted in a decrease of body protein content as well as reduced total enzyme activities of both cellulase and amylase in *D. pulex* after 8 d (Fig. 5 left panel). However, due to the low numbers of replicates only the decrease in total cellulase activity was significant (Kruskal-Wallis,  $p < 0.044$ ). The protein-specific activity of digestive enzymes remained untouched showing rather an increasing trend (Kruskal-Wallis test,  $p > 0.2$ , Fig. 5 left panel). Feeding of UVR-exposed algae enforced the decrease of total amylase activity, but reduced the decrease of body proteins as well as the increase of protein-specific activity of amylase in UVR-exposed *D. pulex*.

As was stated above, fasting control *D. pulex* also showed a decreasing trend of body proteins (Fig. 5 right panel). In contrast to the both direct and indirect UVR-treatments, the total enzyme activity showed a non-significant increase (Kruskal-Wallis,  $p > 0.2$ ) in fasting animals accompanied by a non-significant increase in protein specific activity (Kruskal-Wallis,  $p > 0.24$  or  $0.12$  in amylase and cellulase, respectively) after 5 d of starvation.

## Discussion

Planktonic organisms in ponds are exposed to varying doses of solar UVR-radiation with only limited possibilities to escape. Many studies have described the detrimental effects of solar UVR on zooplankton if animals were exposed in shallow water (e. g. ZELLMER 1995, 1998, ZELLMER et al. 2004, LEECH & WILLIAMSON 2001, RAUTIO & KORHOLA 2002). UVR-induced changes to the biochemistry and productivity of phytoplankton have also been reported (BEHRENFELD et al. 1995, VINCENT & NEALE 2000). However, the fact that, in nature, UVR-exposed algae are also grazed by UVR-exposed herbivores, enhancing the possibility of food chain effects, has not often been examined. In this study we examined whether the direct effect of UVR on *D. pulex* is further modified by an interactive effect of UVR on the algal food.

Despite the differences in spectral composition and intensity between solar UVR and artificial UV sources some of the results of field and standardized laboratory experiments are comparable. In both approaches repeated sublethal UVR-irradiation of *D. pulex* resulted in increased mortality (Figs 2, 3). Also, in field studies, UVR-induced mortality has recently been shown to be preceded by damage to the intestine of *D. pulex* (ZELLMER et al. 2004). The results of our laboratory experiments confirm these findings: UVR-treated *D. pulex* showed an increase in the length of the green-colored portion of the gut, increased percentages of *D. pulex* with damaged guts and increased mortality compared to control animals with no UVR-treatment (Fig. 3). If *D. pulex* were fed on UVR-treated algae these negative effects increased significantly suggesting a food chain effect of UVR, both under field and laboratory conditions. In order to get a more detailed view of the impact of UVR on the digestive process of *D. pulex* we studied food transportation through the gut, amylase and cellulase enzyme activity in UVR-treated and control animals, both with control food and UVR-treated food.

The rate at which algae is transported through the gut of *D. pulex* determines the time available for digestion and assimilation. Control animals fed control food provided a reference for "normal" gut passage time (Fig. 4 a). If control *D. pulex* instead were fed UVR-treated algae, food transportation basically stopped once the midgut was filled with algae (Fig. 4 b). Even after 30 min of grazing the proportion of *D. pulex* with entirely filled guts did not exceed 13.3 % (compared to 90 % after 15 min with control food). The midgut has been identified as the major site of absorption (PETERS 1987). Here food is solubilized and the solutes are absorbed by the epithelium lining the intestine. Our results indicate a longer retention time of UVR-treated algae in this part of the digestive system of *D. pulex* compared to individuals fed control algae. Several authors have reported reduced digestibility of UVR-exposed algae (VAN DONK & HESSEN 1995, VAN DONK et al. 2000) and a high percentage of

undigested cells passing intact through the gut. Longer retention times in the midgut may counteract this phenomenon (but see BOERSMA & WILTSHIRE 2006).

The direct effect of UVR-treatment on *D. pulex* with respect to food transportation through the gut was tested by exposing them repeatedly to three different doses of UVR (10, 15 or 20 min) prior to the grazing experiment (Fig. 4a). UVR-irradiation of *D. pulex* resulted in an increasing, dose-dependant, percentage of animals with an entirely filled green gut compared to controls. In addition, as in control *D. pulex*, the time needed to fill 100 % of the gut with green algae was shorter with increasing UVR-exposure of the animals. This reduction in residence time of algae in the midgut in UVR-treated *D. pulex* is likely to affect the digestion and assimilation process negatively. These conclusions are reinforced by the increased percentage of UVR-treated animals with damaged guts or guts that showed partly or even completely empty intestines. All of these observations suggest a malfunction in the digestive process which should lower the nutritional benefit of the food to *D. pulex*. *Daphnia* exposed to UVR may therefore suffer from some degree of malnutrition comparable to the early stages of fasting animals. Observations on fasting animals, not exposed to UVR, also showed higher percentages of animals with damaged and empty guts compared to fed animals, resembling those effects of the UVR-treated animals (ZELLMER, unpubl.).

Enzymes such as cellulase and amylase play an important role in the digestion of storage and structural polysaccharides contained in algae cells. The cellulases in invertebrates are believed to originate predominantly from symbiotic bacteria or from ingestion of microbial enzymes but may also be produced by the animals (WILDISH & POOLE 1970). Transmission electron microscopy of *Daphnia magna* intestines did not reveal a gut flora and the cellulose utilization demonstrated in individuals of this species feeding on sterile food (SCHOENBERG et al. 1984). However, while evidence is lacking at present, microbial symbionts in microcrustacea cannot be ruled out (SCHOENBERG et al. 1984).

Our experiments show that protein content in the body is reduced if *D. pulex* or their food had been UVR-treated (Fig. 5, left panel). This situation corresponds to malnutrition leading to reduction of growth and using of some body proteins as energy source (Fig. 5, right panel). The total activity of both enzymes were somewhat reduced, but overall enzyme function is not seriously damaged as is clear from the increasing trend of protein-specific activity. The UVR changes are similar to those induced by short-term starvation with the difference, that the decrease of total digestive enzyme activity was not observed after starvation. It may be expected that digestive enzyme production and secretion in continuously feeding invertebrates, such as *Daphnia*, did not vary due to short-term starvation similarly to soil detritivorous oribatid mites

(HUBERT & ŠUSTR 2001), unlike discontinuous feeders as exemplified by blood sucking insects whose enzymes may be produced on demand (CHAPMAN 1985). The increase of protein-specific enzyme activity may be interpreted as a reaction compensating energy malnutrition by more efficient digestion of usually permanently available food. The second possible explanation is that the amounts of digestive enzymes are less reduced due to malnutrition than other body proteins.

In summary, our results demonstrate that solar UVR-irradiation on planktonic organisms of shallow habitats may have complex effects on plankton at different levels in the community. Primary producers as well as grazers are directly affected by radiation which does not necessarily kill the organisms. Even at the low UV doses used in the laboratory experiments we observed the same changes in *D. pulex* as were recorded in the field experiments. In addition, we showed that the effects of UVR on *D. pulex* are enhanced if both the zooplankton and its food are exposed to UVR. We showed that this damage occurs at different levels of the digestive process with food related effects only (food transportation rate) as well as food and grazer related effects (gut damage and enzyme activity). The grazer related effects were similar to those determined in fasting animals. Thus, sublethal UVR treatment may to *D. pulex* result in comparable situations as low food or even fasting conditions, depending on the degree of damage caused in the animal.

Earlier we showed, in standardized laboratory experiments (ZELLMER 1996), that the reproductive output as well as reproductive success in the UVR-stressed daphnids is reduced. Such effects influence the population on a community level even if acute mortality due to the UVR-stress does not occur. In the natural pond populations of *Daphnia* from sub-arctic ponds of Lapland gut damage has only rarely been observed during routine sampling during the last 7 years with the exception of 2003, an extremely hot and dry year (ZELLMER, unpubl.). In 2003 water levels were extremely low allowing UVR to penetrate to the bottom of the ponds. In mid July, 10 % of the sampled animals showed the same symptoms of gut damage as had been recorded in the experimental animals described here. Therefore, *D. pulex* appear to be able to counteract the damages caused by UVR to a certain degree if the exposure time is limited perhaps because repair mechanisms can be brought into play. Enzymatic activity such as amylase and cellulase activity seem to be relatively insensitive to UV stress imposed here and we conclude that they may therefore not be suitable biomarkers of UVR stress in this context.

We conclude that increasing levels of UVR especially in early spring when DOC levels are low and many species are reproducing may affect the biotic communities of shallow subarctic ponds at different levels of the food chain and in subtle ways possibly leading to shifts in community structure to more tolerant species or genotypes.



### Acknowledgements

We wish to thank MARLIES STROBEL for excellent assistance with the laboratory experiments. We thank MELANIE DAMMHAHN for excellent field assistance in Kilpisjärvi and MARGIT SCHÖNBERGER for species identification. ANTERO JÄRVINEN and the team of the Kilpisjärvi Biological Research Station in Finland provided excellent service and assistance in conducting the field experiments. We are grateful for valuable comments on earlier versions of this manuscript by KLAUS HUMBECK and two anonymous referees. This research was funded by a grant of the Deutsche Forschungsgemeinschaft (I. D. Z.), a grant of the Ecosystem Health network (I. D. Z. and V. S.), by the Research Plan No. Z6066911 of the ISB AC CR (V. S.) and by Environment Canada's National Water Research Institute (M. T. A.).

### References

- ARTS, M. T. & RAI, H. (1997): Effects of enhanced ultraviolet-B radiation on the production of lipid, polysaccharide, and protein in three freshwater algal species. – *Freshwat. Biol.* **38**: 597–610.
- ARTS, M. T., ROBERTS, R. D., KASAI, F., WAISER, M. J., TUMBER, V. P., PLANTE, A. J., RAI, H. & DE LANGE, H. J. (2000): The attenuation of ultraviolet radiation in high dissolved organic carbon waters of wetlands and lakes on the northern Great Plains. – *Limnol. Oceanogr.* **45**: 292–299.
- BEHRENFELD, M. J., LEAN, D. R. S. & LEE, H. (1995): Ultraviolet B radiation effects on inorganic nitrogen uptake by natural assemblages of oceanic plankton. – *J. Phycol.* **31**: 25–36.
- BOERSMA, M. & WILTSHIRE, K. H. (2006): Gut passage of phosphorus-limited algae through *Daphnia*: do they take up nutrients in the process? – *Arch. Hydrobiol.* **167**: 489–500, this issue.
- CHAPMAN, R. F. (1985): Coordination of digestion. – In: KERKUT, G. A. & GILBERT, L. I. (eds): *Comprehensive of insect physiology, biochemistry and pharmacology*, Vol. 4. – Pergamon Press, Oxford. pp. 213–240.
- HEBERT, P. D. N. & EMERY, C. J. (1990): The adaptive significance of cuticular pigmentation in *Daphnia*. – *Functional Ecol.* **4**: 703–710.
- HUBERT, J. & ŠUSTR, V. (2001): The effect of starvation on the metabolic rate and microanatomy of *Galumna elimata* (Acari: Oribatida). – *Eur. J. Entomol.* **98**: 265–275.
- LAURION, I., LAMI, A. & SOMMARUGA, R. (2002): Distribution of mycosporine-like amino acids and photoprotective carotenoids among freshwater phytoplankton assemblages. – *Aquat. Microb. Ecol.* **26**: 283–294.
- LEECH, D. & WILLIAMSON, C. E. (2001): In situ exposure to ultraviolet radiation alters depth distribution of *Daphnia*. – *Limnol. Oceanogr.* **46**: 416–420.
- PERIN, S. & LEAN, D. R. S. (2004): The effects of ultraviolet-B radiation on freshwater ecosystems on the Arctic: Influence from stratospheric ozone depletion and climate change. – *Environ. Rev.* **12**: 1–70.
- PETERS, R. H. (1987): Metabolism in *Daphnia*. **45**: *Daphnia*. – Peters R. H. and De Bernardi R., *Mem. Inst. Ital. Idrobiol.*, pp. 193–243.

- RAUTIO, M. & KORHOLA, A. (2002): Effects of ultraviolet radiation and dissolved organic carbon on the survival of subarctic zooplankton. – *Polar Biol.* **25**: 460–468.
- RAUTIO, M., KORHOLA, A. & ZELLMER, I. D. (2003): Vertical distribution of *Daphnia longispina* in a shallow subarctic pond: Does the interaction of ultraviolet radiation and *Chaoborus* predation explain the pattern? – *Polar Biol.* **26**: 659–665.
- SCHOENBERG, S. A., MACCUBBIN, A. F. & HODSON, R. E. (1984): Cellulose digestion by freshwater microcrustacea. – *Limnol. Oceanogr.* **29**: 1132–1136.
- ŠUSTR, V. & CHALUPSKÝ, J. (1996): Activity of digestive enzymes in two species of potworms (Oligochaeta, Enchytraeidae). – *Pedobiologia* **40**: 255–259.
- TARTAROTTI, B., LAURION, I. & SOMMARUGA, R. (2001): Large variability in the concentration of mycosporine-like amino acids among zooplankton from lakes located across an altitude gradient. – *Limnol. Oceanogr.* **46**: 1546–1552.
- URBÁŠEK, F. & STARY, J. (1994): The activity of some enzymes in the guts of five oribatid species (Acari, Oribatida). – *Pedobiologia* **38**: 250–253.
- VAN DONK, E., DE LANGE, H. J. & FAFENG, B. A. (2000): Impact of UV-radiation on grazing activity of *Daphnia middendorffiana* and use of *Chlamydomonas reinhardtii* as UV-biodosimeter in an arctic freshwater lake (Spitsbergen, Norway). – *Verh. Internat. Verein. Limnol.* – **27**: 2031–2035.
- VAN DONK, E. & HESSEN, D. O. (1995): Reduced digestibility of UV-B stressed and nutrient limited algae by *Daphnia magna*. – *Hydrobiologia* **307**: 147–151.
- VINCENT, W. F. & NEALE, P. J. (2000): Mechanisms of UV damage to aquatic organisms. – In: DEMERS M. S. & VERNET, M. (eds.): *The effects of UV radiation in the marine environment*. – Cambridge University, U. K. Press, pp. 149–176.
- WILDISH, D. J. & POOLE, N. J. (1970): Cellulase activity in *Orchestia gammarella* (Pallas). – *Comp. Biochem. Physiol.* **13**: 323–365.
- ZELLMER, I. D. (1995): UVB-tolerance of alpine and arctic *Daphnia*. – *Hydrobiologia* **307**: 153–159.
- (1996): The impact of food quantity on UVB-tolerance and recovery from UVB-damage in *Daphnia pulex*. – *Hydrobiologia* **319**: 87–92.
- (1998): The effect of natural UVA and UVB on subarctic *Daphnia pulicaria* in its natural habitat. – *Hydrobiologia* **379**: 55–62.
- ZELLMER, I. D., ARTS, M. T., ABELE, D. & HUMBECK, K. (2004): Evidence of sublethal damage in *Daphnia* (Cladocera) during exposure to solar UV radiation in subarctic ponds. – *J. Arctic, Antarctic and Alpine Res.* **36**: 370–377.

Submitted: 26 January 2006; accepted: 8 April 2006.

Environment Canada Library, Burlington



3 9055 1017 5681 4


PRINTED IN CANADA  
IMPRIME AU CANADA



ON RECYCLED PAPER  
SUR DU PAPIER RECYCLÉ

**National Water Research Institute**  
**Environment Canada**  
**Canada Centre for Inland Waters**  
P.O. Box 5050  
867 Lakeshore Road  
Burlington, Ontario  
L7R 4A6 Canada

**National Hydrology Research Centre**  
11 Innovation Boulevard  
Saskatoon, Saskatchewan  
S7N 3H5 Canada



**NATIONAL WATER  
RESEARCH INSTITUTE**  
**INSTITUT NATIONAL DE  
RECHERCHE SUR LES EAUX**

**Institut national de recherche sur les eaux**  
**Environnement Canada**  
**Centre canadien des eaux intérieures**  
Case postale 5050  
867, chemin Lakeshore  
Burlington, Ontario  
L7R 4A6 Canada

**Centre national de recherche en hydrologie**  
11, boul. Innovation  
Saskatoon, Saskatchewan  
S7N 3H5 Canada



Environment Canada  
Environnement Canada

**Canada**