Proposed Maximum Residue Limit

Santé

Canada

PMRL2019-35

Ametoctradin

(publié aussi en français)

26 November 2019

This document is published by the Health Canada Pest Management Regulatory Agency. For further information, please contact:

Publications Pest Management Regulatory Agency Health Canada 2720 Riverside Drive A.L. 6607 D Ottawa, Ontario K1A 0K9

Internet: canada.ca/pesticides hc.pmra.publications-arla.sc@canada.ca Facsimile: 613-736-3758 Information Service: 1-800-267-6315 or 613-736-3799 hc.pmra.info-arla.sc@canada.ca



ISSN: 1925-0835 (print) 1925-0843 (online)

Catalogue number: H113-24/2019-35E (print version)

H113-24/2019-35E-PDF (PDF version)

© Her Majesty the Queen in Right of Canada, represented by the Minister of Health Canada, 2019

All rights reserved. No part of this information (publication or product) may be reproduced or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, or stored in a retrieval system, without prior written permission of Health Canada, Ottawa, Ontario K1A 0K9.

Under the authority of the <u>Pest Control Products Act</u>, Health Canada's Pest Management Regulatory Agency (PMRA) has concluded that the addition of a new use on greenhouse lettuce to the product label of Zampro Fungicide, containing technical grade ametoctradin and dimethomorph, is acceptable. The specific use approved in Canada is detailed on the label of Zampro Fungicide, <u>Pest Control Products Act Registration</u> Number 30321.

The evaluation of this ametoctradin application indicated that the end-use product has value and the human health and environmental risks associated with the new use is acceptable.

Before registering a pesticide for food use in Canada, the PMRA must determine the quantity of residues that are likely to remain in or on the food when the pesticide is used according to label directions and that such residues will not be a concern to human health. This quantity is then legally established as a maximum residue limit (MRL). An MRL applies to the identified raw agricultural food commodity as well as to any processed food product that contains it, except where separate MRLs are specified for the raw agricultural commodity and a processed product made from it.

Consultation on the proposed MRL for ametoctradin is being conducted via this document (see Next Steps, the last section of this document). A summary of the greenhouse trial data used to support the proposed MRL can be found in Appendix I. The MRL consultation for dimethomorph, the other active ingredient present in Zampro Fungicide, is being consulted under a separate action.

To comply with Canada's international trade obligations, consultation on the proposed MRL is also being conducted internationally by notifying the <u>World Trade Organization</u>, as coordinated by the <u>Canada's Notification Authority and Enquiry Point</u>.

The proposed MRL, to replace an MRL already established for ametoctradin, is as follows.

 Table 1
 Proposed Maximum Residue Limit for Ametoctradin

| Common Name | Residue Definition | MRL (ppm) ¹ | Food Commodity |
|----------------|---|------------------------|----------------|
| Ametoctradin | 5-ethyl-6-octyl[1,2,4]triazolo[1,5- <i>a</i>]pyrimidin-7-amine | 80^{2} | Leaf lettuce |

 $[\]frac{1}{1}$ ppm = parts per million

MRLs established in Canada may be found using the <u>Maximum Residue Limit Database</u> on the <u>Maximum Residue Limits for Pesticides</u> webpage. The database allows users to search for established MRLs, regulated under the *Pest Control Products Act*, both for pesticides or for food commodities.

² This MRL is proposed to replace the currently established MRL of 40 ppm for this commodity.

International Situation and Trade Implications

MRLs may vary from one country to another for a number of reasons, including differences in pesticide use patterns and the locations of the crop field trials used to generate residue chemistry data.

Table 2 compares the MRL proposed for ametoctradin in Canada with the corresponding American tolerance and Codex MRL. American tolerances are listed in the <u>Electronic Code of Federal Regulations</u>, 40 CFR Part 180, by pesticide. A listing of established Codex MRLs is available on the Codex Alimentarius <u>Pesticide Index</u> webpage, by pesticide or commodity.

Table 2 Comparison of the Canadian MRL, American Tolerance and Codex MRL (where different)

| Food Commodity | Canadian MRL (ppm) | American Tolerance (ppm) | Codex MRL (ppm) |
|----------------|--------------------|---|--------------------------|
| Leaf lettuce | 80 | 40 (Vegetable, leafy, except Brassica, group 4, except spinach) | 50 (Leafy vegetables) |

Next Steps

The PMRA invites the public to submit written comments on the proposed MRL for ametoctradin up to 75 days from the date of publication of this document. Please forward your comments to Publications (see the contact information on the cover page of this document). The PMRA will consider all comments received before making a final decision on the proposed MRL. Comments received will be addressed in a separate document linked to this PMRL. The established MRL will be legally in effect as of the date that it is entered into the Maximum Residue Limit Database.

The Codex Alimentarius Commission is an international organization under the auspices of the United Nations that develops international food standards, including MRLs.

Appendix I

Summary of Field Trial Data Used to Support the Proposed Maximum Residue Limit

Residue data for ametoctradin in greenhouse lettuce were submitted to support the domestic use of Zampro Fungicide on greenhouse lettuce.

Maximum Residue Limit

The recommendation for the maximum residue limit (MRL) for ametoctradin was based upon the submitted greenhouse trial data, and the guidance provided in the OECD MRL Calculator. Table A1 summarizes the residue data used to calculate the proposed MRL for leaf lettuce.

Table A1 Summary of Field Trial Data Used to Support the MRL

| Commodity | Application Method/ Total Application Rate (g a.i./ha) ¹ | Preharvest Interval (days) | Lowest Average Field Trial Residues (ppm) | Highest Average Field Trial Residues (ppm) |
|--------------|---|----------------------------------|--|---|
| Leaf lettuce | Foliar/ 901–942 | 0 | 15.3 | 46.2 |

¹ g a.i./ha = grams of active ingredient per hectare

Following the review of all available data, the MRL as proposed in Table 1 is recommended to cover residues of ametoctradin. Residues of ametoctradin in leaf lettuce at the proposed MRL will not pose an unacceptable risk to any segment of the population, including infants, children, adults and seniors.